The Beneficial Effect of Low-Dose Aspirin on the Survival of Pathological Advanced-Stage Oral Squamous Cell Carcinoma Patients

Sheng-Dean Luo

Kaohsiung Chang Gung Memorial Hospital

Shao-Chun Wu

Kaohsiung Chang Gung Memorial Hospital

Wei-Chih Chen

Kaohsiung Chang Gung Memorial Hospital

Ching-Nung Wu

Kaohsiung Chang Gung Memorial Hospital

Tai-Jan Chiu

Kaohsiung Chang Gung Memorial Hospital

Yao-Hsu Yang

Chiayi Chang Gung Memorial Hospital

Shau-Hsuan Li

Kaohsiung Chang Gung Memorial Hospital

Fu-Min Fang

Kaohsiung Chang Gung Memorial Hospital

Tai-Lin Huang

Kaohsiung Chang Gung Memorial Hospital

Chang-Chun Hsiao

Chang Gung University

Chang-Han Chen (✉ chench7@gmail.com)

National Chi Nan University

Research Article

Keywords: oral squamous cell carcinoma, aspirin, overall survival rate, disease-specific survival rate, pathological cancer stages

Posted Date: February 15th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-193987/v1>

Abstract

Oral squamous cell carcinoma (OSCC) remains one of the most challenging clinical problems in the field due to its high rate of locoregional and distant metastases. However, studies that assess the association between aspirin use and survival in patients with OSCC are limited. Moreover, patients that recruited from those studies might have tumors that arose from different anatomic regions of the head and neck, including the oral cavity, oropharynx, etc. Since tumors within these distinct anatomic regions are unique in the context of epidemiology and tumor progression, we sought to evaluate the association of aspirin use with squamous cell carcinomas located within the oral cavity only. In this 10-year cohort study, we evaluated aspirin use and survival rates in relation to clinical characteristics as well as duration of aspirin use in patients with OSCC. Our findings suggest that OSCC patients with aspirin use for more than 180 days showed improved overall and disease-specific survival rates. Aspirin also improves survival in patients across various stages of OSCC. Cox regression models indicated that aspirin use was associated with a good prognosis. In conclusion, this evidence indicates that aspirin may be potentially used as an adjuvant therapy for OSCC.

Introduction

Oral squamous cell carcinoma (OSCC) is a common malignant tumor in the oral cavity areas including the lip, tongue, palate, etc^{1,2}. It accounts for approximately 80–90% of all cases of oral cancer^{1,2}. According to a report of the World Health Organization (WHO) on oral cancer, the incidence rates of oral cancer are high, and it remains responsible for the top three most common cancers in certain Asian-Pacific countries^{3,4}. Although its prevalence is low in Western countries, the incidence rates of oral cancer driven by human papillomavirus (HPV) infections and some lifestyle habits have increased gradually but significantly and become a public health concern^{5–10}.

Adopting unhealthy lifestyle habits such as tobacco and alcohol consumption is the primary cause of oral cancer in Western countries^{8–10}. In Asian-Pacific countries, on top of tobacco and alcohol consumption, betel (areca) nut chewing is also known as one of the major factors contributing to the risk of developing oral cancer^{11,12}. Indeed, accumulating evidence has demonstrated that adopting these unhealthy lifestyle habits, as well as HPV infection and poor oral hygiene, have significant associations with chronic inflammation that can, in turn, act as a driving force in increasing the risk of oral cancer development^{13–15}.

Aspirin is a well-known nonsteroidal anti-inflammatory drug (NSAID) that has been commonly used to reduce pain and inflammation from a spectrum of inflammatory disorders¹⁴. Given that the anti-inflammatory properties of aspirin are correlated with inhibition of tumor progression across multiple cancer types, it may exert similar effects on OSCC since overwhelming chronic inflammation has a pivotal role in the development of this cancer^{13,14}. In particular, aspirin exhibits as an effective drug for blockage of COX-2 and PGE2. This makes it more plausible that aspirin use will benefit patients with

OSCC due to a higher expression of COX-2 in patients with OSCC¹⁴. For these reasons, our interest in investigating whether aspirin can improve the clinical outcomes of OSCC patients was piqued. In addition, studies of the association between aspirin use and survival in patients with OSCC are lacking. To that end, we used the database from Chang Gung Memorial Hospital over a 10-year follow-up period involving 1,525 participants to analyze the correlation of aspirin use with survival rates in relation to clinical characteristics as well as duration of aspirin use in patients with OSCC.

Methods

Patient Recruitment

This cohort study was reviewed and approved by the Institutional Review Board (IRB) of the Kaohsiung and Chiayi branches of Chang Gung Memorial Hospital. According to these protocols, informed consent was waived due to the nature of the study design and IRB regulations. All methods were performed in accordance with the relevant guidelines and regulations. All experimental protocols were approved by the Kaohsiung and Chiayi branches of Chang Gung Memorial Hospital. The IRB approval protocol numbers are 201801348B0 and 201700253B0C602. This cohort comprises a total of 7,763 patients who were diagnosed with oral cavity cancer in Chang-Gung Memorial Hospital, Taiwan between January 2007 and December 2017. Exclusion criteria were clinical and pathological AJCC stage IVc or missing data on pathological stages of cancer (pathological AJCC or TN staging system), aspirin use for less than 180 days, oral cavity cancer with morphology codes other than squamous cell carcinoma (ICD-9-CM 140, 141, 143–145), and individuals who did not receive surgery or any treatment. In total, 1,525 patients were analyzed in our study.

Statistical analyses

Categorical variables including gender, comorbidities, lifestyle risk factors, pathological AJCC stages of cancer, etc. were tested by either a two-sided Fisher's exact test or a Pearson's chi-squared test. The normally and non-normally distributed continuous data were analyzed using Student's 2-tailed t-tests and Mann–Whitney U tests, respectively. In order to minimize the confounding effect of groups that are comparable due to non-randomized allocations, a Mahalanobis 1:4 propensity score-matched (PSM) study group was performed using the Greedy method with a 0.25 caliper-width. We used NCSS 10 software (NCSS Statistical Software, Kaysville, UT, USA) with the setting mentioned above to test if a balanced covariate distribution between aspirin users and non-users was achieved by using a logistic regression model with the following covariates: gender, age, pathological AJCC stages of cancer and treatment. We excluded one unmatched aspirin user. The standardized mean difference (SMD) was less than 0.1 after 1:4 PSM, suggesting a good balance of covariate distribution was achieved between aspirin users and non-users (Table S1). The Kaplan–Meier method was used to evaluate the effects of aspirin use in the primary outcomes (OS and DSS) after adjusting for potential confounding variables such as gender, age, and pathological AJCC stages of cancer. A univariate analysis and Cox proportional-

hazards model were used to evaluate any covariates that could affect the survival of OSCC patients. All statistical analyses were performed using SPSS Statistics V22.0 software for Windows.

Results

Demographic and clinical characteristics of patients diagnosed with OSCC

The flow chart in Fig. 1 demonstrates the strategy used to recruit OSCC patients for this cohort study that were admitted to Taiwan Chang Gung Memorial Hospital between 2007 and 2017. A total of 6,324 OSCC patients who had neoplasms affecting regions within the oral cavity such as the tongue, buccal area, etc. were recruited after applying the exclusion criteria. OSCC patients with a history of aspirin use were propensity score matched (PSM) in a 1:4 ratio to those without aspirin use: 305 were aspirin users and 1,220 were non-users after excluding one unmatched pair, i.e., 1 aspirin users and 4 non-users. The demographics and clinical characteristics of these 1,525 patients (aspirin users and non-users) are presented in Table 1. The age, gender, pathological AJCC stages of cancer, cancer recurrence, treatment, and lifestyle risk factors of OSCC patients with aspirin use appeared similar to those of non-users. A statistically significant difference was seen between the aspirin users and non-users with regard to mortality, cause of death, and comorbidities (Table 1).

Table 1
Demographic and clinical characteristics of the cohort study (n = 1525).

Variables	N(%) /median (IQR)	Low-dose Aspirin		p-value
		Non-users	Users ≥ 180days	
Sex				
Female	78(5.11%)	58(4.75%)	20(6.56%)	0.2010
Male	1447(94.89%)	1162(95.25%)	285(93.44%)	
Age				
Year	58(52–65)	58(52–65)	58(53–65)	0.7997
Pathological Stages of Cancer (AJCC) ^a				
I & II	892(58.49%)	711(58.28%)	181(59.34%)	0.7355
III & IV ^b	633(41.51%)	509(41.72%)	124(40.66%)	
Cancer Recurrence				
No	1222(80.13%)	967(79.26%)	255(83.61%)	0.0890
Yes	303(19.87%)	253(20.74%)	50(16.39%)	
Mortality				
Alive	1059(69.44%)	816(66.89%)	243(79.67%)	< 0.0001
Death	466(30.56%)	404(33.11%)	62(20.33%)	
Causes of Death				
Alive	1059(69.44%)	816(66.89%)	243(79.67%)	< 0.0001
Death due to OSCC	264(17.31%)	235(19.26%)	29(9.51%)	
Other-Causes of Death	202(13.25%)	169(13.85%)	33(10.82%)	
Treatments				
Surgery	857(56.2%)	681(55.82%)	176(57.7%)	0.5528
Surgery & RT or CCRT	668(43.8%)	539(44.18%)	129(42.3%)	
Lifestyle Risk Factors				
Smoking				
No	400(26.47%)	322(26.68%)	78(25.66%)	0.7187

Variables	N(%) /median (IQR)	Low-dose Aspirin		<i>p</i> -value
		Non-users	Users $\geq$ 180days	
Yes	1111(73.53%)	885(73.32%)	226(74.34%)	0.5293
Betel nuts consumption				
No	637(42.16%)	504(41.76%)	133(43.75%)	0.7753
Yes	874(57.84%)	703(58.24%)	171(56.25%)	
Alcoholic beverages				0.7753
No	665(44.01%)	529(43.83%)	136(44.74%)	
Yes	846(55.99%)	678(56.17%)	168(55.26%)	
Comorbidities				
Atrial fibrillation (flutter)				< 0.0001
No	1483(97.25%)	1203(98.61%)	280(91.8%)	
Yes	42(2.75%)	17(1.39%)	25(8.2%)	
DM				< 0.0001
No	1171(76.79%)	1023(83.85%)	148(48.52%)	
Yes	354(23.21%)	197(16.15%)	157(51.48%)	
Hypertension				< 0.0001
No	1082(70.95%)	987(80.9%)	95(31.15%)	
Yes	443(29.05%)	233(19.1%)	210(68.85%)	
Hyperlipidemia				< 0.0001
No	1221(80.07%)	1099(90.08%)	122(40.0%)	
Yes	304(19.93%)	121(9.92%)	183(60.0%)	
Abbreviations: IQR interquartile range; RT radiotherapy; CCRT concurrent chemoradiotherapy; DM diabetes mellitus. ^a Pathological AJCC Cancer Staging 7 th Edition; ^b Stages IVa and IVb only.				

Univariate and multivariate analyses of predictive variables for OSCC survival

The univariate Cox regression analysis indicated that pathological AJCC stages of cancer (HRstages I & II vs. stages III & IV = 4.22, 95% CI = 3.23–5.52, $p < 0.0001$), pathological T staging system (HRT1, 2 & 3 vs. T4 = 3.11, 95% CI = 2.43–3.98, $p < 0.0001$), pathological N staging system (HRN0 vs. N1, 2 & 3 = 4.25, 95% CI = 3.34–5.42, $p < 0.0001$), treatment (HRsurgery vs. surgery & RT or CCRT = 4.31, 95% CI = 3.27–5.68, $p < 0.0001$), aspirin use HRaspirin non-users vs. users = 0.44, 95% CI = 0.30–0.64, $p < 0.0001$), hyper-tension (HRno vs. yes = 0.68, 95% CI = 0.51–0.91, $p = 0.0084$), and hyperlipidemia (HRno vs. yes = 0.32, 95% CI = 0.21–0.48, $p < 0.0001$) were significant independent prognostic factors of survival. From the perspective of controlling for confounders, we further used three different multivariate Cox regression models (A, B, and C) to assess the influence of covariates on OSCC survival rates. In Model A (Table 2), which adjusted for covariates including age, gender, pathological AJCC stages of cancer, treatment, and aspirin use, the significant independent prognostic factors of survival were pathological AJCC stages of cancer (HRstages I & II vs. stages III & IV = 2.38, 95% CI = 1.69–3.35, $p < 0.0001$), treatment (HRsurgery vs. surgery & RT or CCRT = 2.55, 95% CI = 1.79–3.62, $p < 0.0001$), and aspirin use (HRaspirin non-users vs. users = 0.42, 95% CI = 0.29–0.63, $p < 0.0001$). In Model B (Table 3), which adjusted for covariates including age, gender, pathological TN stage of cancer, treatment and aspirin use, the significant independent prognostic factors of survival were pathological T staging system (HR T1, 2 & 3 vs. T4 = 1.57, 95% CI = 1.20–2.07, $p = 0.0012$), pathological N staging system (HRN0 vs. N1, 2 & 3 = 2.63, 95% CI = 1.99–3.47, $p < 0.0001$), treatment (HRsurgery vs. surgery & RT or CCRT = 2.30, 95% CI = 1.63–3.25, $p < 0.0001$), and aspirin use (HRaspirin non-users vs. users = 0.44, 95% CI = 0.30–0.64, $p < 0.0001$). In Model C (Table S2), which adjusted for covariates including age, gender, pathological AJCC stages of cancer, treatment, aspirin use and all comorbidities, the significant independent prognostic factors of survival were pathological AJCC stages of cancer (HRstages I & II vs. stages III & IV = 2.34, 95% CI = 1.66–3.30, $p < 0.0001$), treatment (HRsurgery vs. surgery & RT or CCRT = 2.60, 95% CI = 1.83–3.70, $p < 0.0001$), aspirin use (HRaspirin non-users vs. users = 0.59, 95% CI = 0.38–0.92, $p = 0.0191$), diabetes mellitus (HRno vs. yes = 1.65 (95% CI = 1.20–2.28, $p = 0.0022$) and hyperlipidemia (HRno vs. yes = 0.33, 95% CI = 0.20–0.53, $p < 0.0001$). Atrial fibrillation was not an independent prognostic factor for survival in either the univariate (HRno vs. yes = 0.80, 95% CI = 0.35–1.79, $p = 0.5779$) or multivariate Cox regression Model C (HRno vs. yes = 1.03, 95% CI = 0.45–2.40, $p = 0.9376$). Hypertension was no longer an independent prognostic factor for survival in multivariate Cox regression Model C (HRno vs. yes = 0.98, 95% CI = 0.71–1.34, $p = 0.8841$). All the univariate and multivariate Cox regression models used in the analyses indicated that age and gender were not independent prognostic factors for survival (Table 2, 3 and S2). Overall, all of our Cox regression models suggested that OSCC patients who used aspirin were associated with good prognoses, showing a decreased hazard of death that varied from 41–58% compared to non-users (Table 2, 3 and S2).

Table 2

Univariate and multivariate Cox proportional hazards of prognostic factors for OSCC survival, Model A: adjusted for pathological stages of cancer (AJCC) **and other covariates as shown in the table.**

Variables	Cohort n=1525	Hazard Ratio (95%CI)			
		Univariate	p-value	Multivariate	p-value
Gender					
Female	78(5.11%)	1	0.0843	1	0.3474
Male	1447(94.89%)	1.94(0.91-4.11)		1.44(0.67-3.07)	
Age					
Years	1525(100.00%)	1.00(0.99-1.01)	0.7096	1.01(1.00-1.02)	0.1167
Pathological Stages of Cancer (AJCC) ^a					
I & II	892(58.5%)	1	<0.0001	1	<0.0001
III & IV ^b	633(41.5%)	4.22(3.23-5.52)		2.38(1.69-3.35)	
Treatments					
Surgery	857(56.2%)	1	<0.0001	1	<0.0001
Surgery & RT or CCRT	668(43.8%)	4.31(3.27-5.68)		2.55(1.79-3.62)	
Aspirin Use					
No	1220(80.0%)	1	<0.0001	1	<0.0001
Yes	305(20.0%)	0.44(0.30-0.64)		0.42(0.29-0.63)	
Abbreviations: 95% CI 95% confidence interval; RT radiotherapy; CCRT concurrent chemoradiotherapy.					
^a Pathological AJCC Cancer Staging 7 th Edition;					
^b Stages IVa and IVb only;					

Table 3

Univariate and multivariate Cox proportional hazards of prognostic factors for OSCC survival, Model B: adjusted for pathological TN staging system and other covariates as shown in the table.

Variables	Cohort n=1525	Hazard Ratio (95%CI)			
		Univariate	p-value	Multivariate	
					p-value
Gender					
Female	78(5.11%)	1	0.0843	1	0.3384
Male	1447(94.89%)	1.94(0.91-4.11)		1.45(0.68-3.10)	
Age					
Years	1525(100.00%)	1.00(0.99-1.01)	0.7096	1.01(1.00-1.02)	0.0745
Pathological T staging ^a					
T1, 2 & 3	1219(79.93%)	1	<0.0001	1	0.0012
T4	306(20.07%)	3.11(2.43-3.98)		1.57(1.20-2.07)	
Pathological N staging ^b					
N0	1140(74.75%)	1	<0.0001	1	<0.0001
N1, 2 & 3	385(25.25%)	4.25(3.34-5.42)		2.63(1.99-3.47)	
Treatments					
Surgery	857(56.2%)	1	<0.0001	1	<0.0001
Surgery & RT or CCRT	668(43.8%)	4.31(3.27-5.68)		2.30(1.63-3.25)	
Aspirin Use					
No	1220(80.0%)	1	<0.0001	1	<0.0001
				0.44(0.30-0.64)	

Yes	305(20.0%)	0.44(0.30-0.64)
-----	------------	-----------------

Abbreviations: 95% CI 95% confidence interval; RT radiotherapy; CCRT concurrent chemoradiotherapy.

^a Pathological T Staging 7th Edition;

^b Pathological N Staging 7th Edition.

Survival analyses

There were 62 (20.33%) aspirin-using OSCC patients that died during the study period, while 404 (33.11%) deaths were of non-users (Table 1). Figures 2A and B show the Kaplan–Meier curves for overall survival (OS) and disease-specific survival (DSS) at 10 years in OSCC patients with and without 180-day aspirin use. Results of these analyses indicated that aspirin use is associated with improved OS and DSS within a 10-year follow-up period. Figure S1 shows no significant difference in DSS among the patients who took aspirin for fewer than 90 days or within 90–179 days, while OSCC patients who took aspirin consistently had better survival rates compared to non-users. We also evaluated the survival benefit of aspirin use in OSCC patients by comparing the cancer-specific (dead due to OSCC) and other-cause mortality (other causes of death) between aspirin users and non-users across different stages of cancer classified according to the pathological AJCC or TN staging system (Table 4). According to these analyses, the survival benefit conferred by aspirin use was observed in various stages of cancer, including pathological AJCC stages III & IV ($p < 0.0001$), T1, 2 & 3 ($p = 0.0051$), T4 ($p = 0.0124$) and N1, 2 & 3 ($p = 0.0001$) (Table 4).

Table 4

Comparing mortality rates between aspirin users (≥ 180 days) and non-users in OSCC patients (n = 1525) according to *different types of staging systems*.

Cancer Staging System	Causes of Death	N(%)	Low-dose Aspirin		p-value	
			Non-users n = 1220	Aspirin Users ≥ 180days n = 305		
Pathological Stages of Cancer (AJCC) ^a						
I & II	Alive	711(79.71%)	560(78.76%)	151(83.43%)	0.3283 ^e	< 0.0001 ^k
	Death due to OSCC	75(8.41%)	64(9.00%)	11(6.08%)		
	Other Causes of Death	106(11.88%)	87(12.24%)	19(10.50%)		
III& IV ^b	Alive	348(54.98%)	256(50.29%)	92(74.19%)	< 0.0001 ^f	
	Death due to OSCC	189(29.86%)	171(33.60%)	18(14.52%)		
	Other Causes of Death	96(15.17%)	82(16.11%)	14(11.29%)		
Pathological T staging ^c						
T1, 2 & 3	Alive	905(74.24%)	696(72.27%)	209(81.64%)	0.0051 ^g	< 0.0001 ^l
	Death due to OSCC	161(13.21%)	141(14.64%)	20(7.81%)		
	Other Causes of Death	153(12.55%)	126(13.08%)	27(10.55%)		
T4	Alive	154(50.33%)	120(46.69%)	34(69.39%)	0.0124 ^h	
	Death due to OSCC	103(33.66%)	94(36.58%)	9(18.37%)		
	Other Causes of Death	49(16.01%)	43(16.73%)	6(12.24%)		
Pathological N staging ^d						
N0	Alive	873(76.58%)	689(75.71%)	184(80.00%)	0.2179 ⁱ	< 0.0001 ^m
	Death due to OSCC	125(10.96%)	107(11.76%)	18(7.83%)		
	Other Causes of Death	142(12.46%)	114(12.53%)	28(12.17%)		

Cancer Staging System	Causes of Death	N(%)	Low-dose Aspirin		p-value
			Non-users n = 1220	Aspirin Users ≥ 180days n = 305	
N1, 2 & 3	Alive	186(48.31%)	127(40.97%)	59(78.67%)	0.0001 ^j
	Death due to OSCC	139(36.10%)	128(41.29%)	11(14.67%)	
	Other Causes of Death	60(15.58%)	55(17.74%)	5(6.67%)	
^a Pathological AJCC Cancer Staging 7th Edition;					
^b Stages IVa and IVb only;					
^c Pathological T Staging 7th Edition;					
^d Pathological N Staging 7th Edition;					
^{e-j} Aspirin non-users versus users;					
^k Stages I & II versus III & IV;					
^l T1, 2 & 3 versus T4;					
^p N0 versus N1, 2 & 3.					

Discussion

Some studies suggest that aspirin use has beneficial effects on head and neck squamous cell carcinoma (HNSCC)^{16–19}, while other studies do not show this^{20,21}. HNSCC is a collective term for carcinoma arising within the anatomic region of the head and neck, including the oral cavity, oropharynx, hypopharynx, nasopharynx and larynx. Although these tumors are all classified as squamous cell carcinomas, tumors that arise from the different anatomic regions of the head and neck differ in their epidemiology, tumor progression and therapeutic approach^{1,12,21–24}. Thus, we paid particular attention to recruiting patients whose squamous cell carcinomas arose within the oral cavity, including the tongue, buccal area, etc. (Figure S2).

A meta-analysis demonstrates that aspirin use confers a survival benefit in a dose-dependent manner, with a higher dosage of aspirin producing a greater reduction in the risk of colorectal cancer²⁵. In addition, aspirin use has also shown benefits in reducing cancer in a duration-dependent manner across

various cancers^{17,26–28}. In our study, low-dose aspirin (100 mg/day) was prescribed and taken by patients after diagnosis of OSCC. We were able to access and trace the records for the duration of aspirin treatment of these OSCC patients by using the database from Chang Gung Memorial Hospital within a 10-year follow-up period. We classified the duration of aspirin use into three groups and examined whether the survival benefit of aspirin use appeared in a duration-dependent manner. The DSS rate of patients with OSCC who received aspirin for at least 180 days was significantly higher than that of non-users. In contrast, aspirin use had no significant effect on DSS in patients who only received aspirin for either less than 90 days or 90–179 days compared with non-users. Our results extend previous findings^{16,26} suggesting that over a 10-year follow-up period, aspirin use at a low dosage (100 mg) per day for at least 180 days has a survival benefit in patients with OSCC.

With the goal of providing more insight into whether aspirin has the potential to improve survival in OSCC patients, we included the analyses of the associations between aspirin use and mortality rates in different stages of cancer according to the pathological AJCC or TN classification of malignant tumors. Patients who used aspirin for at least 180 days after diagnosis of OSCC showed significant improvement in 10-year OS and DSS compared with non-users, with more marked benefit in late-stage OSCC (stages III & IV) (Table 4). Patients with T (T1, 2 & 3 and T4) and N (N1, 2 & 3) categories who took aspirin after diagnosis of OSCC were associated with increased survival rates (Table 4), while no significant survival differences were observed between aspirin users and non-users with OSCC who were classified as having an early stage of cancer (pathological AJCC stages I & II) and N0 (Table 4). Thus, our results suggest that OSCC patients who had advanced tumors, tumors spread to more lymph nodes, or regional metastases were more likely to benefit from aspirin use.

It is not surprising that aspirin use has beneficial effects for patients who have the aforementioned tumor characteristics. The link between aspirin and cancer metastasis has been widely explored across various cancers^{29–31}. Recently, a study demonstrated that aspirin prevents metastasis by disrupting the activation of endothelial cells, vasoconstriction, and the recruitment of monocytes to tumor sites through inhibition of the COX-1/TXA2 pathway³². The importance of this study along with the others is underscored by the fact that there is a phase III clinical trial, ADD-ASPIRIN, launching now that aims to evaluate the efficacy of aspirin in preventing metastases after patients have been diagnosed with breast, colorectal, gastroesophageal or prostate cancers. Hence, there is a clear possibility that aspirin has great potential to be an effective adjuvant cancer therapy for OSCC patients as well³³.

Conclusions

Our study is an observational retrospective cohort study in which we looked at historical data for a group of OSCC patients with or without aspirin use. A limitation of such a retrospective cohort study is that the OSCC patients who were identified as aspirin users in this study were mostly those who showed signs and symptoms of cardiovascular diseases. For example, the OSCC patients who had risks for heart attack or stroke would be prescribed aspirin as part of the overall cancer treatment plan at the time they

were diagnosed with OSCC. On the other hand, OSCC patients who had very little to no risk for heart attack or stroke would be less likely to take aspirin during their cancer treatment. For this reason, it was expected (as we noted) that our baseline patient characteristics would reflect this clinical setting, with prevalence of comorbidities being higher in aspirin users compared to non-users. With this limitation in mind, comorbidities were additionally adjusted in univariate and multivariate Cox regression analyses for controlling confounding effect. When comparing the Cox regression analyses of models with and without comorbidities as variables, aspirin use, pathological stages of cancer and treatment methods were still associated with OSCC survival even after adjusting for comorbidities. These results suggest that it is less likely that comorbidities would affect the outcome of our study.

Similarly, as a result of the inherent constraints of a retrospective cohort study with historical data, we could not assess the association of aspirin use with bleeding events in OSCC patients due to the lack of data on diagnoses of upper and lower gastrointestinal bleeding. Bleeding from a life-threatening condition such as a hemorrhagic stroke needs immediate emergency care^{34,35}, meaning that we would receive this record in our database. However, as we looked at our study's records of OSCC patients with a history of aspirin use, we did not identify any cases of hemorrhagic stroke that linked to aspirin use in OSCC patients. Numerous studies suggest that the benefit of aspirin use in some cancers is both dose- and duration-dependent^{17,25–28}. We were able to classify the association of survival benefit with duration of aspirin use but unable to examine if there would be a greater survival benefit with an even shorter duration and higher dose of aspirin use. Since the use or longer duration of low-dose aspirin is still associated with a risk of bleeding³⁶, the generalizability of our study might be perfected if we were able to identify whether the survival benefits of aspirin outweigh its potential risks in OSCC patients. This would require jointly examining the survival benefit of aspirin use in OSCC patients based on risk of bleeding, dose, and duration.

The results of this study demonstrate that during a 10-year follow-up period, aspirin use at a low dosage (100 mg) per day for at least 180 days confers a survival benefit on patients with OSCC. Given the poor prognosis of this disease, a treatment like aspirin that could not only improve the OSCC survival rate but is also inexpensive would be highly appreciated.

Declarations

Acknowledgments

We thank Jo-Chi Chin, Huei-Yi Kuo, the Health Information and Epidemiology Laboratory at the Chiayi Chang Gung Memorial Hospital and the Biostatistics Center at Kaohsiung Chang Gung Memorial Hospital for statistics work.

Conflict of Interest Disclosures

We have no *conflicts of interest* to declare.

Funding

This research was funded by Kaohsiung Chang Gung Memorial Hospital, Taiwan, CFRPG8H0391 and CMRPG8H1011.

References

1. Johnson, N. W., Jayasekara, P. & Amarasinghe, A. A. Squamous cell carcinoma and precursor lesions of the oral cavity: epidemiology and aetiology. *Periodontol 2000*.**57** (1), 19–37 <https://doi.org/10.1111/j.1600-0757.2011.00401.x> (2011). **doi**
2. Pires, F. R. *et al.* Oral squamous cell carcinoma: clinicopathological features from 346 cases from a single oral pathology service during an 8-year period. *J Appl Oral Sci*.**21** (5), 460–467 <https://doi.org/10.1590/1679-775720130317> (2013). **doi**
3. **American Cancer Society. Global Cancer Facts & Figs. 3rd Edition.** American Cancer Society. 2015;**800:1–64.**
4. Ferlay, J. *et al.* Global cancer observatory: cancer today. Lyon, France: International Agency for Research on Cancer 2018 **Dec 14.**
5. Taberna, M., Inglehart, R. C. & Pickard, R. K. *et al.* Significant changes in sexual behavior after a diagnosis of human papillomavirus-positive and human papillomavirus-negative oral cancer. *Cancer*.**123** (7), 1156–1165 <https://doi.org/10.1002/cncr.30564> (2017). **doi**
6. **IARC Working Group on the Evaluation of Carcinogenic Risks to Humans, IARC Working Group on the Evaluation of Carcinogenic Risks to Humans.** Meeting, World Health Organization, International Agency for Research on Cancer. Human papillomaviruses. World Health Organization; 2007.
7. Jalouli, J. *et al.* Human papilloma virus, herpes simplex virus and epstein barr virus in oral squamous cell carcinoma from eight different countries. *Anticancer Res*.**32** (2), 571–580 (2012).
8. Macpherson, L. M. D. Raising awareness of oral cancer from a public and health professional perspective. *Br Dent J*.**225** (9), 809–814 <https://doi.org/10.1038/sj.bdj.2018.919> (2018). **doi**
9. Chaturvedi, P., Singh, A., Chien, C. Y. & Warnakulasuriya, S. Tobacco related oral cancer. *BMJ*.**365**, l2142 **Published 2019 Jun 5** <https://doi.org/10.1136/bmj.l2142> (2019). **doi**
10. **GBD 2016 Alcohol Collaborators. Alcohol use and burden for 195 countries and territories, 1990–2016: a systematic analysis for the Global Burden of Disease Study 2016 [published correction appears in Lancet. 2018 Sep 29;392(10153):1116] [published correction appears in Lancet. 2019 Jun 22;393(10190):e44].** *Lancet*. **2018;392(10152):1015–1035.** **doi:10.1016/S0140-6736(18)31310-2**
11. Adel, M., Liao, C. T. & Lee, L. Y. *et al.* Incidence and Outcomes of Patients With Oral Cavity Squamous Cell Carcinoma and Fourth Primary Tumors: A Long-term Follow-up Study in a Betel Quid Chewing

- Endemic Area. *Med. (Baltim)*.**95** (12), e2950 <https://doi.org/10.1097/MD.0000000000002950> (2016). **doi**
12. Krishna Rao, S. V., Mejia, G., Roberts-Thomson, K. & Logan, R. Epidemiology of oral cancer in Asia in the past decade—an update (2000–2012). *Asian Pac J Cancer Prev*.**14** (10), 5567–5577 <https://doi.org/10.7314/apjcp.2013.14.10.5567> (2013). **doi**
 13. Shafique, K., Mirza, S. S. & Vart, P. et al. Areca nut chewing and systemic inflammation: evidence of a common pathway for systemic diseases. *J Inflamm (Lond)*.**9** (1), 22 **Published 2012 Jun 7**<https://doi.org/10.1186/1476-9255-9-22> (2012). **doi**
 14. Nasry, W. H. S., Rodriguez-Lecompte, J. C. & Martin, C. K. Role of COX-2/PGE2 Mediated Inflammation in Oral Squamous Cell Carcinoma. *Cancers (Basel)* 2018;**10**(10):**348**. **Published 2018 Sep 22**. **doi**:10.3390/cancers10100348
 15. Perera, M., Al-Hebshi, N. N. & Perera, I. et al. Inflammatory Bacteriome and Oral Squamous Cell Carcinoma. *J Dent Res*.**97** (6), 725–732 <https://doi.org/10.1177/0022034518767118> (2018). **doi**
 16. Hedberg, M. L., Peyser, N. D. & Bauman, J. E. et al. Use of nonsteroidal anti-inflammatory drugs predicts improved patient survival for PIK3CA-altered head and neck cancer. *J Exp Med*.**216** (2), 419–427 <https://doi.org/10.1084/jem.20181936> (2019). **doi**
 17. Lin, Y. S., Yeh, C. C. & Huang, S. F. et al. Aspirin associated with risk reduction of secondary primary cancer for patients with head and neck cancer: A population-based analysis. *PLoS One*.**13** (8), e0199014 **Published 2018 Aug 22**<https://doi.org/10.1371/journal.pone.0199014> (2018). **doi**
 18. Iovoli, A. J., Hermann, G. M. & Ma, S. J. et al. Association of Nonsteroidal Anti-inflammatory Drug Use With Survival in Patients With Squamous Cell Carcinoma of the Head and Neck Treated With Chemoradiation Therapy. *JAMA Netw Open*.**3** (6), e207199 **Published 2020 Jun 1**<https://doi.org/10.1001/jamanetworkopen.2020.7199> (2020). **doi**
 19. Wilson, J. C., Murray, L. J., Hughes, C. M., Black, A. & Anderson, L. A. Non-steroidal anti-inflammatory drug and aspirin use and the risk of head and neck cancer. *Br J Cancer*.**108** (5), 1178–1181 <https://doi.org/10.1038/bjc.2013.73> (2013). **doi**
 20. Becker, C., Wilson, J. C., Jick, S. S. & Meier, C. R. Non-steroidal anti-inflammatory drugs and the risk of head and neck cancer: A case-control analysis. *Int J Cancer*.**137** (10), 2424–2431 <https://doi.org/10.1002/ijc.29601> (2015). **doi**
 21. Tang, L. *et al.* Association of nonsteroidal anti-inflammatory drugs and aspirin use and the risk of head and neck cancers: a meta-analysis of observational studies. *Oncotarget*.**7** (40), 65196–65207 <https://doi.org/10.18632/oncotarget.11239> (2016). **doi**
 22. Vigneswaran, N. & Williams, M. D. Epidemiologic trends in head and neck cancer and aids in diagnosis. *Oral Maxillofac Surg Clin North Am*.**26** (2), 123–141 <https://doi.org/10.1016/j.coms.2014.01.001> (2014). **doi**
 23. Götz, C. *et al.* Impact of HPV infection on oral squamous cell carcinoma. *Oncotarget*.**7** (47), 76704–76712 <https://doi.org/10.18632/oncotarget.12501> (2016). **doi**

24. Wise-Draper, T. M. *et al.* Future directions and treatment strategies for head and neck squamous cell carcinomas. *Transl Res*.**160** (3), 167–177 <https://doi.org/10.1016/j.trsl.2012.02.002> (2012). **doi**
25. Bosetti, C., Santucci, C., Gallus, S. & Martinetti, M. Aspirin and the risk of colorectal and other digestive tract cancers: an updated meta-analysis through 2019. *Ann Oncol*.**31** (5), 558–568 <https://doi.org/10.1016/j.annonc.2020.02.012> (2020). **La Vecchia Cdoi**
26. Macfarlane, T. V., Murchie, P. & Watson, M. C. Aspirin and other non-steroidal anti-inflammatory drug prescriptions and survival after the diagnosis of head and neck and oesophageal cancer. *Cancer Epidemiol*.**39** (6), 1015–1022 <https://doi.org/10.1016/j.canep.2015.10.030> (2015). **doi**
27. Rothwell, P. M. *et al.* Effect of daily aspirin on risk of cancer metastasis: a study of incident cancers during randomised controlled trials. *Lancet*.**379** (9826), 1591–1601 [https://doi.org/10.1016/S0140-6736\(12\)60209-8](https://doi.org/10.1016/S0140-6736(12)60209-8) (2012). **doi**
28. Luo, S. D., Chen, W. C. & Wu, C. N. *et al.* Low-Dose Aspirin Use Significantly Improves the Survival of Late-stage NPC: A Propensity Score-Matched Cohort Study in Taiwan. *Cancers (Basel)*.**12** (6), 1551 **Published 2020 Jun 12**<https://doi.org/10.3390/cancers12061551> (2020). **doi**
29. Rothwell, P. M., Wilson, M. & Elwin, C. E. *et al.* Long-term effect of aspirin on colorectal cancer incidence and mortality: 20-year follow-up of five randomised trials. *Lancet*.**376** (9754), 1741–1750 [https://doi.org/10.1016/S0140-6736\(10\)61543-7](https://doi.org/10.1016/S0140-6736(10)61543-7) (2010). **doi**
30. Elwood, P. C., Morgan, G. & Pickering, J. E. *et al.* Aspirin in the Treatment of Cancer: Reductions in Metastatic Spread and in Mortality: A Systematic Review and Meta-Analyses of Published Studies. *PLoS One*.**11** (4), e0152402 **Published 2016 Apr 20**<https://doi.org/10.1371/journal.pone.0152402> (2016). **doi**
31. Algra, A. M. & Rothwell, P. M. Effects of regular aspirin on long-term cancer incidence and metastasis: a systematic comparison of evidence from observational studies versus randomised trials. *Lancet Oncol*.**13** (5), 518–527 [https://doi.org/10.1016/S1470-2045\(12\)70112-2](https://doi.org/10.1016/S1470-2045(12)70112-2) (2012). **doi**
32. Lucotti, S., Cerutti, C. & Soyer, M. *et al.* Aspirin blocks formation of metastatic intravascular niches by inhibiting platelet-derived COX-1/thromboxane A2. *J Clin Invest*.**129** (5), 1845–1862 **Published 2019 Mar 25**<https://doi.org/10.1172/JCI121985> (2019). **doi**
33. Coyle, C., Cafferty, F. H. & Rowley, S. *et al.* ADD-ASPIRIN: A phase III, double-blind, placebo controlled, randomised trial assessing the effects of aspirin on disease recurrence and survival after primary therapy in common non-metastatic solid tumours. *Contemp Clin Trials*.**51**, 56–64 <https://doi.org/10.1016/j.cct.2016.10.004> (2016). **doi**
34. Gorelick, P. B. & Weisman, S. M. Risk of hemorrhagic stroke with aspirin use: an update. *Stroke*.**36** (8), 1801–1807 <https://doi.org/10.1161/01.STR.0000174189.81153.85> (2005). **doi**
35. Huang, W. Y. *et al.* Frequency of Intracranial Hemorrhage With Low-Dose Aspirin in Individuals Without Symptomatic Cardiovascular Disease: A Systematic Review and Meta-analysis [published online ahead of print, 2019 May 13]. *JAMA Neurol*.**76** (8), 906–914 <https://doi.org/10.1001/jamaneurol.2019.1120> (2019). **doi**

36. García Rodríguez, L. A., Martín-Pérez, M., Hennekens, C. H., Rothwell, P. M. & Lanas, A. Bleeding Risk with Long-Term Low-Dose Aspirin: A Systematic Review of Observational Studies. *PLoS One*.**11** (8), e0160046 **Published 2016 Aug 4**<https://doi.org/10.1371/journal.pone.0160046> (2016). **doi**

Figures

Figure 1

Flowchart showing the inclusion and exclusion criteria of the OSCC patients recruited to engage in this cohort study. A total of 7,763 patients who were diagnosed with OSCC were recruited in this study. After matching aspirin users with non-users based on 1:4 propensity score analysis, there was a final inclusion of 1,525 OSCC patients were recruited for data analysis in this study. a Clinical and Pathological AJCC Cancer Staging 7th Edition; b Clinical AJCC Cancer Staging 7th Edition; c Pathological AJCC Cancer Staging 7th Edition.

Figure 2

Kaplan–Meier survival curve of OS and DSS rates between aspirin users (≥ 180 days) and non-users. (A). The median OS rate of non-users was around 8 years and the estimated 10-year OS rate was 45.44%. Although median survival was not reached for aspirin users (≥ 180 days), estimated 10-year OS rate was 55.16%. (B). The estimated 10-year DSS rate of non-users was 65.67%, whereas aspirin users (≥ 180 days) was 86.67%.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Supplementalresultswithtitlepage.docx](#)