Theoretical Studies on The Mechanism of Pd²⁺-Catalyzed Regioselective C-H Alkylation of Indole With MesI-CH₂-CF₃-OTf

Yao Shi

Nanjing Agricultural University - Weigang Campus: Nanjing Agricultural University

Hongsheng Ma

Nanjing Agricultural University - Weigang Campus: Nanjing Agricultural University

Jiaxuan Shao

Nanjing Agricultural University - Weigang Campus: Nanjing Agricultural University

chao deng (✉ chaodeng@njau.edu.cn)

Nanjing Agricultural University - Weigang Campus: Nanjing Agricultural University

<https://orcid.org/0000-0002-0899-3305>

Research Article

Keywords: Reaction mechanism, Indole, C4-H selectivity, Pd²⁺-catalyst, DFT studies

Posted Date: February 16th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-194046/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

The reaction mechanism of Pd^{2+} -catalyzed regioselective C–H alkylation of indole with $\text{MesICH}_2\text{CF}_3\text{OTf}$ has been investigated by the density functional theory calculations. The reaction mechanism mainly contains four steps—C–H activation, oxidative addition, reductive elimination and ligands substitution. From our calculations, we find that the C–H activation step was realized by the acetate anion (^-OAc) assisted CMD process and the transition state of C–H activation process is a square planar configuration. Moreover, the calculation results suggest that the regioselectivity of C–H bond alkylation of indole with $\text{MesICH}_2\text{CF}_3\text{OTf}$ can be ascribed to the different stability of the CMD transition states in C–H activation step and different acidities of C–H bonds.

Introduction

Indole and its derivatives as one kind of privileged and significant N-heterocycles have been widely found in a variety of natural products and drugs.¹ Direct C–H bond functionalization of indoles is often the most efficient way to construct all kinds of substituted indole compounds. Therefore, the methods for regioselective synthesis of C–H bond functionalized indole compounds is of great important.² Among them, the C4–H substituted indoles as the backbone of ergot alkaloids are attractive synthetic targets.³ However, in the past few years there is a big challenge for C4–H bond selective functionalization of indoles comparing with other positions of indoles despite its extensive applications.⁴ In addition, in recent years the introduction of fluoroalkylated groups into organic compounds has become very popular due to it often exhibits potential pharmaceutical applications.⁵

Recently, Shi group reported a $\text{Pd}(\text{OAc})_2$ catalyzed C–H bond fluoroalkylation of indole with 2,2,2-trifluoroethyl(mesityl)iodonium triflate salt ($\text{MesICH}_2\text{CF}_3\text{OTf}$),⁶ The interesting aspect here is that the isolated products are occurred at the C4 position of indoles. To the best of our knowledge, the mechanism of $\text{Pd}(\text{OAc})_2$ catalyzed C4–H bond alkylation of indoles with $\text{MesICH}_2\text{CF}_3\text{OTf}$ has not been studied. In order to account for the observed regioselectivity and understand the detailed reaction mechanism for this reaction. Herein, we perform the theoretic calculations to address the C4 versus C2 selective C–H bond alkylation of indole, which should be valuable for experimental chemists to design better selective C–H bond functionalization of N-heterocycles.

Computational Methods

All theoretic calculations were performed with Gaussian 16 software package.⁷ Structure optimization of reactants, products, intermediates, and transition states was carried out at M06 level of density functional theory in the gas phase,⁸ this DFT method is reasonable and has been applied in numerous theoretical studies of transition metal catalyzed C-H functionalization reactions.⁹ Frequency analysis was also carried out at the same level of theory to identify the optimized structures as local minimum or transition states. Intrinsic coordinate reaction (IRC) calculations were performed to examine the connectivity of every transition state with relevant intermediates.¹⁰ The 6-31G (d) basis set was adopted for C, H, O, N, F and S atoms, and the 6-31G (d, p) basis set was selected to describe hydrogen atoms located on the indole ring. The effective core potential (ECP) Lanl2dz¹¹ and $\zeta f=1.472$ polarization functions¹² were used for Pd atom.

In order to take the solvation and dispersion effects into account, the energetic results were further refined by single-point energy calculations (based on the gas phase obtained geometries) using the same functional in conjunction with SMD solvation model.¹³ In the M06-SMD calculations, dichloromethane (DCM) was used as the solvent according to the experimental conditions, the 6-311++G (d, p) basis set was employed for C, H, O, N, F and S atoms and the SDD¹⁴ and $\zeta f=1.472$ polarization functions were selected for Pd atom. The solvation-corrected Gibbs free energies (298.15K) are used to discuss the energetics for this reaction.

Results And Discussion

General aspects

For the reaction of Pd(OAc)₂ catalyzed C-H alkylation of indole with MesICH₂CF₃OTf, there are two major possibilities: C2-H alkylation and C4-H alkylation. Unexpectedly, the Pd(II)-catalyzed C4-H alkylation of indole with MesICH₂CF₃OTf was experimentally obtained. In order to explain the Pd(II)-catalyzed regioselective C-H alkylation of indole with MesICH₂CF₃OTf, the detailed reaction mechanism shown in Scheme 1 has been provided by us.⁶ Firstly, the mechanism of C4-H alkylation (Path 1A) occurs through the coordination of the indole substrate with active catalyst Pd(OAc)₂ to form intermediate **A**. Then the C4-H bond activation gives a six-membered metallacycle intermediate **B**. Subsequently, the alkylation reagent MesICH₂CF₃OTf comes in and take place an oxidative addition process to afford a Pd(IV) intermediate **C**, which was followed by a reductive elimination (RE) process to yield the intermediate **D**. Finally, the intermediate **D** yields C4-H alkylated products and recovers the Pd(II) catalyst through ligands substitution. The mechanism for C2-H alkylation of indole (Path 1B) is similar to that of C4-H alkylation.

Mechanism of C4-H alkylation

The free energy profile for C4-H alkylation of indole with MesICH₂CF₃OTf was calculated and shown in Fig. 1 (Path 1A). According to the previous studies on Pd(OAc)₂ catalyzed C-H bond cleavage,¹⁵ the most common mechanism for C-H bond activation is concerted metalation-deprotonation(CMD). The reaction

begins with the coordination of indole substrate with the active catalyst $\text{Pd}(\text{OAc})_2$ through the oxygen atom of directing group to form a 16-electron intermediate **2a**, in which one acetate anion (^-OAc) is bidentate and this step is endergonic by 6.1 kcal/mol. In order to retain a vacant coordination position for the CMD process, species **2a** undergoes a facile isomerization to yield the precursor complex (**2a'**) for C-H bond activation. Subsequently, C4-H bond activation occurs at the C4 position of indole through the CMD process with the assistance of acetate anion, the transition state for this step is a six-membered ring **TS2a** and the activation barrier of the C-H bond cleavage is 21.0 kcal/mol. In **TS2a**, the C4-H bond cleavage and C-Pd bond formation occurs simultaneously. Meanwhile, in transition state **TS2a** the configuration of the metal center is square planar.

From species **3a**, it extrudes an acetic acid into the reaction system to give a new 16-electron six-membered cyclic palladium species **4a**, the energy released in the whole C-H activation step is 13.5 kcal/mol. Meanwhile, it's worth noting that this step is irreversible, as the reverse reaction (34.5 kcal/mol) calls for much more energy than the forward reaction (21.0 kcal/mol). Subsequently, the oxidative reagent $\text{MesI}\text{CH}_2\text{CF}_3\text{OTf}$ comes in and undergoes an oxidative addition process to afford the intermediate **5a**. After that the reductive elimination process takes place to give species **6a** via the three-membered ring transition state **TS5a** by spanning a barrier of 13.0 kcal/mol (**4a**→**TS5a**). Especially, these two steps are energetically favorable and exergonic by 16.2 kcal/mol. Finally, the active catalyst $\text{Pd}(\text{OAc})_2$ and C4-H bond alkylated product are formed through ligands substitution with HOAc.

Overall, the energy released in C4-H bond alkylation of indole with $\text{MesI}\text{CH}_2\text{CF}_3\text{OTf}$ catalyzed by $\text{Pd}(\text{OAc})_2$ is 37.3 kcal/mol. And the C-H bond activation process is the rate-determining step of this reaction, which activation energy is 21.0 kcal/mol.

Mechanism of C2-H alkylation

As mentioned in the Introduction, another possible pathway to produce the hypothetical C2-H bond alkylated product was also been considered as shown in Fig. 2. Firstly, the coordination of directing group ketone in indole substrate with active catalyst $\text{Pd}(\text{OAc})_2$ affords a square planar 16-electron intermediate **2b**, followed by C-H activation at the C2 position of indole, which undergoes a CMD process through the five-membered ring transition state **TS2b** to afford a five-membered ring intermediate **3b** followed by dissociation of AcOH, the activation barrier for this step is 23.8 kcal/mol. Subsequently, one molecule of substrate $\text{MesI}\text{CH}_2\text{CF}_3\text{OTf}$ goes through an oxidative addition process attaching to the Pd^{2+} center to produce intermediate **5b**. From intermediate **5b**, the reductive elimination occurs to the Pd(IV) metal center via the transition state **TS5b** to give a square planar configuration intermediate **6b** by overcoming an activation barrier of 20.1 kcal/mol (**4b**→**TS5b**). Finally, the intermediate **6b** take place ligands substitution of AcOH with TfOH to give the C2-H bond alkylated product **P2** (Fig. 2).

According to mechanism shown in Fig. 2, the rate-determining step for the C2-H alkylation of indole with $\text{MesI}\text{CH}_2\text{CF}_3\text{OTf}$ is the C2-H bond activation with an overall activation barrier of 23.8 kcal/mol. Meanwhile, the overall reaction is exergonic by 36.6 kcal/mol.

Comparison between C2-H and C4-H alkylation of indole with MesICH₂CF₃OTf

Based on the above calculated results (Fig. 1 and Fig. 2), It evidently reveal the fact that the C-H bond activation step is the rate-determining step for these two pathways and the overall activation barrier for C2-H alkylation of indole is 23.8 kcal/mol), which is higher than C4-H bond alkylation of indole (21.0 kcal/mol). Thus the C4-H bond alkylation of indole is kinetically more favorable than C2-H alkylation. The reason for this scenario can be explained by comparing the stabilities of the transition states in CMD process. The transition state (**TS2a**) for C4-H bond alkylation is a six-membered ring TS and the deprotonated hydrogen atom located on phenyl ring. In contrast, the transition state (**TS2b**) in C2-H bond alkylation is a five-membered ring structure and the departing proton located on pyrrole ring, the electron density of pyrrole ring is much greater than phenyl ring. Additionally, we also contend that the relative stabilities of deprotonated intermediates can affect this reaction selectivity through our computational data, the relative free energy of **3a** in C4-H activation is -10.3 kcal/mol, which is much lower than **3b** (-0.2 kcal/mol) in C2-H activation. Therefore, the overall activation barrier for C4-H bond alkylation is lower than C2-H bond alkylation. Meanwhile, the overall energy of C4-H bond alkylation of indole with MesICH₂CF₃OTf is exergonic by 37.3 kcal/mol, which is larger than C2-H bond alkylation (36.6 kcal/mol). Thus the C4-H bond alkylation of indole in Path 1A is clearly more favorable than the C2-H bond alkylation process in both kinetically and thermodynamically. These calculated results are in good consistent with experimental results.

Conclusions

In this work, the detailed reaction mechanism of Pd(II)-catalyzed regioselective C-H alkylation of indole with MesICH₂CF₃OTf is scrupulously explored by our computational studies. Four main steps are contained in the entire catalytic cycle: C-H activation, oxidative addition, reductive elimination and ligands substitution. What's more, we reckon that the acetate anion (⁻OAc) assisted CMD process plays a crucial role in the C-H bond activation after our theoretic research. And our computational results have verified that the relative stabilities of the CMD transition states in C-H bond activation process and different acidities of C-H bonds can control the regioselectivity of C-H bond alkylation of indole. On the one hand, the activation barrier of C4-H bond alkylation (21.0 kcal/mol) is lower than C2-H alkylation (23.8 kcal/mol), It results in the process of C4-H bond alkylation is more feasible to occur. On the other hand, the overall energy released in the process of C2-H bond alkylation (36.6 kcal/mol) of indole with MesICH₂CF₃OTf is lower than C4-H bond alkylation (37.3 kcal/mol). In a word, the C4-H bond alkylation of indole in Path 1A is more favorable than the C2-H bond alkylation process in both kinetically and thermodynamically. We anticipate our efforts could provide experimentalists with a rational perspective that design the substrates and extend the scope of N-heterocycles selective C-H functionalization.

Declarations

Supporting Information Electronic supplementary information available at

<https://doi.org/10.1007/xxxxxx>.

Author's contributions Y. Shi: investigation. H. Ma: writing original draft. J. Shao: data collection. C. Deng: supervision.

Funding We thank the Bioinformatics Center of Nanjing Agricultural University and the Opening Project of PCOSS, Xiamen University (201909).

Compliance with ethical standards

Conflict of interest The authors declare that they have no competing interests.

Consent for publication Written informed consent for publication was obtained from all participants.

References

1. For books and reviews, see: (a) R.-J. Sundberg, *The Chemistry of Indoles*, Academic Press, New York, **1970**; (b) Y. Ban, Y. Murakami, Y. Iwasawa, M. Tsuchiya and N. Takano, *Res. Rev.* **1988**, *8*, 231. (c) R.-J. Sundberg, *Indoles*, Academic Press, San Diego, **1996**; (d) U. Pindur and T. Lemster, *Curr. Med. Chem.* **2001**, *8*, 1681. (e) A. Ramirez and S. Garcia-Rubio, *Curr. Med. Chem.* **2003**, *10*, 1891. (f) A.-J. Kochanowska-Karamyan and M.-T. Hamann, *Chem. Rev.* **2010**, *110*, 4489. (g) G.-R. Humphrey and J.-T. Kuethe. *Chem. Rev.* **2006**, *106*, 2875. (h) S. Cacchi and G. Fabrizi. *Chem. Rev.* **2011**, *111*, 215.
2. W. Gribble, J. C. Badnock, *Heterocyclic Scaffolds II: Reactions and Applications of Indoles*, Springer, Berlin, **2010**.
3. Lanke, K. R. Bettadapur, and K. R. Prabhu. *Org. Lett.* **2016**, *18*, 5496.
4. (a) Q. Liu, Q. Li, Y. Ma and Y. Jia, *Lett.*, **2013**, *15*, 4528. (b) J. Liang, W. Hu, P. Tao and Y. Jia, *J. Org. Chem.*, **2013**, *78*, 5810. (c) Q. L. Xu, L. X. Dai and S. L. You, *Chem. Sci.*, **2013**, *4*, 97.
5. (a) P. Stjernlof, T. Elebring, J. Nilsson, B. Andersson, S. Lagerkvist, K. Svensson, A. Ekman, A. Carlsson and H. Wikström, *Med. Chem.*, **1994**, *37*, 3263. (b) T. Konno, J. Chae, T. Ishihara and H. Yamanaka, *J. Org. Chem.*, **2004**, *69*, 8258. (c) A. Hall, A. Billinton, S. H. Brown, A. Chowdhury, G. M. Giblin, P. Goldsmith, D. N. Hurst, A. Naylor, S. Patel, T. Scoccitti and P. J. Theobald, *Bioorg. Med. Chem. Lett.*, **2008**, *18*, 2684. (d) Y. Qiao, T. Si, M. H. Yang and R. A. Altman, *J. Org. Chem.*, **2014**, *79*, 7122. (e) K. Matcha and A. P. Antonchick, *Angew. Chem., Int. Ed.*, **2014**, *53*, 11960.
6. J. Borah and Z. Shi. *Chem. Commun.*, **2017**, *53*, 3945.
7. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, G. Scalmani, V. Barone, G. A. Petersson, H. Nakatsuji, X. Li, M. Caricato, A. V. Marenich, J. Bloino, B. G. Janesko, R. Gomperts, B. Mennucci, H. P. Hratchian, J. V. Ortiz, A. F. Izmaylov, J. L. Sonnenberg, D. Williams-Young, F. Ding, F. Lipparini, F. Egidi, J. Goings, B. Peng, A. Petrone, T. Henderson, D. Ranasinghe, V. G. Zakrzewski, J. Gao, N. Rega, G. Zheng, W. Liang, M. Hada, M. Ehara, K. Toyota, R. Fukuda, J. Hasegawa, M. Ishida, T. Nakajima, Y. Honda, O. Kitao, H. Nakai, T. Vreven, K. Throssell, J. A.

- Montgomery, Jr., J. E. Peralta, F. Ogliaro, M. J. Bearpark, J. J. Heyd, E. N. Brothers, K. N. Kudin, V. N. Staroverov, T. A. Keith, R. Kobayashi, J. Normand, K. Raghavachari, A. P. Rendell, J. C. Burant, S. S. Iyengar, J. Tomasi, M. Cossi, J. M. Millam, M. Klene, C. Adamo, R. Cammi, J. W. Ochterski, R. L. Martin, K. Morokuma, O. Farkas, J. B. Foresman, and D. J. Fox, Gaussian, Inc., Wallingford CT, **2016**.
8. (a) Y. Zhao and G. Truhlar, *J. Phys. Chem. A* **2006**, *110*, 13126. (b) Y. Zhao and D.-G. Truhlar, *Theor. Chem. Acc.* **2008**, *120*, 215. (c) D.-G. Truhlar, *J. Am. Chem. Soc.* **2008**, *130*, 16824. (d) Y. Zhao and D.-G. Truhlar, *J. Chem. Theory Comput.* **2009**, *5*, 324.
9. (a) F. Ye, S. Qu, L. Zhou, C. Peng, C. Wang, J. Cheng, M.-L. Hossain, Y. Liu, Y. Zhang, Z.-X Wang, and J. Wang. *Am. Chem. Soc.* **2015**, *137*, 4435. (b) Y-Y. Jiang, X. Man and S. Bi, *Sci. China Chem.* **2016**, *59*, 1448. (c) Y. Dang, X. Deng, J. Guo, C. Song, W. Hu and Z. Wang. *J. Am. Chem. Soc.* **2016**, *138*, 2712. (d) H. Zhang, H.-Y. Wang, Y. Luo, C. Chen, Y. Cao, P. Chen, Y.-L. Guo, Y. Lan and G. Liu, *ACS Catal.* **2018**, *8*, 2173. (e) L. Zhu, Y.-Y Jiang, X. Fan, P. Liu, B-P. Ling, S. Bi. *Organometallics* **2018**, *37*, 2222. (f) J. Wang, B. Ling, P. Liu, Y. Liu, Y.-Y. Jiang, and S. Bi. *ACS Omega* **2020**, *5*, 15446. (g) C. Deng, Y. Sun, Y. Ren and W. Zhang. *Dalton Trans.* **2019**, *48*, 168. (h) Y. Wang, Y. Shi, J. Shao and C. Deng. *Int. J. Quantum Chem.* **2020**, doi.org/[10.1002/qua.26475](https://doi.org/10.1002/qua.26475).
10. Fukui, *J. Phys. Chem.* **1970**, *74*, 4161. (b) K. Fukui, *Acc. Chem. Res.* **1981**, *14*, 363.
11. P-J. Hay and W-R. Wadt, *Chem. Phys.* **1985**, *82*, 270. (b) P-J. Hay and W-R. Wadt, *J. Chem. Phys.* **1985**, *82*, 284. (c) C-E. Check, T-O. Faust, J-M. Bailey, B-J. Wright, T-M. Gilbert and L-S. Sunderlin, *J. Phys. Chem. A* **2001**, *105*, 8111.
12. A-W. Ehlers, M. Böhme, S. Dapprich, A. Gobbi, A. Höllwarth, V. Jonas, K-F. Köhler, R. Stegmann, A. Veldkamp and G. Frenking, *Phys. Lett.* **1993**, *208*, 111.
13. A-V. Marenich, C-J. Cramer and D-G. Truhlar, *Phys. Chem. B.* **2009**, *113*, 6378.
14. (a) P. J. Hay, W. R. Wadt, *Chem. Phys.* **1985**, *82*, 299. (b) G. Igel-Mann, H. Stoll, H. Preuss. *Mol. Phys.* **1988**, *65*, 1321. (c) D. Andrae, U. Häußermann, M. Dolg, H. Stoll, H. Preuß, *Theoret. Chim. Acta* **1990**, *77*, 123. (d) Y. Zhao and D. G. Truhlar, *Acc. Chem. Res.*, **2008**, *41*, 157.
15. (a) S. Zhang, L. Shi, and Y. Ding. *Am. Chem. Soc.* **2011**, *133*, 20218. (b) Y-Y. Jiang, X. Man and S. Bi, *Sci. China Chem.* **2016**, *59*, 1448. (c) C. Deng, J-X. Zhang, and Z. Lin. *ACS Catal.* **2018**, *8*, 2498. (d) P. Hu, S. Wang, Y. Wang, C. Deng and W. Zhang. *Dalton Trans.*, **2018**, *47*, 14644.

Figures

Figure 1

Free energy diagram for the C4-H alkylation of indole with MesI-CH₂CF₃OTf. The solvation-corrected relative free energies are given in kcal/mol.

Figure 2

Free energy diagram for the C2-H alkylation of indole with MesI $\text{CH}_2\text{CF}_3\text{OTf}$. The solvation-corrected relative free energies are given in kcal/mol.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Onlinefloatimage3.png](#)