

Integrating the prevention of mother-to-child transmission of HIV after AIDS denialism in South Africa: Perspectives of experts and health care workers - A qualitative study

Jean Claude MUTABAZI (✉ mutajeanc@yahoo.fr)

Université de Montréal

Corie Gray

Curtin University

Lorrein Muhwava

University of Cape Town, Department of Medicine

Helen Trottier

Université de Montréal

Lisa Jayne Ware

University of the Witwatersrand

Shane Norris

University of the Witwatersrand

Katherine Murphy

University of Cape Town, Department of Biological Sciences

Naomi Levitt

University of Cape Town, Department of Medicine

Christina Zarowsky

Université de Montréal

Research article

Keywords: integration, health systems, primary health care, Prevention of Mother-to-Child Transmission of HIV, PMTCT, MTCT, HIV/AIDS, South Africa

Posted Date: April 2nd, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-19420/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published on June 26th, 2020. See the published version at <https://doi.org/10.1186/s12913-020-05381-5>.

Abstract

Background Integrating vertical programmes into routine health services under complex socio-political and health system conditions is a priority and a challenge. The successful rollout of Prevention of Mother-to-Child Transmission programmes (PMTCT) in sub-Saharan Africa has decreased Human Immunodeficiency Virus (HIV), reduced child mortality and improved maternal health. In South Africa, PMTCT is now integrated into existing health services and this experience could serve as a relevant example for integrating other programmes into comprehensive primary care. We explored the perspectives of both experts and frontline health workers (FHCWs) in South Africa on PMTCT integration into primary health care (PHC) in the context of post-AIDS denialism using a Complex Adaptive Systems framework.

Methods We conducted a total of 20 in-depth semi-structured interviews; 10 with experts (PMTCT and health system experts, policymakers, researchers and activists) and 10 with FHCWs including clinic managers, nurses and midwives. All interviews were conducted in person, audio-recorded and transcribed. Three investigators collaborated in coding transcripts and used an iterative approach for thematic analysis.

Results Experts and FHCWs agreed on the importance of integrated PMTCT services. Experts reported a slow and partial integration of PMTCT programs into PHC following its initial rollout as a stand-alone program in the aftermath of the AIDS denialism period. Experts and FHCWs diverged on the challenges associated with integration of PMTCT. Experts highlighted bureaucracy, HIV stigma and discrimination and a shortage of training for staff as major barriers to PMTCT integration. In comparison, FHCWs emphasized high workloads, staff turnover and infrastructure issues (such as lack of rooms, small spaces, etc.) as their main challenges to integration. Both experts and FHCWs suggested that working with community health workers, particularly in the post-partum period, helped to address cases of loss to follow-up of women and their babies and to improve linkages to protein-chain reaction (PCR) testing and immunisation.

Conclusions Despite organised efforts in South Africa, experts and FHCWs reported multiple barriers for the full integration of PMTCT programmes in PHC, especially postpartum. Our results suggest opportunities to address operational challenges towards more integrated PMTCT and other health services in order to improve maternal and child health.

Background

Health systems operate in conditions of increasing complexity, in which health care providers interact with patients and their families in the context of multiple diseases or health problems dealt with using modern sophisticated technologies and other inputs (1–3). Within this complex context, interventions to address particular diseases are scaled-up and health systems have no option other than to adapt to these underlying components, that are crucial for successful implementation (4,5). Integrating stand-alone

interventions into comprehensive services at local facilities has evolved as systems and regions including sub-Saharan Africa (SSA) have faced the challenges of limited health service accessibility, critical socio-economic conditions and the double burden of communicable and non-communicable (6–8). Integration of the prevention of mother-to-child transmission (PMTCT) of human immunodeficiency virus (HIV) programme in South Africa can serve as a case study of this adaptation under complex conditions.

The transmission of HIV from mother to child can be reduced by more than 95% with effective interventions during pregnancy, labor, delivery and breastfeeding (9,10). These interventions are known collectively as PMTCT (9,10). The PMTCT cascade encompasses services aiming to virtually eliminate mother-to-child transmission (MTCT) of HIV, reduce child mortality and improve maternal health (11,12). Since first reported in the 1980s, the proportion of infections by MTCT has decreased from an estimated 10% of new HIV cases to less than 2% (13,14). The rollout of PMTCT has been particularly successful where the burden is greatest in low and middle income countries (LMICs) including in SSA and in South Africa (13,14).

The acknowledgement of the impact of perinatal HIV on maternal and child health (MCH) (14,15) saw PMTCT services receive significant attention from various actors at local, national and international levels. These stakeholders included organizations such as the United States President's Emergency Plan for AIDS Relief (PEPFAR) and the Global Fund to fight against AIDS, TB and Malaria (GFATM), United States Agency for International Development (USAID), Centers for Disease Control (CDC) and international non-governmental organizations (NGOs), different governmental initiatives and local health systems (16). This support allowed for the integration of PMTCT into existing health care services (17), with the ambition of achieving virtual elimination of MTCT (18,19).

Over the last two decades, PMTCT has increasingly been rolled-out in primary health care (PHC) in order to maximise impact and benefits to maternal and child health services. PMTCT has been integrated into existing services within PHC or community outreach programmes to prevent other diseases such as tuberculosis, diarrhea, depression and others (20,21). The impact of PMTCT on the improvement of MCH in SSA has been well documented, especially in improving service availability, accessibility and utilization (22,23). Unlike the situation for other health problems, this integration of PMTCT in PHC has provided the opportunity to improve preventive services and manage other common health issues within countries (21), rather than shifting scarce resources away in vertical programmes (24). It is acknowledged that well-resourced programmes, like PMTCT, have had a role to play in improving the quality and ensuring the sustainability of other services offered in the same clinics (25). However, achieving such encouraging results, and in particular sustaining them through national rather than donor funding and management, has increased pressure on some facilities that face shortages in terms of human resources and medical supplies, or that lack required knowledge and training to deliver full service packages among existing staff members (26).

AIDS denialism and PMTCT implementation in South Africa

The history of AIDS denialism and the challenges of initially launching PMTCT programmes in South Africa have been extensively documented (27–29). This history is important to understanding the context of PMTCT rollout in South Africa.

Despite now having the world's largest antiretroviral therapy (ART) programmes and well-established PMTCT rollout (30), South Africa's initial response to HIV was stalled due to political interference in health policy during the time of Thabo Mbeki's presidency from 1999–2008. Mbeki subscribed to the AIDS denialist view that all retroviruses are harmless and that other influences, such as drug abuse, poverty, and antiretroviral medications cause AIDS (31,32). Under his presidency, the government instituted policies denying antiretroviral drugs to AIDS patients and withdrew support from clinics that had started using AZT to prevent mother-to-child transmission of HIV. In 1998, the Treatment Action Campaign (TAC) was founded, campaigning for increased access to treatment, care and support for people living with HIV and measures to reduce new HIV infections. In 2001, TAC initiated legal proceedings against the government's policy of limiting treatment for the purpose of PMTCT. The use of ARV/AZT only resumed in 2002 when the constitutional court ruled that the denial of treatment was an infringement of the fundamental right to healthcare (Sect. 27 & 28) (33). In this time period an estimated 100,000 infants were believed to be infected (30,34).

However, it wasn't until 2008, with a new government elected, that the rapid implementation of PMTCT and ART services was prioritised (35). In the period between the first Nevirapine regimen in 2002 and the national PMTCT plan in 2008 at least 330,000 preventable deaths were estimated to have occurred, with 35,000 babies born with HIV. The delay in the rollout of PMTCT caused inestimable financial and social loss, with HIV infection rates well above other SSA countries that had initiated PMTCT in the early 2000s (36,37).

Since then, substantial achievements in PMTCT have been seen in South Africa. HIV testing coverage of pregnant women is now close to 100% and PMTCT services were provided in almost 98% of all health facilities by 2015 (38). In 2011, MTCT decreased to 2.7%, with ART coverage reaching 87% in 2012 (35,38). These significant results were achieved following changes in National PMTCT to initiate treatment for all women living with HIV regardless of CD4 count, drawing on the experience and leadership developed by key actors during the AIDS denialism and PMTCT pilot period. A timeline of South Africa's PMTCT Guidelines is summarized in the adapted table below (Table 1) (30).

Table 1
Adapted timeline of national PMTCT guidelines, South Africa (30).

PMTCT Guidelines	Year
National PMTCT programme begins after Constitutional Court judgement against the Government: • Single dose nevirapine to the mother in labour, and to the infant within 72 hours of birth.	2002
• AZT to mother from 28 weeks gestation; single dose nevirapine to mother in labour and to infant within 72 hours of birth and HAART for pregnant women.	2004
• AZT from 14 weeks gestation: single dose nevirapine plus tenofovir/3TC in labour; infant prophylaxis with nevirapine for 6 weeks if mother on HAART or formula feeding, or until the end of all breastfeeding if mother not eligible for HAART.	2010
• All pregnant women eligible for HAART, irrespective of CD4 count. Infant prophylaxis with nevirapine for 6 weeks. Women initiating HAART with CD4 < 350 and no other indication for HAART, to stop treatment after all breastfeeding has ceased.	2013
• All pregnant and breastfeeding women eligible for lifelong HAART.	2015
AZT: zidovudine; HAART: highly active antiretroviral therapy	

There are however few reports of research findings regarding the process, successes and failures of PMTCT implementation (19,39,40). Understanding the perspectives, experiences or practices of different health system and PMTCT specialists, alongside frontline health workers who have been involved in daily PMTCT activities at PHC level, could facilitate the further integration of PMTCT services and inform efforts to integrate other vertical programmes into PHC. The objective of this study was to explore these perspectives in the context of post-AIDS denialism in South Africa and document important considerations for the integration of other comprehensive services in maternal and child health care.

Methods

Study framework

For this study, we used an adapted Complex Adaptive Systems (CAS) framework to describe features explaining key components of integration of PMTCT (41). There are four nested levels that can be considered for health care system changes and improvement (42,43). These include: individual patients; care teams (family members, nurses, doctors, pharmacists, etc.); organisation (clinics, hospitals, infrastructure, quality assurance, etc.); and the larger system or environment (public and private regulators, insurance companies, research funders, legislative and policy frameworks, etc.) (42,43). Many local, national and international actors, including individual or country initiatives and bilateral or multilateral organizations, also exist in PMTCT programmes.

CAS has been proposed as a theory to both understand, and work towards, integrated health care (44–46). Health systems are perceived as “complex” because they imply a wide variety of components,

“adaptive” due to the ability to adjust or change over time and “system” because of its group of connections or interdependence (47). Though there is no true consensus on the set of characteristics that define CAS (48), Paul Cilliers (49) proposed a list of important CAS characteristics, later summarised by Maguire et al. (50). These characteristics can be applied to the South African health system and to PMTCT intervention and include: unpredictable, multiple with circular causality, self-organised cooperative synergistic, obey asymmetric statistics, modular, robust, open systems, non-linear and adaptive (these framework features are described in the supplemental material 1). Considering the complexity of the process and environmental/contextual elements, the following framework to understand the integration of PMTCT in PHC was proposed for this study (Fig. 1).

Study design and sample

Health system or PMTCT experts, policymakers, researchers and activists at international, national and provincial levels in South Africa were purposively recruited. Recruitment occurred between June 2016 and October 2017. Eligibility criteria included experience in design or implementation of PMTCT in South Africa and other regional countries or at international level, and fluency in English. Twelve experts were contacted by email and recruited for this study, but two were unable to attend the interview, leaving a final sample of 10.

Frontline health care workers (FHCWs) included clinic managers, PMTCT nurses and midwives in clinics randomly selected from the Public Health Department of Cape Town or Western Cape province, South Africa. Recruitment occurred alongside the recruitment of experts. Eligibility included the following: managers of the local facilities offering PMTCT services, nurses or midwives delivering PMTCT services or dealing with PMTCT related problems in the clinics.

Data collection

Participants were able to complete the face-to-face interview at their workplace or any other suggested quiet setting. A self-administered questionnaire was used to collect sociodemographic information including age, sex, years of experience and institution or centre of affiliation before the interview. Interviews were conducted by JCM. Interviews ranged from 30 to 60 minutes. A semi-structured interview guide was developed through a review of various PMTCT policies and implementation guidelines from both WHO and the South African Government, by JCM and revised by CZ. This included open-ended questions on the experience of PMTCT service integration at PHC. The interview guide was modified over the course of the study, using an iterative process informed by the content of previous interviews. For example, questions regarding health information systems and PMTCT data and related statistics were added to understand how information and datasets are shared among various health professionals and institutions, and how monitoring and evaluation of PMTCT services were carried out. The interviewer used reflective probes to encourage respondents to clarify and expand on their statements. Interviews were ongoing until no new information was obtained and thematic saturation was achieved. Participants were offered no compensation.

Data analysis

All interviews were audiorecorded and a verbatim transcription was conducted by a hired assistant. Using an iterative approach to thematic analysis (51,52), two investigators (JCM and CZ) independently read de-identified transcripts, identifying meaningful segments within the responses. All interviews were coded by JCM in collaboration with CG and CZ, and all discrepancies in the coding process were negotiated between these three investigators. The CAS framework was first applied in the process of identifying relevant categories and later in the interpretation of the themes that emerged from the transcripts. General themes and subthemes summarizing perspectives of the two groups of participants (experts and FHCWs) were discussed among members of the research team, and we collectively selected the representative quotes presented in this article. Atlas.ti software was used to assist data analysis and management. Characteristics of participants were analyzed using SPSS version 24.0.

Ethics approval

The study was approved by the Human Research Ethics Committee at the Faculty of Health Sciences, University of Cape Town (HREC REF: 946/2014) and the Centre de Recherche du Centre Hospitalier de l'Université de Montréal (2018–6690, 17.044 - ID). Written informed consent was obtained from all participants.

Results

Participant characteristics

Twenty participants were interviewed; 10 experts and 10 frontline health workers. The experts or key informants included national and international HIV policy makers and researchers while the frontline health workers included three PHC clinic managers and seven PHC clinic nurses. The majority of interviewees were female ($n = 16, 80\%$), with an age range of 29–61 years (mean age 45 years) (Table 2).

Themes and categories

The key themes that emerged from interviews were grouped into seven categories. These categories were then mapped onto three components of the adapted CAS framework, organised by the occurrence of events throughout PMTCT rollout during and after AIDS denialism. They were classified as: “system” (experiences in trials and PMTCT adoption during AIDS denialism era); “adaptive” (gradual move towards PMTCT acceptance and integration, increased awareness and community participation, commitment to PMTCT services integration, community health workers' involvement, clinical training and retraining for nurses) and “complex” (persistent barriers to PMTCT integration). This classification with illustrative quotes is depicted in Table 3.

Table 3
Key findings based on proposed CAS components

CAS component and key findings	Illustrative quotes
<p>System 1. Experiences in PMTCT trials during after AIDS denialism era</p>	<p>“We had to get special permission to roll out the programme, and... when there was a court instruction to roll it out; but then they were still controlling the roll-out and we were all, like, pilot projects” – Expert 2. “I think over the years what has happened is the pressure on the ground that has resulted in people changing their perspective and focus around PMTCT” – Expert 1. “I’m just thinking back now, in general and for Cape Town in particular,...So basically, we received global funding, and in 2006, we appointed quite a number of PMTCT coordinators, as well as PMTCT registered professional nurses within the PHC setting...” – Expert 4. “It was only AZT and Nevirapine that was given at that time. Then as time goes by then we changed” – FHCW 1. “.....she started to run the PMTCT programme about 15 years ago, lots of patients were lost before that and statistics were quite high when it came to PCRs. They weren’t followed up appropriately. So, I think after the PMTCT programme started, there was somebody actually taking charge, invested in the patient’s wellbeing, and they were being followed up, and if any other issues arose besides PMTCT, it could also be managed more appropriately and be referred” – FHCW 2. “I believe there was a problem with a theft on medication. As a result now, the nurses that are doing ANC, they don’t dispense medication” – FHCW 3.</p>
<p>Adaptive 2. Commitment to PMTCT integration 3. Gradual move towards PMTCT acceptance and integration in the clinics 4. Increased awareness and community participation 5. Community health workers’ involvement 6. Clinical training and retraining for nurses</p>	<p>“And in some settings, actually referring the patient involved having them moving from the facility to another to obtain the care for other conditions; and you’ve got to think of integration in terms of, at least, the making available within the same facility of other services” – Expert 6. “...transferring clients to specialists may cause gaps and some women become missing, even for a short distance..., they disappear within 50 meters to where you send them. They fall in hole in that walk and with this we can’t defragment the system” – Expert 10. “It’s a logical move for midwives and other health workers who are working in maternity, antenatal, and postnatal and delivery services to treat mothers who are infected with HIV in a holistic way” – Expert 3. “It’s quite good, because we try to integrate services.....So we do the PCRs when the mothers are here for immunisation, I mean, it can be sooner done at once” – FHCW 3. “You want your baby to be immunised here; then we don’t want you to go up and down, you must also be here and your baby, so that we can see both of youBut the mother and the baby must be on the same facility” FHCW 4. “...because they see mother and child, then they also check which medication did the child get, or is the child a high risk infant or not? Does the mother still remain in care? All those check ups, PAP smears, they also check immunisations, deworming, nutrition,...everything possible” – FHCW 5. “...PMTCT figures almost synonymous with antenatal care, because the chances of being HIV infected are so high, relatively, that you almost don’t see one without the other” – Expert 8. “So most of our nurses, our midwives are now trained, so which means they’re offering integrated antenatal and antiretroviral treatment (PMTCT) and they’re also doing TB screening” – Expert 4. “Western Cape was the only province to start integration of PMTCT and other ANC services and others did the same later. Different stakeholders used to meet once per</p>

CAS component and key findings	Illustrative quotes
	<p>week to discuss the progress and issues and everyone came together to assess how it was managed” – Expert 9.</p> <p>“The community is more open-minded now of what is expected of them once they are pregnant...So, it was an eye-opener, also for the nurses, because and the community as well; because now they know, once they are pregnant, they are eager to know their status” – FHCW 6.</p> <p>“...within the facility, it’s now good that we realised that there is a need for that continuity of care.....when we started this project of Child Health” – FHCW 1.</p> <p>“I can say one in five years, we have only one child, one in five years that is positive. She didn’t come to the clinic when she was pregnant, and yet she was positive. You see, it can take one in five years for one child due to PMTCT; and we are not happy with that” – FHCW 4.</p> <p>“It has so many adverse consequences affecting key public health indicators, such as maternal and under five and infant mortality, that it became a priority health problem for the health services to solve” – Expert 3.</p> <p>“...bringing HIV care back to the community, now increasingly having what they call community-based club, and you are going to see for instance more room for those people, to deal with more urgent issues, because they are overcrowded” – Expert 6.</p> <p>“...community support also plays a key role in supporting moms throughout this whole cascade. Peer support is important” – Expert 10.</p> <p>“We have got programmes like MomConnect where they are enrolled and they get all the support, like where they are sent sms’s and stuff” – FHCW 6.</p> <p>“Mothers to Mothers services which is well supporting the mothers who are pregnant and those after delivery, so they work hand in hand with antenatal labour ward as well as Child Health for continuity of care” FHCW 1.</p> <p>“So, when we talk to them we get the social worker involved as well, then we will try and make a plan to see if there are not other family members because, I mean, any support, it doesn’t just have to be the partner, support in general is good” – FHCW 2.</p> <p>“CHWs are part of health care system to link clinics to the communities and enforce adherence among their other roles. They need to be trained” – Expert 9.</p> <p>“...if they were in place, in many communities, it would be relatively easy to send a message to those community health workers and teams” – Expert 3.</p> <p>“They (CHWs) can go to the houses and they can reinforce adherence...and the idea is that they go into that household and look at all the problems. They look at the teenagers, they counsel them about contraception, and look at the mother that’s struggling with breastfeeding and at the father that’s smoking” – Expert 5.</p> <p>“I will say they are helpful, because we use them as the bridge just to get to the communities, because they go the extra mile in this way that they go inside those houses; then they will go there and give talk...So we work hand in hand with them” – FHCW 7.</p> <p>“...but mostly our community health workers are working with TB and HIV,... Only if, for example, we need to recall our babies for immunisation.... then they will recall those mothers for us. When we are doing some bloods screening and there is an abnormality, they can’t get that person from the phone, and then somebody has to go and do a home visit ” – FHCW 3.</p> <p>“After delivery, CHWs maybe they visit them (women) at home” – FHCW 8.</p> <p>“We trained nurses and they are delivering ART and none believed that in those days, training changes everything...” – Expert 10.</p> <p>“...When everyone diagnosed with HIV was to be initiated to treatment and nurses had to be train on ART initiation. NIMART was created and trained them” – Expert 9.</p> <p>“We said that the NIMART nurse who’s got a dispensing licence can also dispense the medication in the MOU, to make sure that the mom gets more of an integrated service...” – Expert 4.</p>

CAS component and key findings	Illustrative quotes
	<p>“In South Africa, for special training, they have to do the NIMART training, and then they can dispense the drugs and look after the patients and the counselling...” – Expert 5.</p> <p>“We wanted all the midwives to be NIMART trained, so they would be able to initiate, so they don’t send the patients from pillar to post. Then they get tired and don’t start their medication because they must wait in other queues” – FHCW 1.</p> <p>“...just to send the staff to the trainings on PMTCT, on BANC...yes, I see who is running short, so that I send them for training through NIMART training programme” – FHCW 4.</p> <p>“I do BANC and most of the time... I don’t have the HIV course, so I must run around and asking those sisters there for my patients all the time; and if they would send me for HIV training that would be resolved” – FHCW 8.</p> <p>“...because, the staff have to be trained, and especially on the PMTCT guidelines when they change again, so, they change a lot and there should be uniformity” – FHCW 3.</p>
<p>Complex 7. Persistent barriers to PMTCT integration</p>	<p>“If we have an electronic system that talks to each other and the patient has got a single number, we can track her...it’s not installed in all the computers and not everyone is trained how to use it...Other health problems like Hypertension is not integrated into the PMTCT services” – Expert 5.</p> <p>“I think that there was resistance to change. I remember there was a time, just before we introduced the same day initiation. When we spoke to nurses about it, they said it’s impossible, we can’t do it; and so there are attitudinal issues and they needed support and training and encouragement. So people don’t really want to change necessarily, but when it’s policy, you say, well we don’t have a choice” – Expert 7.</p> <p>“Some mothers book late, and so don’t benefit from the services available. Other mothers convert after they’ve been initially tested and they’re especially at high risk, so identifying and getting the mothers onto treatment is one of the big challenges; and then keeping mothers adherent is another challenge” – Expert 3.</p> <p>“We wanted all the midwives to be NIMART trained, so they would be able to initiate, so they don’t send the patients from pillar to post. Then they get tired and don’t start their medication because they must wait in other queues” – FHCW 1.</p> <p>“...for stigma purpose, because they know in our communities if you breastfeed then you are regarded as negative....So we have that challenge, breastfeeding for two weeks then stop, swap to formula feeding” – FHCW 7.</p> <p>“I believe there was a problem with a theft on medication. As a result now, the nurses that are doing ANC, they don’t dispense medication. So now those mothers, they queue twice” – FHCW 3.</p> <p>“Our waiting area is very congested. We don’t even separate them, you know, like kids as such and mothers..., and also there’s poor ventilation in there,... and then they come here and actually, those kids may leave this place more sicker” – FHCW 3.</p>

The following section discusses the results and demonstrates the sequence and complexity of integrating PMTCT services within PHC in South Africa.

1. Experiences in PMTCT trials and its adoption during AIDS denialism era

From the perspectives of HIV experts who were involved in the initial PMTCT trials, during the era of AIDS denialism, doctors and the other health care professionals, community groups and activists representing people living with HIV were the first to push for access to HIV treatment. These early actors in the campaign for treatment for mothers and children had tremendous and diverse experiences of how PMTCT was rolled-out during and after AIDS denialism. Several experts interviewed had participated in the design and implementation of PMTCT in local pilot sites. FHCWs interviewed had not been involved in the early stage of PMTCT rollout.

Experts, who to some extent played a key role in advocacy for HIV treatment or against AIDS denialism, recalled the introduction of PMTCT. The majority discussed how unsupportive the Department of Health (DoH) was, and the difficulty in obtaining approvals at local clinics. Six experts who were involved in the initial PMTCT trials in South Africa described how the PMTCT intervention was vertically rolled-out in selected pilot sites prior to and following the constitutional court ruling in August 2002.

“In South Africa, when we started the PMTCT Programme, it was really a programme implemented in 18 pilot sites, and then based on a court order, we had to scale it up nationally, and at that time, it was very much a vertical programme” – Expert 1.

Though campaigns to prevent MTCT had legal support, it took time for the DoH and all health care providers to fully cooperate with the teams that were delivering HIV interventions. Researchers and activists who were involved in the scaling-up of PMTCT interventions recounted how they personally sought support from health care providers, clinic committees and from pregnant women involved. Bringing together these different groups of stakeholders was essential in gaining support for implementing complex interventions such as PMTCT:

“We would first meet with the clinic... Every clinic had a clinic board made up of lay people and community leaders in that district; and so, the first strategy was you get those people to be on your side” – Expert 2.

In contrast, the majority of the FHCWs knew very little about how PMTCT services were first implemented into PHC in South Africa. One clinic manager assumed that PMTCT had been integrated from the beginning.

“It’s always been integrated, so I don’t want to say something that I don’t have experience of. I don’t want to talk based on assumptions” – FHCW 1.

Another clinic manager who worked at the pilot site was unable to recall what happened when PMTCT services were offered for the first time in her facility. Like others with the same experience, stories were short, imprecise and with few details on how PMTCT services were initiated and later integrated into PHC.

“We started by piloting it in 1999. We were the pilot site. I must try to think now where I was working in that time of the PMTCT; because I remember these babies, they were HIV affected. You know, it was that

way.... We didn't even have registers, at that time, because we didn't even know how big the problem was at the time" – FHCW 2.

2. Gradual move towards PMTCT acceptance and integration in the clinics

Most experts involved in this study reported that they were among the teams of researchers and activists that were pioneering this intervention and worked to mobilise buy-in and support for PMTCT at an individual and organisational levels. They opposed the government's stand, and worked with clinic boards and community leaders in every district for the acceptance and initiation of PMTCT, as illustrated in the following quote:

"It was in the beginning of AIDS denialism, and so the way we entered into the clinic, was we would first meet with the clinic board, and if the clinic board says, we sanction this, it's very hard for the clinic staff not to allow you into the hospital" – Expert 2.

According to the experts who witnessed the whole evolution of PMTCT scale-up in South Africa, despite the Constitutional Court ruling in favour of PMTCT rollout, research teams, activists and non-government organisations (NGOs) continued to face challenges in efforts to implement the intervention in most of the provinces. Those championing the rollout encountered resistance from FHCWs who worried about HIV transmission to themselves and that the intervention would add to already heavy workloads. Meanwhile, HIV-related health care needs were increasing in the country, especially among pregnant women attending clinics for antenatal care, which put healthcare workers under increasing pressure to respond. Health care research teams were also inspired by the civic-minded attitude of those women who willingly participated in early PMTCT trials to support efforts to rollout services to more women. These women were hailed as champions:

"... When there was a court instruction to roll it [PMTCT] out; we were all, like, pilot projects. We were doing rollouts, but it was always very controlled.... it was a placebo-controlled study, and the women said to us, "we don't mind being on placebo, as long as you promise us that if this drug works, you will roll it out, and you will give it to all other [women]... So, we will sacrifice ourselves for the future women". So we promised the women that if it worked, we would do whatever it took to make sure that other women in their situation [received the intervention]..." – Expert 2.

All FHCWs interviewed spoke unreservedly about the benefits of the PMTCT programme, reporting that they had progressively embraced new initiatives and expressed their full support for PMTCT integration into PHC services. They saw how PMTCT integration at a facility level helped to deliver services to community, and how this had become crucial for the successful delivery of other services for maternal and child health. A researcher who participated in PMTCT rollout from the very beginning and a manager working at the PHC level discussed how integration of PMTCT services in local health facilities benefited other services.

“..Mother to child transmission is such a big problem. It has so many adverse consequences affecting key public health indicators, such as maternal and under five and infant mortality, that it became a priority health problem for the health services to solve. So the natural home for PMTCT is within the main stream services” – Expert 6.

“With the integration, it’s the best thing that ever happened, and also where we are now, compared to ten years ago, it’s like a big improvement; from the infection perspective-wise, of cross-infection and stuff, from mother to child” – FHCW 1.

Despite strong support from women and FHCWs, the rollout faced a number of challenges. Addressing HIV related stigma and increasing programme acceptance required support from community-based NGOs. NGOs included those with a special focus on treatment like Médecins Sans Frontières (MSF) and those with activism and advocacy aims like Treatment Action Campaign (TAC). Experts and FHCWs who worked with NGOs on a daily basis in their local facilities, praised them as key players in PMTCT intervention acceptability, in retaining women, or in boosting support from families of women attending PMTCT with their babies after delivery. Four FHCWs and three experts discussed how NGOs have helped from the very beginning - from awareness campaigns, assisting in the training of lay health workers and equipping clinics, as well as providing support and follow-up with women in their communities. NGOs using community health workers (CHWs) and lay counselors continue to play a vital role in working with national health department to reinforce governmental initiatives dealing with MTCT and increasing women retention.

“...So it has improved a lot based on other NPOs [non-profit organisational staff] that assist us if we have a problem when we can’t find, a patient, then mothers trace them, or the NPOs trace them” – FHCW 3.

3. Commitment to PMTCT services integration

Participants discussed knowledge of health care services integration and their experience in the process of integrating PMTCT cascades in PHC. Experts considered PMTCT integration as the ‘right process’ to bring together clinical activities. All experts believed that PMTCT fitted well within other maternal and child health (MCH) services and felt that it had been smoothly integrated into other available services in the clinics such as tuberculosis treatment, sexually transmitted infections and mental health services.

“It’s linking better with the MCH programmes... On the ground, I think, at facility level, there always is integration” – Expert 1.

FHCWs who had been involved in service delivery at the clinics understood how challenging it had been to integrate various services in their facilities. However, they recognised the importance of PMTCT integration in daily service provision at the clinics. In striving to deliver a package of comprehensive services, PMTCT became the core component among the services offered at their local facilities. For example, one of the FHCWs shared how she worked to ensure all services were integrated for the convenience of women.

“So, what I did, I tried to integrate everything, and to make sure that whoever is working here is going to give a one-stop-shop... They [the woman] are going to get all the services they are supposed to get, instead of going there and there and there” – FHCW 1.

4. Increased awareness and community participation

In addition to combined efforts from both national and provincial health systems, local authorities and NGOs to increase PMTCT awareness, community involvement was also essential to increasing uptake and sustainability of the programme. PMTCT mobilisation and integration campaigns have therefore included community-members to share first-hand messages and experiences to peers, relatives and neighbours and ensure the intervention is widely known and accepted. The following two groups were, based on our interviews, identified to have carried out important advocacy and treatment messages about PMTCT and boosted its successful implementation: mothers-to-mothers, and male partners of women who accessed PMTCT.

Mother-to-mother services

Barriers to accessing services, treatment initiation and adherence to it, were acknowledged by both experts and FHCWs. Mother-to-mother (M2M) was perceived as the core peer-support organisation to offer services that helped to address these challenges. First initiated and registered as an NGO partnering with other governmental, multilateral and individual organisations, M2M started in South Africa employing trained mothers living with HIV to provide psychosocial support to pregnant women and mothers of babies diagnosed with HIV, to promote retention and to encourage disclosure (53). Strong support for mother-to-mother’ initiatives was expressed by both experts and FHCWs:

“Mother-to-mother initiatives have been fantastic for making sure moms come back and take the medication...community support also plays a key role in supporting moms throughout this whole cascade. Peer support is important” – Expert 3.

“Mothers to Mothers services which are supporting the HIV mothers who are pregnant and those after delivery, work hand in hand with antenatal labour ward as well as Child Health for continuity of care” – FHCW 3.

Participants also discussed how the M2M peer support group meetings were forums in which women discussed appropriate solutions in responding to HIV infection in their communities, as well as issues they experience, such as difficulties with treatment adherence, social support and poverty.

Male involvement in PMTCT

Active involvement of males in supporting pregnant women living with HIV was regarded as important by both experts and FHCW. However, experts described how antenatal and postnatal health services in the country are ‘womanised’ - a situation that did not foster male involvement in services. Both experts and FHCWs emphasized the need for more efforts to bring in partners to services to improve PMTCT.

“Male integration is something else we didn’t do well and it needs to be improved a bit...and our health services are female oriented and male staff and partners of our women clients are to (should) be more involved” – Expert 3.

FHCWs found male support and collaboration difficult to gain:

“Other contributions were involving other... allowing people to be involved from outside, especially the men, because our African men don’t go to antenatal....to promote the partners to come with” – FHCW 3.

5. Community health workers’ involvement

While there is a long history of CHW involvement in PHC, the advent of HIV and of treatment for HIV and tuberculosis in South Africa led to a major increase in the numbers and visibility of CHWs such as lay counsellors. Many were employed by NGOs and funded by external donors but working in close collaboration with public health facilities (54,55). For experts, CHWs were a cornerstone for linking communities and health systems, addressing stigma and discrimination at family and community level, and for supporting medication adherence. They emphasised the need to establish or enhance community-based services (CBS) with CHWs engagement and full collaboration with health systems.

“They (CHWs) are more available than other health workers in the clinics....and if you go back to the HIV story again, I mean, using CHWs is all about bringing HIV care back to the community as they are in charge of what they call community-based club,.... and this gives more room for those nurses and others in clinics to deal with more urgent issues, because they are overcrowded now. It (the club) gives them that repeated contact and a lot of those redundant tasks can obviously be shifted nicely to a community health worker who is trained and accessible, and then that eases the pressure on the clinic” – Expert 4.

Experts valued the contributions of CHWs, especially in responding to HIV and in facilitating PMTCT implementation. Increased training and appropriate payments for CHWs were suggested by experts, to ensure that they are motivated and are able to support communities in partnership with local health facilities. Experts experience was that, despite intentions that NGOs provide adequate training, supervision and payment, this was done so inconsistently.

“That systems of peers and CHWs, they need to be properly trained and paid, that’s my opinion...When they are trained and offer their services, they have to be paid” – Expert 3.

FHCWs also recognised the importance of CHWs at different fronts to link local health facilities and the communities. Three FHCWs stated they had reservations about the involvement of CHWs but acknowledged that they are useful in ensuring follow-up of women and continuous PCR testing and immunisation for babies. Some CHWs were thought not to have the required education to support community members. This meant that continuous training and supervision by the designated clinic nurse or midwife was required to ensure that the right information and quality services were delivered by CHWs as suggested by one clinic manager:

“...because I need first somebody to train them, or maybe somebody senior to them to listen to what they are saying, because people they start to ask questions. We also see that as an option and then some of them don't even have grade 12, so, those kinds of things. So we try to control it...” – FHCW 2.

However, most FHCWs described collaboration with CHWs as critical to the success of the PMTCT programme and facilitating its reach into the surrounding communities. CHWs supported the delivery of health education for a range of health issues, located mothers for appointments if they could not be reached via phone and organised HIV/AIDS support groups at antenatal clinics.

“I will say they are helpful, because we use them as the bridge just to get to the communities, because they go the extra mile in this way that they go inside those houses; then they will go there and give talk... So we work hand in hand with them” – FHCW 5.

6. Clinical training and retraining for nurses

Given the challenge of doctors providing treatment to a large number of people living with HIV, South Africa has adopted a decentralised approach, allowing other health professionals to prescribe treatment. The Nurse Initiated and Managed Anti-Retroviral Therapy (NIMART) training programme was established as an effective task-shifting strategy, following the conclusion of the Streamlining Tasks and Roles to Expand Treatment and Care for HIV (STRETCH) study in South Africa in 2010 by Fairall L. et al. (56). Its role was to train nurses to not only test HIV, but prescribe ART support to pregnant women living with HIV. It was regarded as successful by both experts and FHCWs but they described several challenges in the roll out of this programme. The shortage of NIMART trained nurses and midwives was thought to result in high workload for those trained and was linked to long waiting hours in the queues and poor-quality services.

“Well, I think in the facilities where you've got enough NIMART trained nurses, there isn't a problem...” – Expert 5.

“We wanted all the midwives to be NIMART trained, so they would be able to initiate, so they don't send the patients from pillar to post. Then they get tired and don't start their medication because they must wait in other queues” – FHCW 3.

“Because I do BANC [basic antenatal care] and most of the time... I don't have the HIV course, so I must run around and asking those sisters there for my patients all the time; if they would send me for HIV, then it's going to be okay...” – FHCW 6.

Frequent changes in PMTCT and ART guidelines also presented challenges in ensuring all staff members understood and were implementing updated guidelines.

“I think it is still challenging, particularly when we have to implement new guidelines, because we've got to go through a whole process of training and retraining. And the trouble is with the whole HIV and PMTCT

field, is evidence evolves and guidelines change rapidly or consecutively in close succession, so we need to keep retraining groups of people” – Expert 1.

In particular, several experts and FHCWs mentioned changes in the guidelines about breastfeeding. While HIV mothers were previously recommended to formula feed, mothers who have a suppressed viral load are now being asked to breastfeed consistently. This has resulted in confusion for both staff members and mothers.

“The other challenge was consistent messaging, because you know, different people were saying different things, and there was a lot of confusion about infant feeding” – Expert 1.

7. Persistent barriers to PMTCT integration

Like other complex health programmes, PMTCT rollout has faced challenges in South Africa (Table 4). Experts mentioned attitudinal issues from facility-based health care workers and lack of accountability and bureaucracy at all levels of management in the health system. They also discussed how poor communication and inconsistencies in the fidelity to the clinical algorithm and practice guidelines reduced the overall quality of care, in some facilities.

Table 4
Persistent barriers to PMTCT integration

According to experts	Common to both experts and FHCWs	According to FHCWs
<ul style="list-style-type: none"> • Bureaucratic slowness • Lack of managerial accountability • Poor quality of care including suboptimal fidelity to algorithms • Poor training of staff • Attitudinal issues • Discrimination, stigma • Lack of national • Lack of health information systems and issues relating to electronic records 	<ul style="list-style-type: none"> • Loss to follow-up • Socio-economic issues for women • Women migration • Lack of unique patient identifiers • Resistance to change 	<ul style="list-style-type: none"> • Small working spaces and other infrastructure related challenges like poor ventilation, congestion, etc., • Shortage and work overload • High staff turnover • Guideline changes • Too much paperwork for managers

“I think it’s due to a lack of accountability and a lack of good management at all levels...And currently, there’s no great communication between primary level and hospital level generally, referral both ways” – Expert 6.

Experts suggested that a single health information system has the potential to improve efficiency in patient identification, follow up and reporting and take its integration to a new stage.

“So, we know patients are nomadic... we need linked electronic patient records, and the problem is for HIV care for example in this city, they have three different systems” – Expert 5.

FHCWs also expressed frustration regarding the lack of communication between clinics and their referral hospitals and the lack of a national-wide system of unique identifiers for all patients.

“If we transfer them, they must take over everything, but the problem is, if the patient went to a hospital, even if she went to our Khayelitsha District hospital, if they give them medication there, it doesn't appear, because we don't know where they captured them. So they appear to us as defaulters, up until we must phone them” – FHCW 8.

FHCWs discussed how their heavy workload, inadequate training, staff shortages and high staff turnover affect the quality and outcomes of service delivery in their health settings.

“It was not all smooth sailing, due to staff turnover....I had, like almost, six, seven nurses who were basic antenatal care [BANC] trained, who can do all for the pregnant mother... from the time that their patient is pregnant until the baby is born, and still look after the baby, but then with the staff turnover, they left and it was tough, terrible! I've got only two people who are BANC trained, and these people, it's not like the only thing that they must do. They were doing other things as well, and not only focussing on this [PMTCT]” – FHCW 1.

Discussion

Political support of AIDS denialism at national level resulted in the deaths of many mothers and children, and delayed South Africa's adoption of PMTCT until 2002 (30). Since then, South Africa has been leading the development of PMTCT integrated into PHC in order to address broader maternal and child health care. This study adds to the limited literature on what has enabled, and challenged, the integration of PMTCT, from the perspectives of both FHCWs and experts.

The interviews highlighted that commitment and ground level collaborations between different stakeholders, including policy, research and activist groups, FHCWs and women from the broader community were essential to navigating around the political climate to ensure the integration of PMTCT services. Considering health system and community elements and their interactions is key to the success of interventions or programmes and should be looked at in terms of degree of interrelatedness and actors from health system structures, their understanding and functions (57) as demonstrated in the study conceptual framework. While the rollout of PMTCT was met with resistance early on from FHCWs, they slowly joined the efforts to integrate, agreeing with experts on the importance of PMTCT integration within existing services, especially those related to MCH. As PMTCT care was incorporated into well-equipped facilities where trained doctors and nurses were available, NIMART training contributed to a decentralised approach with a special focus on women and children from general HIV care. As the task-shifting model aimed to overcome issues of low doctor/patient ratio, PMTCT integration was seen as relatively successful by all participants. However, consistent with what happens in other countries,

FHCWs argued that beyond focused training on PMTCT algorithms, workload and infrastructure constraints have impeded work towards full integration and eliminating MTCT in South Africa (58–60). Apart from NIMART that has not been offered to all nurses due to many not reaching the qualification threshold, PMTCT protocols have continued to evolve and require compulsory both off-and on-the-job training – which not all FHCWs are able to attend, resulting in PMTCT integration in some local facilities being limited.

The need to involve other groups outside of the health system from the outset for training, treatment, advocacy and community mobilisation (such as MSF, TAC, M2M or other well-equipped NGOs) to reinforce PMTCT services integration was acknowledged by all study participants. Such collaborative initiatives were required from the very beginning to manage the health needs of pregnant women and their babies (61). NGOs and partnering local community groups supplemented overwhelmed FHCWs to implement various programmes at local health facilities. Despite the criticism of competition for funding rather than cooperation with national, provincial and district health departments (62), this experience of collaboration was positively regarded by all study participants. Mother to mother services were highlighted by both experts and FHCWs as a way to reduce MTCT and strengthen community-based support. Participants credit M2M efforts and services with establishing links between PMTCT mothers and their local PHC facilities, and maintaining them through the many elements of the PMTCT cascade (63). A study conducted in Zimbabwe by Shroufi A. et al. (53) has similarly found that M2M increases retention in PMTCT, owing to their community-based approach. The three M2M core objectives of 1) reducing the number of babies infected with HIV during or after delivery, 2) promoting the health of pregnant women and new mothers and increasing their opportunities to access and effectively use health and life-sustaining ART, and 3) supporting disclosure and fighting stigma and discrimination (64) have been generally achieved in the SA context (65–68).

Participants in this study explained how and why the CHWs are increasingly accepted into PHC. CHWs have emerged as key players in response to HIV/AIDS, tuberculosis and the Ebola virus, among other health problems - particularly in overwhelmed health systems such as South Africa's (69–72). In case of PMTCT services, the CHWs' role has evolved to improve health for both mothers and their infants, as well as for other family members in the households in different areas of health priority such as disease prevention, hygiene, nutrition, family planning, social support and domestic abuse (73–77). Even though critical, investment in CHWs in South Africa has to date been through NGOs partnerships and have not yet been formally institutionalised nor thoroughly documented despite being considered as a key element of PHC re-engineering strategy, launched almost a decade ago (72,78,79). This situation does not motivate nor facilitate the development of CHWs, who are mostly from a low social-economic background, in their commitment to advance health care services in communities and at the local facilities. Continuous in-service training, adequate supervision, proper financial incentives and other benefits could bridge the gaps in CHW's knowledge and ability to work closely with PHCs, issues that currently can raise ethical questions in relation to quality of care (80).

Both experts and FHCWs were enthusiastic about the positive results already achieved by PMTCT but were dismayed by HIV stigma and discrimination, small and insufficient spaces and other infrastructure related challenges in the facilities. They did however diverge regarding other health system challenges that impede the full integration of PMTCT within PHC and consequently the virtual elimination of MTCT in South Africa that is a clear strategic aim of all PMTCT efforts. Experts insisted on bureaucratic slowness and lack of managerial accountability, poor quality of care, including suboptimal fidelity to algorithms and remnant attitudinal issues in the facilities as obstacles. FHCWs expressed concerns with work overload, high staff turnover, loss to follow-up of women, shortage and poor training of staff. These challenges, among others, are well documented to have historically affected the quality of health care that is a constitutional obligation in South Africa (81,82) and have indeed slowed down the progress with regard to PMTCT integration and virtual MTCT elimination in the country (26,83). Participants from both groups suggested approaches and initiatives to address these complex common issues faced by health systems in South Africa and in other LMICs as illustrated in this study conceptual framework. These measures include more spending in infrastructures, adequate training of workforce in various capabilities, use of electronic unique patient identifiers and health information technology, social assistance to facilitate transportation, fight against hunger, and a number of social factors leading to stigma and discrimination that still result in non-disclosure and low adherence rates (84–88). This would be a tremendous step towards fully integrated PMTCT especially in the wake of introduction of WHO recommended option B + adopted in South Africa, in which all positive tested women are initiated to ART regardless of their CD4 cell count (84,85). Once achieved, full integration of PMTCT would serve as a PHC model to adequately deal with multimorbidity epidemics among ageing populations, including women living with HIV (89–91). With more investment from stakeholders and consistent changes in bureaucracy and greater accountability from National Department of Health, PMTCT integration within other basic health services might be revamped and more fully reported on in South Africa (30,92).

Strengths and limitations of the study

- Included both experts and FHCWs who had different perceptions. Diversity in number of years involved, including those who had been involved in pre-2008 PMTCT services.
- Recaptured the history of PMTCT implementation at local PHC facilities and documents the lost lessons or failures that could be remembered in the integration process of other interventions.
- Social desirability may have influenced some responses and those experiencing more challenges may not be captured.
- Experts were from different areas, including policy, research, advocacy among others.

Conclusion

While the integration of PMTCT into PHC has been hailed as a success, this research identified ongoing challenges in the integration process for HIV/PMTCT from both the perspectives of experts and FHCWs.

Existing issues in bureaucracy and accountability presented barriers to full integration of PMTCT. For FHCWs, concerns of heavy workload and infrastructure constraints, and ongoing issues with training and high staff turnover created challenges in the care of both mother and child. South Africa has prioritised the virtual MTCT elimination – addressing such challenges in the integration of PMTCT into PHC will enable South Africa to achieve this.

Abbreviations

AIDS	Acquired Immunodeficiency Syndrome
ART	Anti-Retroviral Therapy
AZT	Zidovudine
CDC	Centers for Disease Control
FHCWs	Frontline Health Care Workers
GFATM	Global Fund to fight against AIDS, TB and Malaria
HAART	Highly Active Anti-Retroviral Therapy
HIV	Human Immunodeficiency Virus
IQR	Inter Quartile Range
LMICs	Low- and Middle-Income Countries
MCH	Maternal and Child Health
MSF	Médecins Sans Frontières
MTCT	Mother-To-Child Transmission
M2M	Mother-to-mother
NGOs	Non-Governmental Organizations
NIMART	Nurse Initiated and Managed Anti-Retroviral Therapy
PCR	Protein-Chain Reaction
PEPFAR	President’s Emergency Plan for AIDS Relief
PHC	Primary Health Care
PMTCT	Prevention of Mother-To-Child Transmission

SD	Standard Deviation
SSA	Sub-Saharan Africa
TAC	Treatment Action Campaign
USAID	United States Agency for International Development
WHO	World Health Organisation

Declarations

Ethics approval and consent to participate

The study was approved by the Human Research Ethics Committee at the Faculty of Health Sciences, University of Cape Town (HREC REF: 946/2014) and the Centre de Recherche du Centre Hospitalier de l'Université de Montréal (2018-6690, 17.044 - ID). Written informed consent was obtained from all participants.

Consent for publication

Not applicable

Competing interests

JCM, CG, LM, LJW, SN, KM, NL, HT and CZ declare no competing interests.

Funding

JCM holds a PhD scholarship of her supervisor Christina Zarowsky from the Canadian Institutes of Health Research (CIHR) under the "Team Grant – Implementation Research in the Prevention and Treatment of Type II Diabetes in Low- and Middle-Income Countries" competition. This funding is for the following randomized trial: integrated health system intervention aimed at reducing type 2 diabetes risk in women after gestational diabetes in South Africa (IINDIAGO). HT holds a salary award (chercheur-boursier) from the "Fond de la recherche en santé- du Québec (FRQ-S) and a salary award (New Investigator Salary Award) from Canadian Institutes of Health Research (CIHR).

Authors' contributions

JCM, CZ, and HT designed the study. JCM conducted interviews, data analysis and wrote the first manuscript under the supervision of CZ and HT. CG, LM, LW, SN, KM, NL, HT and CZ critically revised versions of the manuscript. All authors read and approved the final manuscript for publication.

Availability of data and materials

The datasets analysed during the current study are not publicly available to preserve participant anonymity.

Acknowledgements

We would like to thank the participants for their contribution to this study.

References

1. Sturmborg JP, Martin C. Handbook of Systems and Complexity in Health. Springer Science & Business Media; 2013. 941 p.
2. Cohn J. Leading healthcare in complexity. Nurs Leadersh (Tor Ont). déc 2014;27(4):52-64.
3. Greenhalgh T, Papoutsis C. Studying complexity in health services research: desperately seeking an overdue paradigm shift. BMC Med [Internet]. 20 juin 2018 [cité 14 févr 2020];16. Disponible sur: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6009054/>
4. Hawe P, Shiell A, Riley T. Theorising interventions as events in systems. Am J Community Psychol. juin 2009;43(3-4):267-76.
5. Borgermans L, Marchal Y, Busetto L, Kalseth J, Kasteng F, Suija K, et al. How to Improve Integrated Care for People with Chronic Conditions: Key Findings from EU FP-7 Project INTEGRATE and Beyond. Int J Integr Care [Internet]. [cité 14 févr 2020];17(4). Disponible sur: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5854097/>
6. Temu F, Leonhardt M, Carter J, Thiam S. Integration of non-communicable diseases in health care: tackling the double burden of disease in African settings. Pan African Medical Journal [Internet]. 05 2014 [cité 14 févr 2020];18(202). Disponible sur: <https://www.panafrican-med-journal.com/content/article/18/202/full/>
7. Amon JJ. Preventing HIV infections in children and adolescents in sub-Saharan Africa through integrated care and support activities: a review of the literature. African Journal of AIDS Research. 1 janv 2002;1(2):143-9.
8. Ekman B, Pathmanathan I, Liljestrang J. Integrating health interventions for women, newborn babies, and children: a framework for action. The Lancet. 13 sept 2008;372(9642):990-1000.

9. WHO | Mother-to-child transmission of HIV [Internet]. WHO. [cité 1 mars 2017]. Disponible sur: <http://www.who.int/hiv/topics/mtct/en/>
10. Prevention of mother-to-child transmission (PMTCT) of HIV | AVERT [Internet]. [cité 1 mars 2017]. Disponible sur: <https://www.avert.org/professionals/hiv-programming/prevention/prevention-mother-child>
11. Dionne-Odom J, Welty TK, Westfall AO, Chi BH, Ekouevi DK, Kasaro M, et al. Factors Associated with PMTCT Cascade Completion in Four African Countries. *AIDS research and treatment*. 2016;2016.
12. USAID. Prevention of Mother-to-Child Transmission of HIV (PMTCT) [Internet]. AIDSFree. [cité 21 sept 2017]. Disponible sur: <https://aidsfree.usaid.gov/resources/pkb/biomedical/prevention-mother-child-transmission-hiv-pmtct>
13. Paintsil E, Andiman WA. Update on successes and challenges regarding mother-to-child transmission of HIV. *Curr Opin Pediatr*. févr 2009;21(1):94-101.
14. Cock KMD, Fowler MG, Mercier E, Vincenzi I de, Saba J, Hoff E, et al. Prevention of Mother-to-Child HIV Transmission in Resource-Poor Countries: Translating Research Into Policy and Practice. *JAMA*. 1 mars 2000;283(9):1175-82.
15. Kuhn L, Sinkala M, Thea DM, Kankasa C, Aldrovandi GM. HIV prevention is not enough: child survival in the context of prevention of mother to child HIV transmission. *J Int AIDS Soc*. 11 déc 2009;12:36.
16. Idele P, Hayashi C, Porth T, Mamahit A, Mahy M. Prevention of Mother-to-Child Transmission of HIV and Paediatric HIV Care and Treatment Monitoring: From Measuring Process to Impact and Elimination of Mother-to-Child Transmission of HIV. *AIDS Behav*. 2017;21(Suppl 1):23-33.
17. Wiegert K, Dinh T-H, Mushavi A, Mugurungi O, Kilmarx PH. Integration of Prevention of Mother-to-Child Transmission of HIV (PMTCT) Postpartum Services with Other HIV Care and Treatment Services within the Maternal and Child Health Setting in Zimbabwe, 2012. *PLoS One* [Internet]. 10 juin 2014 [cité 22 sept 2017];9(6). Disponible sur: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4051591/>
18. Gourlay A, Birdthistle I, Mburu G, Iorpenda K, Wringe A. Barriers and facilitating factors to the uptake of antiretroviral drugs for prevention of mother-to-child transmission of HIV in sub-Saharan Africa: a systematic review. *J Int AIDS Soc* [Internet]. 19 juill 2013 [cité 22 sept 2017];16(1). Disponible sur: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3717402/>
19. Bhardwaj S, Carter B, Aarons GA, Chi BH. Implementation Research for the Prevention of Mother-to-Child HIV Transmission in Sub-Saharan Africa: Existing Evidence, Current Gaps, and New Opportunities. *Curr HIV/AIDS Rep*. juin 2015;12(2):246-55.
20. Grimwood A, Fatti G, Mothibi E, Eley B, Jackson D. Progress of preventing mother-to-child transmission of HIV at primary healthcare facilities and district hospitals in three South African provinces. *South African Medical Journal*. 27 janv 2012;102(2):81-3.
21. Tolle MA. A package of primary health care services for comprehensive family-centred HIV/AIDS care and treatment programs in low-income settings. *Trop Med Int Health*. juin 2009;14(6):663-72.

22. Mutabazi JC, Zarowsky C, Trottier H. The impact of programs for prevention of mother-to-child transmission of HIV on health care services and systems in sub-Saharan Africa - A review. *Public Health Rev.* 1 déc 2017;38(1):28.
23. Fokam J, Santoro M-M, Musolo C, Chimbiri I, Chikwenga G, Deula R, et al. Impact of a Holistic Intervention on Pmtct Uptake Within Sub-Saharan Africa: Evidence from 'Save the Families for Africa' in Malawi. *BMJ Global Health.* 1 févr 2017;2(Suppl 2):A49-A49.
24. Pfeiffer J, Montoya P, Baptista AJ, Karagianis M, Pugas M de M, Micek M, et al. Integration of HIV/AIDS services into African primary health care: lessons learned for health system strengthening in Mozambique - a case study. *J Int AIDS Soc.* 20 janv 2010;13:3.
25. Chi BH, Bolton-Moore C, Holmes CB. Prevention of mother-to-child HIV transmission within the continuum of maternal, newborn, and child health services. *Curr Opin HIV AIDS.* sept 2013;8(5):498-503.
26. Sprague C, Chersich MF, Black V. Health system weaknesses constrain access to PMTCT and maternal HIV services in South Africa: a qualitative enquiry. *AIDS Res Ther.* 3 mars 2011;8:10.
27. Fassin D, Schneider H. The politics of AIDS in South Africa: beyond the controversies. *BMJ.* 1 mars 2003;326(7387):495-7.
28. Buckler A. Denialism and the Struggle against AIDS in South Africa. Nattrass N, éditeur. *Journal of Southern African Studies.* 2008;34(3):722-4.
29. Wang J. AIDS denialism and « The humanisation of the African ». *Race & Class.* 1 janv 2008;49(3):1-18.
30. Burton R, Giddy J, Stinson K. Prevention of mother-to-child transmission in South Africa: an ever-changing landscape. *Obstet Med.* mars 2015;8(1):5-12.
31. Duesberg PH, Nicholson JM, Rasnick D, Fiala C, Bauer HH. HIV-AIDS hypothesis out of touch with South African AIDS—A new perspective. *Population.* 2009;45:50.
32. Kalichman SC, Eaton L, Cherry C. "There is no Proof that HIV Causes AIDS": AIDS Denialism Beliefs among People Living with HIV/AIDS. *J Behav Med.* déc 2010;33(6):432-40.
33. Cullinan K, Thom A. *The virus, vitamins and vegetables: the South African HIV/AIDS mystery.* Jacana Media; 2009.
34. Heywood M. Preventing mother-to-child HIV transmission in South Africa: Background strategies and outcomes of the treatment action campaign case against the Minister of Health. *S Afr J on Hum Rts.* 2003;19:278.
35. Barron P, Pillay Y, Doherty T, Sherman G, Jackson D, Bhardwaj S, et al. Eliminating mother-to-child HIV transmission in South Africa. *Bulletin of the World Health Organization.* 2013;91:70-4.
36. Chigwedere P, Seage GR, Gruskin S, Lee T, Essex M. Estimating the Lost Benefits of Antiretroviral Drug Use in South Africa. *J AIDS Journal of Acquired Immune Deficiency Syndromes.* 1 déc 2008;49(4):410-5.

37. Nattrass N. AIDS and the Scientific Governance of Medicine in Post-Apartheid South Africa. *Afr Aff (Lond)*. 1 avr 2008;107(427):157-76.
38. National Department of Health. National consolidated guidelines for the prevention of mother-to-child transmission of HIV (PMTCT) and the management of HIV in children, adolescents and adults. National Department of Health Pretoria; 2015.
39. Aizire J, Fowler MG, Coovadia HM. Operational issues and barriers to implementation of prevention of mother-to-child transmission of HIV (PMTCT) interventions in Sub-Saharan Africa. *Curr HIV Res. mars 2013;11(2):144-59*.
40. Okoli JC, Lansdown GE. Barriers to successful implementation of prevention-of-mother-to-child-transmission (PMTCT) of HIV programmes in Malawi and Nigeria: a critical literature review study. *Pan Afr Med J [Internet]*. 15 oct 2014 [cité 26 sept 2017];19. Disponible sur: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4345230/>
41. Martínez-García M, Hernández-Lemus E. Health systems as complex systems. *American Journal of Operations Research*. 2013;3(01):113.
42. Reid PP, Compton WD, Grossman JH, Fanjiang G. A Framework for a Systems Approach to Health Care Delivery [Internet]. National Academies Press (US); 2005 [cité 24 mai 2019]. Disponible sur: <https://www.ncbi.nlm.nih.gov/books/NBK22878/>
43. Ferlie EB, Shortell SM. Improving the Quality of Health Care in the United Kingdom and the United States: A Framework for Change. *Milbank Q*. juin 2001;79(2):281-315.
44. Edgren L. The meaning of integrated care: a systems approach. *International Journal of Integrated Care [Internet]*. 23 oct 2008 [cité 21 oct 2015];8(4). Disponible sur: <http://www.ijic.org/index.php/ijic/article/view/URN%3ANBN%3ANL%3AUI%3A10-1-100498>
45. Lars Edgren, Keith Barnard. Complex adaptive systems for management of integrated care. *Leadership in Health Services*. 27 janv 2012;25(1):39-51.
46. Nugus P, Carroll K, Hewett DG, Short A, Forero R, Braithwaite J. Integrated care in the emergency department: a complex adaptive systems perspective. *Soc Sci Med*. déc 2010;71(11):1997-2004.
47. Begun JW, Zimmerman B, Dooley K. Health care organizations as complex adaptive systems. *Advances in health care organization theory*. 2003;253:288.
48. McDaniel RR, Lanham HJ, Anderson RA. Implications of complex adaptive systems theory for the design of research on health care organizations. *Health Care Manage Rev*. 2009;34(2):191-9.
49. Cilliers P. Complexity and postmodernism: Understanding complex systems. routledge; 2002.
50. Maguire S, McKelvey B, Mirabeau L, Oztas N. Complexity science and organization studies. In: Clegg SR, Hardy C, Lawrence TB, Nord WR. 2^e éd. London: The SAGE handbook of organization studies; 2006. 921 p.
51. Srivastava P, Hopwood N. A Practical Iterative Framework for Qualitative Data Analysis. *International Journal of Qualitative Methods*. 1 mars 2009;8(1):76-84.

52. Nowell LS, Norris JM, White DE, Moules NJ. Thematic Analysis: Striving to Meet the Trustworthiness Criteria. *International Journal of Qualitative Methods*. 1 déc 2017;16(1):1609406917733847.
53. Shroufi A, Mafara E, Saint-Sauveur JF, Taziwa F, Viñoles MC. Mother to Mother (M2M) Peer Support for Women in Prevention of Mother to Child Transmission (PMTCT) Programmes: A Qualitative Study. *PLoS One* [Internet]. 5 juin 2013 [cité 18 janv 2018];8(6). Disponible sur: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3673995/>
54. Clarke M, Dick J, Lewin S. Community health workers in South Africa: Where in this maze do we find ourselves? *SAMJ: South African Medical Journal*. sept 2008;98(9):680-1.
55. Tsolekile LP, Schneider H, Puoane T. The roles, training and knowledge of community health workers about diabetes and hypertension in Khayelitsha, Cape Town. *Curationis*. 2018;41(1):e1-8.
56. Fairall L, Bachmann MO, Lombard C, Timmerman V, Uebel K, Zwarenstein M, et al. Task shifting of antiretroviral treatment from doctors to primary-care nurses in South Africa (STRETCH): a pragmatic, parallel, cluster-randomised trial. *Lancet*. 8 sept 2012;380(9845):889-98.
57. Kannampallil TG, Schauer GF, Cohen T, Patel VL. Considering complexity in healthcare systems. *Journal of Biomedical Informatics*. 1 déc 2011;44(6):943-7.
58. Davies NECG, Homfray M, Venables EC. Nurse and manager perceptions of nurse initiated and managed antiretroviral therapy (NIMART) implementation in South Africa: a qualitative study. *BMJ Open* [Internet]. 15 nov 2013 [cité 19 janv 2018];3(11). Disponible sur: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3831110/>
59. Jones M, Cameron D. Evaluating 5 years' NIMART mentoring in South Africa's HIV treatment programme: Successes, challenges and future needs. *S Afr Med J*. 22 2017;107(10):839-42.
60. Sanne I, Orrell C, Fox M, Conradie F, Ive P, Zeinecker J, et al. Nurse management is not inferior to doctor management of antiretroviral patients: The CIPRA South Africa randomised trial. *Lancet*. 3 juill 2010;376(9734):33-40.
61. Uwimana J, Zarowsky C, Hausler H, Jackson D. Engagement of non-government organisations and community care workers in collaborative TB/HIV activities including prevention of mother to child transmission in South Africa: opportunities and challenges. *BMC Health Serv Res*. 2 août 2012;12:233.
62. Mate KS, Ngubane G, Barker PM. A quality improvement model for the rapid scale-up of a program to prevent mother-to-child HIV transmission in South Africa. *Int J Qual Health Care*. 1 sept 2013;25(4):373-80.
63. SOUTH AFRICA| m2m.org [Internet]. mothers2mothers | m2m.org. [cité 10 janv 2018]. Disponible sur: <https://www.m2m.org/where-we-work/south-africa/>
64. Besser M. Mothers 2 mothers. *South African Journal of Obstetrics and Gynaecology*. 1 nov 2006;12(3):122-8.
65. Baek C, Mathambo V, Mkhize S, Friedman I, Apicella L, Rutenberg N. Key findings from an evaluation of the mothers2mothers program in KwaZulu-Natal, South Africa. 2007;

66. Lazarus R, Struthers H, Violari A. Starting HIV-positive babies on antiretroviral treatment: Perspectives of mothers in Soweto, South Africa. *Journal of Pediatric Health Care*. 2010;24(3):176-83.
67. Schunder T, Bagchi-Sen S, Canty M. Uneven Geographies and Treatment Challenges of People Living with HIV and AIDS: Perspectives from Geography. In: *Transforming Global Health*. Springer; 2020. p. 245-60.
68. Nieuwoudt SJ, Ngandu CB, Manderson L, Norris SA. Exclusive breastfeeding policy, practice and influences in South Africa, 1980 to 2018: A mixed-methods systematic review. *PloS one*. 2019;14(10).
69. Mwai GW, Mburu G, Torpey K, Frost P, Ford N, Seeley J. Role and outcomes of community health workers in HIV care in sub-Saharan Africa: a systematic review. *J Int AIDS Soc [Internet]*. 10 sept 2013 [cité 21 déc 2017];16(1). Disponible sur: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3772323/>
70. Rachlis B, Naanyu V, Wachira J, Genberg B, Koech B, Kamene R, et al. Community Perceptions of Community Health Workers (CHWs) and Their Roles in Management for HIV, Tuberculosis and Hypertension in Western Kenya. *PLOS ONE*. 22 févr 2016;11(2):e0149412.
71. Gautier L, Hougbedji KA, Uwamaliya J, Coffee M. Use of a community-led prevention strategy to enhance behavioral changes towards Ebola virus disease prevention: a qualitative case study in Western Côte d'Ivoire. *Global Health Research and Policy*. 22 déc 2017;2:35.
72. Schneider H, Hlophe H, van Rensburg D. Community health workers and the response to HIV/AIDS in South Africa: tensions and prospects. *Health Policy Plan*. mai 2008;23(3):179-87.
73. Futterman D, Shea J, Besser M, Stafford S, Desmond K, Comulada WS, et al. Mamekhaya: A pilot study combining a cognitive behavioral intervention and mentor mothers with PMTCT services in South Africa. *AIDS Care*. sept 2010;22(9):1093-100.
74. Shetty AK, Marangwanda C, Stranix-Chibanda L, Chandisarewa W, Chirapa E, Mahomva A, et al. The feasibility of preventing mother-to-child transmission of HIV using peer counselors in Zimbabwe. *AIDS Research and Therapy*. 1 août 2008;5(1):17.
75. Bhutta ZA, Lassi ZS, Pariyo G, Huicho L. Global experience of community health workers for delivery of health related millennium development goals: a systematic review, country case studies, and recommendations for integration into national health systems. *Global health workforce Alliance*. 2010;1(249):61.
76. Lewin S, Munabi-Babigumira S, Glenton C, Daniels K, Bosch-Capblanch X, van Wyk BE, et al. Lay health workers in primary and community health care for maternal and child health and the management of infectious diseases. *Cochrane database of systematic reviews*. 2010;(3).
77. Perry HB, Zulliger R, Rogers MM. Community health workers in low-, middle-, and high-income countries: an overview of their history, recent evolution, and current effectiveness. *Annu Rev Public Health*. 2014;35:399-421.
78. Pillay Y, Barron P. The implementation of PHC re-engineering in South Africa. *Public Health Association of South Africa*. 2011;1-6.

79. National Planning Commission. National Development Plan 2030: Our future-make it work. 2012; Disponible sur: <http://www.psetresearchrepository.dhet.gov.za/document/national-development-plan-2030-our-future-make-it-work>
80. Ormel H, Kok M, Kane S, Ahmed R, Chikaphupha K, Rashid SF, et al. Salaried and voluntary community health workers: exploring how incentives and expectation gaps influence motivation. *Human Resources for Health*. 19 juill 2019;17(1):59.
81. Maphumulo WT, Bhengu BR. Challenges of quality improvement in the healthcare of South Africa post-apartheid: A critical review. *Curationis* [Internet]. 29 mai 2019 [cité 19 févr 2020];42(1). Disponible sur: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6556866/>
82. Stuckler D, Basu S, McKee M. Health Care Capacity and Allocations Among South Africa's Provinces: Infrastructure–Inequality Traps After the End of Apartheid. *Am J Public Health*. 1 janv 2011;101(1):165-72.
83. Goga A, Chirinda W, Ngandu NK, Ngoma K, Bhardwaj S, Feucht U, et al. Closing the gaps to eliminate mother-to-child transmission of HIV (MTCT) in South Africa: Understanding MTCT case rates, factors that hinder the monitoring and attainment of targets, and potential game changers. *South African Medical Journal*. 2018;108(3):17-24.
84. Chersich MF, Newbatt E, Ng'oma K, de Zoysa I. UNICEF's contribution to the adoption and implementation of option B+ for preventing mother-to-child transmission of HIV: a policy analysis. *Globalization and Health*. 1 juin 2018;14(1):55.
85. Etoori D, Kerschberger B, Staderini N, Ndlangamandla M, Nhlabatsi B, Jobanputra K, et al. Challenges and successes in the implementation of option B+ to prevent mother-to-child transmission of HIV in southern Swaziland. *BMC Public Health* [Internet]. 20 mars 2018 [cité 17 oct 2018];18. Disponible sur: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5859825/>
86. Rodriguez VJ, LaCabe RP, Privette CK, Douglass KM, Peltzer K, Matseke G, et al. The Achilles' heel of prevention to mother-to-child transmission of HIV: Protocol implementation, uptake, and sustainability. *SAHARA J*. 19 sept 2017;14(1):38-52.
87. Mahy M, Stover J, Kiragu K, Hayashi C, Akwara P, Luo C, et al. What will it take to achieve virtual elimination of mother-to-child transmission of HIV? An assessment of current progress and future needs. *Sexually transmitted infections*. 2010;86(Suppl 2):ii48-55.
88. Goga AE, Dinh T-H, Essajee S, Chirinda W, Larsen A, Mogashoa M, et al. What will it take for the Global Plan priority countries in Sub-Saharan Africa to eliminate mother-to-child transmission of HIV? *BMC Infect Dis* [Internet]. 16 sept 2019 [cité 29 janv 2020];19(Suppl 1). Disponible sur: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6745780/>
89. Trickey A, May MT, Vehreschild J-J, Obel N, Gill MJ, Crane HM, et al. Survival of HIV-positive patients starting antiretroviral therapy between 1996 and 2013: a collaborative analysis of cohort studies. *The Lancet HIV*. 1 août 2017;4(8):e349-56.
90. Luijckx HD, Lagro-Janssen ALM, van Weel C. Multimorbidity and the primary healthcare perspective. *J Comorb*. 21 avr 2016;6(2):46-9.

91. Sturmberg JP, Botelho RJ, Kissling B. Integrated Multimorbidity Management in Primary Care: Why, What, How, and How To? J Comorb. 1 janv 2016;6(2):114-9.
92. Nicol E, Dudley L, Bradshaw D. Assessing the quality of routine data for the prevention of mother-to-child transmission of HIV: An analytical observational study in two health districts with high HIV prevalence in South Africa. International Journal of Medical Informatics. 1 nov 2016;95:60-70.

Figures




Figure 1

Adapted study framework towards integrated PMTCT in PHC (41–43)