First-in-human AAV Gene Therapy for Tay-Sachs Disease

Miguel Sena-Esteves (✉ Miguel.Esteves@umassmed.edu)

Department of Neurology, University of Massachusetts Medical School, Worcester, MA, USA.

Terence Flotte

University of Massachusetts Medical School <https://orcid.org/0000-0002-8255-0588>

Oguz Cataltepe

University of Massachusetts Medical School

Ajit Puri

University of Massachusetts Medical School

Ana Rita Batista

University of Massachusetts Medical School

Richard Moser

University of Massachusetts Medical School

Diane McKenna-Yasek

University of Massachusetts Medical School

Catherine Douthwright

University of Massachusetts Medical School

Gwladys Gernoux

University of Massachusetts Medical School <https://orcid.org/0000-0002-1268-7001>

Meghan Blackwood

UMass Medical School

Christian Mueller

Department of Pediatrics and Gene Therapy Center University of Massachusetts Medical School

Xuntian Jiang

Washington University School of Medicine <https://orcid.org/0000-0001-9048-7294>

Scot Bateman

University of Massachusetts Medical School

Spiro Spanakis

University of Massachusetts Medical School

Julia Parzych

University of Massachusetts Medical School

Allison Keeler

UMass Medical School

Aly Abayazeed

University of Massachusetts Medical School

Saurabh Rohatgi

University of Massachusetts Medical School

Laura Gibson

University of Massachusetts Medical School

Robert Finberg

UMass Medical School

Bruce Barton

UMass Medical School

Zeynep Vardar

University of Massachusetts Medical School <https://orcid.org/0000-0002-7510-9524>

Mohammed Shazeeb

University of Massachusetts Medical School

Matthew Gounis

University of Massachusetts Medical School

Florian Eichler

Massachusetts General Hospital

Robert Brown

University of Massachusetts Medical School

Douglas Martin

Auburn University <https://orcid.org/0000-0002-3444-8407>

Heather Gray-Edwards

University of Massachusetts Medical School

Article

Keywords: case report, Tay Sachs disease, gene therapy, GM2 gangliosidosis, AAV, 25 adeno-associated virus, clinical trial

Posted Date: February 18th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-195847/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Nature Medicine on February 10th, 2022. See the published version at <https://doi.org/10.1038/s41591-021-01664-4>.

Case Report: First-in-human AAV Gene Therapy for Tay-Sachs Disease

Terence R. Flotte M.D.^{1,7*}, Oguz Cataltepe M.D.^{1,2}, Ajit Puri M.D.⁴, Ana Rita Batista PhD^{3,7}, Richard Moser M.D.², Diane McKenna-Yasek RN³, Catherine Douthwright PhD³, Gwladys Gernoux PhD^{1,7}, Meghan Blackwood^{1,7}, Christian Mueller PhD^{1,7}, Xuntian Jiang PhD⁹, Scot Bateman M.D.¹, Spiro Spanakis M.D.⁵, Julia Parzych M.D.⁵, Allison M. Keeler PhD^{1,7}, Aly Abayazeed M.D.⁴, Saurabh Rohatgi M.D.⁴, Laura Gibson M.D.^{1,6}, Robert Finberg M.D.⁶, Bruce A. Barton PhD⁸, Zeynep Vardar M.D.⁴, Mohammed Salman Shazeeb PhD⁴, Matthew Gounis PhD⁴, Florian S. Eichler M.D.¹⁰, Robert H Brown Jr. D. Phil, M.D.^{3,7}, Douglas R. Martin PhD¹¹, Heather L. Gray-Edwards PhD, DVM^{4,7}, Miguel Sena-Esteves^{3,7} PhD.*

¹Departments of Pediatrics, ²Neurosurgery, ³Neurology, ⁴Radiology, ⁵Anesthesiology, ⁶Internal Medicine, ⁷Horae Gene Therapy Center and Weibo Li Institute for Rare Diseases Research, ⁸Population and Quantitative Health Sciences, University of Massachusetts Medical School; ⁹Department of Medicine and Cardiovascular Division, Washington University School of Medicine, St. Louis, MO; ¹⁰Department of Neurology, Massachusetts General Hospital and Harvard Medical School, Boston, MA; ¹¹Scott-Ritchey Research Center, Department of Anatomy, Physiology and Pharmacology, College of Veterinary Medicine, Auburn University, Auburn, AL

*Correspondence should be addressed to Miguel Sena-Esteves (Miguel.esteves@umassmed.edu, 508-856-4412) or Terence R. Flotte (Terry.Flotte@umassmed.edu, 506-856-2107), Horae Gene Therapy Center, University of Massachusetts Medical School, 368 Plantation Street, Worcester, MA 01605, USA

Keywords: case report, Tay Sachs disease, gene therapy, GM2 gangliosidosis, AAV, adeno-associated virus, clinical trial

Abstract

Tay-Sachs Disease (TSD) is an inherited neurological disorder caused by deficiency of hexosaminidase A (HexA). Preclinical work demonstrated safety and efficacy of CNS gene therapy using AAVrh8-HEXA/HEXB. Here we describe an expanded access trial in two patients with infantile TSD (IND 18225).

Case TSD-001 demonstrated neurodevelopmental regression by 8 months of age and severe seizures by 1 year was treated at 30 months. An equimolar mix of AAVrh8-HEXA and AAVrh8-HEXB (now AXO-AAV-GM2) was administered intrathecally (IT), with 75% of the dose (1×10^{14} vg) delivered to the cisterna magna and 25% at the thoraco-lumbar junction. The second patient (TSD-002) was treated at 7 months of age with 4.2×10^{13} vg by a combination of bilateral thalamic (0.18 mL; 1.5×10^{12} vg per thalamus), and IT infusion (3.9×10^{13} vg). Both patients underwent immunosuppression with sirolimus, corticosteroids, and rituximab.

Injection procedures were well tolerated and have shown no vector-related adverse events to date. CSF HexA activity nearly doubled from baseline and remained stable. In TSD-002 (now 16 months of age), MRI showed stabilization of disease by 3 months post-injection; there now appeared to temporarily deviate from the natural history of infantile TSD but declined again 6 months post-treatment. TSD-001 (now 4.5 years of age) remains seizure-free on the same anti-convulsant therapy as pre-therapy, but TSD-002 developed seizures between 13 and 17 months post-treatment (by 2 years of age).

Administration of AXO-AAV-GM2 by IT and thalamic injections was safe, HexA activity increased in CSF and ongoing myelination was apparent in the younger patient treated at an early symptomatic stage. This study provides early safety and proof-of-concept in humans for treatment of TSD patients by AAV gene therapy.

Introduction

Tay-Sachs disease (TSD) and Sandhoff disease (SD) are recessively inherited and caused by mutations in *HEXA* and *HEXB* genes, respectively, encoding heterodimeric β -N-acetylhexosaminidase A enzyme (HexA). Severity correlates with residual activity and infantile patients (<0.1% of normal) are diagnosed in the first 12-14 months of life¹. Disease symptoms include inability to sit, difficulty swallowing, seizures, and progressive loss of acquired developmental milestones. Presently there is no effective treatment.

Adeno-associated virus (AAV) gene therapy has shown remarkable success in treating neurological disorders including SMA type 1², and AADC deficiency^{3,4}. However global CNS diseases, including TSD and SD, require widespread distribution of therapeutic proteins. Combined AAV delivery to the thalamus and cerebral spinal fluid (CSF) is an effective approach to achieve broad CNS distribution through axonal transport between interconnected structures and CSF flow⁵⁻¹⁸. In mouse, sheep and cat models of these diseases, combination of intraparenchymal and CSF delivery of AAV vectors showed remarkable therapeutic effects^{19,20}. These preclinical studies demonstrated that axonal transport is an evolutionarily conserved mechanism for widespread distribution of lysosomal enzymes in large brains^{21,22}. Here we describe the initial results of an expanded access clinical trial with two infantile TSD children treated by a similar approach using a combination of AAVrh8-HEXA and AAVrh8-HEXB vectors through a combination of thalamic and CSF delivery. This is the first human use of AAV gene therapy for TSD and provides the basis for future studies.

Materials and Methods

Approvals

The expanded access protocol was reviewed by the University of Massachusetts Medical School IRB, IBC, Radiation Safety Committee and Western IRB. External safety oversight was provided

by an independent Data Safety Monitor. The study was approved by the Food and Drug Administration (IND 18225).

Immune Suppressive Regimen

Patients received a regimen²³ that included rituximab (375 mg/m²), IV infusion of methylprednisolone (10 mg/kg) and sirolimus (1 mg/m²). Prednisone (2mg/kg/day) was administered for 90 days and sirolimus was maintained at 3-7 ng/mL for 180 days, both followed by a one-month taper. B-cell levels decreased to <1% of total lymphocytes. Intravenous immunoglobulin (IVIG) was given as needed to maintain serum levels between 700-1,000 mg/dL.

Dosing

The study agent consists of an equimolar mix of AAVrh8-HEXA and AAVrh8-HEXB vectors (now known as AXO-AAV-GM2). TSD-001 was treated at 30 months only by intrathecal (IT) administration due to severe thalamic degeneration. The dose 1×10^{14} vg (vector genomes) was split between cisterna magna (75%; 9mL) and thoracolumbar junction (25%; 3mL) using a SL-10 microcatheter in the intrathecal space²⁴. TSD-002 was treated at 7 months by bilateral thalamic injections followed by IT delivery as above, for a total combined dose of 4.2×10^{13} vg. The trajectory and depth for thalamic injections were based on MRI and CT, and a robotic platform (ROSA, Zimmer Biomet, Warsaw, IN) was used for cannula (Smartflow, MRI Interventions, Irvine, CA) placement. Vector was infused with a micropump (Harvard Apparatus, Holliston, MA) at 3 μ L/min to a total of 180 μ L (1.54×10^{12} vg) per thalamus.

Results

Case Histories

TSD-001 (female) showed developmental delays at 5-6 months, an exaggerated startle reflex at 8 months, macrocephaly, seizures and abnormal myelination at 12 months. Coincident with seizure onset (14 months), TSD-001 was diagnosed with infantile TSD (HEXA mutations: 4-BP insertion c.1274-1278 and c.82C>T p.Gln28X).

TSD-002 (female) was clinically well at enrollment (6 months) having acquired normal milestones, including tripod sitting, but showing lower extremity weakness and cherry red spots. TSD-002 was diagnosed shortly after birth with infantile TSD (HEXA mutations: 1·75kb deletion of exons 11-13; p.Val381X), because of two preceding TSD siblings that exhibited neurodevelopmental delays at 7 months of age, exaggerated startle and in one case death before 3 years.

General Safety

Study procedures were well-tolerated, resulting in no neurologic deficits or other significant acute procedural complications, with exception of a low-grade, transient post-operative fever occurring in the first 48 hours (TSD-002), without any identifiable source. All other adverse events (AEs) were judged to be unrelated to the study (supplementary Table S1). Transaminases were higher than the non-TSD reference, but remained within the range observed TSD²⁵ throughout the study (Figure 1A, B). A very brief (<2 week) deviation from baseline (but remaining within the TSD range) was observed after the operative procedures and was deemed to be due to a combination of anesthesia medications and underlying liver dysfunction in TSD (supplementary Table S2). There were two serious adverse events (SAEs), both unrelated to either vector or infusion procedures. TSD-001 was hospitalized 9 months after treatment (3 months after immune suppression) due to a viral pneumonia, subsequently complicated by ventilator-associated bacterial pneumonia and gastroesophageal reflux-associated aspiration. TSD-001 recovered and was discharged. TSD-002 was hospitalized 4 months after treatment with a febrile urinary tract infection from *Klebsiella pneumoniae* that responded to antibiotics.

Immune Suppression and Immune Responses

Neither patient showed detectable increases in anti-capsid IgG from baseline to 6 months (Figure 1C). This contrasts with a marked increase in anti-capsid IgG serum levels in an amyotrophic lateral sclerosis patient after intrathecal AAV gene therapy (Figure 1C; assay positive control). TSD-001 showed a rise in anti-capsid neutralizing antibodies (NAb), peaking at week 2 and again at 6 months after treatment (Figure 1D). Increases in NAb titers coincided with IVIG treatment

(IVIG not tested for anti-AAV antibodies). Mild T-cell responses to AAVrh8 capsid were detected at months 3·5 and 6 in TSD-001, and up until month 1 in TSD-002 (Figure 1E-F). Positive ELISpot assays in TSD-002 occurred when trough sirolimus levels dropped below 4 ng/mL. Neither patient showed T-cell responses to HEXA or HEXB (Figure 1G-H).

Bioefficacy: Protein and Enzyme Activity in CSF and serum.

In both patients, baseline CSF HexA activity was ~0·3 nmol/hr/mL and increased to 0·5-0·6 nmol/hr/mL after treatment (Figure 2A). Baseline serum HexA activity was ~0·3 nmol/hr/mL in both patients (Figure 2A-B), and increased slightly at all time points (except at 6 months in TSD-002). Western blot of CSF showed faint bands corresponding to HEXA and HEXB proteins (Figure 2D). HEXA became detectable at 6-months for TSD-001 and 3-months for TSD-002 (Figure 2D). Various GM2 ganglioside species in CSF (Figure 1C) were increased in TSD-001 and to a lesser degree in TSD-002, suggesting that GM2 levels may be directly proportional to disease state. There was a disparity between GM2 species, where GM2(22:0) and GM2(24:0) plateaued after treatment whereas remaining species continued to increase.

Radiologic Assessments

Post-injection MRI images showed successful thalamic targeting in TSD-002, without any detectable deleterious effects (Figure 3A). Fluid signal averaged 474 mm³ per thalamus, ~2·6 times the infused volume. At the time of treatment, TSD-001 had advanced cortical atrophy, mild ventricular enlargement and diffusely affected white matter (Figure 3B), but after treatment no further deterioration was apparent on MRI. At baseline, TSD-002 MRI was suggestive of mild dysmyelination that appeared unchanged at 3 months (Figure 3B). At 6 months, areas of increased myelination were noted in several sites, including the posterior corpus callosum. DTI showed minimal changes over time, except in the anterior corpus callosum of TSD-001 (Figure 3C, supplementary Figure S1). Volumetric analyses from TSD-001 suggested stabilization of corpus callosum volumes after treatment. Total brain volume of TSD-002 increased over the 6-

month period, a finding also noted in individual brain structures (Figure 3C; supplementary Figure S2).

Neurologic and Neurodevelopmental Outcomes

CHOP INTEND scores reflected the advanced disease stage of TSD-001 and the early stage in TSD-002. TSD-001 remained stable with a score of ~20 (Figure 2E). Presently, TSD-001 is 4 years and 6 months old has been seizure-free since treatment on the same anticonvulsant therapy (Keppra) and non-adjusted dose that was in place pre-therapy when seizures were active. Electroencephalograms, auditory brainstem responses, and retinal exams did not change significantly through the first 6 months after therapy, after which time data were not available due to health-related travel restrictions.

TSD-002's score remained ~60 until 6 months, when it decreased to 52. At 13 months of age, TSD-002 was still sitting, an important milestone for infantile TSD¹, but with some truncal stability loss and exaggerated startle response. At 16 months of age, TSD-002 could sit for 5 seconds, retained visual tracking and response to auditory stimuli, but this was ability was lost by 24 months of age. Seizures developed by 24 months of age (17 months post-therapy). As with the first patient, electroencephalograms, auditory brainstem responses, and retinal exams did not change significantly through the first 6 months after therapy, after which time data were not available.

Conclusions

This is the first in human test of AAV gene therapy in TSD; this trial achieved a mild increment in CSF HEXA activity, showed good general safety in both patients, and demonstrated feasibility of bilateral thalamic injections in a 7-month-old infant. There were no vector related SAEs, and mild AEs after treatment were judged to be related to procedural aspects of the protocol, including a brief, mild post-operative fever. Transaminase levels remained within the historical range

reported for TSD²⁵. No anti-transgene immune responses were observed, and anti-capsid immune responses were short-lived and mild.

Post-treatment CSF HexA activity in the range observed here could alter the trajectory of TSD, since patients with 0.5% of normal HexA activity exhibit a late infantile/ juvenile phenotype²⁶. HexA levels in CSF may not accurately represent brain tissue activity, but nonetheless MRI of TSD-002 showed some degree of disease burden stabilization and increased myelination in several structures, in agreement with some attenuation of clinical disease progression. If sustained, this would represent a meaningful benefit when compared to siblings and natural course of infantile TSD. This impact occurred in TSD-002 patient despite receiving half the dose of TSD-001, pointing to the value of early treatment. A retrospective study of infantile Tay-Sachs disease showed that 98% of patients developed seizures with an onset at 16.8 ± 5.5 months of age that are often refractory to antiepileptic drugs¹. The observation that TSD-001 has remained free of seizures at 4.5 years of age (24 months post-therapy), in spite of continuing the same anti-convulsant therapy (Keppra) at a dose which had not been adjusted since gene therapy, suggests a positive benefit of the gene therapy intervention. Although this is encouraging, this effect did not persist in TSD-002, where seizures were noted by 24 months of age (17 months after therapy). This is the first report of intrathalamic gene therapy in patients. The technique holds potential for treatment of other diseases, because axonal transport enables global CNS distribution^{21,22}. The thalamus is a high-risk target because of its critical functions as an integration and relay center of the cerebral cortex. Therefore, to minimize risk to the patient, the thalamic injection volume in TSD-002 (180 μ L) was roughly equivalent to that safely injected in non-human primates (150 μ L; data not shown). The dose/volume used corresponds to $\sim 1/10$ of that projected to be most efficacious in pre-clinical experiments. Since the thalamic volume is estimated at $\sim 4,500$ microliters²⁷, and infusate spread was ~ 474 microliters, a substantial increase in dose/volume will be needed to achieve maximal therapeutic benefit. Animal studies showed a 10-fold higher thalamic injection volume is safe; however, thalamic injury in patients is of concern. Since TSD

is a fatal disease, risk associated with injecting larger volumes is warranted and necessary to achieve transformative therapeutic outcomes.

In summary, expanded access use of AXO-AAV-GM2 was safe after intrathecal and bilateral thalamic delivery, and accompanied by a modest increase in CSF HEXA activity. Clinical findings in TSD-002 who was treated at a younger age, suggest a possible deviation from the natural history of TSD despite receiving a lower dose than indicated by preclinical experiments. This expanded access trial provides the basis for future trials of AXO-AAV-GM2 in patients with TSD and SD.

References

1. Bley, A.E., *et al.* Natural history of infantile G(M2) gangliosidosis. *Pediatrics* **128**, e1233-1241 (2011).
2. Mendell, J.R., *et al.* Single-Dose Gene-Replacement Therapy for Spinal Muscular Atrophy. *The New England journal of medicine* **377**, 1713-1722 (2017).
3. Muramatsu, S., *et al.* A phase I study of aromatic L-amino acid decarboxylase gene therapy for Parkinson's disease. *Molecular therapy : the journal of the American Society of Gene Therapy* **18**, 1731-1735 (2010).
4. Hwu, W.L., *et al.* Gene therapy for aromatic L-amino acid decarboxylase deficiency. *Science translational medicine* **4**, 134ra161 (2012).
5. Passini, M.A., *et al.* Intraventricular brain injection of adeno-associated virus type 1 (AAV1) in neonatal mice results in complementary patterns of neuronal transduction to AAV2 and total long-term correction of storage lesions in the brains of beta-glucuronidase-deficient mice. *Journal of virology* **77**, 7034-7040 (2003).
6. Gray, S.J., Nagabhushan Kalburgi, S., McCown, T.J. & Jude Samulski, R. Global CNS gene delivery and evasion of anti-AAV-neutralizing antibodies by intrathecal AAV administration in non-human primates. *Gene therapy* **20**, 450-459 (2013).

- 232 7. Samaranch, L., *et al.* Strong cortical and spinal cord transduction after AAV7 and AAV9
233 delivery into the cerebrospinal fluid of nonhuman primates. *Human gene therapy* **24**, 526-
234 532 (2013).
- 235 8. Hinderer, C., *et al.* Intrathecal gene therapy corrects CNS pathology in a feline model of
236 mucopolysaccharidosis I. *Molecular therapy : the journal of the American Society of Gene*
237 *Therapy* **22**, 2018-2027 (2014).
- 238 9. Meyer, K., *et al.* Improving Single Injection CSF Delivery of AAV9-mediated Gene Therapy
239 for SMA: A Dose-response Study in Mice and Nonhuman Primates. *Molecular therapy :*
240 *the journal of the American Society of Gene Therapy* **23**, 477-487 (2015).
- 241 10. Miyanojara, A., *et al.* Potent spinal parenchymal AAV9-mediated gene delivery by subpial
242 injection in adult rats and pigs. *Molecular therapy. Methods & clinical development* **3**,
243 16046 (2016).
- 244 11. Cearley, C.N., *et al.* Expanded Repertoire of AAV Vector Serotypes Mediate Unique
245 Patterns of Transduction in Mouse Brain. *Molecular therapy : the journal of the American*
246 *Society of Gene Therapy* (2008).
- 247 12. Salegio, E.A., *et al.* Axonal transport of adeno-associated viral vectors is serotype-
248 dependent. *Gene therapy* **20**, 348-352 (2013).
- 249 13. Passini, M.A., Lee, E.B., Heuer, G.G. & Wolfe, J.H. Distribution of a lysosomal enzyme in
250 the adult brain by axonal transport and by cells of the rostral migratory stream. *The Journal*
251 *of neuroscience : the official journal of the Society for Neuroscience* **22**, 6437-6446 (2002).
- 252 14. Kells, A.P., *et al.* Efficient gene therapy-based method for the delivery of therapeutics to
253 primate cortex. *Proceedings of the National Academy of Sciences of the United States of*
254 *America* **106**, 2407-2411 (2009).
- 255 15. Baek, R.C., *et al.* AAV-mediated gene delivery in adult GM1-gangliosidosis mice corrects
256 lysosomal storage in CNS and improves survival. *PloS one* **5**, e13468 (2010).

16. Cearley, C.N. & Wolfe, J.H. A single injection of an adeno-associated virus vector into nuclei with divergent connections results in widespread vector distribution in the brain and global correction of a neurogenetic disease. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **27**, 9928-9940 (2007).
17. Dodge, J.C., et al. Gene transfer of human acid sphingomyelinase corrects neuropathology and motor deficits in a mouse model of Niemann-Pick type A disease. *Proceedings of the National Academy of Sciences of the United States of America* **102**, 17822-17827 (2005).
18. Foust, K.D., Flotte, T.R., Reier, P.J. & Mandel, R.J. Recombinant adeno-associated virus-mediated global anterograde delivery of glial cell line-derived neurotrophic factor to the spinal cord: comparison of rubrospinal and corticospinal tracts in the rat. *Human gene therapy* **19**, 71-82 (2008).
19. Cachon-Gonzalez, M.B., et al. Effective gene therapy in an authentic model of Tay-Sachs-related diseases. *Proceedings of the National Academy of Sciences of the United States of America* **103**, 10373-10378 (2006).
20. Cachon-Gonzalez, M.B., et al. Gene transfer corrects acute GM2 gangliosidosis--potential therapeutic contribution of perivascular enzyme flow. *Molecular therapy : the journal of the American Society of Gene Therapy* **20**, 1489-1500 (2012).
21. Gray-Edwards, H., et al. AAV gene therapy in a sheep model of Tay-Sachs disease. *Human gene therapy* (2017).
22. Rockwell, H.E., et al. AAV-mediated gene delivery in a feline model of Sandhoff disease corrects lysosomal storage in the central nervous system. *ASN Neuro* **7**(2015).
23. Corti, M., et al. B-Cell Depletion is Protective Against Anti-AAV Capsid Immune Response: A Human Subject Case Study. *Molecular therapy. Methods & clinical development* **1**(2014).

24. Taghian, T., *et al.* A Safe and Reliable Technique for CNS Delivery of AAV Vectors in the Cisterna Magna. *Molecular therapy : the journal of the American Society of Gene Therapy* (2019).
25. Schneck, L., Maisel, J. & Volk, B.W. The Startle Response and Serum Enzyme Profile in Early Detection of Tay-Sachs' Disease. *The Journal of pediatrics* **65**, 749-756 (1964).
26. Mahuran, D.J. Biochemical consequences of mutations causing the GM2 gangliosidoses. *Biochim Biophys Acta* **1455**, 105-138 (1999).
27. Tutunji, R., *et al.* Thalamic volume and dimensions on MRI in the pediatric population: Normative values and correlations: (A cross sectional study). *Eur J Radiol* **109**, 27-32 (2018).

Figure legends

Figure 1. *Longitudinal safety and immunological studies.* Serum levels of liver enzymes A) alanine aminotransferase (ALT) and B) aspartate aminotransferase (AST) enzymes in TSD-001 and TSD-002 patients before treatment (Pre-TX), weeks 1, 2, and 3 (W1, 2, 3) and months 1, 2, 3 and 6 (M1, 2, 3, 6). Shaded bars in A) and B) represent the normal interval for both assays. Dashed line in B) indicates the maximum value reported in TSD patients²⁵. C) Total anti-AAVrh8 IgG in serum quantified by ELISA. In C) the gray squares and lines represent anti-capsid IgG levels in serum of an ALS patient treated by IT AAV gene therapy. D) Neutralizing antibody titers to AAVrh8 capsid quantified by a transduction assay. Peripheral blood mononuclear cells (PBMCs) were isolated from TSD-001 and TSD-002 patients before treatment and at various time points thereafter to assess T-cell responses by interferon gamma ELISpot assays using stimulation with pools of overlapping peptides spanning E, F) AAVrh8 VP1, or G, H) human HEXA and HEXB proteins. Unstimulated or CD3/CD28-stimulated PBMCs were included in all assays as negative and positive controls. A-D) TSD-001 data is represented by blue circles and connecting lines, while TSD-002 data is represented by red triangles and connecting lines.

Figure 2. *Longitudinal biochemical and neurological function outcome measures.* HexA activity was measured in A) cerebral spinal fluid (CSF) and B) serum of TSD-001 (blue bars) and TSD-002 (red bars) patients and non-TSD individuals (gray bars). CSF was collected on treatment day prior to infusion (D0), months 3 and 6 (M3, 6). Serum was collected at weeks 1, 2, 3 (W1-3), months 2, 3, and 6 (M2, M3, M6). The increase in HexA activity levels in CSF and serum post-treatment was statistically significant using a mixed effects model to adjust for repeated measures within individual. C) Mass spectrometry quantification of GM2 ganglioside species in CSF at day 0, months 3 and 6. TSD-001 data is represented by blue circles and connecting lines. TSD-002 data is represented by red triangles and connecting lines. D) Western blot analysis of HEXA and HEXB protein expression in CSF of TSD-001 and TSD-002 patients at day 0 (D0), months 3 (M3) and 6 (M6). Transthyretin (TTR), which is the most abundant protein in CSF, was used as a loading control. Arrowheads indicate the location of the bands corresponding to the HEXA and HEXB proteins. E) CHOP-INTEND scores for patients TSD-001 and TSD-002 from 2 days prior to treatment until 6 months or 12 months post-treatment.

Figure 3. *MRI findings in TSD patients treated with rAAVrh8.HEX/HEXB vectors.* A) Thalamic targeting in TSD-002. T1 weighted coronal (left panel) and T2 weighted (T2) coronal and axial (middle and right panels) MRIs immediately after bilateral thalamic injection. Note injection locations and fluid distribution (yellow arrows). B) Longitudinal MRIs from TSD-001 and TSD-002 at the time of injection and at 3 and 6 months after treatment. T2 MRI of TSD-001 shows some unchanged cortical atrophy and ventricular enlargement and diffusely affected white matter over the 6 month period. Color map (CM) of the fractional anisotropy (FA) and radial diffusivity (RD) suggest a combination of hypomyelination and myelin loss. The pretreatment MRI for TSD-002 shows incomplete myelination during infancy as posterior limbs of the internal capsule were not myelinated and lacked parietal and occipital myelination. After treatment, the degree of

myelination, as shown by the darkening on T2 imaging (B) and FA metrics (C), suggest stabilization over time and potential improvement in the posterior corpus callosum.

Author Contributions

TRF, RHB, OC, RM, AP, SB, SS, JP, AA, AR, LG, RF, and FSE were part of the medical team. DMY, CD and AMK were part of the clinical trial coordination team, data collection and analysis. ARB and MSE produced the AAVrh8 vectors. GW, MB and CM performed all immunological assays and respective data interpretation. ARB performed all enzyme assays and western blots, as well as respective data analysis and interpretation. BAB performed the statistical analyses. AA, SR, ZV, MSS, MG were responsible for analysis and interpretation of MRI data. TRF, RHB, DRM, HGE, and MSE wrote the manuscript.

Acknowledgments

This work was supported in part by the BlueGenes Foundation. We are also grateful for the exceptional support from the National Tay-Sachs and Allied Diseases Association and the Cure Tay-Sachs Foundation during the pre-clinical development of this AAV gene therapy.

Figure 2

A

TSD-001

TSD-002

B

C

Supplementary Table S1 - Adverse Events

	ID#	Adverse event (reported term)	AE Start date	AE Stop date	Severity (mild, moderate, severe)	Outcome (resolved, not resolved, resolved with sequelae, fatal)	Causality to gene therapy (not related, unlikely, possible, probable, definitely)	Causality to neurosurgical procedures (not related, unlikely, possible, probable, definitely)	Causality to immunosuppressive regimen (not related, unlikely, possible, probable, definitely)	Other possible cause of the AE (disease under study, concomitant medication, protocol related procedure, etc.)	SAE (Y/N)	SAE criteria (death, life-threatening, initial/prolong hospitalization (dates), congenital anomaly/birth defect, medically significant)	Action taken to drug (dose changed, drug interrupted, drug withdrawn, NA,)	Treatment given
1	TSD-001	Red raised rash during rituximab infusion	11/9/18	11/9/18	moderate	resolved	not related	not related	related	NA	No	NA	NA	Benadryl and temporarily stopped infusion, restarted at slower rate
2	TSD-001	Rash on perineum and neck	11/27/18	1/2/19	mild	resolved	not related	not related	likely	NA	No	NA	NA	antifungal cream and powder
3	TSD-001	white coating on tongue	11/29/18	12/29/18	mild	resolved	not related	not related	likely	NA	No	NA	NA	antifungal medication
4	TSD-001	Bruising at IV and blood draw site on foot	11/14/18	12/10/18	mild	resolved	not related	not related	not related	protocol related procedure	No	NA	NA	None
5	TSD-001	oral lesion on inner lower lip	2/23/19	3/5/19	mild	resolved	not related	not related	likely	NA	No	NA	NA	Acyclovir
6	TSD-001	oral lesion on right inner lower lip	4/24/19	5/7/19	mild	resolved	not related	not related	likely	NA	No	NA	NA	Acyclovir
7	TSD-001	nasal stuffiness	6/7/19	6/22/19	mild	resolved	not related	not related	not related	NA	No	NA	NA	None
8	TSD-001	cough	6/7/19	6/24/19	mild	resolved	not related	not related	not related	NA	No	NA	NA	None
9	TSD-001	Upper respiratory congestion	UN/jul/2019	UN/jul/2019	moderate	resolved	not related	not related	not related	pneumonia	No	NA	NA	unknown
10	TSD-001	Pneumonia	8/2/19	UNAUG/2019	severe	resolved	not related	not related	not related	NA	Yes	hospitalization	NA	antibiotics and intubation
11	TSD-001	Labored breathing	8/2/19	8/2/19	severe	resolved	not related	not related	not related	pneumonia	Yes	hospitalization	NA	intubation
12	TSD-001	darky coloring	8/2/19	8/2/19	severe	resolved	not related	not related	not related	pneumonia	Yes	hospitalization	NA	intubation
1	TSD-002	post-op pain	6/11/19	6/14/19	moderate	resolved	unlikely	definitely; thalamic injections	not related	NA	No	NA	NA	acetaminophen/oxycodone
2	TSD-002	fever	6/11/19	6/11/19	moderate	resolved	unlikely	unknown	not related	unknown	No	NA	NA	acetaminophen
3	TSD-002	fever	6/12/19	6/12/19	moderate	resolved	possible	unknown	not related	unknown	No	NA	NA	acetaminophen,blood and urine cultures
4	TSD-002	cough	6/13/19	6/24/19	mild	resolved	not related	not related	not related	unknown	No	NA	NA	none
5	TSD-002	fever	9/9/19	9/10/19	mild	resolved	unlikely	not related	not related	anesthesia and intubation	No	NA	NA	acetaminophen
6	TSD-002	higher fever	9/10/19	9/11/19	moderate	resolved	unlikely	not related	not related	urinary tract infection	Yes	hospitalization	NA	antibiotics
7	TSD-002	urinary tract infection	9/10/19	9/16/19	moderate	resolved	unlikely	not related	not related	unknown	No	NA	NA	antibiotics

TSO-002	Lab Test	Ref Range & Units	Screening	Day-7	Day-2	Day 1	Week 2	Week 3	Month 1	Unschedule d Visit 1	Unschedule d Visit 2	Month 2	Month 3	Month 6	
Date:			5/21/19	6/3/19	6/10/19	6/12/19	6/26/19	7/2/19	7/11/19	7/22/19	7/29/19	8/8/19	9/6/19	11/21/19	
Chemistry	Sodium	135-145 mmol/L	139	--	--	136	--	--	--	--	--	--	136	139	
	Potassium	3.5-5.3 mmol/L	4.4	--	--	4.3	--	--	--	--	--	--	4.4	3.4	
	Chloride	97-110 mmol/L	105	--	--	102	--	--	--	--	--	--	109	109	
	CO2	24-32 mmol/L	24	--	--	23	--	--	--	--	--	--	16	ND	
	Anion Gap	15-may	10	--	--	11	--	--	--	--	--	--	11	ND	
	Glucose	70-99 mg/dL	78	--	--	86	--	--	--	--	--	--	180	65	
	Creatinine	0.20-0.73 mg/dL	0.25	--	--	0.21	--	--	--	--	--	--	<0.20	0.24	
	Calcium	8.7-10.7 mg/dL	10.2	--	--	9.7	--	--	--	--	--	--	8.5	10	
	Total Protein	5.5-7.5 g/dL	6.1	--	6.4	5.8	7.6	7	7.3	--	--	--	7	5.6	
	Albumin	3.2-5.2 g/dL	4.5	--	4.4	4.1	4.1	4.2	4.1	--	--	--	4.1	3.2	
	Bilirubin, Total	0.1-1.2 mg/dL	0.3	--	0.3	0.3	0.3	0.3	0.3	--	--	--	0.2	0.3	
	Alkaline Phosphatase	30-300 U/L	227	--	<0.1	109	94	88	96	--	--	--	92	101	
	AST	10-40 U/L	80	--	130	80	160	125	128	--	--	--	146	126	
	ALT	10-40 U/L	19	--	80	19	26	26	21	--	--	--	23	184	
	BUN	7-23 mg/dL	6	--	18	5	6	--	--	--	--	--	6	12	
	WBC	5.5-15.5 10 ³ /uL	7.2	--	--	8.6	--	9.6	--	--	--	--	--	8.6	--
	RBC	3.90-5.50 10 ⁶ /uL	4.34	--	--	3.69	--	4.34	--	--	--	--	--	4.86	--
	Hemoglobin	11.1-15.3 g/dL	11.9	--	--	10.1	--	11.9	--	--	--	--	--	12.6	--
	Hematocrit	34-40.0 %	34.6	--	--	30.1	--	35	--	--	--	--	--	37.9	--
	MCV	75-87 fL	79.7	--	--	81.5	--	80.5	--	--	--	--	--	78	--
MCH	24.0-30.0 pg	27.4	--	--	27.5	--	27.5	--	--	--	--	--	26	--	
MCHC	29.0-36.0 g/dL	34.3	--	--	33.7	--	34.2	--	--	--	--	--	33.3	--	
RDW	11-15.5 %	11.4	--	--	13	--	12.8	--	--	--	--	--	12	--	
Hematology	Platelets	140-400 10 ³ /uL	340	--	--	330	--	340	445	--	--	--	342	--	
	MPV	7.6-11.6 fL	7.7	--	--	8.1	--	8.1	--	--	--	--	7.7	--	
	Neutrophil %	%	24.7	--	--	62.5	--	22.3	--	--	--	--	--	39.9	
	Lymphocyte %	%	65.3	--	--	24.4	--	64.3	--	--	--	--	--	43.4	
	Monocyte %	%	6.8	--	--	11.8	--	12.7	--	--	--	--	6.8	--	
	Eosinophil %	%	2.8	--	--	0.7	--	0.4	--	--	--	--	5.1	--	
	Basophil %	%	0.4	--	--	0.6	--	0.4	--	--	--	--	0.6	--	
	Neutrophil #	1.50-8.50 10 ³ /uL	1.8	--	--	5.4	--	2.1	--	--	--	--	3.4	--	
	Lymphocyte #	2.0-6.0 10 ³ /uL	4.7	--	--	2.1	--	6.2	--	--	--	--	3.7	--	
	Monocyte #	0.2-1.0 10 ³ /uL	0.5	--	--	1	--	1.2	--	--	--	--	0.9	--	
	Eosinophil #	0.0-0.6 10 ³ /uL	0.2	--	--	0.1	--	0	--	--	--	--	0.4	--	
	Basophil #	0.0-0.3 10 ³ /uL	0	--	--	0.1	--	0	--	--	--	--	0.1	--	
	Smear review		normal	--	--	vacuolated neutrophils	--	platelet estimate increased	--	--	--	--	--	no	--
	Immunogenic	IGG	533-1078 mg/dL	--	277	--	present	--	platelet estimate increased	577	--	--	746	672	366
		IgA	24-121 mg/dL	14	--	--	--	--	--	--	--	--	--	--	--
		hs CRP	<10 mg/L	--	--	--	--	--	--	--	--	--	--	--	--
%CD19		6-23%	21	<1	--	--	--	--	<1	<1	--	<1	<1	<1	
%CD20		6-13%	24	<1	--	--	--	--	<1	<1	--	<1	<1	<1	
GGT		<25 U/L	--	--	21	--	23	--	--	--	--	--	--	--	
Creatine Kinase		<134 U/L	--	--	--	--	--	--	--	--	--	--	--	--	
ID labs	CH50	31-60 U/mL	--	--	--	--	--	160	--	--	--	--	--	--	
	Sirostimus	2.0-18.0 ng/mL	--	10	--	--	--	2.6	2.9	4.2	6	4.7	--	--	
	IgG/IgA/IgM level	mcg/mL	--	--	--	--	--	--	--	--	--	--	--	--	
	HSV 1 IgG Type Specific	<90 negative	1.32	--	--	--	--	--	--	--	--	--	--	--	
	HSV 2 IgG Type Specific	<90 negative	--	--	--	--	--	--	--	--	--	--	--	--	
	Hep B surface antibody	>0-0.10 mIU/mL	non-reactive	--	--	--	--	--	--	--	--	--	--	--	
	Varicella Zoster	>165 positive	<135.00	--	--	--	--	--	--	--	--	--	--	--	
	Toxoplasma gondii antib	<7.2 IU/mL negative	<7.20	--	--	--	--	--	--	--	--	--	--	--	
	CMV antibody (IgG)	<60 U/mL negative	<0.2	0.72	--	--	--	--	--	--	--	--	--	--	
	Color	--	--	--	--	--	colorless	--	--	--	--	--	colorless	colorless	
CSF	Appearance	--	--	--	--	clear	--	--	--	--	--	clear	clear	--	
	Total Nucleated Cells	--	--	--	--	0	--	--	--	--	--	--	3	9	
	RBC	--	--	--	--	1	--	--	--	--	--	--	0	1	
	Total Cells counted	--	--	--	--	--	--	--	--	--	--	--	89	89	
	Narrative	--	--	--	--	--	--	--	--	--	--	--	11	11	
	Protein	--	--	--	--	19	--	--	--	--	--	--	23	30	
	Glucose	--	--	--	--	57	--	--	--	--	--	--	78	54	
CSF	LDH	--	--	--	--	102	--	--	--	--	--	--	97	--	
	Gram Stain	--	--	--	--	negative	--	--	--	--	--	--	negative	negative	
	Culture	--	--	--	--	negative	--	--	--	--	--	--	negative	negative	

A

B

C

Supplementary Figure S1: Diffusion tensor imaging (DTI) results. A) Mean diffusivity (MD) maps of TSD-002 brain indicating regions of interest (ROI) for calculation of DTI parameters. MD quantifications over time are displayed below the images for various ROIs in the brain for both TSD-001 and TSD-002. B) Axial diffusivity (AD) and C) radial diffusivity (RD) measurements are also shown for the same brain ROIs over time for both patients. In TSD-002 both optic tracts show stable FA values and increased average diffusivity values. AD values remain stable even when FA decreases, as shown in the anterior corpus callosum for TD-001.

Supplementary Figure S2: volumetric calculations of lateral ventricular size and size of the lentiform nucleus. TSD-001 has larger ventricles consistent with cortical atrophy whereas the relevance of TSD-002 ventricular size remains unclear and could be associated with normal brain development.

Supplementary Materials and Methods

Hex A enzyme activity

Cerebral spinal fluid or serum from patients and non-TSD controls were used either neat or diluted 1:10 to 1:100 in 0.01 M Phosphate Citrate Buffer pH 4.4 with 0.1% Triton X100. The assay for Hexosaminidase A activity was conducted in 96 well plate format using the artificial substrate 4-methylumbelliferyl 6-sulfo-2-acetamido-2-deoxy- β -D-glucopyranoside (MUGS, Toronto Research Chemicals, Toronto, Canada) and a standard curve of 4-methylumbelliferone (Sigma, St. Louis, MO). Samples were incubated at 37°C for up to 4 hrs and reactions stopped by addition of 0.2 mL of 1M sodium carbonate, pH 10.7. Fluorescence was measured using a Synergy-HTX multi-mode reader (BioTek, Winooski, VT) using 360 and 460 nm excitation and emission filters, respectively.

Statistical analysis of changes in HexA activity

A mixed effects model was used to estimate the statistical significant of HexA increased activity in CSF and serum after treatment compared to baseline, adjusting for repeated measures within individual. We used months as the time metric in the models. Estimates were obtained from the least-squares means generated from the model. All analyses were performed using SAS 9.4 (SAS Institute, Inc. Cary, NC).

Western Blot

CSF samples (30 μ L each) were separated in a 4-20% polyacrylamide SDS-PAGE gel (Bio-Rad) and then transferred to 0.2 μ m pore nitrocellulose (GE Healthcare) membrane using a wet transfer system. Primary antibodies used were rabbit polyclonal anti-HexA (1:100; [aa431-480], LS-B6959-100, LSBio), rabbit monoclonal anti-HexB (1:200; [EPR7978], ab140649, Abcam), and rabbit polyclonal anti-human prealbumin (1:500; A0002, Dako). All primary antibodies were incubated at 4°C for 16-18 hr. Detection was performed by chemiluminescence using Clarity

Western ECL Substrate (Bio-Rad) and images were acquired using the ChemiDoc system (Bio-Rad).

Mass Spectrometry quantification of GM2 ganglioside in CSF

C-MS/MS analysis was conducted on a Shimadzu Prominence UFLC system coupled with an Applied Biosystems/MDS Sciex 000QTRAP mass spectrometer using multiple reaction monitoring (MRM) and two Valco switching valves. The CSF samples were injected directly to the LC-MS/MS system. A quality control sample was prepared by pooling 20 mL from each study sample and injected every three study samples to monitor the instrument performance. The relative quantification data were measured as peak areas of analytes and reported as total counts. All the samples were analyzed in one batch so that relative comparisons could be made accurately without run-to-run variation. The coefficient of variation of each lipid in the quality control sample was less than 15%, indicating that the mass spectrometric response was reproducible within the batch.

Immunologic Assays

Neutralizing antibody assay

Sera sampled prior to dosing, Days 7, 14, 21, 28, Months 2, 3 and 6 were heat inactivated for 30 min at 56°C. AAVrh8-CMV-LacZ (10^9 gc/well) was incubated with 2-fold serial dilutions of samples for 1h at 37°C, 5% CO₂. The mixture was added to Huh7 cells (1×10^5 cells) previously infected with an adenovirus (100 particles/well) and incubated for 18-22h at 37°C, 5% CO₂. Cells were washed and developed with Galactor-Star kit (Applied Biosystems). Luminescence was measured with a luminometer. The NAb titers are expressed as the highest dilution that inhibited β -galactosidase expression by at least 50% compared to a negative mouse serum control.

Anti-AAVrh8 IgG ELISA

Patient sera were collected prior to dosing, Days 7, 14, 21, 28, Months 2, 3 and 6. Anti-AAVrh8 IgG antibodies were detected by enzyme-linked immunosorbent assay (ELISA). Plates were coated with 5e9 vg of rAAVrh8 vector in coating buffer (carbonate-bicarbonate 0.5M, pH=9.6) at 4°C overnight. The plates were blocked with PBS-Tween 0.1% for 1 hour at 37°C. Patient sera were serially diluted and incubated for 2 hours at 37°C. Plates were then incubated with an anti-human IgG-HRP antibody (Bethyl Labs A80-104P) for 1 hour at room temperature. Substrate (Tetramethylbenzidine (TMB), BD) was added and plates were incubated 7 minutes at room temperature. The reaction was stopped using 2N H₂SO₄. Plates were read in a spectrometer (VersaMax, Molecular Devices) at 450nm.

IFN γ ELISpot assay

PBMCs were isolated prior to dosing, Days 14, 21, Months 1, 2, 3 and 6. The IFN γ -ELISpot assay was performed according to the manufacturer's recommendations (Human IFN- γ ELISpot^{BASIC}, MABTech). Briefly, PBMCs were stimulated *in vitro* for 24-48hrs with overlapping peptides spanning either the AAVrh8 VP1 capsid, the HexA or HexB sequences and divided into 2 to 3 pools (2 μ g/ml, 15-mers overlapping by 10 aa, Mimotopes, Australia). A negative control consisted of unstimulated cells (medium only), and CD3/CD28 stimulation were used as a positive control for cytokine secretion. Spot number was determined using an ELISpot reader ELR068IFL (AID) and analyzed with AID ELISpot Reader Software V6.0. Responses were considered positive when the number of spot-forming units (SFU) per 1e6 cells were >50 and at least 3-fold higher than the control condition.

MRI and image analysis

Brain MRI was performed using a Signa HDxt 1.5T scanner (GE Healthcare, Chicago, IL) equipped with a HD 8-channel head coil (GE Healthcare). T1-weighted images were acquired

using a 3D fast spoiled gradient echo (FSPGR) sequence (TR/TE = 13/4 ms, flip angle = 15°, axial direction slice thickness = 1.2 mm, acquisition matrix = 256 × 256, NEX = 0.7, FOV = 22 cm). T2-weighted images were acquired using a 3D fluid-attenuated inversion recovery (FLAIR) sequence (TR/TE = 6000/133 ms, inversion time = 1851 ms, sagittal direction slice thickness = 1.8 mm, acquisition matrix = 224 × 224, NEX = 1, FOV = 19 cm). Diffusion tensor imaging (DTI) was also performed with the following parameters: TR/TE = 8000/116 ms, 15-gradient directions, b-values = 0 and 1000 s/mm², slice thickness = 5 mm, acquisition matrix = 128 × 128, NEX = 1, FOV = 19 cm.

Volumetric analysis

Amira (V2019.1, Thermo Fisher Scientific, Waltham, MA) was used to perform volume calculations of the different brain regions at each time point for each patient. A combination of 3D T1- and 3D T2-weighted MR images were used for segmenting the following regions: whole brain (without ventricles), cerebellum, thalamus, corpus callosum, caudate, ventricles, and lentiform nucleus. For the whole brain and ventricle regions, a semi-automated technique using a combination of thresholding and edge detection was used to capture the respective regions in each coronal slice and manual corrections were performed thereafter before rendering the 2D slices into a 3D volume. The remaining structures were segmented manually due to isointense signal of the structures with surrounding tissue. All segmentations were performed using both the axial and sagittal orientations at the same time while carefully paying attention to tissue boundaries. All segmentations were verified using age-appropriate atlases under the guidance of an experienced neuroradiologist (SR).

DTI analysis

DTI analysis was performed using DSI Studio (<http://dsi-studio.labsolver.org>). For each patient at each time point, fractional anisotropy (FA), axial diffusivity (AD), mean diffusivity (MD), and

radial diffusivity (RD) maps were generated to visualize the changes in the brain over time. To further quantify the changes in sub-regional structures, the following regions of interest were selected: anterior and posterior corpus callosum, dorsal brainstem, left and right centrum semiovale, superior cerebellar peduncle, left and right proximal occipital optic tractus, and left and right distal occipital optic tractus. FA, MD, AD, and RD quantification were made for each of the aforementioned structures and tracked over time for each patient.

Figures

Figure 1

Longitudinal safety and immunological studies. Serum levels of liver enzymes A) alanine aminotransferase (ALT) and B) aspartate aminotransferase (AST) enzymes in TSD-001 and TSD-002 patients before treatment (Pre-TX), weeks 1, 2, and 3 (W1, 2, 3) and months 1, 2, 3 and 6 (M1, 2, 3, 6).

Shaded bars in A) and B) represent the normal interval for both assays. Dashed line in B) indicates the maximum value reported in TSD patients²⁵. C) Total anti-AAVrh8 IgG in serum quantified by ELISA. In C) the gray squares and lines represent anti-capsid IgG levels in serum of an ALS patient treated by IT AAV gene therapy. D) Neutralizing antibody titers to AAVrh8 capsid quantified by a transduction assay. Peripheral blood mononuclear cells (PBMCs) were isolated from TSD-001 and TSD-002 patients before treatment and at various time points thereafter to assess T-cell responses by interferon gamma ELISpot assays using stimulation with pools of overlapping peptides spanning E, F) AAVrh8 VP1, or G, H) human HEXA and HEXB proteins. Unstimulated or CD3/CD28-stimulated PBMCs were included in all assays as negative and positive controls. A-D) TSD-001 data is represented by blue circles and connecting lines, while TSD-002 data is represented by red triangles and connecting lines.

Figure 2

Longitudinal biochemical and neurological function outcome measures. HexA activity was measured in A) cerebral spinal fluid (CSF) and B) serum of TSD-001 (blue bars) and TSD-002 (red bars) patients and non-TSD individuals (gray bars). CSF was collected on treatment day prior to infusion (D0), months 3 and 6 (M3, 6). Serum was collected at weeks 1, 2, 3 (W1-3), months 2, 3, and 6 (M2, M3, M6). The increase in HexA activity levels in CSF and serum post treatment was statistically significant using a mixed effects

model to adjust for repeated measures within individual. C) Mass spectrometry quantification of GM2 ganglioside species in CSF at day 0, months 3 and 6. TSD-001 data is represented by blue circles and connecting lines. TSD-002 data is represented by red triangles and connecting lines. D) Western blot analysis of HEXA and HEXB protein expression in CSF of TSD-001 and TSD-002 patients at day 0 (D0), months 3 (M3) and 6 (M6). Transthyretin (TTR), which is the most abundant protein in CSF, was used as a loading control. Arrowheads indicate the location of the bands corresponding to the HEXA and HEXB proteins. E) CHOP-INTEND scores for patients TSD-001 and TSD-002 from 2 days prior to treatment until 6 months or 12 months post-treatment.

Figure 3

MRI findings in TSD patients treated with rAAVrh8.HEX/HEXB vectors. A) Thalamic targeting in TSD-002. T1 weighted coronal (left panel) and T2 weighted (T2) coronal and axial (middle and right panels) MRIs immediately after bilateral thalamic injection. Note injection locations and fluid distribution (yellow arrows). B) Longitudinal MRIs from TSD-001 and TSD-002 at the time of injection and at 3 and 6 months after treatment. T2 MRI of TSD-001 shows some unchanged cortical atrophy and ventricular enlargement

and diffusely affected white matter over the 6 month period. Color map (CM) of the fractional anisotropy (FA) and radial diffusivity (RD) suggest a combination of hypomyelination and myelin loss. The pretreatment MRI for TSD-002 shows incomplete myelination during infancy as posterior limbs of the internal capsule were not myelinated and lacked parietal and occipital myelination. After treatment, the degree of myelination, as shown by the darkening on T2 imaging (B) and FA metrics (C), suggest stabilization over time and potential improvement in the posterior corpus callosum.