

Obesity prevention in the early years: a mapping study of national policies in England from a behavioural science perspective

Helen Croker (✉ h.croker@ucl.ac.uk)

University College London <https://orcid.org/0000-0002-7247-6599>

Simon Russell

University College London

Aswathikutty Gireesh

University College London

Aida Bonham

King's College London

Corinna Hawkes

City University

Helen Bedford

University College London

Susan Michie

University College London

Russell M. Viner

University College London

Research

Keywords: Policy, mapping, obesity prevention, childhood obesity, diet, physical activity, behavioural science

Posted Date: March 30th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-19653/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Version of Record: A version of this preprint was published at PLOS ONE on September 30th, 2020. See the published version at <https://doi.org/10.1371/journal.pone.0239402>.

Abstract

Background

Evidence indicates that early life is a critical time for determining future risk of obesity. A sharper policy focus on pregnancy and early childhood would be useful to improve obesity prevention efforts. This study aimed to systematically identify and categorise the policy levers being used in England that have potential to influence early lifecourse (pregnancy and 0-5 years) and identify how these policies interface with energy balance behaviours. The objective is to identify gaps and where further policy actions could most effectively focus.

Methods

A behavioural science approach was taken using the Capability-Opportunity-Motivation-Behaviour (COM-B) model and Behaviour Change Wheel (BCW) framework. The key determinants of energy balance in the early years were identified from the Foresight Systems Map. Policy actions were scoped systematically from available literature, including any health or non-health policies which could impact on energy balance behaviours. The Foresight variables and policy actions were considered in terms of COM-B and the BCW to determine approaches that would be likely to be effective for obesity prevention and treatment. Existing policies were then overlaid across the map of key risk factors to identify gaps in obesity prevention and treatment provision.

Results

A wide range of policy actions were identified (n=115) to address obesity-relevant risk factors. The most common policy actions were educational and guidelines relating to environmental restructuring (i.e. changing the physical or social context). Scope for strengthening policies relating to the food system (e.g. the market price of food) and psychological factors contributing to obesity were identified. Policies acted via all aspects of the COM-B model, but there was scope for improving policies to increase capability through skills acquisition and both reflective and automatic motivation.

Conclusions

There is substantial policy activity to address early years obesity but much is focused on education. Scope exists to strengthen actions relating to upstream policies which act on food systems and those targeting psychological factors contributing to obesity risk.

Background

Rates of overweight and obesity in children have increased worldwide since the 1980s, and despite a slowing recently, it remains an important public health issue with serious consequences for health (1, 2). In the UK, approximately 10% of children aged 4–5 years and 20% of those aged 10–11 years were living with obesity in 2018/19 (3). Children in the UK are also developing obesity at a younger age and therefore are likely to accumulate greater levels of overweight across their lifetimes (4). Behaviour-changing interventions to treat children with overweight and obesity can reduce body mass index (BMI) but effects tend to be small (5) and effective prevention programmes have generally proved elusive (6). There is growing evidence for the importance of pregnancy and maternal factors, including maternal weight, general health and the environment, for programming weight gain in offspring across the life-course (7, 8). Pre and postnatal growth, and their interaction, are important for establishing risk factors for obesity (7). Children from disadvantaged backgrounds have higher rates of obesity and, in England, these inequalities have increased over the past decade (9). Early life is likely to be critical; a recent systematic review showed consistent associations between rapid growth in the first two years of life and later obesity, with the strongest effects in children of lower socioeconomic status (SES) and ethnic minority groups (10). Evidence also suggests that children of lower SES have greater exposure to several early life risk factors for obesity, including pre-pregnancy maternal obesity and diabetes, low birth weight and poorer early life nutrition (11). This suggests that a sharper policy focus on pregnancy and early childhood could be useful for improving the effectiveness of obesity prevention efforts and for addressing health inequalities.

The complex nature of obesity, with its numerous interacting contributing factors, requires a comprehensive, coordinated and sustained policy response (12). Policy levers have been described as “instruments that can be adjusted by governments to achieve system-wide change” (13). Adjusting or applying a policy lever should significantly impact on the system, and in this case influence behaviours or determinants of behaviour related to child obesity (14). Policy actions are available through the ‘layers of influence on health’, from the individual, through family, social and community networks, to wider socioeconomic and environmental conditions and up to national level (15). Therefore, there are a range of periods and settings where policy levers may potentially be applied during the early lifecourse. An overview of national policy to tackle obesity in England demonstrated the changing nature of government policy with a shift from ‘top-down’ regulation to more collaborative approaches and a strong focus on individual responsibility and locally-led actions (16). They however found limited evidence of measurable success and highlighted the challenges of evaluating policy.

Some previous work has been done to map obesity-related policies in England, this has been limited to interventions specifically promoting healthier ready to eat meals (17) and local level policies (18). These identified a range of policies in place with a focus on ‘downstream’ action (i.e. those primarily relying on individual level change). There has also been mapping work undertaken of obesity policies across countries which included

England (19). However, a comprehensive mapping of existing policies in England to intervene during the critical early years part of the lifecourse, and whether they target appropriate behaviours or behavioural influences has not been conducted to the authors' knowledge. An improved understanding of the current policy landscape and whether actions are appropriately targeted has potential to provide insight on whether resources are being appropriately allocated. The aim of this study was to use a behavioural science approach to identify and categorise the policy levers being used in England that have potential to influence obesity across the early lifecourse (here considered to be pregnancy and 0–5 years) and identify how these interface with energy balance behaviours, by which we mean behaviours that are related to either energy intake or expenditure. The objective is to identify gaps and areas where further policy actions should focus.

Methods

To meet the overarching aim, the objectives of this study were to: 1) identify and describe the key factors relevant to obesity prevention in the early years; 2) identify and describe English national policy in the early years for obesity prevention; 3) systematically map the identified policies onto the key factors related to obesity prevention in early life; and 4) identify gaps and where further policy actions could be most effectively focused. A behavioural science approach was taken using the Capability-Opportunity-Motivation-Behaviour (COM-B) model and the Behaviour Change Wheel (BCW) framework, shown in Fig. 1 (20, 21). The COM-B model identifies the factors required to bring about desired behaviour change and the BCW highlights the approaches likely to be effective to support this. According to the model, for a behaviour to occur, there must be sufficient Capability (physical and psychological); Opportunity (physical and social); and Motivation (reflective and automatic). Definitions of the COM-B and BCW components are provided in Table 1. Coding guidelines and data extraction templates were developed for each stage of the work (see Supplementary File). The methods used to address each aim are described below.

Reproduced with permission (21)

Table 1
COM-B and Behaviour Change Wheel (BCW) components

COM-B components	
Physical capability	Physical skill, strength, or stamina
Psychological capability	Knowledge or psychological skills, strength or stamina to engage in the necessary mental processes
Physical opportunity	Opportunity afforded by environment involving time, resources, location, cues, physical 'affordance'
Social opportunity	Opportunity afforded by interpersonal influences, social cues and cultural norms that influence the way we think about things, e.g. the words and concepts that make up our language
Automatic motivation	Automatic processes involving emotional reactions, desires (wants and needs), impulses, inhibitions, drive states and reflex responses
Reflective motivation	Reflective processes involving plans (self-conscious intentions) and evaluations (beliefs about what is good and bad)
Behaviour Change Wheel intervention types	
Education	Increasing knowledge or understanding
Persuasion	Using communication to induce positive or negative feelings or stimulate action
Incentivisation	Creating an expectation of reward
Coercion	Creating an expectation of punishment or cost
Training	Imparting skills
Restriction	Using rules to reduce the opportunity to engage in the target behaviour (or to increase the target behaviour by reducing the opportunity to engage in competing behaviours)
Environmental restructuring	Changing the physical or social context
Modelling	Providing an example for people to aspire to or imitate
Enablement	Increasing means/reducing barriers to increase capability (beyond education and training) or opportunity (beyond environmental restructuring)
Behaviour Change Wheel policy categories	
Communication/marketing	Using print, electronic, telephonic or broadcast media
Guidelines	Creating documents that recommend or mandate practice. This includes all changes to service provision
Fiscal measures	Using the tax system to reduce or increase the financial cost
Regulation	Establishing rules or principles of behaviour or practice
Legislation	Making or changing laws
Environmental/social	Designing and/or controlling the physical or social environment
Service provision	Delivering a service
(Taken from Michie et al, 2014)	

To address objective 1 (identify and describe the key factors relevant to obesity prevention in the early years), the Foresight Systems map was used to identify key mechanisms influencing energy balance, providing insight into the complex relationships between determinants of obesity (22). The factors are organised into broad groups which include those relating to an individual's physiology, physical activity, biology, diet, and psychology, the physical activity environment, and the food production system. In the centre of the model is 'energy balance' and the map shows how a wide range of variables influence this. The variables most strongly connected to the core were the focus of this study ('key variables' and 'first tier variables'). These are considered leverage points for obesity policy as they have strong connections with the core (energy balance) and many connections with the wider system (23). Such 'upstream' interventions are expected to have greater influence than those 'downstream'. Variables relevant to children were used, including behaviours occurring via parent actions. These were described using COM-B, to indicate how each variable could act to influence energy balance. Variables were independently coded by two researchers (HC and SR), the rate of agreement was recorded and discrepancies were discussed to reach consensus.

To address objective 2 (identify and describe national policy in the early years for obesity prevention), a systematic online search of current national government policies in England relating to child obesity prevention in the early years was undertaken (see the Supplementary File for the search strategy). Key policy documents relating specifically to obesity were included, and policies relating more generally to child health, including

pregnancy and maternity care, child care settings, primary schools and leisure facilities. Policies in other areas which may influence early years were also included: health-related (e.g. dental health) and social (e.g. travel, housing). Policy actions have been defined as ‘the specific actions put into place by any level of government and associated agencies to achieve the public health objective’ (24). Therefore, we included interventions as well as policies providing they were judged to be ‘concrete and actionable’ (i.e. not aspirational/ vague). These were systematically described by coding the intervention types and policy categories identified in the BCW. Each policy was independently coded by two researchers (HC and AB), the rate of agreement recorded and any discrepancies discussed to reach consensus. The searches were undertaken in May 2018. The Childhood Obesity Plan: a call for action, Chap. 2 (COP2) was published in June 2018 and there coded at a later date (25). Coding was undertaken by HC and AG as a consensus exercise. Policies were also coded according to the mode of influence, this could be direct action on the child (e.g. change in food provided to the child in a child care setting), direct action via the parent (e.g. policy resulting in parent changing food provided to child), indirect action via the parent (e.g. policy resulting in parent changing own behaviour resulting in modelling of healthier behaviours), or a combination.

To address objective 3 (systematically map the identified policies onto the key factors related to obesity prevention in early life), the matrix developed by Michie et al (2014) was used, see Table 2 (21). This indicates the intervention types most likely to be useful for bringing about change in each COM-B construct. The coding generated for objectives 1 and 2 were used; the Foresight variables were examined sequentially and relevant policies that could potentially address them were identified; variables were considered in terms of each applicable COM-B construct. This was done by two researchers (HC and AB) as a consensus exercise; coding of the COP2 was additionally undertaken by HC and AG.

Table 2- Matrix of links between COM-B and intervention types

(Taken from Michie et al, 14)

COM-B components	Intervention types								
	Education	Persuasion	Incentivisation	Coercion	Training	Restriction	Environmental restructuring	Modelling	Enablement
Physical capability					X				X
Psychological capability	X				X				X
Physical opportunity					X	X	X		X
Social opportunity						X	X	X	X
Automatic motivation		X	X	X	X		X	X	X
Reflective motivation	X	X	X	X					

X indicates the intervention types most applicable for bringing about change in each COM-B construct

To address objective 4 (identify potential gaps and opportunities for policy actions to be developed or implemented), the mapping work generated for objective 3 was used. Coherence between the key influences of energy balance and current policies was examined (i.e. presence of policies to address key factors influencing energy balance). We also examined whether there were opportunities for developing or implementing policy. This is where there are factors known to influence obesity but for which we were unable to identify any policies to address, or where the policies were not targeted in the most effective way according to the BCW.

Results

The coding of the Foresight variables using the COM-B model resulted in an agreement between coders of 71%, considered moderate, and agreement for the coding of policies using the BCW was 90%, considered very high (26). The full coding can be found in the Supplementary File.

Objective 1: Identify and describe the key factors relevant to obesity prevention in the early years

The codes relating to COM-B assigned to the Foresight variables varied widely, so that there was a good spread of potential influences on children’s energy balance, these are shown in detail in the Supplementary File and summarised in Fig. 2.

Objective 2: identify and describe national policy in the early years for obesity prevention

A total of 106 specific policy actions were identified in the initial search, this included specific policy initiatives (e.g. the 'National Child Measurement Programme' and 'Change for Life') and national clinical and public health guidelines produced by the National Institute for Health and Care Excellence (NICE). Four additional policy actions were identified but not included as they either fed into other policies (update of Nutrient Profile Model), were planned for the future (Healthy Rating Scheme for primary schools; suite of digital applications for healthy eating) or were not a specific policy action (healthy marketing strategy). Some of the policies had more than one specific policy action (e.g. measurement and feedback elements of the 'National Child Measurement Programme' and a wide range of actions at different time points in the 'Healthy Child Programme'); the number of unique policies was 79. When the COP2 was included, an additional nine policy actions were identified. In total, 115 specific policy actions were identified and 88 unique policies. The remainder of the results relate to these 115 policy actions. The coding of the policy actions is described fully in the Supplementary File, but examples of how they relate to the intervention types in the BCW, as well as the mode of action, is provided in Table 3. The most common intervention type was education, with environmental restructuring and modelling approaches also commonly used. Coercion, restriction and incentivisation approaches were rarely used. As well as policies impacting directly on the child, many were via adult-focused initiatives which provide educational opportunities for parents and modelling opportunities for children (via parent behaviour change). The number of policy actions assigned to each intervention type and policy option is shown in Figs. 3 and 4 respectively.

Table 3
Examples of how identified policy actions relate to BCW intervention types

BCW intervention type	Examples of policy actions relating to each intervention type		
	Direct action on child	Direct action via parent	Indirect action via parent
Education Increasing knowledge or understanding	Food teaching in primary schools framework; School Food Plan; NICE guidance- obesity prevention- schools; resources to support NCMP in schools; Cycling and walking investment strategy (walk to school project)	Healthy Start Scheme; Healthy Child Programme; National Child Measurement Programme feedback; Change4Life resources; 5aday logos	Healthy Child Programme; One You NHS campaign
Persuasion Using communication to induce positive or negative feelings or stimulate action	None identified	Promotions of breastfeeding within Healthy Child Programme; Clearer food labelling; consistent calorie labelling; Change4Life resources; NICE physical activity guidelines	One You campaign online resources; physical activity infographic
Incentivisation Creating an expectation of reward	Cycling and walking investment strategy (walk to school project- children's challenge);	Change4Life vouchers for healthy food	None identified
Coercion Creating an expectation of punishment or cost	None identified	None identified	None identified
Training Imparting skills	Funding for bikeability training; Cooking in the national curriculum; Sports skills within national curriculum	NICE guidelines including imparting skills regarding breastfeeding, cooking, positive parenting	NICE guideline on obesity prevention (NHS interventions for adults)
Restriction Using rules to reduce the opportunity to engage in the target behaviour (or to increase the target behaviour by reducing the opportunity to engage in competing behaviours)	Advertising restrictions in children's media	Ban on sales of energy drinks to children; proposed ban of promotion of unhealthy foods and drinks	None identified
Environmental restructuring Changing the physical or social context	Sugar drinks levy; Sugar and Calorie Reduction Programmes; provision of opportunities to be active at school (breaktimes and PE); Primary PE and Sports Premium; provision of healthier foods in school; Food in Early Years Settings; lowered maximum protein content of formula milk; Nursery Milk Scheme; universal free school meals for KS1; Daily mile	Sporting Future (improved local leisure facilities); local transport plans to encourage active travel; NICE guidance on postnatal care and maternal and child nutrition; consistent calorie labelling	NICE obesity prevention guidelines; NICE physical activity guidelines on walking and cycling
Modelling Providing an example for people to aspire to or imitate	Creation of healthy food environments in public sector settings (including leisure centres); staff modelling recommended in Food in Early Years Settings; School Fruit and Vegetable Scheme;	Healthy Child Programme (peer support for breastfeeding); positive images within Change4Life; Baby Friendly accreditation	One You Campaign, NHS weight loss plan, various NICE guidelines
Enablement Increasing means/reducing barriers to increase capability (beyond education and training) or opportunity (beyond environmental restructuring)	Universal free school meals for KS1; What works in schools and colleges to increase physical activity?	Vouchers within Healthy Start Scheme; behavioural elements and vouchers with Change4Life; prompts for behaviour change with Start4Life;	Support for behaviour change and access to resources in NICE guidelines; NHS weight loss plan

Objective 3: Systematically map the identified policies onto the key factors related to obesity prevention in early life

The results of the mapping work as a whole are shown in the HEAT map (Fig. 2), which indicates where policy actions were identified for Foresight variable as per the BCW. Of a possible 287 opportunities for policy, over half had at least one policy action (n = 157; 54.7%), but we were unable to identify any actions for the remainder (n = 128; 44.6%) or there were uncertainties over the coding (n = 2; 0.7%). A summary of how well the identified policies mapped onto the COM-B theoretical model and the intervention types from the BCW is shown in Table 4. For the COM-B constructs, 'physical capability' was the best covered and 'physical opportunity' and 'social opportunity' were reasonably well covered. The constructs with the least identified policy actions were 'psychological capability', 'automatic motivation' and 'reflective motivation'. For the intervention types from the BCW, there was wide variation in coverage. There were many policy actions based on education and environmental restructuring, with more than 80% of the opportunities for these approaches having policies in place. There were a reasonable number of policy actions in place to address modelling and enablement (with over 60% of opportunities having policies in place), fewer for training, restriction, persuasion and incentivisation (less than 60% of opportunities) and no policy actions based on coercion.

Table 4
Policy actions in place by COM-B constructs and intervention types (across all Foresight variables)

COM-B construct	Number with any mapped policies	%
Physical capability Physical skill, strength, strength or stamina	13/14	93%
Social opportunity Opportunity afforded by interpersonal influences, social cues and cultural norms that influence the way we think about things, e.g. the words and concepts that make up our language	42/60	70%
Physical opportunity Opportunity afforded by environment involving time, resources, location, cues, physical 'affordance'	38/60	63%
Psychological capability Knowledge or psychological skills, strength or stamina to engage in the necessary mental processes	13/24*	54%
Reflective motivation Reflective processes involving plans (self-conscious intentions) and evaluations (beliefs about what is good and bad)	22/52	42%
Automatic motivation Automatic processes involving emotional reactions, desires (wants and needs), impulses, inhibitions, drive states and reflex responses	29/77*	38%
BCW intervention type		
Environmental restructuring Changing the physical or social context	37/41	90%
Education Increasing knowledge or understanding	17/21	81%
Modelling Providing an example for people to aspire to or imitate	18/26	69%
Enablement Increasing means/reducing barriers to increase capability (beyond education and training) or opportunity (beyond environmental restructuring)	36/56	64%
Persuasion Using communication to induce positive or negative feelings or stimulate action	13/24	54%
Restriction Using rules to reduce the opportunity to engage in the target behaviour (or to increase the target behaviour by reducing the opportunity to engage in competing behaviours)	13/30	43%
Training Imparting skills	17/41**	41%
Incentivisation Creating an expectation of reward	6/24	25%
Coercion Creating an expectation of punishment or cost	0/24	0%
*There was uncertainty regarding the coding for one of the policy actions within each of these categories, these were therefore not coded as a mapped policy action		
** There was uncertainty regarding the coding for two of the policy actions within each this category, these were therefore not coded as mapped policy actions		

The HEAT map (Fig. 2) can also be used to examine the policy actions available for individual Foresight variables. The nutrition-related variables with the most policy actions in place were food literacy, food abundance, food exposure, palatability of food, breastfeeding and weaning, energy density (at least 60% of opportunities had identified policies). For example, actions for food literacy included policies for clearer food labelling and education within the Healthy Child Programme. There were a reasonable number of policies (40–60% of opportunities had identified policies) to address de-skilling, convenience of food, portion size, purchasing power, perceived inconsistency of science messages, and tendency to graze. For example, nutrient-based standards in food outlets to target food convenience. There is much overlap with many of the policy actions; for example, healthier foods in schools could help to address food abundance, food exposure, palatability of food offerings, and energy density. For physical activity, the variables with the most policy actions (at least 60%) were degree of physical education, innate activity in childhood, recreational activity, transport activity, functional fitness, with a reasonable number (40–60%) present for de-skilling. For example, guidance for schools to target physical education and NICE guidance re public open spaces and schools to target recreational activity. The variables with limited identified policy actions were stress, food advertising, self-esteem, side-effects of drugs, demand for indulgence, and market price of food (less than 30% of the opportunities for policies had identified policy actions). There were however some efforts to target these variables with marketing restrictions identified as key actions in the COP2.

Objective 4: Identify potential gaps and opportunities for policy actions to be developed or implemented

As identified from the above mapping work, the COM-B constructs with the least identified policy actions were 'psychological capability', 'automatic motivation' and 'reflective motivation'. The HEAT map (Fig. 2) indicates which intervention types specifically were lacking for these. In relation to increasing psychological capability, there were opportunities to strengthen policies through training and enablement. The Foresight variables that had gaps for these included stress, food literacy, inconsistent messaging, 'demand for indulgence', and de-skilling. In relation to increasing automatic motivation, there were, in particular, opportunities to strengthen policies based on persuasion, incentivisation, coercion, training, and enablement. In relation to reflective motivation, there were opportunities to particularly strengthen policies based on incentivisation and coercion. A number of Foresight variables had particular gaps relating to motivation, including stress, food advertising, self-esteem, 'demand for indulgence', tendency to graze, and market price of food. The Foresight variable 'market price of food' was notable in that no policies were identified which targeted it.

Discussion

This scoping and mapping exercise identified where there were policies in place in England to target risk factors for obesity, and whether the methods employed were appropriate according to a behavioural science perspective, and where there was scope for additional actions. A substantial amount of policy activity was identified aiming to address childhood obesity and strong coverage of policies to target many of the energy balance-related risk factors from the Foresight systems map. A total of 115 relevant policy actions were identified and over half of the potential opportunities for addressing these risk factors had appropriate actions in place. This indicates that Government has implemented many actions in England to address early years obesity.

The mapping work in the current study was able to provide specific information about whether the policy action in place to target the Foresight variables used appropriate approaches, as identified by COM-B and the BCW. There were policy actions targeting all of aspects of the model but we also identified gaps. The COM-B model identifies two types of capability- physical and psychological. Physical capability refers to physical skill, strength or stamina and is best acted upon with training or enablement, whilst psychological capability refers to knowledge and psychological skills and is best acted upon with education, training and enablement. The majority of Foresight variables identified as being amenable to change through increasing physical capability related to physical activity (e.g. transport activity) and we identified good coverage of policy actions targeting these across most of the relevant Foresight variables. The majority of the Foresight variables related to psychological capability were psychological (e.g. stress, food literacy) or dietary (e.g. portion size) and we identified numerous policy actions targeting this based on education but gaps for actions based on training and enablement.

Within the COM-B model, opportunity comprises physical and social aspects. Physical opportunity relates to time, resources, locations and cues, whilst social opportunity relates to interpersonal influences, social cues and social norms. Both can be targeted using restriction, environmental restructuring and enablement; physical opportunity can also be increased with training and social opportunity can be increased with modelling (i.e. setting a 'good' example). The majority of the Foresight variables across several domains (diet, physical activity, diet, economic) were related to opportunity and we identified reasonable coverage of policy actions across all of these. The Foresight variable 'market price of food' was a notable exception with no policy actions identified for either physical or social opportunity. The COM-B model goes onto identify reflective motivation (including self-conscious intentions and making evaluations) and automatic motivation (including emotional responses, impulses, and inhibitions). Both reflective and automatic motivation can be increased using multiple approaches from the BCW. The majority of the Foresight variables across several domains (diet, physical activity, diet, economic, physiology) were related to either reflective or automatic motivation. We identified numerous gaps in the coverage of policy actions to target these Foresight variables, with less than half having any actions. In particular, there were few policy actions relating to motivation for the following Foresight variables: stress, food advertising, self-esteem, demand for indulgence, tendency to graze, and market price of food.

Looking across the components in COM-B and the Foresight variables, the most common policy approach (as per the BCW) that we identified was education, along with a focus on guidelines targeting environmental restructuring and policies encouraging modelling opportunities (indirectly acting on the child via parent/carer behaviour change). We identified opportunities to further develop policy actions focused on enablement, persuasion, incentivisation, restriction, and coercion. For example; restriction is a possible approach for increasing physical and social opportunity, so policies based on restriction could be developed to promote physical activity (including both recreational and transport activity). Potential examples of this are restrictions on car use near schools to promote active school journeys or adding restrictions to tablets to limit their use to encourage active recreational time. Another example; since incentivisation is a possible approach for increasing reflective and automatic motivation, consideration could be given to developing policies based on incentivisation in relation to the Foresight variable portion size, an example could be incentivising purchasing of smaller packaged snack foods. Together, this highlights the focus on education and indicates that there are opportunities to build upon efforts for upstream change. In particular, there are numerous opportunities for further developing policies which act via increases psychological capability, reflective motivation and automatic motivation. Strengthening policies which increase the latter, such as people's desires, emotions and inhibitions may be particularly powerful as they go beyond a reliance on people's self-conscious 'choices'.

There were a many policies addressing environmental change, with regards to both the food and physical activity environment. However, despite the UK's good record of developing evidence-based policy guidelines, implementation of guidelines (especially public health policies) has often been poor (27). A study of implementation, using the Food Environment Policy Index (Food EPI) to map out and rate policies targeting childhood obesity in England, included a rating of implementation by experts (28). Implementation was rated highest for monitoring (of obesity, risk factors, diet), nutrient declarations on labels, access to information, availability of dietary guidelines, school food standards, and population level targets. Implementation was rated lowest for food subsidies, planning policies to encourage fruit and vegetables, and systems-based approaches. This supports our findings that upstream policies are particularly challenging to implement. A focus on strengthening existing policy recommendations to facilitate implementation, especially those targeting upstream actions, may be useful. An example of such an upstream action is the price of food which can have a huge impact on people's purchasing decisions, with less healthy foods typically costing less and being consumed more by lower SES groups than healthier foods (29); a lack of activity here may result in other policies having limited influence.

The emphasis on education and the limited action targeting automatic motivation indicates a reliance on policies primarily focusing on individual level change. A recent study found that, for addressing obesity, governments from developed countries tended to concentrate on policy levers addressing individual-level change rather than the environment, even in countries (such as England) with a strong policy focus on childhood obesity (19). A recent systems-mapping exercise examined how local authorities in England address obesity using the 'Action Mapping Tool' (18). Consistent with our findings, this work found that whilst only a small proportion of the causes of obesity were coded as 'individual lifestyle factors', nearly 60% of the actions around obesity targeted individuals. This suggests that an individual-orientated approach is a common theme throughout both national and local obesity policy. Interventions based only on individual choice have limitations. They require families to perceive change as important and be in a position to make such changes. This is likely to be challenging for many families but particularly difficult in families from lower SES backgrounds; this may act to further widen the health inequalities apparent with obesity (30). A recent review indicated that all intervention types risk widening health inequalities but complex interventions which are targeted at multiple levels (systems, community, individual) and in multiple settings (school, health, population) appear to have less negative effects, and fiscal measures may even bring benefits (31). Successive UK governments' policies highlight obesity as a serious problem; however there is a political tension between state and individual responsibility. Health choices are assumed to be the individual's responsibility to control even though the behaviours leading to excess weight gain are acknowledged to be greatly influenced by the environment (32). Consistent with the findings in the current study, previous government policy documents have focused on information provision to change behaviour (33). One example comes from an analysis of the pledges within the Public Health Responsibility Deal which found that most pledges focused on providing information for consumers, rather than structural changes (e.g. reformulation) (34). Consumer views echo this, with analyses of online reactions to news stories about obesity policy finding either contradictory views around responsibility (35, 36) or that blame is attributed to the individual (37, 38). This discourse is at odds with the evidence for the important role of environmental factors in contributing to obesity (39). Of note, a greater emphasis on restricting of unhealthy food advertising was observed with the mapping of COP2 policy actions suggesting a move to more upstream action.

Strengths and limitations

This study systematically identified national policies on childhood obesity using an authoritative system analysis of risk factors for obesity (Foresight) and took a behavioural science approach to first describe risk factors and policies, and then to conduct mapping work. We believe that this is the first time that a comprehensive mapping of obesity policies has been conducted using this approach. This allowed behavioural targets and policies to be systematically described in detail, allowing the identification of gaps and opportunities for further policy development. These gaps and opportunities were specifically characterised (according to the type of intervention, the method and the target) providing explicit information to inform the strengthening of current policy and future policy development. The work has potential to be built on and could be applied at a local level and used to inform needs assessments.

Our work is subject to a number of limitations. England is a populous country with high childhood obesity levels and a history of strong public health actions on obesity, thus findings are not necessarily generalisable to other countries. We focused on national-level policy and recognise that in most countries, including England, local or community policy actions may also be in place. We were unable here to include policy activity by the 152

upper-tier local authorities in England, but the approach used in the current study could be applied in that context. This analysis recorded actions and recommendations, not how well they were being implemented, which was beyond the scope of the study. Estimating the expected impact of policy actions on behaviour and weight was also beyond the scope of this work but could be useful since the number of policies available does not necessarily correspond to their expected impact. In particular, insight into the implementation of NICE guidelines would be useful, especially for public health guidelines where there may not be the same accountability as the clinical guidelines, which are included in service commissioning processes. Tools to support implementation include the Food EPI, an established method using expert consensus to provide policy ratings and identifies gaps and policy priorities (28) and surveillance plans (19). The Foresight systems map was used to identify risk factors for childhood obesity as it is a comprehensive review of the evidence; however it was developed in 2007 so may not capture recent research. In addition, this is a fast moving field, the Childhood Obesity Plan (Chap. 3) was published as part of the Prevention Green Paper after the mapping work was completed and therefore not included (40). There is also considerable interest in the role of the pre-conception period for later obesity risk (41); however, the life-course stage for this work was restricted to pregnancy and early life to ensure that the study was feasible. The policy scoping was done via online searches, it is possible that there are additional policy actions not identified with this approach.

Conclusion

This work generated a systematic map of current national government policy on obesity in England for the early years, with the purpose of identifying additional policy opportunities across the system. We identified a substantial amount of activity aiming to address early years obesity but scope for strengthening actions, especially upstream policies (acting on the environment and systems) and those targeting psychological factors contributing to obesity (stress, self-esteem, use of food for non-nutritive purposes). It is important to further consider implementation and likely impact of policy action. We found that using the COM-B model and BCW was useful for characterising the risk factors and existing policies, allowing for a detailed exploration of the current policy landscape and identification of gaps.

List Of Abbreviations

COM-B Capability-Opportunity-Motivation-Behaviour (COM-B) model

BCW Behaviour Change Wheel

SES Socioeconomic status

COP2 The Childhood Obesity Plan: a call for action, Chap. 2

NICE National Institute for Health and Care Excellence

Food EPI Food Environment Policy Index

Declarations

- Ethics approval and consent to participate- not applicable
- Consent for publication- permission to use the image for Figure 1 was gained from Susan Michie (one of the authors)
- Availability of data and materials- The authors declare that the information supporting the findings of this study are available within the article [and its supplementary information files]
- Competing interests- The authors declare that they have no competing interests
- Funding- This report is independent research funded by the National Institute for Health Research (NIHR) Policy Research Programme (Policy Research Unit: Obesity/ PR-PRU-0916-21001). The views expressed in this publication are those of the authors and not necessarily those of the NIHR or the Department of Health and Social Care.
- Authors' contributions- RMV and CH conceived the study. HC designed the study methods with input from SR, CH, HB, SM, RMV. HC, SR, AG, AB undertook the data coding, mapping and interpretation. HC wrote the first draft of the manuscript. All authors provided input into the draft and read and approved the final draft.
- Acknowledgements- Thank you to Heidi Yu Spurrell and Rosita Ibrahim who additionally assisted with coding.

References

1. Ng M, Fleming T, Robinson M, Thomson B, Graetz N, Margono C, et al. Global, regional, and national prevalence of overweight and obesity in children and adults during 1980-2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet*. 2014;384(9945):766-81.
2. Sharma V, Coleman S, Nixon J, Sharples L, Hamilton-Shield J, Rutter H, et al. A systematic review and meta-analysis estimating the population prevalence of comorbidities in children and adolescents aged 5 to 18 years. *Obesity Reviews*. 2019;20(10):1341-9.
3. Lifestyles Team ND. National Child Measurement Programme, England 2018/19 School Year. 2019.

4. Johnson W, Li L, Kuh D, Hardy R. How Has the Age-Related Process of Overweight or Obesity Development Changed over Time? Co-ordinated Analyses of Individual Participant Data from Five United Kingdom Birth Cohorts. *PLOS Medicine*. 2015;12(5):e1001828.
5. Mead E, Brown T, Rees K, Azevedo LB, Whittaker V, Jones D, et al. Diet, physical activity and behavioural interventions for the treatment of overweight or obese children from the age of 6 to 11 years. *Cochrane Database Syst Rev*. 2017;6:Cd012651.
6. Wang Y, Cai L, Wu Y, Wilson RF, Weston C, Fawole O, et al. What childhood obesity prevention programmes work? A systematic review and meta-analysis. *Obes Rev*. 2015;16(7):547-65.
7. Matthews EK, Wei J, Cunningham SA. Relationship between prenatal growth, postnatal growth and childhood obesity: a review. *Eur J Clin Nutr*. 2017;71(8):919-30.
8. Ratnasingham A, Eiby YA, Dekker Nitert M, Donovan T, Lingwood BE. Review: Is rapid fat accumulation in early life associated with adverse later health outcomes? *Placenta*. 2017;54:125-30.
9. Public Health England. Changes in children's body mass index between 2006/07 and 2015/16. England: Public Health England; 2017.
10. Andrea SB, Hooker ER, Messer LC, Tandy T, Boone-Heinonen J. Does the association between early life growth and later obesity differ by race/ethnicity or socioeconomic status? A systematic review. *Ann Epidemiol*. 2017;27(9):583-92.e5.
11. Cameron AJ, Spence AC, Laws R, Hesketh KD, Lioret S, Campbell KJ. A Review of the Relationship Between Socioeconomic Position and the Early-Life Predictors of Obesity. *Curr Obes Rep*. 2015;4(3):350-62.
12. Gortmaker SL, Swinburn BA, Levy D, Carter R, Mabry PL, Finegood DT, et al. Changing the future of obesity: science, policy, and action. *Lancet (London, England)*. 2011;378(9793):838-47.
13. Grace FC, Meurk CS, Head BW, Hall WD, Carstensen G, Harris MG, et al. An analysis of policy levers used to implement mental health reform in Australia 1992-2012. *BMC Health Services Research*. 2015;15(1):479.
14. Williams A. Getting health reform right: a guide to improving performance and equity by Marc J. Roberts, William Hsaio, Peter Berman and Michael R. Reich. Oxford University Press, Oxford, New York, 2004. No. of pages: 332. ISBN 0-19-516232-3. *Health Economics*. 2005;14(4):433-4.
15. Dahlgren G, Whitehead M. WHOLIS E89384 European strategies for tackling social inequities in health: Levelling up Part 22007.
16. Jebb SA, Aveyard PN, Hawkes C. The evolution of policy and actions to tackle obesity in England. *Obesity Reviews*. 2013;14(S2):42-59.
17. Hillier-Brown FC, Summerbell CD, Moore HJ, Wrieden WL, Adams J, Abraham C, et al. A description of interventions promoting healthier ready-to-eat meals (to eat in, to take away, or to be delivered) sold by specific food outlets in England: a systematic mapping and evidence synthesis. *BMC Public Health*. 2017;17(1):93.
18. Nobles J, Christensen A, Butler M, Radley D, Pickering K, Saunders J, et al. Understanding how local authorities in England address obesity: A wider determinants of health perspective. *Health Policy*. 2019;123(10):998-1003.
19. Esdaile E, Thow AM, Gill T, Sacks G, Golley R, Love P, et al. National policies to prevent obesity in early childhood: Using policy mapping to compare policy lessons for Australia with six developed countries. *Obesity Reviews*.0(0).
20. Michie S, van Stralen MM, West R. The behaviour change wheel: a new method for characterising and designing behaviour change interventions. *Implementation science : IS*. 2011;6:42-.
21. Michie S. The behaviour change wheel : a guide to designing interventions / Susan Michie, Lou Atkins, Robert West. Atkins L, West R, editors: Sutton : Silverback Publishing; 2014.
22. Butland B. Tackling obesities: future choices. Project report. . In: Government Office for Science DoIuAs, editor. 2nd ed. London2007.
23. Vandebroek P, Goossens J, Clemens M. Tackling Obesities: Future Choices – Building the Obesity System Map. Londib: Government Office for Science; 2007.
24. Hawkes C, Jewell J, Allen K. A food policy package for healthy diets and the prevention of obesity and diet-related non-communicable diseases: the NOURISHING framework. *Obesity Reviews*. 2013;14(S2):159-68.
25. Department of Health and Social Care, Global Public Health Directorate, Obesity Food and Nutrition. Childhood obesity: a plan for action, Chapter 2 England; 2018.
26. McHugh ML. Interrater reliability: the kappa statistic. *Biochemia medica*. 2012;22(3):276-82.
27. Poobalan AS, Aucott LS, Ahmed S, Smith WCS. Analysis of the UK recommendations on obesity based on a proposed implementation framework. *BMC Public Health*. 2010;10(1):17.
28. Watson F, Taylor A, Rayner M, Lobstein T, Hinks R. Priority actions for addressing the obesity epidemic in England. *Public Health Nutrition*. 2018;21(5):1002-10.
29. Darmon N, Drewnowski A. Contribution of food prices and diet cost to socioeconomic disparities in diet quality and health: a systematic review and analysis. *Nutr Rev*. 2015;73(10):643-60.
30. Beauchamp A, Backholer K, Magliano D, Peeters A. The effect of obesity prevention interventions according to socioeconomic position: a systematic review. *Obesity Reviews*. 2014;15(7):541-54.
31. Venturelli F, Ferrari F, Broccoli S, Bonvicini L, Mancuso P, Bargellini A, et al. The effect of Public Health/Pediatric Obesity interventions on socioeconomic inequalities in childhood obesity: A scoping review. *Obesity Reviews*.0(0).

32. Ulijaszek SJ, McLennan AK. Framing obesity in UK policy from the Blair years, 1997–2015: the persistence of individualistic approaches despite overwhelming evidence of societal and economic factors, and the need for collective responsibility. *Obesity Reviews*. 2016;17(5):397-411.
33. Vallgård S. English obesity policies: To govern and not to govern. *Health Policy*. 2015;119(6):743-8.
34. Knai C, Petticrew M, Durand MA, Eastmure E, James L, Mehrotra A, et al. Has a public–private partnership resulted in action on healthier diets in England? An analysis of the Public Health Responsibility Deal food pledges. *Food Policy*. 2015;54:1-10.
35. Gregg R, Patel A, Patel S, O'Connor L. Public reaction to the UK government strategy on childhood obesity in England: A qualitative and quantitative summary of online reaction to media reports. *Health Policy*. 2017;121(4):450-7.
36. Chambers SA, Traill WB. What the UK public believes causes obesity, and what they want to do about it: a cross-sectional study. *J Public Health Policy*. 2011;32(4):430-44.
37. Thomas-Meyer M, Mytton O, Adams J. Public responses to proposals for a tax on sugar-sweetened beverages: A thematic analysis of online reader comments posted on major UK news websites. *PLOS ONE*. 2017;12(11):e0186750.
38. Hertwig R, Mata J. Public Beliefs About Obesity Relative to Other Major Health Risks: Representative Cross-Sectional Surveys in the USA, the UK, and Germany. *Annals of Behavioral Medicine*. 2018;52(4):273-86.
39. James WP. The fundamental drivers of the obesity epidemic. *Obes Rev*. 2008;9 Suppl 1:6-13.
40. Kuunders TJM, van Bon-Martens MJH, van de Goor IAM, Paulussen TGWM, van Oers HAM. Towards local implementation of Dutch health policy guidelines: a concept-mapping approach. *Health Promot Int*. 2018;33(4):635-47.
41. Poston L. Obesity in pregnancy; Where are we, where should we go? *Midwifery*. 2017;49:4-6.

Figures




Figure 1

Behaviour Change Wheel

COM-B component Intervention type Foresight Variable	Physical capability		Psychological capability			Physical opportunity				Social opportunity				Automatic motivation						Reflective motivation				
	Tr	En	Ed	Tr	En	Tr	Re	Env	En	Re	Env	Mo	En	Per	Inc	Co	Tr	Env	Mo	En	Ed	Per	Inc	Co
Stress			Green	Red	Red									Green	Red	Red	Red	Red	Red	Red				
Food advertising																								
Food literacy			Green	Red	Green																			
Self-esteem			Green	Orange	Green																Green	Green	Red	Red
Inconsistent messages			Green	Red	Red																Green	Green	Red	Red
Degree physical edn	Green	Green							Red													Green	Green	Red
Food abundance																								
Food exposure																								
Innate activity	Green	Green																						
Recreation activity	Green	Green																						
Transport activity	Green	Green																						
Side effects drugs																								
Functional fitness	Green	Green																						
Palatability food																								
Breastfeeding/ weaning																								
Demand indulgence																								
Tendency to graze																								
De-skilling	Green	Red																						
Convenience of food																								
Portion size																								
Purchasing power																								
Energy density food																								
Market price food																								

Figure 2

HEAT MAP showing Foresight variables with mapped policy action Intervention types: Tr-training; En-enablement; Ed-education; Re-restriction; Env-environmental restructuring; Mo-modelling; Per-persuasion; Inc-incentivisation; Co-coercion; Mapping: Grey- not applicable (COM-B construct not relevant for that variable); Green- policy actions identified; Red- no policy actions identified, Orange -uncertain




Figure 3

Policy actions coded by 'intervention type'




Figure 4

Policy actions coded by 'policy option'

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupplementaryFile.xlsx](#)