

# The regeneration of *Acer rubrum* L. "October Glory" through embryonic callus

**Chongwen Dai**

southwest university college of resources and environment <https://orcid.org/0000-0001-8333-6957>

**Yang-yang YAN**

institute of mountain hazards and environment

**Yumin Liu** (✉ [yuminliu@163.com](mailto:yuminliu@163.com))

Resources and Environment Department of Southwest University Three Gorges Res-ervoir area laboratory of ecological environment, ministry of education Southwest University

<https://orcid.org/0000-0002-3558-7097>

**Ya-min LIU**

southwest university

**Yuan-wei DENG**

southwest university

**Hong-yu YAO**

southwest university

---

## Research article

**Keywords:** *Acer rubrum* L. "October Glory ", tissue culture, embryogenic callus, regeneration

**Posted Date:** June 4th, 2020

**DOI:** <https://doi.org/10.21203/rs.3.rs-19994/v2>

**License:**   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

**Version of Record:** A version of this preprint was published at BMC Plant Biology on July 2nd, 2020. See the published version at <https://doi.org/10.1186/s12870-020-02496-z>.

# Abstract

**Background:** Tissue culture and rapid propagation technology is an important way to solve the difficulties of plant propagation. This experiment aims to explore the appropriate conditions at each stage of the red maple's tissue culture process and to obtain plantlets, thus providing a theoretical basis for the establishment of the red maple's tissue culture system.

**Results:** The results showed that the stem segment is the most suitable explant for inducing embryogenic callus. The MS (Murashige&Skoog) +0.8 mg/L TDZ (Thidiazuron) +1.0 mg/L 6-BA (6-Benzylaminopurine) +0.5 mg/L IAA(Indole-3-acetic acid) +35 g/L sucrose+7.5 g/L semi-fixed medium was the best for callus formation. When selecting type  $\square$  callus as embryonic callus induction material, MS+0.6 mg/L TDZ+0.5 mg/L 6-BA +2.0 mg/L IAA +35 g/L sucrose+7.5 g/L semi-fixed medium can get embryonic callus. The optimal medium for adventitious bud induction is MS+1.0 mg/L TDZ+3.0 mg/L 6-BA+0.2 mg/L NAA (1-Naphthaleneacetic acid)+1.2 mg/L IAA+35 g/L sucrose+7.5 g/L semi-fixed medium. The induction rate of adventitious roots in MS+0.6 mg/L TDZ+1.0 mg/L 6-BA+3 mg/L NAA+35 g/L sucrose+7.5 g/L semi-fixed medium was the highest, reaching 76%.

**Conclusions:** In the course of our research, we found that PGRs play an important role in the callus induction stage, and the effect of TDZ is particularly obvious; The callus cells grow and proliferate according to the "S" growth curve, and can be sub-cultured when the highest growth point is reached to maintain the rapid proliferation of the callus cells and to avoid inactivation of callus caused by tight niche.

## Background

Landscape industry has been recognized as the "eternal sunrise industry". As a kind of ideal colored-leaf tree species, red maple has a promising market and is one of the most popular trees in modern landscape design planning in China (Li Y et al., 2014). However, due to environmental factors, the introduction of colored-leaf tree species is often accompanied by the problem of unstable leaf color and variation of progeny traits.

Introduced in China in 2011, October Glory (*Acer rubrum* L., red maple), originating in the northeastern United States, is an excellent cultivar, due to its beauty, colorful leaves, wet resistance, cold resistance, drought tolerance (Townsend et al., 1974; Schaub et al., 2003), ecological characteristics (Abrams et al., 1990; Li L et al., 2014), and broad ability to adapt to a variety of soil types. After domesticating and cultivating the red maple, we selected this cultivar as it is well adapted to the climate conditions in southwest China (especially the bud mutations we selected) and has rapid growth along with stable leaf color. However, its low seed maturation rate and unstable leaf color traits of its progeny make it difficult to promote planting through seed propagation. The use of cuttings and other vegetative propagation methods are limited by the season and quantity of material. In addition, it is difficult to satisfy mass

production requirements by small-scale cutting propagation. Therefore, it is of practical significance to establish the rapid propagation system of red maple.

Plant transformation regeneration systems mainly depend on whether they can form mature embryonic cells (Onay et al., 2011). Therefore, the formation of callus embryo becomes the most important link. Types of explants, osmotic pressure, plant growth regulators (PGRs), environmental conditions, these factors play a vital role in the formation and maturation of embryonic callus (Asad H et al., 2010; Bahrami Sirmandi H et al., 2009; Sheikh Beig Goharrizi MA et al., 2010). Among them, the interaction between the medium and the PGRs is the basis for the establishment of embryonic cells. (Skoog et al., 1957; Wang et al., 2012). When a PGR stimulates explants and calluses, the internal intrinsic hormones produce feedback (Bosco D et al., 2010) and induce a callus to form an embryonic callus (Friml J et al., 1957; Benishelkh et al., 2015). In many cases, specific combinations and proportions of auxin and cytokinin can effectively induce embryonic callus or directional differentiation of organs (Saikat G et al., 2018; Mazri M A et al., 2018). Somatic embryogenesis technology plays an important role in the preparation of artificial seeds and the genetic improvement of garden plants. This technology has been widely used in the breeding of garden plants that propagate slowly or have problems with propagation, and are of great significance for the propagation and conservation of rare plants. (Kozai T et al., 1986; Moradi Sh et al., 2017). At present, the tissue culture and rapid propagation system of red maple is still being perfected, and yet there are no reports about red maple's embryonic callus and adventitious organ formation.

To provide a theoretical basis and technical support for the production of *Acer rubrum* L. October Glory, this experiment explores the influence of culture medium and plant growth-regulating substances on callus induction in the procession of embryogenesis and plantlet regeneration and screened out the appropriate medium formulas in different stages.

## Results

### Callus induction

During the period of callus induction, the callus induced by different types of explants showed significant differences. With the newly germinated twigs (approximately 15 d after germination, without buds) as the explants and after the initial 25-d culture, there was substantial swelling at the incision of the explant (Fig. 1B). A mixture of various types of calluses was formed when the twigs (approximately 15 d after germination, with buds) were cultured for 25 d as explants (Fig. 1C). However, the callus propagation was relatively slower than that of the twigs when the hard branches (the spring sprouting branches collected in October of the same year) were taken as explants and cultured for 25d (Fig. 1A). When the tender leaves were taken as explants in the initial culture to the 7th day, a reddish-brown, yellow, loose and green callus started at the edge of the blade. After the 25-d culture, the leaves began to curl from the edge to the petiole, forming multiple loose callus (Fig. 1D). The leaves were covered by a large number of callus after being sub-cultured, and the callus were mostly reddish-brown, green and yellowish curled. Compared to twigs and hard branch, the blade callus induction during the early stage was relatively slow, and the

callus were mostly granular and loose. But in later stage, the callus proliferation rate of blade was significantly faster than that of twigs and hard branch, and showed a looser callus state than before.

There were 10 explant callus types in this stage. The type I callus (indicated by an arrow in Fig. 1A, I) was a white and loose callus that was fragile when there was successive transfer of the culture by explant induction on the lenticels. The induction of the adventitious roots stage can be directly induced from the callus. The type II callus (indicated by an arrow in Fig. 1A, II) appeared to grow longer in hard branches and was pale yellow and translucent with water damage. After successive generations, the callus turned brown and died. In the microscopic examination, the cells were fewer and more irregular in shape, and the cells were highly vacuolated. The type III callus (indicated by an arrow in Fig. 1B, III) appeared in the stem section and had a yellow, spherical, compact shape, with a rapid reproductive rate. The type IV callus (indicated by an arrow in Fig. 1B, IV) appeared in the twig stem section and was yellow-green and globular, multiplying faster and easily forming an embryonic callus. The type V callus (indicated by an arrow in Fig. 1C, V) was green with a spherical, compact shape. After successive transfers, the callus appeared similar to the type of callus that is white on the outside and pale yellow on the inside. The type VI callus (indicated by an arrow in Fig. 1C, VI) was bottle-green and globular, with the fastest reproduction rate, and its later form was not easy to change. In the microscopic examination, the cells were larger and elliptical, with dense cytoplasm, which was between the embryonal and non-embryonic cells. The type VII callus (indicated by an arrow in Fig. 1C, VII) was pale yellow and loose. After the subculture, this type browned and died. The type VIII callus (indicated by an arrow in Fig. 1C, VIII) was milk-white flaxen in a loose state and was more on the edge of the callus formation. After the subculture, the reproduction rate was slow and easily exhausted. The type IX callus (indicated by an arrow in Fig. 1D, IX) formed in the blade explants and had green water damage, and its callus cells died easily. The type X callus (indicated by an arrow in Fig. 1D, X) formed in the blade explants and was a green, dense granule, and it was a reddish-brown callus, although, at times, it did not induce embryonic callus formation.

During the initial induction stage, the explants broke the normal propagation mode and built a new adaptive mechanism. Several experiments have shown that this mechanism is closely related to the type of explants, the type of culture medium and the type and content of PGR. The orthogonal experiment results showed that the callus induction rate in the MS + 0.8 mg/L TDZ + 1.0 mg/L 6-BA + 0.5 mg/L IAA medium was the highest, as high as 82% (Fig. 2). The effect of culture conditions on callus induction was TDZ > medium > IAA > 6-BA, the medium (P = 0.049) and TDZ (P = 0.037) had significant effects on callus induction (P < 0.05), while 6-BA (P = 0.17) and IAA (P = 0.338) had no significant effect (P > 0.05).

### **Somatic embryo induction and somatic embryogenesis**

In the process of embryonic callus induction, type VI callus with faster reproduction rate and good growth was selected as the embryonic callus induction material. After a 56-d culture, the callus gradually changed into white somatic embryos (Fig. 3D). In the early stage, the surface embryo cells were increasing, but the proliferation rate was relatively slow (Fig. 3B). After the secondary 5-week culture, the callus was transferred to a medium without PGR. The microscopic examination revealed that the cells

were small and elliptical, with dense cytoplasm, presenting typical embryonic cell characteristics (Fig. 3C). The process is achieved under PGR regulation, while embryogenesis is composed of proembryogenic masses (PEM), which may indicate that PEM formation is the normal reaction of the explants to PGR. Although it is difficult to strictly distinguish between direct or indirect somatic embryogenesis, embryogenesis is guaranteed to undergo somatic embryo formation when the PEM continues to develop and form an indeterminate organ (Fig. 3D). In this study, the somatic embryo was derived from the spherical tissue block, and this spherical tissue embryo was the equivalent cleavage polymorphism that formed the multicolumn embryonal cell embryo phenomenon presented in Fig. 3-D. Unlike somatic embryos, meristematic tissues can be independently oriented to form roots or stems (which will also grow from within meristematic tissues), and somatic embryos grow almost without exception on the surface of callus tissue, and are easily separated from surrounding cells.

When the callus has not been sub-cultured for a long time, its surface gradually becomes dark brown. At this time, the embryogenic callus cell disappeared, and the callus was almost dead, which occurred from outside to inside (Fig. 3a). After the callus was cut, fit the longitudinal section to the surface of the MS + 2.0 mg/L TDZ + 1.0 mg/L 6-BA + 0.5 mg/L NAA + 35 g/L sucrose + 7.5 g/L AGAR semi-fixed medium for subculture. After a 25-d culture, the brown explants turned green (Fig. 3b). When this reactivated callus was further cultured, we found that the original healing surface appeared white with green and brown healing (Fig. 3b). When this callus continued to be inoculated in the embryonic induction medium of MS + 0.6 mg/L TDZ + 0.5 mg/L 6-BA + 2.0 mg/L IAA + 35 g/L sucrose + 7.5 g/L AGAR semi-fixed culture medium, the callus gradually changed to a green dense globule. On the surface of the callus, some cells were found to have embryonal cell traits (Fig. 3d).

### **Adventitious bud induction**

When the mature embryogenic callus were cut and inoculated into the adventitious bud induction medium, the green adventitious bud gradually formed (Fig. 4A). After the subculture, embryonic callus began to swell, and the adventitious bud elongated (Fig. 4B). After 30 days of continuous cultivation, the adventitious buds begin to unfold (Fig. 4C). The stem branches were clearly visible at 40 d (Fig. 4D).

It was difficult to distinguish between the inner embryonic cells and non-embryonic cells. However, embryonic cell orientation development is closely related to the in vitro culture environment and can be achieved by changing the in vitro culture conditions, especially PGRs. The results of the orthogonal test showed that when the exogenous growth regulator was 1.0 mg/L TDZ + 3.0 mg/L 6-BA + 0.2 mg/L NAA + 1.2 mg/L IAA, the adventitious bud induction rate was the highest (Fig. 4), reaching 52% (Fig. 5). The effect of different PGRs on adventitious bud induction rates was different, and the PGR influence sequence was 6-BA > IAA > TDZ > NAA. Moreover, 6-BA ( $P = 0.022$ ), IAA ( $P = 0.027$ ) and TDZ ( $P = 0.046$ ) had the most significant effects on callus induction, and the impact of NAA ( $P = 0.0678$ ) was not significant.

### **Induction of adventitious root**

Root induction is the last key link in the success of the rapid propagation of red maple, and the thickness and quantity of the adventitious roots affect the survival rate of the subsequent acclimation and transplants. When the embryonic callus type VI and embryonic callus with adventitious bud were inoculated on adventitious root induction medium and cultured for 15 d, the green embryonic callus gradually turned white and generated many adventitious roots (Fig. 6C). Furthermore, after cultivation for 20 d, the adventitious roots of the embryonic callus with buds germinated, and the plant leaves increased (Fig. 6D).

This study found that the adventitious roots can be induced directly from the stem and can also be induced through an embryonal callus (Fig. 6). The orthogonal experiment showed that the induction rate was as high as 76% (Fig. 7), when the MS + 0.6 mg/L TDZ + 1.0 mg/L 6-BA + 3 mg/L NAA + 35 g/L sucrose + 7.5 g/L AGAR semi-fixed medium were used. Moreover, with an increase in the NAA concentration, the induction rate of adventitious roots increased. The main effect of NAA on the induction of the adventitious roots process was the most significant ( $P = 0.023$ ), while 6-BA ( $P = 0.442$ ) and TDZ ( $P = 0.816$ ) had no significant influence on the adventitious root process.

### **Effects of different training methods on survival rate of plantlets**

Among the 9 treatments, it can be seen that the survival rate of T2t2 group (namely the group under 6-d culture in closed flasks and 2-d in opened flasks) was the highest, reaching 86.7% (Fig. 8). And the survival rate of transplanted plantlets increased first and then decreased with the increase of bottle closing time.

### **Effects of different substrates on survival rate of transplanted plantlets**

The plantlets of T2t2 group were used as experimental materials. The survival rate of plantlets under different substrate treatments was significantly different, and the average survival rate of plantlets under 1:1 mixture of sand and humus treatment was the highest, reaching 81.1% (Fig. 9).

## **Discussion**

### **The effect of PGRs in callus induction stage**

Endogenous hormones naturally exist in plants; a low concentration of these compounds controls the whole process of cell division and elongation (Huetteman C A, et al., 1993). Hormones also play an important role in establishing and maintaining plant polarity (Gahan P B, et al., 2007), the most notable of which is the maintenance of apical dominance. In this study, we explored callus induction and embryo regeneration of the red maple. In the early stages of induction and propagation, the explants and callus were more dependent on PGRs, especially the cytokinin TDZ. As cytokinins TDZ was applied in a site, such as leaves, bark, evolution will make the processing units for a single amino acid active library. The surrounding amino acids were moved to the library, thereby promoting the synthesis of RNA and proteins and enhancing the activity of certain enzymes (Pasternak et al., 2000; Hunter et al., 2003). This also

explains the phenomenon in this experiment: when cytokinins, especially TDZ, were added to the medium, the apical dominance of the shoot segment with buds was inhibited, the lateral buds germinated into a single stem branch, and the proliferation of callus cells was promoted. In tissue culture (as well as intact plants and plant organs), cytokinin can also shorten the metaphase of cell division and thus accelerate cell division and proliferation (Jouanneau et al., 1970). (Jouanneau et al., 1970). Therefore, PGR plays an important role in callus induction and propagation.

### **The regeneration of somatic embryos**

Some embryonic cells can be formed gradually when the non-embryonic cells mutate through several divisions. If the secondary embryonic callus induction medium is present, embryonic callus can continue to form. In contrast, if cultured on the same medium, the embryonic cells will eventually be covered by non-embryonic cells, causing the embryonic callus to disappear. This result is illustrated by the phenomenon of callus degeneration after several successive generations of this experiment. In this experiment, it was found that when the culture medium, at the early stage of tissue culture, provides an adequate nutrition for the complete life cycle, the embryonic cells begin to multiply and differentiate into adventitious organs. If this supply is sustained, the number of cells will multiply in the "S" growth curve., but with the production of cells and the consumption of nutrients from the medium, the supply will continue to shrink and the cells will start to develop in the reverse direction, namely to develop into non-embryonic cells. This result is due to the restrictions of the culture medium space environment and nutrient supply (Hvoslef-Eide A K et al., 2005). Therefore, after the "S" curve reaches the highest point, the rate of cell proliferation will continue to decline, and the ecological niche of the cell population will be sharply constricted. Even the embryonic cells that have been formed will no longer have a survival advantage (Kallendrusch S et al., 2017). Then, the cellular self-protection mechanism is activated (Ghanbarali S et al., 2016), and non-embryonic cells will cover the cell population. Therefore, it is thought that cell reproduction will continue to be carried out by successive generations before reaching the highest point of the "S" curve. In this way, the sustained proliferation of the cells can be maintained by the nourishing of the new culture medium.

The explanation is derived from "competition" (Reinert et al., 1968), which is the disappearance of cellular embryos. After multiple cell divisions, if the medium cannot supply enough energy to maintain the normal division of embryonic cells, the cells will return to their best state of preservation, that is, the state of non-embryonic cells. We call this "the backward development of cells. This kind of self-protection mechanism will benefit the cell life cycle when there is a deteriorated environment (Li Junming et al., 2005). If the environment changes again to the appropriate conditions, the embryonic cells will be activated again, and develop into new plants under suitable conditions.

This study explored the activation process of this browning callus. The results showed that during the 30-day culture without renewing the medium, the callus growth began to slow down on the 18th day. After cultivation for 24 d, the white and green callus gradually turned brown. After a culture of 40 d, the callus was mostly brown. After the brown callus were sub-cultured in the new medium, green and white callus

gradually formed from the original explants at the 20d, and this time was significantly delayed compared with the initial induction time of the explant. This study also found that the callus induced from leaves was usually loose and light yellow, while that of twigs and branches was mostly white and green, and the formation of the former needed more time. This result indicates that the explants first break the original propagation mode before forming callus. Compared with twigs and branches, leaves have higher specialization process, thus their original propagation pattern is more difficult to break through.

### **The formation of adventitious organs**

Typically, every normal cell in a plant has the totipotency to develop into a complete plant but only in the best environment. In an appropriate medium, calluses can maintain long-term familiar stimulation and regenerate stems and branches, such as rice callus in 2% sucrose and 3% mannitol medium, and can maintain their root and embryo callus formation of bark for a long time (Kishor et al., 1986). Hormone, as the most effective factor in the induction of embryogenic callus, can play different roles under different combinations and proportions of hormone and the interaction with other factors. **Abscisic acid** (ABA), which is often used as a growth inhibitor in tissue culture, can significantly promote the maturation and germination of embryonic cells in tissue culture of some plants (Fang HY et al., 2013), and the addition of ABA to the medium can effectively promote the synthesis and accumulation of nutrients in embryonic cells (Arnold SV et al., 2002). In tissue culture of *Saposhnikovia divaricata*, normally differentiated embryonic cells can only be obtained by adding ABA (MA J et al., 2004); For Persian walnuts, ABA promotes the maturation and germination of somatic embryos and the growth of both buds and roots (Bayat Sh et al., 2009; Vahdati K et al., 2006; Vahdati K et al., 2008). When ABA is combined with other hormones or other factors such as cold treatment, the plant conversion rate can be significantly improved (Bahrami Sirmandi H et al., 2010). The formation of adventitious organs requires the combination of auxin and cytokinin. Our results showed that when the ratio of NAA to TDZ was greater than 3:1, the bud induction mechanism was activated. The formation of adventitious buds is more complicated than that of adventitious roots. However, when the ratio increased to 5:1, the adventitious bud induction rate began to decrease. This result may be caused by the polar transport of auxin. If the callus was cultured in this kind of medium, it will lead to the death of embryonic cells (Ruffoni B et al., 2013). In the process of low concentration to high concentration, its enhancement effect is increasing continuously. However, when reaching a certain threshold, higher concentrations of auxin also exhibit inhibitory effects (Aryakia et al., 2013) because most auxin has the physiological effect of the bell-shaped active curve (Thorpe, 2006). In this study, auxin NAA was effectively absorbed by red maples, especially at 4 mg/L with a high root induction rate of 90%. Callus proliferation in the culture medium containing NAA was slow but the addition of NAA can induce the formation of adventitious roots. However, it is necessary to induce the adventitious root of the blade callus in the embryo callus stage. In this experiment, the rooting rate of explant was greater than that of the callus. Moreover, when the NAA concentration increased, the callus induced by stems was obviously stagnant. As a highly differentiated organ, the leaf cannot directly enter the rooting state and gradually become dry. Thus, it is more difficult to break the inherent propagation mechanisms during the process of inducing the formation of adventitious organs from highly developed plant organs.

# Conclusion

In this study, we screened the most suitable callus type for embryonic callus induction and selected suitable media formulations in various stages including callus induction, Somatic embryo induction and somatic embryogenesis, adventitious buds and adventitious root induction. In the course of our research, we found that PGRs play an important role in the callus induction stage, and the effect of TDZ is particularly obvious; The callus cells grow and proliferate according to the "S" growth curve, and can be sub-cultured when the highest growth point is reached to maintain the rapid proliferation of the callus cells and to avoid inactivation of callus caused by tight niche.

## Methods

### Source of plants

Plant materials were obtained from the forestry test base of Southwest University, and were formally identified by Associate Professor Liu Yu-min. The plants (Southwest University deposition number SWU-OG-03) are kept in the base (not a voucher specimen but the whole living plants) and can be obtained with the permission of Associate Professor Liu.

### Treatment and training conditions of test materials

Four types of materials were selected as explants in this experiment: tender leaves, twigs (approximately 15 d after germination) with buds, twigs without buds and hard branch (the spring sprouting branches collected in October of the same year). First, the explants were rinsed with tap water for 5 min, and then they were added to a 500 ml beaker. Then, the explants were soaked for 5 min in detergent water and washed with a hair brush, and surface stains were removed. Then, the explants were rinsed with tap water for 20 min, cut into 1-2 cm sections and moved to a sterile counter. The leaves were rinsed with a small flow of water, and all the injured and browning leaves were discarded when the surface was washed. The pretreated explants were disinfected with 75% alcohol on a sterile operating table for 30 seconds and treated with 0.1% HgCl<sub>2</sub> for 8 min. The materials were rinsed 6 times with sterile water, and then the cutting sites were excised. The culture medium (PH5.8-6.0) used in the experiments was sterilized at 121 °C for 20min;Hormones were sterilized by filtration.AD7000 digital camera (Nikon Corporation, Tokyo, Japan) using manual settings (F = 4.5, shutter 320 s, iso = 500) was used to photograph the callus regularly and observe and record the changes.

### The induction of callus

An orthogonal experiment was designed with medium DKW, N6, MS, / WPM, and the growth regulator IAA(0.5, 1.0, 1.5, 2.0 mg/L),6-BA(0.5, 1.0, 1.5, 2.0 mg/L) and TDZ (0.2, 0.4, 0.6, 0.8 mg/L), with a total of 16 treatments (Fig.2). Each group was treated with 10 bottles of 5 to 6 explants. The culture bottles were placed in an artificial climate box and cultured under conditions of humidity 75%, 25 ° C, 14h of light (1500lx) and 10h of darkness. The changes of explants were observed and recorded every 2 days.

## **Embryo callus induction and somatic embryogenesis**

During the stages of callus induction and proliferation, the MS + 0.6 mg/L TDZ + 0.5 mg/L 6-BA + 2.0 mg/L IAA + 35 g/L sucrose + 7.5 g/L AGAR semi-fixed culture medium effectively induced the formation of somatic embryo, and the somatic embryo induction rate was as high as 41 per gram. Based on the above results, the light yellow and white, loose callus was selected to induce somatic embryo formation in the medium (pH = 5.8). After 5 weeks, the somatic embryo was transferred to the MS medium free of growth regulating substances and cultured until the embryo matured. Culture conditions were 85% humidity, (25±2) °C, 16 h of light (1500 lx) and 8 h of darkness. During the culture, the changes of callus and somatic embryo were observed and recorded once every two days.

## **Induction and germination of adventitious bud**

To obtain adventitious buds, the embryonic callus was used as the experimental material, and an orthogonal experiment was designed with TDZ (0.5, 1.0, 1.5, 2.0 mg/L), 6-BA (1.0, 2.0, 2.5, 3.0mg/L), NAA (0, 0.1, 0.2, 0.5 mg/L), and IAA(0.6, 0.8, 1.0, 1.2 mg/L), with a total of 16 treatments (Fig. 5). Culture conditions were 85% humidity, (25 + 2) °C , 18 h of light (3000 lx) and 2 h of darkness(20h cycle). During the culture, the changes of callus and the germination of adventitious bud were observed and recorded once every two days.

## **The induction and germination of the adventitious root**

Based on the above test results, the obtained embryonic callus and embryonic callus with buds were used as experimental materials for adventitious root induction. An orthogonal experiment was designed with NAA(1.0, 2.0, 3.0 mg/L), 6-BA (0.5, 1.0, 2.0 mg/L), and TDZ (0.6, 0.8, 1 mg/L), with a total of 9 treatments (Fig. 5). Culture conditions were 85% humidity, (25 + 2) °C, 12 h of light (1500 lx) and 12 h of darkness. During the culture, the changes of callus and the germination of adventitious root were observed and recorded once every two days.

## **Training of plantlets**

After the plantlets were moved to the nursery room at room temperature of 20 °C, they were first cultivated in closed flasks for T1=3d, T2=6, T3=12d respectively, and then the lids were opened to continue cultivating for t1=1d, t2=2d, t3=3d, with a total of 9 treatments, 30 plantlets per treatment. After the cultivation, the plants were transplanted, and the survival rate was counted 30 days after transplanting.

## **Screening of transplanting substrate**

The plantlets obtained from the best training method were transplanted into sand, humus soil, and a 1:1 mixture of sand and humus respectively, and the survival rate was counted 30 days after transplanting.

## **Data Statistics and Analysis**

The callus induction rate (%) = callus/explants number x 100%.

The somatic embryo induction rate (%) = the number of somatic embryo/callus mass x 100%.

The adventitious bud germination rate (%) = the number of adventitious bud/ the total number of callus x 100%.

The adventitious root germination rate (%) =the number of adventitious root/ the total number of callus x 100%.

The average survival rate (%) =the number of viable plantlets/ the total number of plantlets x 100%.

The data was analyzed using SPSS version 20.0, and Duncan's new multiple range test was used for multiple comparisons.

## Abbreviations

2,4-D	2,4-Dichlorophenoxyacetic acid
6-BA	6-Benzylaminopurine
ABA	Abscisic acid
DKW	Driver and Kuniyuki walnut medium
IAA	Indole-3-acetic acid
IBA	Indole-3-Butyric acid
MS	Murashige & Skoog
N6	Chu's N-6 Medium
NAA	1-Naphthaleneacetic acid
PGRs	Plant growth regulators
TDZ	Thidiazuron
WPM	Woody Plant medium

## Declarations

**Ethical approval and consent to participate** This article does not contain any studies with human participants or animals performed by any of the authors.

**Consent for publication** Not applicable.

**Availability of data and materials** The data used to support the findings of this study are available from the corresponding author upon request.

**Competing interests** The authors declare that they have no competing interests

**Funding** This study were supported by Project supported by the Key Project of Forestry Science and Technology of Chongqing Municipality(cfa2018lykj05);Fundamental Research Funds for the Central Universities (SWU119066);the Application Development Plan of Chongqing Municipality (cstc2014yykfb80015). The funding bodies didn't play a role in the design of the study and collection, analysis, and interpretation of data and in writing the manuscript.

**Authors' contributions** YY designed the experiment, DC,DY and YH performed the experiment. DC analyzed the data and was the major contributor in writing the manuscript.LYM(Yu-min) and LYM(Ya-min) revised the manuscript. All authors read and approved the manuscript.

**Acknowledgments** We thank all those who helped us during the discussion and assistance. Also thanks to Dr. Krysta of AJE, which has polished the language. We hve benefited from their help to finish the paper.

## References

- Abrams M D, Kubiske M E. Photosynthesis and water relations during drought in *Acer rubrum* L. genotypes from contrasting sites in central Pennsylvania. *Functional Ecology*, 1990, 4(6):727-733.
- Arnold S V , Sabala I , Bozhkov P , et al. Developmental pathways of somatic embryogenesis. *Plant Cell, Tissue and Organ Culture*, 2002, 69(3):233-249.
- Aryakia E, Naghavi M R, Farahmand Z, et al. Evaluating allelopathic effects of some plant species in tissue culture media as an accurate method for selection of tolerant plant and screening of bioherbicides. *Journal of Agricultural Science & Technology*, 2015, 17(4):1011-1023.
- Asad H, Lotfi M, and Vahdati K (2010) Development of an embryogenic suspension culture in *Cucumis melo*. *Acta Horticulturae*. 871: 553-558.
- Bahrami Sirmandi H and Vahdati K. Effect of carbohydrate source and polyethylene glycol on maturation and germination of somatic embryos in walnut (*Juglans regia* L.). *Acta Horticulturae*.2009, 839:165-172.
- Bahrami Sirmandi H and Vahdati K. Enhancement of Maturation and Germination of Somatic Embryos of Persian Walnut (*Juglans regia* L.) by Gellan Gum, Cold and PGR. *Acta Horticulturae*. 2010, 861:333-344.
- Baya Sh, Ebrahimzadeh H, Vahdati K, Mirmasoumi M. Somatic embryo maturation and germination of Persian walnut (*Juglans regia* L.). *Acta Horticulturae*. 2009,812:313-318.

- Benishelkh A A G, Tijani Y, Askira N K, et al. Prevalence and Management of Laboratory Microbial Contaminants in Plant Tissue culture. *Anatomical Record*, 2015, 127(3):551-567.
- Bosco D D, Sinski I, Ritschel P S, et al. Residues of 2,4-Dichlorophenoxyacetic Acid in Plant-Tissue Culture Glassware: Their Effects and A Decontamination Protocol. *International Journal of Biotechnology & Biochemistry*, 2010.
- Fang HY,Zhu H,Yao JX et al. Abscisic acid and Tissue Culture of Medicinal Plant.*China journal of Chinese materia medica*.2013,38(01):14-18.
- Friml J, Vieten A, Sauer M, et al. Efflux-dependent auxin gradients establish the apical-basal axis of *Arabidopsis*. *Nature*, 2003, 426(6963):147-53.
- Gahan P B. Totipotency and the cell cycle// *Protocols for Micropropagation, of Woody Trees and Fruits*. 2007:3-14.
- Ghanbarali S, Abdollahi M R, Zolnorian H, et al. Optimization of the conditions for production of synthetic seeds by encapsulation of axillary buds derived from minituber sprouts in potato ( *Solanum tuberosum* ). *Plant Cell Tissue & Organ Culture*, 2016(3):1-10.
- Huetteman C A, Preece J E. Thidiazuron: a potent cytokinin for woody plant tissue culture. *Plant Cell Tissue & Organ Culture*, 1993, 33(2):105-119.
- Hunter C, Sun H, Poethig R S. The *Arabidopsis*, Heterochronic Gene ZIPPY, Is an ARGONAUTE, Family Member. *Current Biology*, 2003, 13(19):1734.
- Hvoslef-Eide A K, Preil W. *Liquid Culture Systems for in vitro, Plant Propagation*[M]. Springer Netherlands, 2005.
- Jouanneau J P. Renouvellement des protéines et effet spécifique de la kinétine sur des cultures de cellules de Tabac[J]. *Physiologia Plantarum*, 1970, 23(2):232–244.
- Kallendrusch S, Merz F, Bechmann I, et al. Long-Term Tissue Culture of Adult Brain and Spleen Slices on Nanostructured Scaffolds. *Advanced Healthcare Materials*, 2017.
- Kishor P B K, Reddy G M. Regeneration of plants from long-term cultures of *Oryza sativa*, L. *Plant Cell Reports*, 1986, 5(5):391.
- Kozai T, Fujiwara K, Watanabe I. *Fundamental Studies on Environments in Plant Tissue Culture Vessels: (1) Relation between the Culture Medium Composition and Water Potential of Liquid Culture Media*. *J.agric.meteorol*, 1986, 42:1-6.
- Li Jun-ming꠫Zhu Deng-yun. *Plant tissue culture*. 3rd ed. Beijing꠫China Agricultural University Press;2005.


- Li L, Liu Y M, Wang M et al. Physiological response mechanism of three kinds of *Acer rubrum* L. under continuous high temperature and drought stress. *Acta Ecologica Sinica*.2014;34 (22) : 6471-6480.
- Li Y, He, B H, Yu C et al. American red maple "brandy" tissue culture and rapid propagation technology . *Journal of southwest normal university (natural science edition)*, 2014 (8) : 36-42.
- Ma J, Qiao Q, Xiao YP et al. Occurrence and Control of Abnormal Somatic Embryos in *Saposhnikovia divaricata* Tissue Culture. 2005(03):552-556.
- Mazri M A , Meziani R , Belkoura I , et al. A combined pathway of organogenesis and somatic embryogenesis for an efficient large-scale propagation in date palm (*Phoenix dactylifera* L.) cv. Mejhoul[J]. *Biotech*, 2018, 8(4):215.
- Moradi Sh, Dianati Daylami Sh, Arab M, Vahdati K (2017) Direct somatic embryogenesis in *Epipactis veratrifolia*, a temperate terrestrial orchid. *The Journal of Horticultural Science and Biotechnology*. 92(1):88–97.
- Onay A, Yildirim H, Tokatli Y O, et al. Plant tissue culture techniques—Tools in plant micropropagation. *Current Opinion in Biotechnology*.2011;22(5):S130-S130.
- Pasternak T, Miskolczi P, Ayaydin F, et al. Exogenous auxin and cytokinin dependent activation of CDKs and cell division in leaf protoplast-derived cells of alfalfa. *Plant Growth Regulation*.2000;32(2-3):129-141.
- Reinert J, Backs D. Control of Totipotency in Plant Cells growing in vitro. *Nature*.1968;220(5174):1340-1341.
- Saikat G, Jitendriya P. In vitro biotechnological advancements in Malabar nut ( *Adhatoda vasica* Nees): Achievements, status and prospects.[J]. *Journal, genetic engineering & biotechnology*,2018,16(2).
- Schaub M, Skelly J M, Steiner K C, et al. Physiological and foliar injury responses of *Prunus serotina*, *Fraxinus americana*, and *Acer rubrum* seedlings to varying soil moisture and ozone. *Environmental Pollution*.2003;124(2):307-320.
- Sheikh Beig Goharrizi MA, Vahdati K, Bahrami Sirmandi H, Kalantari K, Leslie CA. Enhancement of maturation and germination of somatic embryos in Persian walnut (*Juglans regia* L.) using osmolites, hormones and cold treatments. *African Journal of Food Science*. 2010, 4:735-743.
- Skoog F, Miller C O. Chemical regulation of growth and organ formation in plant tissues cultured in vitro. *Symposia of the Society for Experimental Biology*, 1957, 11(21):118.
- Thorpe T A. History of Plant Tissue Culture// *Plant Cell Culture Protocols*. Humana Press, 2006:pp 9-32.
- Townsend A M, Dochinger L S. Relationship of seed source and developmental stage to the ozone tolerance of *acer rubrum*, seedlings. *Atmospheric Environment*, 1974, 8(9):957-964.

Vahdati K, Jariteh M, Niknam V, Mirmasoumi M and Ebrahimzadeh H (2006) Somatic embryogenesis and embryo maturation in Persian walnut. *Acta Horticulturae*. 705:199-205.

Vahdati K, Bayat Sh., Ebrahimzadeh H., Jariteh M. and Mirmasoumi M. Effect of exogenous ABA on somatic embryo maturation and germination in Persian walnut (*Juglans regia* L.). *Plant Cell Tissue and Organ Culture*. 2008, 93:163–171.


Wang Q M, Wang L. An evolutionary view of plant tissue culture: somaclonal variation and selection. *Plant Cell Reports*, 2012, 31(9):1535-47.

## Figures




**Figure 1**

Callus of *Acer rubrum* L. explants in different periods A: Cultured 15 d, In the incision site and lenticel, explants produce  $\square$ ,  $\square$  callus; B: Cultured 30 d, explant calluses were  $\square$  and  $\square$  callus; C: Cultured 30 d, the stem section with a bud  $\square$ ,  $\square$ ,  $\square$  and mixed type  $\square$  callus; D: Leaf explant culture for 25 d,  $\square$  and  $\square$  calluses




**Figure 2**

## The influence of medium type and PGRs on callus induction



**Figure 3**


Somatic embryo induction and embryonic cell regeneration of blastocyst A: After three generations, the green ball callus was observed to contain a small number of embryonic cells. B: After five successive generations, the callus differentiated the embryonic callus cells. C: The embryonic callus after 6 successive generations; D: After 7 generations, the body embryo maturation and the callus showed a polyembryonal stalk. a: In the early stage, the explants of stem grafts were not in sync, which caused browning and even partial cell death; b: The brown callus was cultured over 25 d, and the brown callus was now white. c: After two successive generations of 60 d, a dark green, spherically dense callus was formed. d: The brown callus was cultured in the 45-d secondary culture, with embryogenic cells in the periphery



**Figure 4**


The adventitious bud induction of embryonic callus A: On the basis of mature embryo callus, the adventitious bud germinated after being cultured 7-9d. B: After 2 subcultures, the elongation of

adventitious bud was increased; C: Continue to culture 30d, new adventitious buds formed like ①; D: After a 40d subculture, bud ① unfolded to form branches.




**Figure 5**

The influence of kinds and concentrations of different growth regulators on adventitious buds




**Figure 6**

The adventitious root induction of embryonic callus A: After culturing type ① callus in adventitious root induction medium for 23 d, the embryonic callus gradually became white and loose and formed many infinitive roots. B: After 60 d of continuous cultivation ② the adventitious roots of embryonic callus without adventitious buds germinated in large quantities. C: After the callus that had been differentiated to form adventitious buds was cultured for 25 days, the green embryonic callus gradually changed to white, and a large number of adventitious roots were generated on it. D: The amount of adventitious roots and leaves thrived after 20 d




**Figure 7**

The influence of kinds and concentrations of different growth regulators on adventitious root



**Figure 8**

Effects of different training methods on survival rate of plantlets



**Figure 9**

Effects of different substrates on survival rate of transplanted plantlets