CircRNA TADA2A relieves idiopathic pulmonary fibrosis by inhibiting proliferation and activation of fibroblasts

Juan Li

Zhengzhou University First Affiliated Hospital

Ping Li

Zhengzhou University First Affiliated Hospital

Guojun Zhang

Zhengzhou University First Affiliated Hospital

Pan Qin

Zhengzhou University First Affiliated Hospital

Da Zhang

Zhengzhou University First Affiliated Hospital

Wei Zhao (✉ WeiZhao7271@yeah.net)

Zhengzhou University First Affiliated Hospital

Research

Keywords: circRNA TADA2A, miR-526b, miR-203, caveolin, fibroblasts, idiopathic pulmonary fibrosis

Posted Date: June 22nd, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-20000/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Cell Death & Disease on July 1st, 2020. See the published version at <https://doi.org/10.1038/s41419-020-02747-9>.

Abstract

The excessive activation and proliferation of lung fibroblasts are responsible for the abundant deposition of extracellular matrix (ECM) in idiopathic pulmonary fibrosis (IPF), while its specific mechanism is still unknown. This study focuses on the role of circRNA (circ) TADA2A in functional abnormalities of lung fibroblasts and aims to elaborate its regulatory mechanism. In the present study, circTADA2A was down-regulated in both IPF primary human lung fibroblasts and human IPF fibroblastic cell lines. Functionally, the overexpression of circTADA2A repressed the activation and proliferation of normal human fibroblastic cell line induced by several fibrogenic growth factors. Using fluorescence in situ hybridization (FISH), luciferase reporter assays, and RNA pull-down, circTADA2A was confirmed to function as sponges of miR-526b and miR-203, thus releasing the expression of Caveolin (Cav)-1 and Cav-2. The overexpression of circTADA2A suppressed lung fibroblasts activation via Cav-1 and reduced lung fibroblasts proliferation via Cav-2. *In vivo* experiments also confirmed that the overexpression of circTADA2A decreased fibrogenic responses induced by bleomycin in lung fibrosis mice. Collectively, circTADA2A repressed lung fibroblasts activation via miR-526b/Cav-1 and reduced lung fibroblasts proliferation via miR-203/Cav-2, thus inhibiting the excessive deposition of ECM and relieving IPF.

Introduction

Idiopathic pulmonary fibrosis (IPF) is the most common form of interstitial lung disease, with high morbidity and lacking medical therapies to improve the survival rate. The representative characteristic of IPF is the abundant deposition of extracellular matrix (ECM) in the alveolar parenchyma, which will sequentially irreversibly alter normal lung architecture and ultimately result in respiratory failure [1]. As a major producer of ECM, the significant role of activated fibroblasts in the development of IPF is apparent.

Fibroblasts serve a crucial role in tissue remodeling and fibrosis. In normal wound healing, activated fibroblasts secrete ECM to provide a scaffold for the restoration of normal lung architecture and eventually undergo apoptosis to restrain the excessive production of ECM [2]. Nevertheless, in the lung tissues of patients with IPF, fibroblasts present several dysfunctions, including apoptosis resistance, abundant proliferation and sustained activation [3], which lead to the massive accumulation of ECM and give rise to fibrosis. Though the mechanism of the excessive activation and proliferation of fibroblasts hasn't been expounded explicitly, several fibrogenic growth factors [platelet-derived growth factor (PDGF), insulin-like growth factor 1 (IGF-1) and transforming growth factor- β 1 (TGF- β 1)] have been proved to contribute to this process during the development of IPF [4]. If the functional abnormalities of fibroblasts which located in or around pulmonary fiber lesion can be restrained, it will be of great significance to the clinical treatment of IPF.

Circular RNA (circRNA) is a new class of endogenous noncoding RNA, generating in the back-splicing process of pre-mRNA. The majority of circRNAs are highly conserved across species and play regulatory roles in the onset and progression of numerous human diseases. A recent study revealed that several

circRNAs were dysregulated in the plasma of IPF patients [5], indicating circRNAs may play a potential role in the development of IPF. Besides, emerging evidence showed that circRNAs owned the ability to regulate fibroblasts proliferation and activation via acting as the sponges of microRNAs (miRNAs), binding to them directly and removing their suppressive effect on mRNA expression [6]. Zhu et al. [7] found that the overexpression of circNFIB notably decreased the proliferation of primary mice cardiac fibroblasts via sponging miR-433. Tang et al. [8] reported circRNA_000203 exhibited its promotion effect on the activation of mice cardiac fibroblasts through targeting miR-26b-5p. Inspired by these findings, we speculated that circRNA could also function as miRNA sponges, regulate proliferation and activation of lung fibroblasts, thus influencing the development of IPF.

Materials And Methods

Isolation and growth of human primary lung fibroblasts

IPF primary human lung fibroblasts (IPF-HLF) were acquired from diagnostic biopsies of IPF patients ($n=5$) and normal primary human lung fibroblasts (N-HLF) were acquired from healthy donor lung tissues failed for transplantation ($n=5$). The diagnosis of IPF was supported by history, physical examination, pulmonary function tests, and typical high-resolution chest computed tomography findings of IPF. The IPF patients ($n=5$) and healthy donors ($n=5$) were all male and aged from 40 to 65 years. Exclusion criteria included current or recent use of immunosuppression; chronic infection such as HIV or hepatitis; known pulmonary hypertension; cardiovascular, renal, or neoplastic disease; and inability to provide informed consent. This study was approved by the Ethics Committees of The First Affiliated Hospital of Zhengzhou University and written informed consent was obtained on all patients prior to the procedure being performed.

The human primary lung fibroblasts were isolated as previously described [9]. Briefly, lung tissues were cut into small pieces and subjected to enzymatic dissociation in Hank's balanced salt solution (containing 600 U/ml collagenase I, 2 U/ml papain, 2 U/ml protease, and 3.8 mM calcium chloride) at 37°C for 1 hour. Next, tissues were ground by glass pipette trituration, and the supernatant was centrifuged at 800g for 5min to collect the cells. Then, cells were redispersed in high-glucose Dulbecco's Modified Eagle Medium (DMEM) containing 10% fetal bovine serum (FBS), 50 U/ml strepto-ECM, and 50 µg/ml penicillin. The phenotype of human primary lung fibroblasts was confirmed by positive immunocytochemistry for vimentin (data not shown). In all experiments, the lung fibroblasts were harvest in passages four to six for the test.

Cell culture

The human IPF fibroblastic cell lines (LL-97A and LL-29) and normal human fibroblastic cell line (LL-24) were obtained from the American Type Culture Collection (USA), maintained in Kaighn's Modification of Ham's F-12 Medium (F-12K Medium) containing 15% FBS and cultured with 5% CO₂ at 37°C. The hyperproliferation and activation of normal human lung fibroblasts were performed as previously

described [4]. Briefly, LL-24 cells were starved for 48 hours and incubated with fetal calf serum (FCS; 2% or 5%), PDGF-BB (30 or 60 ng/ml), and IGF-1 (100 or 200 ng/ml) for 6 h to promote fibroblasts proliferation and incubated with TGF- β 1 (5 or 10 ng/ml) for 6 h to activate fibroblasts. All growth factors and cytokines were purchased from Solarbio (China).

Cell transfection and infection

Cells were cultured in 6-well plates with a concentration of 4×10^5 cells/well. When the cells were cultured to 70% confluence, cells were transfected with RNAi-vector [si-circTADA2A, si-Caveolin 1 (Cav-1), si-Cav2] or micro RNA inhibitor (miR-526b inhibitor, miR-203 inhibitor) or their relative negative controls (si-control, NC) using Lipofectamine 2000 (Invitrogen, USA). Cells were incubated with 2 ml Opti-MEM medium (GIBCO, USA) containing plasmids (1 μ g) and Lipofectamine 3000 (2.5 μ l). The medium was changed after 6 h, and the RNA extraction was performed at 48 h to verify the transfection efficiency.

To overexpress circTADA2A or Cav1 or Cav2, the adenovirus expressed circTADA2A or Cav1 or Cav2 (Ad-circTADA2A or Ad-Cav1 or Ad-Cav2) were produced by Ribobio (China). The appropriate volume of virus particles calculated by the multiplicity of infection (MOI) was added in the cell culture medium. Forty-eight hours later, virus infection efficiency was monitored by GFP expression using the fluorescence microscope. Ad-GFP was used as a negative control. The sequences of the transfected components were shown in Table 1.

Cell Proliferation Assay

Lung fibroblasts proliferation was detected using the BrdU incorporation assay kit (Sigma Aldrich, USA). Cells were cultured in 6-well plates with a concentration of 4×10^5 cells/well. When the cells reached confluence, cells were incubated with BrdU (10 μ M) for 40 min. After that, cells were fixed in 4% paraformaldehyde (PFA) for 10min and permeabilized with 0.5% Triton X-100 for 15 min. Then, cells were orderly treated with a blocking buffer, a primary antibody against BrdU, secondary antibody, and DAPI. The BrdU-positive cells were counted using the fluorescence microscope (Nikon, Japan).

Quantitative RT-PCR

Total RNAs were isolated from cells or lung tissues of lung fibrosis mice using TRIzol Reagent (Invitrogen, USA). The quality of total RNA samples was evaluated by spectrophotometer and the high-quality RNAs ($1.8 < OD_{260/280} < 2.0$) were inversely transcribed into cDNA using a cDNA synthesis kit (Thermo Fisher). Quantitative RT-PCR was performed to measure circRNAs and miRNAs expressions using the Thunderbird SYBR qPCR mix (Toyobo, Japan). Gene expressions were calculated by the $2^{-\Delta\Delta CT}$ method, the relative expressions of circRNA and miRNA were normalized to GAPDH and U6 respectively. The sequences of qRT-PCR primers were shown in Table 1.

Western blot

The determination of protein levels of collagen 1a1 (COL1A1), collagen 3a1 (COL3A1), α -smooth muscle actin (α -SMA), laminin (LN), fibronectin (FN), Caveolin-1 (Cav1) and Caveolin-2 (Cav2) were done by western blot with total protein purified from cell lysate or lung tissues of lung fibrosis mice by RIPA lysis buffer. Proteins were subjected to 10% sodium dodecyl sulfate-polyacrylamide gel electrophoresis and then transferred to PDVF membrane (ThermoFisher Scientific, USA). After being blocked with 5% skim milk for 30 min, membranes were incubated with primary antibodies (against COL1A1, COL3A1, α -SMA, LN, FN, Cav1 and Cav2; all purchased from Abcam). After the night, the membranes were incubated with the secondary antibodies specifically. Immunoblots were visualized in IBright FL1500 Intelligent Imaging System (ThermoFisher, USA) and GAPDH was used as an internal control.

The primary antibodies used in the experiment were as follows: anti-COL1A1 (1:1000; sc-59772, Santa Cruz Biotechnology, USA), anti-COL3A1 (1:1000; sc-271249, Santa Cruz Biotechnology, USA), anti- α -SMA (1:1000; ab32575, Abcam, UK), anti-LN (1:2000; ab11575, Abcam, UK), anti-FN (1:1000; ab268021, Abcam, UK), anti-Cav1 (1 μ g/ml; ab2910, Abcam, UK), anti-Cav2 (1:5000; ab133484, Abcam, UK), and GAPDH (1:2500; ab9485, Abcam, UK). The secondary antibody used in the experiment were m-IgGk BP-HRP (1:10000; sc-516102, Santa Cruz Biotechnology, USA) and Goat Anti-Rabbit IgG H&L (1:5000; ab205718, Abcam, UK).

Dual-luciferase reporter gene assay

To verify the combination of circTADA2A and miR-526b/miR-203, the sequence of circTADA2A was amplified and inserted into pGL3-basic plasmids. 0.5 μ g plasmid and 20nM miR-526b mimic or miR-203 mimic or mimic-negative control (Pre-NC) were co-transfected in well-grown 293T cells by using lipofectamine 2000 (ThermoFisher, USA). Forty-eight hours after transfection, the cells were lysed and the activities of Renilla luciferase and firefly luciferase were measured with Dual-luciferase Reporter Assay Kit (Promega, China) following the manufacturer's protocol. To verify the combination of miR-526b and Cav1/miR-203 and Cav2. The luciferase activities of Cav1-3'-UTR-mutant (Cav1 UTR WT), Cav2-3'-UTR-mutant (Cav2 UTR WT), and Cav2-3'-UTR-mutant (Cav2 UTR WT) were measured in the same way. The primer sequences were shown in Table 2.

RNA pull-down assay

The combination of circTADA2A and miR-526b/miR-203 was determined by RNA pull-down assay. The circTADA2A probe and its negative control [oligonucleotide probe (Oligo probe)] were constructed by Ribobio (China). LL-29 cells (1.5×10^7) were collected and lysed using 100 μ l lysis buffer. The lysate was then incubated with 50 pmol biotin-labeled circTADA2A probe and 50 μ l streptavidin agarose magnetic beads for 1 h at 4°C. miR-526b and miR-203 in circTADA2A probe pull-down complex were detected by qRT-PCR using the Oligo probe pull-down complex as a negative control. The probe sequences were shown in Table 2.

Fluorescence in situ hybridization (FISH)

LL-29 cells were cultured on the coverslips, fixed with 4% paraformaldehyde for 15 min., incubated with proteinase K, and washed with alcohol solutions. Then, the slides were incubated with hybridization solution for 30 min at 37°C. Cy3-labeled circTADA2A probe and FAM-labeled miR-526b/miR-203 probes were denatured for 8 min at 73°C and hybridized to the slides for 24 h at 42 °C. Then, blocking was performed and 4,6-diamidino-2-phenyl-indole (DAPI) was used to counterstain the cell nuclei. Finally, the images were obtained with a confocal microscope (Carl Zeiss, Germany).

Mouse model of IPF

Male C57BL/6 mice (4-6 weeks old) were purchased from Laboratory Animal Resources, Chinese Academy of Sciences (Beijing, China). The Ad-circTADA2A and its negative control (Ad-vector) were produced by Ribobio (China). Before the establishment of a lung fibrosis mice model, 50 µl saline containing 110^{11} virus particles of Ad-circTADA2A or Ad-vector were injected in mice intratracheally. After 2 days of injection, mice were divided into four groups. In bleomycin (BLM; n=6), BLM+vector (n=6), and BLM+ circTADA2A (n=6) groups, BLM (3 U/kg) in 50 µl saline was administered intratracheally in mice using trachea cannula. In the saline group (n=6), 50 µl saline was administered intratracheally in mice using the same way. Two weeks later, lung function measurements (total lung capacity, lung compliance, and tissue resistance) were detected by Resistance and Compliance Plethysmographs (Yuyan Instruments, China), and then mice were sacrificed for the following experiment. All protocols in this study have been approved by the Ethics Committee of The First Affiliated Hospital of Zhengzhou University.

Lung Histological Examination

For H&E staining, fresh lung tissues were fixed in 4% PFA, embedded in paraffin, and sectioned using an automatic slicing machine (Leica, Germany). The slices were undergone deparaffinating and rehydration and then stained with hematoxylin and eosin (Nanjing Jiancheng Bioengineering Institute, China).

To evaluate the deposition of collagen in lung tissues, Masson trichrome staining was performed as previously described [10]. Briefly, the slices were orderly stained with hematoxylin, ponceau acid fuchsin, and aniline blue. Finally, images were captured using a microscope (Nikon, Japan).

Hydroxyproline Assay

The assessment of hydroxyproline content in lung tissues was performed using a Hydroxyproline Assay Kit (Abcam, USA) under the manufacturer's protocol. The absorbances of samples at 550 nm were obtained utilizing a microplate reader. The results were expressed as µg/mg lung tissues.

Statistical analysis

Experimental results were expressed as mean $\pm$ standard deviation (SD) and analyzed using GraphPad 7.0 Prism. The differences were analyzed by Student t-test or one-way analysis of variance (ANOVA) with the Newman-Keuls post hoc test. Results were considered statistically significant when $P < 0.05$.

Results

CircTADA2A was down-regulated in IPF lung fibroblasts

Li et al. [5] identified 67 circRNAs which were significantly dysregulated in the blood samples of patients with IPF. Among them, 6 dysregulated circRNAs with high absolute fold-change and low P-values (circANKRD42, circCDC27, circZMYM2, circGRHPR, circTADA2A, and circARHGAP26) were selected in our study. As shown in Figure 1A, compared with N-HLF, a significant decrease in circTADA2A expression was observed in IPF-HLF. Then, the human IPF fibroblastic cell lines (LL-97A and LL-29) and normal human fibroblastic cell line (LL-24) were used for further study. The results of qRT-PCR and FISH depicted that circTADA2A was mainly located in the cytoplasm and the abundance of circTADA2A was lower in human IPF fibroblastic cell lines than that of the normal human fibroblastic cell line (Figure 1B). Interestingly, when we added growth factors and cytokines (FCS, PDGF-BB, IGF-1, and TGF- β 1, respectively), which have been reported to driven lung fibroblasts proliferation and activation *in vitro* [4], in the culture medium of LL-24 to stimulate fibroblasts, circTADA2A expression was declined in response to these growth factors and cytokines in a dose-dependent manner (Figure 1C), implying that with the activation of fibroblasts, the circTADA2A expression was gradually decreased.

CircTADA2A inhibited growth factors-driven lung fibroblasts proliferation and activation

In light of the above results, the possibility was hypothesized that circTADA2A played a potential role in IPF progression. To verify this hypothesis, si-circTADA2A or Ad-circTADA2A transfection was implemented to silence or overexpress circTADA2A in LL-24 cells. After that, the potential effect of circTADA2A on the proliferation of lung fibroblasts was assessed. The result of the BrdU incorporation assay implied that si-circTADA2A further elevated LL-24 proliferation induced by FCS, PDGF-BB, and IGF-1, respectively, compared with si-control transfection (Figure 2A). To evaluate the effect of circTADA2A on activation of lung fibroblasts driven by TGF- β 1, a potent stimulator of fibroblasts, the protein levels of fibroblasts activation markers [collagen 1a1 (COL1A1), collagen 3a1 (COL3A1), and α -smooth muscle actin (α -SMA)] and components of the extracellular matrix [laminin (LN) and fibronectin (FN)] secreted by activated fibroblasts were determined. As shown in Figure 2A, the interference of circTADA2A aggravated fibroblasts activation and extracellular matrix secretion. On the contrary, the overexpression of circTADA2A retarded fibroblasts proliferation induced by FCS, PDGF-BB, and IGF-1, and suppressed fibroblasts activated by TGF- β 1 (Figure 2B). Besides, in the human IPF fibroblastic cell line LL-29, the overexpression of circTADA2A reduced the protein levels of COL1A1, COL3A1, α -SMA, LN, and FN (Supplemental Figure 1).

CircTADA2A functioned as sponges of miR-203 and miR-526b

Studies have shown that circRNA could function as miRNA sponges, binding to miRNA directly and removing their suppressive effect on mRNA expression, thus participating in the regulation of gene expression [6]. Using bioinformatics databases (Circular RNA Interactome, <https://circinteractome.nia.nih.gov/>), we found several target miRNAs that had potential binding sites

with circTADA2A. Among them, six miRNAs (miR-203, miR-520f, miR-450b-3p, miR-526b, miR-638, and miR-769-3p) with the highest prediction score were selected in our study. As shown in Figure 3A, the overexpression of circTADA2A distinctly reduced expression levels of miR-203 and miR-526b while not affect expression levels of miR-520f, miR-450b-3p, miR-638, and miR-769-3p in human IPF fibroblastic cell line LL-29. Besides, compared with LL-24 cells, the expression levels of miR-526b and miR-203 were up-regulated in LL-29 cells (Supplemental Figure 2), indicating that miR-526b and miR-203 may take part in the development of IPF. Furthermore, in response to the transfection of si-circTADA2A, the expression levels of miR-203 and miR-526b were ascended in normal human fibroblastic cell line LL-24 (Figure 2B). Next, Dual-luciferase Reporter Assay and RNA pull-down assay were performed to evaluate the endogenous combination of circTADA2A and miR-526b or miR-203. Compared with the negative control (Pre-NC), transfection of both miR-526b-mimic and miR-203-mimic could decrease the luciferase activity of circTADA2A, and the luciferase activity presented further reduction after co-transfection of miR-526b-mimic and miR-203-mimic (Figure 3C). The RNA pull-down assay displayed that abundant miR-526b and miR-203 were observed in the complex pulled down by the circTADA2A probe than that of the Oligo probe (Figure 3D). The results of the FISH assay further proved that circTADA2A and miR-526b/miR-203 were preferentially colocalized in the cytoplasm (Figure 3E).

CircTADA2A released Cav1 and Cav2 expression via sponging miR-526b and miR-203

Cav1 and Cav2 are the major components of the caveolin protein family and expressed widely in lung tissues. It has been reported that Cav1 could suppress lung fibroblasts activation by blocking the TGF- β signaling pathway [11], Cav2 could decrease proliferation [12] and apoptosis resistance [13] of lung fibroblasts. What's more, an online bioinformatics database (microRNA.org) revealed that miR-526b and miR-203 had the putative binding sites with Cav1 and Cav2 respectively. Firstly, the effect of miR-526b on Cav1 was evaluated. The miR-526b inhibitor raised Cav1 protein expression (Figure 4A), and the Dual-luciferase Reporter Assays showed that compared with the negative control (NC), the luciferase activity of Cav1 3'-UTR wild type (WT) was significantly up-regulated in the cells transfected with miR-526b inhibitor. Whereafter, in IPF fibroblastic cell line (LL-29), the overexpression of miR-526b declined the high expression of Cav1 which was accelerated by Ad-circTADA2A (Figure 4B) while not affect Cav2 (data not shown). In normal fibroblastic cell line (LL-24), the silence of miR-526b elevated the low expression of Cav1 which was suppressed by si-circTADA2A (Figure 4C) while not affect Cav2 (data not shown). The above data indicating that circTADA2A up-regulated Cav1 expression through sponging miR-526b. Afterward, following the same experimental procedure, we proved circTADA2A could also up-regulate Cav2 expression through sponging miR-203 (Figure 4D-F).

The inhibitory effects of circTADA2A on lung fibroblasts activation and proliferation were mediated via Cav1 and Cav2

Once demonstrated that circTADA2A could release the expressions of Cav1 and Cav2 via acting as sponges of miR-526b and miR-203, we decided to confirm whether circTADA2A restrained lung fibroblasts proliferation and activation through Cav1 and Cav2. Firstly, in LL-24 cells that have been

transfected with si-circTADA2A, the overexpression of Cav1 reversed the promotion effect of si-circTADA2A, decreased the lung fibroblasts activation and ECM production (Figure 5A), while the overexpression of Cav2 reduced lung fibroblasts proliferation induced by si-circTADA2A (Figure 6A), hinting that circTADA2A inactivated lung fibroblasts via Cav1 and reduced lung fibroblasts proliferation via Cav2. To further verify this finding, TGF- β 1 was added in the culture medium of LL-24 cells to induce cell activation, while FCS, PDGF-BB, and IGF-1 were added to induce cell proliferation. The overexpression of circTADA2A restrained LL-24 cells from activation and proliferation, on the contrary, the silence of Cav1 eliminated the inhibitory effect of Ad-circTADA2A on lung fibroblasts activation (Figure 5B), and the silence of Cav2 reversed the inhibitory effect of Ad-circTADA2A on cell proliferation (Figure 6B).

CircTADA2A alleviated fibrogenic responses in lung fibrosis mouse

To assess the potential therapeutic effect of circTADA2A *in vivo*, mice were infected with Ad-circTADA2A or negative control (vector) before the establishment of a lung fibrosis mouse model induced by BLM, and the qRT-PCR results proved the infections worked (Figure 7D). We observed that the lung function of circTADA2A-treated lung fibrosis mice was improved as compared to vector-treated lung fibrosis mice (Figure 7A). In line with the lung function analysis, analysis of lung sections manifested Ad-circTADA2A could mainly preserve the alveolar structure and greatly reduce collagen deposition in comparison with vector-treated lung fibrosis mice (Figure 7B). Meanwhile, in circTADA2A-overexpressed lung fibrosis mice, the hydroxyproline content was lower than that of vector-treated lung fibrosis mice (Figure 7C). Additionally, Ad-circTADA2A was also able to down-regulated the BLM-induced high expression levels of miR-256b and miR-203 (Figure 7E) and up-regulated the BLM-induced low protein levels of Cav1 and Cav2 (Figure 7F). The above data signified that miR-256b/Cav1 and miR-203/Cav2 also took part in the therapeutic effect of circTADA2A on lung fibrosis mice.

Discussion

The etiology of IPF has been explained by several hypotheses, among them, the accumulation of ECM induced by hyperproliferating and activating fibroblasts are considered to be the core mechanism. The present study, performed in primary human lung fibroblasts (isolated from healthy and IPF lung tissues), lung fibroblasts cell lines and a mouse model of lung fibrosis, identified new regulatory pathways concerning fibroblasts activation and proliferation, involving circTADA2A, miR-526b, miR-203, Cav1, and Cav2. Briefly, we clarified that high expression of circTADA2A suppressed fibroblasts activation by up-regulating Cav1 via targeting miR-526b, reduced fibroblasts proliferation by up-regulating Cav2 via targeting miR-203, thus decreasing ECM deposition and alleviating IPF. Our research forwarded strong evidence for the vital role of circTADA2A in the progression of IPF, providing novel perspectives for IPF treatment.

Whole-transcriptome RNAs (ncRNAs) sequencing revealed the existence of noncoding RNAs. In recent years, dysfunction of ncRNAs has been proved to be associated with various complex human diseases, and increasing researchers paid attention to the roles of ncRNAs in IPF. Savary et al. [14] identified long

noncoding RNA (lncRNA) DNMT3A as a critical downstream effector of TGF- β during the activation of lung fibroblasts. Yang et al. [15] showed that miR-200 family members (200a, 200b, and 200c) reversed fibrogenic responses in IPF via restraining epithelial-mesenchymal transition (EMT). Nevertheless, to date, compared with lncRNA and miRNA, little is known about the role of circRNA in IPF. Through a circRNA microarray, Li et al [5] screened out 67 circRNA presented visibly dysregulated in the plasma of IPF patients, implying the potential role of circRNA in IPF, while the explicit regulatory mechanism of circRNA in IPF is confusing. In our study, we observed a distinct down-regulation of circTADA2A (hsa_circ_0043278) in the primary human lung fibroblasts isolated from IPF lung tissues. Then, we confirmed this reduction of circTADA2A in both human IPF fibroblastic cell lines and the activation process of a human normal fibroblastic cell line (Fig. 1). Further study showed that the overexpression of circTADA2A effectively suppressed the proliferation and activation of lung fibroblasts driven by fibrogenic growth factors (Fig. 2) and expounded the specific mechanism. The study herein investigated the specific role of circRNA in IPF and clearly point out the anti-fibrosis effect of circTADA2A.

CircTADA2A is identified from exons 5 and 6 of the Transcriptional Adaptor 2A (TADA2A) gene, and its role in disease or biological process is rarely reported. In the osteosarcoma cell line, Wu et al.[16] determined that circTADA2A was mainly located in the cytoplasm and exhibited marked stability under RNase R treatment, revealing the potential of circTADA2A on function as miRNA sponges and abolish the downstream effects of these miRNAs on target mRNAs. In our study, through luciferase reporter gene experiments and RNA pull-down assay, we confirmed circTADA2A could act as sponges of miR-526b and miR-203, hinting that except IPF, circTADA2A may take part in the regulatory network of more human diseases related to miR-526b or miR-203. Chi et al. [17] found that miR-203 inhibited cell proliferation, invasion, and migration of non-small-cell lung cancer by targeting RGS17. Interestingly, in our study, we found that miR-203 negatively regulated Cav2 expression, which could inhibit the proliferation of fibroblasts, hinting that miR-203 may promote cell proliferation in lung fibroblasts. In line with our study, Ren et al. [18] also found that miR-203 promoted proliferation in pancreatic cancer cells via degrading salt-inducible kinase 1. As reported, miRNA can bind to the 3'UTR region of target mRNA and represses gene expression [19]. Therefore, the function of the miRNA is determined by its downstream target gene. In different cell types or different disease states, miRNA could target different target genes and exhibited different functions. Besides, the researches of miR-526b were focused on its anticancer effect [20], in the present study, we firstly reported miR-526b was related to the activation of lung fibroblasts by targeting Cav1.

Caveolin protein family is the major protein component of caveolae (invaginations of the plasma membrane) and takes part in the modulation effect of caveolae on signal transduction [21]. Three members of caveolin protein family have been identified: caveolin-1, -2, and -3. Among them, Cav1 is expressed widely in lung tissues and exhibits an inhibitory effect on fibroblasts activation via directly interacting with the TGF- β 1 receptor (TGF- β 1R) and inhibiting the excessive expression of TGF- β 1 [22]. TGF- β 1 is an essential promoter of collagen amassing and fibrogenesis, driving fibroblasts activation via TGF- β /Smad2/3 and PI3K/Akt signaling [23, 24]. An evident decline of Cav1 expression in lung tissues of IPF patients has been observed [25] and the low expression of Cav1 resulted in the activation of

fibroblasts [11]. In line with this, in our study, Cav1 displayed low expression in IPF fibroblastic cell line (Fig. 4B), and the Dual-luciferase reporter gene assay showed that the decline was due to the combination of miR-526b and 3' UTR of Cav1, the overexpression of circTADA2A released Cav1 expression via sponging miR-526b. What's more, the silence of Cav1 reversed the inhibitory effect of Ad-circTADA2A on lung fibroblasts induced by TGF- β 1 (Fig. 5B), confirming that circTADA2A repressed TGF- β 1-induced fibroblasts activation via Cav1. Except Cav1, emerging evidence showed that Cav2 was also involved in the regulation of pulmonary fibrosis. Almeida et al. [12] reported that during the pulmonary fibrosis induced by bleomycin, compared with wild-type mice, the TGF- β level was not altered while the proliferation of cells was promoted in the lung tissues of Cav2^{-/-} mice. In the present study, we found that the overexpression of Cav2 reversed the promotion effect of si-circTADA2A on cell proliferation while not affect fibroblasts activation (Fig. 6A), confirming that Cav2 was only responsible for the inhibitory effect of circTADA2A on fibroblasts proliferation.

Conclusion

In conclusion, the current study elucidated that circTADA2A could repress fibroblasts activation and proliferation via miR-526b/ Cav1 and miR-203/ Cav2 pathway, thus alleviating IPF. Our study clarified the anti-fibrosis effect of circTADA2A for the first time and providing new perspectives for IPF therapy and circRNA investigation.

Declarations

Ethics approval and consent to participate

All procedures of this study were approved by the Ethics Committees of The First Affiliated Hospital of Zhengzhou University and each subject signed informed consent forms.

All protocols in this study have been approved by the Ethics Committee of The First

Consent for publication

Not applicable.

Availability of data and material

All data generated or analyzed during this study are included in this published article.

Conflict of interest

The authors declare no conflict of interest.

Acknowledgments

Not applicable.

Funding

This study was supported by Medical Science and Technology Projects of Henan Province of China (SB201902018; 2018020060); Key Research Project of Higher Education in Henan Province of China (20A320066).

Authors' contributions

Wei Zhao contributed to the conception of the study;

Juan Li performed the experiments and wrote the manuscript;

Ping Li and Guojun Zhang performed the data analysis;

Pan Qin and Da Zhang helped perform the analysis with constructive discussions.

References

1. Burgess JK, et al. The extracellular matrix - the under-recognized element in lung disease? *J Pathol.* 2016;240(4):397–409.
2. White ES, Lazar MH, Thannickal VJ. Pathogenetic mechanisms in usual interstitial pneumonia/idiopathic pulmonary fibrosis. *J Pathol.* 2003;201(3):343–54.
3. Waters DW, et al. Fibroblast senescence in the pathology of idiopathic pulmonary fibrosis. *Am J Physiol Lung Cell Mol Physiol.* 2018;315(2):L162–72.
4. Al-Tamari HM, et al. FoxO3 an important player in fibrogenesis and therapeutic target for idiopathic pulmonary fibrosis. *EMBO molecular medicine.* 2018;10(2):276–93.
5. Li R, et al. Potential regulatory role of circular RNA in idiopathic pulmonary fibrosis. *Int J Mol Med.* 2018;42(6):3256–68.
6. Wilusz JE, Sharp PA. Molecular biology. A circuitous route to noncoding RNA. *Science.* 2013;340(6131):440–1.
7. Zhu Y, et al. Upregulation of Circular RNA CircNFIB Attenuates Cardiac Fibrosis by Sponging miR-433. *Frontiers in genetics.* 2019;10:564–4.
8. Tang CM, et al. CircRNA_000203 enhances the expression of fibrosis-associated genes by derepressing targets of miR-26b-5p, Col1a2 and CTGF, in cardiac fibroblasts. *Sci Rep.* 2017;7:40342.
9. Ghavami S, et al. Autophagy and the unfolded protein response promote profibrotic effects of TGF- β (1) in human lung fibroblasts. *Am J Physiol Lung Cell Mol Physiol.* 2018;314(3):L493–504.
10. Huang H, et al. ANGPTL2 regulates autophagy through the MEK/ERK/Nrf-1 pathway and affects the progression of renal fibrosis in diabetic nephropathy. *American journal of translational research.* 2019;11(9):5472–86.

11. Liu J, et al. Fluorofenidone attenuates TGF-beta1-induced lung fibroblast activation via restoring the expression of caveolin-1. *Shock*. 2015;43(2):201–7.
12. de Almeida CJG, et al. Genetic ablation of caveolin-2 sensitizes mice to bleomycin-induced injury. *Cell Cycle*. 2013;12(14):2248–54.
13. Predescu SA, et al. Mouse Lung Fibroblast Resistance to Fas-Mediated Apoptosis Is Dependent on the Baculoviral Inhibitor of Apoptosis Protein 4 and the Cellular FLICE-Inhibitory Protein. *Frontiers in physiology*. 2017;8:128–8.
14. Savary G, et al. The Long Noncoding RNA DN30S Is a Reservoir of FibromiRs with Major Functions in Lung Fibroblast Response to TGF-beta and Pulmonary Fibrosis. *Am J Respir Crit Care Med*. 2019;200(2):184–98.
15. Yang S, et al. Participation of miR-200 in pulmonary fibrosis. *Am J Pathol*. 2012;180(2):484–93.
16. Wu Y, et al. Circular RNA circTADA2A promotes osteosarcoma progression and metastasis by sponging miR-203a-3p and regulating CREB3 expression. *Mol Cancer*. 2019;18(1):73–3.
17. Chi Y, et al. miR-203 inhibits cell proliferation, invasion, and migration of non-small-cell lung cancer by downregulating RGS17. *Cancer Sci*. 2017;108(12):2366–72.
18. Ren ZG, et al. miR-203 promotes proliferation, migration and invasion by degrading SIK1 in pancreatic cancer. *Oncol Rep*. 2016;35(3):1365–74.
19. Fabian MR, Sonenberg N, Filipowicz W. Regulation of mRNA translation and stability by microRNAs. *Annu Rev Biochem*. 2010;79:351–79.
20. Zhang R, et al. miR-526b-3p functions as a tumor suppressor in colon cancer by regulating HIF-1alpha. *Am J Transl Res*. 2016;8(6):2783–9.
21. Fridolfsson HN, et al., *Regulation of intracellular signaling and function by caveolin*. FASEB journal: official publication of the Federation of American Societies for Experimental Biology, 2014. 28(9): p. 3823–3831.
22. Gvaramia D, et al. Role of caveolin-1 in fibrotic diseases. *Matrix Biol*. 2013;32(6):307–15.
23. Kulkarni AA, et al. PPAR-γ ligands repress TGFβ-induced myofibroblast differentiation by targeting the PI3K/Akt pathway: implications for therapy of fibrosis. *PloS one*. 2011;6(1):e15909–9.
24. Zhao B, et al. Human amniotic epithelial cells attenuate TGF-beta1-induced human dermal fibroblast transformation to myofibroblasts via TGF-beta1/Smad3 pathway. *Cytotherapy*. 2016;18(8):1012–24.
25. Wang XM, et al. Caveolin-1: a critical regulator of lung fibrosis in idiopathic pulmonary fibrosis. *J Exp Med*. 2006;203(13):2895–906.

Tables

Table 1 Sequences of the transfected components or primers used in the experiments

Transfected components	Sequences
si-circTADA2A	5'-ATTCCATTTCACTACTTCAGA-3'
si-Cav1	5'-GCAAAUACGUAAUGUACAAGU-3'
Si-Cav2	5'-GGAGAUUGGGAUACUGUAAUA-3'
miR-526b mimic	5'-CTCTTGAGGGAAGCACTTTCTGT-3'
miR-203 mimic	5'-GUGAAAUGUUUAGGACCACUAG-3'
miR-526b inhibitor	5'-ACAGAAAGTGCTTCCCTCAAGAG-3'
miR-203 inhibitor	5'-CUAGUGGUCCUAAACAUUUCAC-3'
Ad-circTADA2A	F: 5'-CAAGGAGGAGTGTGAGAAGCACT
(for circTADA2A amplification)	R: 5'-GTCTTGGTGCACATTTGATTGGCT-3'
Primers	Sequences
circANKRD42	F: 5'-CTGGACAAGGCCACATAGAGT-3' R: 5'-CAGAGCAGCCAATGAAGACAC-3'
circCDC27	F: 5'-TCTATTAGGGCATGAGTTTGTCTT-3' R: 5'-TCCTTGGTTGTGGAGCTGTC-3'
circZMYM2	F: 5'-GCACCTGACAGCATCTATTACC-3' R: 5'-GACAGTAGAAACGCAGTAAGCAA-3'
circGRHPR	F: 5'-ACAGATACCACCGCCGAAC-3' R: 5'-TCTAGCTCCTTGGCAGGGAT-3'
circTAD2A	F: 5'-AGCCATTCCATTTCACTACT-3' R: 5'-CCACAGTCCATCACAGCTTC-3'
circARHGAP26	F: 5'-CCATGCAAGCTTTGTGCGAA-3' R: 5'-CATACTTCTTTTTGGCTTCA-3'
miR-203	F: 5'-GGGGTGAAATGTTTAGGAC-3' R: 5'-CAGTGCGTGTCTGGAGT-3'
miR-520f	F: 5'-ACACTCCAGCTGGGAAGTGCTTCCTTTTAG-3' R: 5'-CTCAACTGGTGTCTGGAGTCGGCAATTCAGTTGAGAACCCTCT-3'
miR-450b-3p	F: 5'-GATCCCCGGAUGCAAAUUGAUCCCAATTCA-3' R: 5'-AGCTTAAAAAGGAUGCAAA AUGAUCCCAAT-3'
miR-526b	F: 5'-GTCTCTTGAGGGAAGCACT-3' R: 5'-GTGCAGGGTCCGAGGT-3'
miR-638	F: 5'-AGGGATCGCGGGCGGGTGGCGGCCT-3' R: 5'-ATTCTAGAGGCCGAGGCGGCCGACA TGT-3'
miR-769-3p	F: 5'-TCGGCAGGCTGGGATCTCCGGGG-3' R: 5'-GTGCAGGGTCCGAGGT-3'
U6	F: 5'-CTCGCTTCGGCAGCACA-3' R: 5'-AACGCTTCACGAATTTGCGT-3'
GAPDH	F: 5'-TCACTCAAGATTGTCAGCAA-3' R: 5'-AGATCCACGACGGACACATT-3'
LUC-circTADA2A vector	F: 5'-CCGCTCGAGGCAGGATGTAGCCAATCAAAT-3' R: 5'-ATAAGAAATGCGGCCGCACTGAAATGGAATGGCTGTGT-3'
circTADA2A probe	5'-CATCCTGCAGTGAAATGGAATGGC-3'

Figures

Figure 1

CircTADA2A was down-regulated in IPF lung fibroblasts. (A) Circular RNAs (circRNAs) expression levels in IPF primary human lung fibroblasts (IPF-HLF; n=5) and normal primary human lung fibroblasts (N-HLF; n=5) were measured by qRT-PCR. * $P < 0.05$, ** $P < 0.01$ vs N-HLF. (B) Right: CircTADA2A expression in human IPF fibroblastic cell lines (LL-97A and LL-29) and normal human fibroblastic cell line (LL-24) was measured by qRT-PCR. Left: Representative fluorescence in situ hybridization (FISH) images of circTADA2A in LL-97A, LL-29, and LL-24 (blue, DAPI; red spot, positive staining; Scale bar=20 μ m). ** $P < 0.01$ vs LL-24 cells. (C) LL-24 cells were stimulated with fetal calf serum (FCS; 2% or 5%), platelet-derived growth factor-BB (PDGF-BB; 30 or 60 ng/ml), insulin-like growth factor 1 (IGF-1; 100 or 200 ng/ml), and transforming growth factor- β 1 (TGF- β 1; 5 or 10 ng/ml) for 6 h, respectively. CircTADA2A expression was measured by qRT-PCR. * $P < 0.05$, ** $P < 0.01$ vs LL-24 cells without stimulation.

Figure 2

CircTADA2A inhibited growth factors driven fibroblasts proliferation and activation LL-24 cells were transfected with (A) si-circTADA2A or its negative control (si-control) or (B) adenovirus vector (Ad)-circTADA2A or its negative control (Ad-GFP). Cell proliferation was measured by BrdU incorporation assay after stimulation of 5% FCS or 60 ng/ml PDGF-BB or 200 ng/ml IGF-1 for 6 h. The protein expression of collagen 1a1 (COL1A1), collagen 3a1 (COL3A1), laminin (LN), fibronectin (FN) and α-smooth muscle actin (α-SMA) were determined by western blot after stimulation of 10 ng/ml TGF-β1 for 6 h, GAPDH was used as an internal control. *P<0.05, **P<0.01.

Figure 3

CircTADA2A bound to miR-203 and miR-526b (A) LL-29 cells were transfected with Ad-circTADA2A or Ad-GFP, the expression levels of miR-203, miR-520f, miR-450b-3p, miR-526b, miR-638, and miR-769-3p were measured by qRT-PCR. ** $P < 0.01$ vs Ad-GFP. (B) LL-24 cells were transfected with si-circTADA2A or si-control, the expression levels of miR-526b and miR-203 were measured by qRT-PCR. ** $P < 0.01$ vs si-control. (C) HEK 293T (293T) cells were transfected with miR-526b mimic or miR-203 mimic or miR-526b mimic+miR-203 mimic or negative control (Pre-NC), relative luciferase activities of circTADA2A vector were measured using Dual-Luciferase Reporter Assay System. * $P < 0.05$, ** $P < 0.01$ vs Pre-NC. (D) Detection of miR-526b and miR-203 using qRT-PCR in the samples pulled down by the circTADA2A probe and its negative control (Oligo probe). ** $P < 0.01$ vs Oligo probe. (E) Colocalization between circTADA2A (red) and

miR-526b/miR-203 (green) was observed (arrowheads) using FISH in LL-29 cells. The nuclei were stained with DAPI. Scale bar=20 μ m.

Figure 4

CircTADA2A released Cav1 and Cav2 expression via targeting miR-526b and miR-203 (A) Putative binding sites between miR-526b and Cav1 were forecasted by Miranda. The expression of Cav1 was determined by western blot in LL-29 cells that have been transfected with miR-526b inhibitor or its negative control (NC). Luciferase activity of Cav1 3'-UTR wide type (WT) and Cav1 3'-UTR mutation (MUT) in 293T cells that have been transfected with miR-526b inhibitor or NC. The expression of Cav1 was determined by western blot in (B) LL-29 cells that have been transfected with Ad-circTADA2A, miR-526b mimic, or the negative control of miR-526b mimic (Pre-NC) and (C) LL-24 cells that have been transfected with si-

circTADA2A, miR-526b inhibitor or NC. GAPDH was used as an internal control. (D) Putative binding sites between miR-203 and Cav2. The expression of Cav2 and the luciferase activity of Cav2 3'-UTR WT and Cav2 3'-UTR MUT were determined. The expression of Cav2 was determined by western blot in (E) LL-29 cells that have been transfected with Ad-circTADA2A, miR-203 mimic, or Pre-NC and (F) LL-24 cells that have been transfected with si-circTADA2A, miR-203 inhibitor, or NC. GAPDH was used as an internal control. **P<0.01 vs NC.

Figure 5

CircTADA2A inactivated lung fibroblasts via Cav1 LL-24 cells were transfected with (A) si-circTADA2A or Ad-Cav1+si-circTADA2A or Ad-GFP+si-circTADA2A or (B) Ad-circTADA2A or Ad-GFP or Ad-circTADA2A+si-control or Ad-circTADA2A+si-Cav1, and then stimulation with 5% FCS or 60 ng/ml PDGF-BB or 200 ng/ml

IGF-1 or 10 ng/ml TGF-β1 for 6 h. The expression of COL1A1, COL3A1, LN, FN, and α-SMA were measured by western blot, and the cell proliferation was measured by BrdU incorporation assay. **P<0.01.

Figure 6

CircTADA2A inhibited lung fibroblasts proliferation via Cav2 LL-24 cells were transfected with (A) si-circTADA2A or si-circTADA2A+Ad-Cav2 or i-circTADA2A+Ad-GFP or (B) Ad-circTADA2A or Ad-GFP or Ad-circTADA2A+si-control or Ad-circTADA2A+si-Cav2, and then stimulation with 5% FCS or 60 ng/ml PDGF-BB or 200 ng/ml IGF-1 or 10 ng/ml TGF-β1 for 6 h. The expression of COL1A1, COL3A1, LN, FN, and α-SMA were measured by western blot, and the cell proliferation was measured by BrdU incorporation assay. **P<0.01.

Figure 7

CircTADA2A alleviated pulmonary fibrosis in lung fibrosis mice. Mice were divided into four groups: saline (n=6), bleomycin (BLM; n=6), BLM+adenovirus circTADA2A (circTADA2A; n=6), and BLM+adenovirus vector (vector; n=6). Mice were infected with Ad-circTADA2A and Ad-vector, 2 days later, BLM (3 U/kg) in 50 μl saline was administered intratracheally in mice to induce pulmonary fibrosis. (A) Two weeks after BLM stimulation, lung function measurements of mice (total lung capacity, lung compliance, and tissue resistance) were measured, and then the mice were sacrificed and the lung tissues were collected for the following experiments. (B) H&E and Masson staining of each group. Scale bars= 200 μm. (C) Hydroxyproline content of lung homogenates was evaluated using a Hydroxyproline Assay Kit. (D) The expression level of circTADA2A was measured by qRT-PCR. (E) The expression levels of miR-526b and

miR-203 were measured by qRT-PCR. (F) The expression levels of Cav1 and Cav2 were measured by western blot. GAPDH was used as an internal control. **P<0.01.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupplementalFigure1.jpg](#)
- [SupplementalFigure1.jpg](#)
- [SupplementalFigure2.jpg](#)
- [SupplementalFigure2.jpg](#)