DNA Methylation Signatures Associated With Prognosis of Gastric Cancer

Jin Dai

Peking University Cancer Hospital

Akihiro Nishi

University of California, Los Angeles

Zhe-Xuan Li

Peking University Cancer Hospital

Yang Zhang

Peking University Cancer Hospital

Tong Zhou

Peking University Cancer Hospital

Wei-Cheng You

Peking University Cancer Hospital

Wen-Qing Li (✉ wenqing_li@bjmu.edu.cn)

Peking University Cancer Hospital

Kai-Feng Pan

Peking University Cancer Hospital

Research Article

Keywords: bioinformatics, biomarkers, epigenetics, gastric cancer, heterogeneity, methylation, precision medicine, prognosis; survival, The Cancer Genome Atlas

Posted Date: February 22nd, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-200490/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

DNA methylation signatures associated with prognosis of gastric cancer

Jin Dai^{1,2}; Akihiro Nishi²; Zhe-Xuan Li¹, Yang Zhang¹, Tong Zhou¹, Wei-Cheng You¹, Wen-Qing Li^{1,*}, Kai-Feng Pan^{1,*}

¹Key Laboratory of Carcinogenesis and Translational Research (Ministry of Education/Beijing), Department of Cancer Epidemiology, Peking University Cancer Hospital & Institute, Beijing, 100142, People's Republic of China.

²Department of Epidemiology, UCLA Fielding School of Public Health, Los Angeles, CA, 90095, United States.

* These authors contributed equally to this work

Joint Correspondence to:

Wen-Qing Li, Key Laboratory of Carcinogenesis and Translational Research (Ministry of Education/Beijing), Department of Cancer Epidemiology, Peking University Cancer Hospital & Institute, 52 Fucheng Rd, Haidian District, Beijing, 100142, P. R. China, Tel.: +86 10-88196751, Fax: +86 10-88122437, Email: wenqing_li@bjmu.edu.cn

Kai-Feng Pan, Key Laboratory of Carcinogenesis and Translational Research (Ministry of Education/Beijing), Department of Cancer Epidemiology, Peking University Cancer Hospital & Institute, 52 Fucheng Rd, Haidian District, Beijing, 100142, P. R. China, Tel: +86 10-88196701, Fax: +86 10-88122437, Email: pankaufeng2002@yahoo.com.

Short title: DNA methylation signature for GC prognosis

Word Count: Abstract (344); Main text (3559)

Number of Tables: 3; **Number of figures:** 3

Abstract

Background: Few studies have examined prognostic outcomes-associated molecular signatures other than overall survival (OS) for gastric cancer (GC). We aimed to identify DNA methylation biomarkers associated with multiple prognostic outcomes of GC in an epigenome-wide association study.

Methods: Based on the Cancer Genome Atlas (TCGA), DNA methylation loci associated with OS (n=381), disease-specific survival (DSS, n=372), and progression-free interval (PFI, n=383) were discovered in training set subjects (false discovery rates < 0.05) randomly selected for each prognostic outcome and were then validated in remaining subjects (P -values < 0.05). Key CpGs simultaneously validated for OS, DSS, and PFI were further assessed for disease-free interval (DFI, n=247). Gene set enrichment analyses were conducted to explore the Gene Ontology and Kyoto Encyclopedia of Genes and Genomes pathways simultaneously enriched for multiple GC prognostic outcomes. Methylation correlated blocks (MCBs) were identified for co-methylation patterns associated with GC prognosis. Based on key CpGs, risk score models were established to predict four prognostic outcomes. Spearman correlation analyses were performed between key CpG sites and their host gene mRNA expression.

Results: We newly identified DNA methylation of seven CpGs significantly associated with OS, DSS, and PFI of GC, including cg10399824 (*GRK5*), cg05275153 (*RGS12*), cg24406668 (*MMP9*), cg14719951 (*DSC3*), and cg25117092 (*MED12L*), and two in intergenic regions (cg11348188 and cg11671115). Except cg10399824 and cg24406668, five of them were also significantly associated with DFI of GC. Neuroactive ligand-receptor interaction pathway was suggested to play a key role in the effect of DNA methylation on GC prognosis. Consistent with individual CpG-level association, three MCBs involving cg11671115, cg14719951, and

cg24406668 were significantly associated with multiple prognostic outcomes of GC. Integrating key CpG loci, two risk score models performed well in predicting GC prognosis. Gene body DNA methylation of cg14719951, cg10399824, and cg25117092 was associated with their host gene expression, whereas no significant associations between their host gene expression and four clinical prognostic outcomes of GC were observed.

Conclusions: We newly identified seven CpGs associated with OS, DSS, and PFI of GC, with five of them also associated with DFI, which might inform patient stratification in clinical practices.

Keywords: bioinformatics; biomarkers; epigenetics; gastric cancer; heterogeneity; methylation; precision medicine; prognosis; survival; The Cancer Genome Atlas.

Introduction

Gastric cancer (GC) is the fifth most common cancer and third leading cause of cancer deaths worldwide¹. A considerable proportion of GC patients are diagnosed at advanced stages and their prognosis remains poor². Moreover, GC patients with same clinical stage may develop diverse prognostic outcomes due to the epigenetic or genetic host heterogeneities. Hence, identifying molecular signatures to predict GC prognosis would be warranted for tailored clinical procedures. However, established molecular biomarkers which could predict the prognosis of GC are still very limited.

DNA methylation is a covalent chemical modification, which results in the addition of a methyl (CH₃) group at the carbon 5 position of the cytosine ring³. Alterations in DNA methylation, possessing several merits, such as high frequency in tumor, easy detection, and stability in fixed samples over time, have been considered as promising targets for developing prognostic biomarkers⁴. Based on the Cancer Genome Atlas (TCGA), several studies have identified DNA methylation-based biomarkers associated with overall survival (OS) of GC patients, with three of them based on differentially methylated genes between GC tumor and normal tissue⁵⁻⁷ and the other one not appropriately account for multiple hypothesis testing⁸. In addition, although OS is the most clinically relevant endpoint with the least equivocal definition, solely using OS as the prognostic endpoint may not be sufficient to mirror all perspectives of tumor prognosis, as patient deaths may be due to GC or other causes⁹. Disease-specific survival (DSS) therefore would possess greater relevance to GC-specific biology and therapeutic impact. In addition, given the relatively short follow-up period of GC (median follow-up time = 14 months) in TCGA, the number of deaths at the end of follow-up may be insufficient to reveal a difference in deaths among comparison groups^{10,11}. Progression-free interval (PFI) and disease-free interval (DFI) have therefore been considered as better choices of surrogate clinical

endpoints. To our best knowledge, few studies have identified DNA methylation-based biomarkers associated with prognostic outcomes other than OS of GC patients.

In the present epigenome-wide association study, we comprehensively examined the association between DNA methylation and four prognostic outcomes of GC, including OS, DSS, PFI, and DFI of GC patients in TCGA.

Materials and methods

Data sources and preprocessing

DNA methylation profile (IDAT files) from 395 GC patients were obtained using R/Bioconductor package TCGAAbiolinks¹². In TCGA, DNA methylation of cancer tissue from all patients were assayed using the Infinium HumanMethylation450 BeadChip arrays. A TCGA online tool was used to estimate batch effect (<https://bioinformatics.mdanderson.org/BatchEffectsViewer/>), which would be presented if the dispersion separability criterion (DSC) value was greater than 0.5 and the DSC *P*-value was less than 0.05. Data importation, annotation, and quality control were conducted in the R/Bioconductor package minfi¹³. We used R/Bioconductor package SeSAMe to perform background correction and dye-bias correction^{14,15}. In brief, normal-exponential using out-of-band probes (NOOB) background correction was utilized to account for additive error to the measure of signal intensities due to background fluorescence. Non-linear dye-bias correction was performed to control for the different average intensities in the red and green channels of Type II probes. In addition, *P*-value with out-of-band (OOB) array hybridization (POOBAH) method was applied to address artifacts which reflected failed hybridization to target DNA respectively. For probe filtering, we excluded probes for which the detection *P*-value was bigger than 0.05 for more than 10% of the remaining samples, those on the sex chromosomes, those mapped to multiple regions in the genome, and those harboring known single-nucleotide polymorphisms

with a minor allele frequency greater than 5%¹⁶.

Demographical and clinical characteristics of GC patients, including age at initial pathologic diagnosis, gender, ethnicity, histological grade, anatomic region, family history of GC, *Helicobacter pylori* infection, radiation therapy, and the tumor, lymph nodes, and metastasis (TNM) stage were obtained using R/Bioconductor package TCGAbiolinks¹². We further imputed missing ethnicities using R/Bioconductor package SeSAME¹⁵. Four prognostic outcomes, including OS, DSS, PFI, and DFI of GC, were derived from those defined in a previous study¹¹. In short, using the date of diagnosis of primary GC as the beginning date, OS was defined as the period until the date of death from any cause; DSS was defined as the period until the date of GC-specific death; PFI was defined as the period until the date of the first occurrence of a new GC; DFI was defined as the period until the date of the first new GC progression event subsequent to the ascertainment of a patient's disease-free status after their initial GC diagnosis and treatment. The GC-free status of patients must be confirmed in order for them to be qualified for calculating DFI.

Identification of key CpG sites associated with four clinical endpoints

We first examined the association between DNA methylation of each individual CpG site and OS of GC. GC patients with information on OS were randomly divided into training and validation sets (50:50). For CpG sites significantly associated with OS in the training set (false discovery rate (FDR) < 0.05 considering multiple comparisons)¹⁷, we sought to replicate the associations in the validation sets ($P < 0.05$). For those replicated CpG loci, associations with DSS and PFI of GC were then examined. GC patients with information on DSS or PFI were randomly divided into training and validation sets (50:50) for the assessment of each specific outcome. We then selected key CpG sites significantly associated with OS, DSS, and PFI in the respective training and validation sets, which were further assessed for the association with DFI.

Due to the difficulty of ascertaining GC-free status of patients after their initial diagnosis and treatment, a large body of patients, especially for those in stage IV, were excluded in DFI analysis. We therefore did not seek to divide it into training and validation sets for statistical power consideration. Cox proportional hazard analyses were performed to calculate hazard ratios (HRs) and 95% confidence intervals (CIs) for the association between each individual CpG site and clinical endpoint. Our primary analyses were adjusted for age at initial pathologic diagnosis (continuous) and gender (male, female). In secondary analyses, ethnicity (Asian, Black or African American, or White), TNM stage (T1-2, T3-4, or unknown; N0-1, N2-3, or unknown; M0, M1, or unknown), histological grade (G1-2, G3, or unknown), anatomic neoplasm region (antrum/distal, cardia/proximal, fundus/body, gastroesophageal junction, or unknown), radiation therapy (yes, no, or unknown), *H. pylori* infection (yes, no, or unknown), and family history of GC (yes, no, or unknown). We did not find violations of proportional hazard assumptions for all Cox proportional hazard models.

Gene set enrichment analysis

Gene set enrichment analyses (GSEA) were conducted to explore the Gene Ontology (GO) and Kyoto Encyclopedia of Genes and Genomes (KEGG) pathways which simultaneously enriched for multiple GC prognostic outcomes using DNA methylation data^{18,19}. Utilizing functional class scoring approach, all CpG sites were ranked based on their Cox regression *P*-values with GC prognosis. Gene set level *P*-values were computed through permutation using R/Bioconductor package methylGSA²⁰. The number of CpG sites located in one gene was adjusted in the logistic model to avoid length bias. Gene sets were defined as those containing at least 10 genes and at most 500 genes. FDR less than 0.05 was set as the criteria defining significantly enriched GO or KEGG terms.

Identification of methylation correlated blocks (MCBs) based on key CpGs

In addition to individual CpG loci-level analyses, we also examined the association for a cluster of CpG sites located in the same genomic region based on the concept of genetic linkage disequilibrium. To investigate whether key CpG sites had co-methylation patterns with their adjacent CpG sites and whether these patterns were associated with GC prognosis, Pearson correlation analyses were conducted between DNA methylation level of key CpG sites and their adjacent loci positioned within their flanking areas of one kilobase up- or downstream. Pearson correlation coefficients less than 0.5 indicated boundaries of uncorrelated methylation²¹. For adjacent loci that were not separated by boundaries, we continued examining the correlation between DNA methylation level of these loci and their neighboring loci located within their flanking areas. All CpG sites not separated by the boundary were combined into methylation correlated blocks (MCBs)²¹. The mean β values of all CpG sites within one MCB was defined as the methylation value of that MCB. The R/Bioconductor co-MET package was utilized to visualize the co-methylation patterns of identified MCBs²². We then examined the association between those identified MCBs and four clinical endpoints using Cox regression analyses.

Establishment of two risk score models

Incorporating key CpG sites associated with GC prognosis, risk score models to predict the risk of specific prognostic outcomes were constructed, which were defined as the sum of auto-scaled DNA methylation level of each CpG weighted by the regression coefficients obtained from the Cox regression model in the whole dataset:

$$\text{Risk score} = \beta_1 X_1 + \beta_2 X_2 + \beta_3 X_3 + \dots + \beta_n X_n,$$

GC patients were categorized into low-, median-, and high- risk score groups based on the tertile of their risk scores. Kaplan-Meier survival curves were drawn and log-rank tests were performed to compare the survival probabilities among different risk score groups.

Correlation between DNA methylation and mRNA expression

To explore the potential regulating effect of DNA methylation on host gene expression, Spearman correlation analyses were performed between DNA methylation of those key CpG sites and their host gene mRNA expression. Normalized encoding genes' mRNA expression data (legacy data) based on RNA-Seq (Illumina RNA-Seq HiSeq platform) was downloaded and preprocessed using R/Bioconductor package TCGAAbiolinks¹². Both mRNA expression level and DNA methylation level (β value) were auto-scaled before down-stream analysis. P -value < 0.05 was set as the criteria for the significant correlation. For those genes whose DNA methylation was correlated with their mRNA expression, the association of their mRNA expression with four clinical endpoints of GC were then examined using Cox regression analysis.

All analyses were performed and visualized using R version 4.0.2.

Results

Major characteristics of GC patients in TCGA

In the present study, we included GC patients having both DNA methylation profiles and intact follow-up information of OS (n=381), DSS (n=372), PFI (n=383), and DFI (n=247). During the follow-up, 149, 99, 137, and 46 GC patients developed corresponding clinical endpoints of OS (death from any cause), DSS (GC-specific death), PFI (the first occurrence of a new GC event), and DFI (the first new GC progression event), respectively. Major characteristics of included GC patients were shown in Table 1.

Identification of key CpG sites associated with four clinical endpoints

According to TCGA Batch Effect Viewer, no significant batch effect was observed (DSC = 0.307, DSC P -value = 5×10^{-4}). After data preprocessing, 411,408 CpG sites were kept in our analysis. By performing the Cox regression analyses, we identified a total of 488 CpG sites significantly associated with OS (FDR < 0.05 in the training set and P -value < 0.05 in the

validation set). Among them, seven key CpG sites were simultaneously associated with DSS and PFI in both training (FDR < 0.05) and validation sets (P -value < 0.05), including cg10399824 (G protein-coupled receptor kinase 5, *GRK5*), cg05275153 (regulator of G protein signaling 12, *RGS12*), cg24406668 (matrix metalloproteinase 9, *MMP9*), cg14719951 (desmocollin 3, *DSC3*), and cg25117092 (mediator complex subunit 12L, *MED12L/MED12*) and two CpG loci (cg11348188 and cg11671115) located within intergenic regions. Of them, DNA hypermethylation of three CpG sites, including cg11348188 and cg11671115 in intergenic regions, and cg25117092 in gene body was significantly associated with unfavorable OS, DSS, and PFI of GC, whereas DNA hypermethylation of cg05275153, cg14719951, cg10399824, and cg24406668 in gene body was significantly associated with favorable OS, DSS, and PFI of GC. These CpG loci were further assessed for DFI of GC. Except for cg10399824 and cg24406668, DNA methylation level of the other five CpG sites was also significantly associated with DFI. The directions of associations between seven key CpG sites and all four clinical endpoints remained consistent (Table 2).

GSEA based on CpGs within gene body

Since five key CpG sites that had their host gene information were all located in gene body, GSEA was conducted exclusively based on all CpG sites located in gene body to elucidate the biological mechanisms of gene body DNA methylation on GC prognosis. A total of five GO terms were enriched in all four clinical endpoints according to their original P -values, but none remained simultaneously significant for all four endpoints after multiple comparison adjustment. Four KEGG pathways were simultaneously enriched in OS, DSS, and PFI, with one pathway involving neuroactive ligand-receptor interaction remaining consistently significant even after multiple comparison adjustment (Table 3).

Identification of MCBs associated with GC prognosis

Investigating the potential co-methylation patterns focusing on seven key CpG sites, we identified three MCBs, including one consisting of cg11671115 and cg19989498 in the intergenic region, another containing cg14719951 and 12 adjacent CpG sites in *DSC3* gene body, and the third one including cg24406668 and seven adjacent CpG sites in *MMP9* gene body. Consistent with individual CpG-level association, DNA methylation of these three MCBs was significantly associated with multiple prognostic outcomes of GC (Figure 1).

Establishment of two risk score models

Based on DNA methylation of the seven key CpG sites associated with OS, DSS, and PFI of GC, a risk score models was established:

Model 1 = (-0.301) cg05275153 + (-0.436) cg10399824 + (0.375) cg11348188 + (0.348) cg11671115 + (-0.335) cg14719951 + (-0.363) cg24406668 + (0.353) cg25117092

We also established a risk score model based on the five CpG sites associated with DFI:

Model 2 = (-0.301) cg05275153 + (0.375) cg11348188 + (0.348) cg11671115 + (-0.335) cg14719951 + (0.353) cg25117092

For both models, higher scores indicated worse prognosis potential. After adjusting for all available, potential confounders, both risk scores were independently associated with corresponding clinical endpoints, with a HR (95% CI) per one score increase of 1.41 (1.24-1.60) for OS, 1.70 (1.44-1.99) for DSS, 1.49 (1.31-1.70) for PFI, and 1.76 (1.30-2.38) for DFI. Kaplan-Meier survival curves showed that two models performed well on predicting OS (log-rank $P = 3 \times 10^{-7}$), DSS ($P = 3 \times 10^{-8}$), PFI ($P = 3 \times 10^{-8}$), and DFI ($P = 5 \times 10^{-4}$) (Figure 2).

Integrating with all clinicopathologic features, our risk score models achieved the Harrell's concordance index (C-index) of a mean (standard deviation (SD)) of 0.770 (0.019) for OS, 0.794 (0.021) for DSS, 0.743 (0.023) for PFI, and 0.784 (0.030) for DFI, which outperformed in

prediction compared with models solely utilizing either clinicopathologic features or DNA methylation signatures.

Correlation analysis between DNA methylation and mRNA expression

For the above highlighted key CpG sites in the gene body, we examined the associations of their DNA methylation level with host gene mRNA expression. Gene body DNA hyper-methylation of cg14719951 (correlation coefficient = -0.42, P -value $< 2.2 \times 10^{-16}$) and cg10399824 (correlation coefficient = -0.40, P -value $< 2.2 \times 10^{-16}$) was inversely associated with mRNA expression of *DSC3* and *GRK5*, respectively. On the contrary, Gene body DNA hyper-methylation of cg25117092 (correlation coefficient = 0.46, P -value $< 2.2 \times 10^{-16}$) was significantly associated with increased mRNA expression of *MED12L*. We did not find significant associations between the mRNA expression of these three genes and four clinical prognostic outcomes of GC (Figure 3).

Discussion

We comprehensively examined DNA methylation biomarkers associated with multiple prognostic outcomes of GC in an epigenome-wide association analysis based on TCGA. Altered DNA methylation of seven newly reported CpGs was significantly associated with OS, DSS and PFI of GC, including cg10399824 (*GRK5*), cg05275153 (*RGS12*), cg24406668 (*MMP9*), cg14719951 (*DSC3*), and cg25117092 (*MED12L*), and two in intergenic regions (cg11348188 and cg11671115). Except for cg10399824 and cg24406668, five of them were also significantly associated with DFI of GC. Combining the significant CpG sites, two risk score models performed well in predicting GC prognosis. Consistent with individual CpG-level associations, DNA methylation of three MCBs involving cg11671115, cg14719951 and cg24406668 was significantly associated with multiple prognostic outcomes of GC.

Genomic location of DNA methylation sites is crucial for determining their potential biological functions^{4,23}. It is well known that aberrant promoter DNA hypermethylation could suppress gene expression³. The biological mechanisms of aberrant DNA methylation in intergenic regions or gene body, however, remained overall unclear^{4,24}. In the current study, cg11348188 and cg11671115 were located in intergenic regions and the other five highlighted CpG sites were located in gene body. Of them, DNA hypermethylation of *MED12L* was associated with increased mRNA expression, while hypermethylation of *DSC3* and *GRK5* were associated with decreased mRNA expression. Indeed, previous studies have also reported the complex associations between gene body hypermethylation and increased^{4,25} or decreased gene expression^{24,26}. The reason for the opposite directions of findings is unclear. Gene body DNA methylation might increase transcription through blocking the initiation of alternative promoters within gene body or regulating the activities of repetitive DNAs within the transcribed unit^{23,26}. In addition, splicing or elongation of the ordered structure within the transcriptional unit, which was induced by gene body DNA methylation, might also increase transcriptional activity²⁵. However, intragenic regions also contain many functional elements, such as enhancers, transcription factor binding sites, and repetitive elements, activation of which induced by aberrant gene body DNA methylation could affect host gene expression. In our study, DNA hypermethylation of *RGS12* and *MMP9* were not correlated with host gene expression. Interestingly, previous studies on melanoma have reported gene body DNA hypermethylation and increased mRNA expression of *MMP9*, suggesting possible cell-type specificity of DNA methylation pattern^{27,28}.

We placed special interests on *MED12L*, *DSC3*, and *GRK5*, the DNA methylation of which was significantly associated with mRNA expression. Somatic mutations and DNA

methylation may commonly target the same gene associated with cancer prognosis²⁹. *MED12L* somatic mutations have been identified in both familial and sporadic GC as well as several other cancers^{30,31}. Promoter DNA hypermethylation of *DSC3*, which suppressed host gene expression, has been associated with poor OS of colorectal cancer and recurrence of prostate cancer^{32,33}. The potential role of *GRK5* DNA methylation in the progression/prognosis of cancer has been rarely studied. Interestingly, *MED12L*, *DSC3*, and *GRK5* mRNA expression was not associated with any GC prognostic outcome in our study. Therefore, gene body DNA methylation might affect GC prognosis not through regulating host gene expression, supporting that the identification of associations between DNA methylation and mRNA expression may not be prerequisites of developing DNA methylation-based biomarkers⁴. In fact, several commercially available DNA methylation-based biomarkers, including DNA methylation of *APC* (adenomatosis polyposis coli) and *RASSF1* (ras association domain family member 1) as both diagnostic and prognostic biomarkers of prostate cancer and DNA methylation of *SEPT9* (sepin 9) as predictive biomarker of colorectal cancer, were not correlated with host gene expression⁴.

Gene set enrichment analysis revealed that gene body methylation could affect GC prognosis by regulating the neuroactive ligand-receptor interaction pathway, which has been also enriched for a number of long noncoding RNAs associated with OS of GC in a prior study³⁴. Moreover, this pathway has also been enriched for differentially methylated genes between tumor and normal tissue of GC³⁵, indicating a significant role in both carcinogenesis and prognosis of GC. Future studies are warranted to clarify how aberrant DNA methylation affect the neuroactive ligand-receptor interaction pathway.

Although DNA methylation of single CpG dinucleotides could effectively regulate host gene expression and are qualified as cancer prognostic biomarkers³⁶, the pertinent alterations of

DNA methylation are often regional based, with interplays of a number of adjacent CpG sites. Therefore, in addition to identifying the optimal genomic location, defining the optimal number of relevant CpG sites are of great importance for developing DNA methylation-based biomarker assays⁴. In our study, we also examined DNA methylation patterns accounting for the cluster of adjacent CpG loci. The association of DNA methylation of three identified MCBs with GC prognosis was in line with that based on single CpG locus, which further strengthened our results.

Based on the publicly available dataset with a relatively large sample size of GC patients, our study was the first to thoroughly study DNA-methylation based biomarkers associated with multiple prognostic endpoints of GC, including OS, DSS, DFI, and PFI. The association between DNA methylation of seven key CpG loci and GC prognosis was newly reported. The genomic location and the optimal number of adjacent relevant CpG sites were critically evaluated and reported. Two DNA methylation-based risk score models performed well in predicting four clinical outcomes of GC patients. We were able to adjust for the major clinicopathological characteristics in a secondary analysis, which did not change the results materially.

We acknowledged several limitations. First, DNA methylation profiles in TCGA were measured by Infinium 450k microarrays, which might restrict us from identifying other uncovered CpG dinucleotides with biological relevance to GC prognosis. Second, although we tried to adjust for potential confounders, residual confounding might be present. Third, as discovery and validation datasets utilized to identify prognostic CpG sites were both from TCGA, external datasets are warranted to further validate our results in the future. Fourth, our study was an association study based on biostatistical and bioinformatics analyses. Laboratory work is warranted to unravel the biological consequence of these types of DNA methylation.

In conclusion, our study newly identified seven CpG sites associated with OS, DSS, and PFI of GC, with five of them also associated with DFI of GC. Two DNA methylation-based risk score models were established, which may have implications for clinical practices regarding GC patient stratification of prognosis. Exploration in the experiment setting may contribute to our understanding of the underlying molecular mechanism of gene body DNA methylation on GC prognosis and may inspire the development of novel individualized DNA-methylation based therapeutic strategies.

Declarations

Ethics approval and consent to participate: Inform consent was waived by the Institutional Review Board of Peking University Cancer Hospital as this study was based on a publicly accessible dataset.

Consent for publication: Not applicable

Availability of data and materials: The datasets analyzed during the current study are available in the Genomic Data Commons Data Portal of National Cancer Institute

(<https://portal.gdc.cancer.gov/>)

Competing interests: The authors declare no potential conflicts of interest.

Funding: This work was supported by Beijing Talents Foundation (2018000021223ZK01), Michigan Medicine-PKUHS Joint Institute for Translational and Clinical Research (BMU2020JI004), Capital's Funds for Health Improvement and Research (CFH 2020-2-1026), Beijing Municipal Administration of Hospitals' Ascent Plan (DFL20181102).

Authors' contributions: Jin Dai - study concept and design, acquisition of data, statistical analysis, interpretation of data, drafting of the manuscript, critical revision of the manuscript for important intellectual content; Akihiro Nishi, Zhe-Xuan Li, Yang Zhang, Tong Zhou, Wei-Cheng You - critical revision of the manuscript for important intellectual content; Wen-Qing Li - study concept and design, interpretation of data, critical revision of the manuscript for important intellectual content, funding support, study supervision; Kai-Feng Pan - critical revision of the manuscript for important intellectual content, funding support, study supervision.

Acknowledgements: We thank all individuals who participated in this study and donated samples.

Reference

- 1 Bray, F. *et al.* Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA: a cancer journal for clinicians* **68**, 394-424, doi:10.3322/caac.21492 (2018).
- 2 The global, regional, and national burden of stomach cancer in 195 countries, 1990-2017: a systematic analysis for the Global Burden of Disease study 2017. *Lancet Gastroenterol Hepatol* **5**, 42-54, doi:10.1016/s2468-1253(19)30328-0 (2020).
- 3 Das, P. M. & Singal, R. DNA methylation and cancer. *J Clin Oncol* **22**, 4632-4642, doi:10.1200/jco.2004.07.151 (2004).
- 4 Koch, A. *et al.* Analysis of DNA methylation in cancer: location revisited. *Nat Rev Clin Oncol* **15**, 459-466, doi:10.1038/s41571-018-0004-4 (2018).
- 5 Peng, Y., Wu, Q., Wang, L., Wang, H. & Yin, F. A DNA methylation signature to improve survival prediction of gastric cancer. *Clin Epigenetics* **12**, 15, doi:10.1186/s13148-020-0807-x (2020).
- 6 Bai, Y. *et al.* Development and Validation of a Prognostic Nomogram for Gastric Cancer Based on DNA Methylation-Driven Differentially Expressed Genes. *Int J Biol Sci* **16**, 1153-1165, doi:10.7150/ijbs.41587 (2020).
- 7 Liu, D. *et al.* Discovery and validation of methylated-differentially expressed genes in *Helicobacter pylori*-induced gastric cancer. *Cancer Gene Ther*, doi:10.1038/s41417-019-0125-7 (2019).
- 8 Li, C. *et al.* A four-DNA methylation signature as a novel prognostic biomarker for survival of patients with gastric cancer. *Cancer Cell Int* **20**, 88, doi:10.1186/s12935-020-1156-8 (2020).
- 9 Hudis, C. A. *et al.* Proposal for standardized definitions for efficacy end points in adjuvant breast cancer trials: the STEEP system. *J Clin Oncol* **25**, 2127-2132, doi:10.1200/jco.2006.10.3523 (2007).
- 10 Chibaudel, B. *et al.* Alternative end points to evaluate a therapeutic strategy in advanced colorectal cancer: evaluation of progression-free survival, duration of disease control, and time to failure of strategy--an Aide et Recherche en Cancerologie Digestive Group Study. *J Clin Oncol* **29**, 4199-4204, doi:10.1200/jco.2011.35.5867 (2011).
- 11 Liu, J. *et al.* An Integrated TCGA Pan-Cancer Clinical Data Resource to Drive High-Quality Survival Outcome Analytics. *Cell* **173**, 400-416.e411, doi:10.1016/j.cell.2018.02.052 (2018).
- 12 Colaprico, A. *et al.* TCGAbiolinks: an R/Bioconductor package for integrative analysis of TCGA data. *Nucleic acids research* **44**, e71, doi:10.1093/nar/gkv1507 (2016).
- 13 Aryee, M. J. *et al.* Minfi: a flexible and comprehensive Bioconductor package for the analysis of Infinium DNA methylation microarrays. *Bioinformatics* **30**, 1363-1369, doi:10.1093/bioinformatics/btu049 (2014).
- 14 Triche, T. J., Jr., Weisenberger, D. J., Van Den Berg, D., Laird, P. W. & Siegmund, K. D. Low-level processing of Illumina Infinium DNA Methylation BeadArrays. *Nucleic Acids Res* **41**, e90, doi:10.1093/nar/gkt090 (2013).
- 15 Zhou, W., Triche, T. J., Jr., Laird, P. W. & Shen, H. SeSAMe: reducing artifactual detection of DNA methylation by Infinium BeadChips in genomic deletions.

Nucleic Acids Res **46**, e123, doi:10.1093/nar/gky691 (2018).

16 Dai, J. *et al.* Whole Genome Messenger RNA Profiling Identifies a Novel Signature to Predict Gastric Cancer Survival. *Clin Transl Gastroenterol* **10**, e00004, doi:10.14309/ctg.0000000000000004 (2019).

17 Glickman, M. E., Rao, S. R. & Schultz, M. R. False discovery rate control is a recommended alternative to Bonferroni-type adjustments in health studies. *J Clin Epidemiol* **67**, 850-857, doi:10.1016/j.jclinepi.2014.03.012 (2014).

18 Kanehisa, M. *et al.* KEGG for linking genomes to life and the environment. *Nucleic Acids Res* **36**, D480-484, doi:10.1093/nar/gkm882 (2008).

19 Gene Ontology Consortium: going forward. *Nucleic Acids Res* **43**, D1049-1056, doi:10.1093/nar/gku1179 (2015).

20 Ren, X. & Kuan, P. F. methylGSA: a Bioconductor package and Shiny app for DNA methylation data length bias adjustment in gene set testing. *Bioinformatics* **35**, 1958-1959, doi:10.1093/bioinformatics/bty892 (2019).

21 Hao, X. *et al.* DNA methylation markers for diagnosis and prognosis of common cancers. *Proc Natl Acad Sci U S A* **114**, 7414-7419, doi:10.1073/pnas.1703577114 (2017).

22 Martin, T. C., Yet, I., Tsai, P. C. & Bell, J. T. coMET: visualisation of regional epigenome-wide association scan results and DNA co-methylation patterns. *BMC Bioinformatics* **16**, 131, doi:10.1186/s12859-015-0568-2 (2015).

23 Jones, P. A. Functions of DNA methylation: islands, start sites, gene bodies and beyond. *Nat Rev Genet* **13**, 484-492, doi:10.1038/nrg3230 (2012).

24 Kulis, M. *et al.* Epigenomic analysis detects widespread gene-body DNA hypomethylation in chronic lymphocytic leukemia. *Nat Genet* **44**, 1236-1242, doi:10.1038/ng.2443 (2012).

25 Yang, X. *et al.* Gene body methylation can alter gene expression and is a therapeutic target in cancer. *Cancer Cell* **26**, 577-590, doi:10.1016/j.ccr.2014.07.028 (2014).

26 Maunakea, A. K. *et al.* Conserved role of intragenic DNA methylation in regulating alternative promoters. *Nature* **466**, 253-257, doi:10.1038/nature09165 (2010).

27 Falzone, L. *et al.* MMP-9 overexpression is associated with intragenic hypermethylation of MMP9 gene in melanoma. *Aging (Albany NY)* **8**, 933-944, doi:10.18632/aging.100951 (2016).

28 Kristensen, L. S., Raynor, M. P., Candiloro, I. & Dobrovic, A. Methylation profiling of normal individuals reveals mosaic promoter methylation of cancer-associated genes. *Oncotarget* **3**, 450-461, doi:10.18632/oncotarget.480 (2012).

29 Chan, T. A. *et al.* Convergence of mutation and epigenetic alterations identifies common genes in cancer that predict for poor prognosis. *PLoS Med* **5**, e114, doi:10.1371/journal.pmed.0050114 (2008).

30 Majewski, I. J. *et al.* An α -E-catenin (CTNNA1) mutation in hereditary diffuse gastric cancer. *J Pathol* **229**, 621-629, doi:10.1002/path.4152 (2013).

31 Barbieri, C. E. *et al.* Exome sequencing identifies recurrent SPOP, FOXA1 and MED12 mutations in prostate cancer. *Nat Genet* **44**, 685-689, doi:10.1038/ng.2279 (2012).

- 32 Cui, T. *et al.* DSC3 expression is regulated by p53, and methylation of DSC3 DNA is a prognostic marker in human colorectal cancer. *Br J Cancer* **104**, 1013-1019, doi:10.1038/bjc.2011.28 (2011).
- 33 Pan, J. *et al.* Association of DSC3 mRNA down-regulation in prostate cancer with promoter hypermethylation and poor prognosis. *PLoS One* **9**, e92815, doi:10.1371/journal.pone.0092815 (2014).
- 34 Su, X. *et al.* Identification of the Prognosis-Related lncRNAs and Genes in Gastric Cancer. *Front Genet* **11**, 27, doi:10.3389/fgene.2020.00027 (2020).
- 35 Li, H., Liu, J. W., Liu, S., Yuan, Y. & Sun, L. P. Bioinformatics-Based Identification of Methylated-Differentially Expressed Genes and Related Pathways in Gastric Cancer. *Dig Dis Sci* **62**, 3029-3039, doi:10.1007/s10620-017-4740-6 (2017).
- 36 Claus, R. *et al.* Quantitative DNA methylation analysis identifies a single CpG dinucleotide important for ZAP-70 expression and predictive of prognosis in chronic lymphocytic leukemia. *J Clin Oncol* **30**, 2483-2491, doi:10.1200/jco.2011.39.3090 (2012).

Table 1. Major Characteristics of GC patients with multiple clinical prognostic endpoints in TCGA

	OS		DSS		PFI		DFI
	Training	Validation	Training	Validation	Training	Validation	
Total number, No	191	190	186	186	192	191	247
Ethnicity, No. (%)¹							
Asian	42 (22.0)	41 (21.6)	38 (20.4)	42 (22.6)	46 (24.0)	37 (19.4)	62 (25.1)
Black or African American	9 (4.7)	5 (2.6)	6 (3.2)	8 (4.3)	10 (5.2)	5 (2.6)	7 (2.8)
White	140 (73.3)	144 (75.8)	142 (76.3)	136 (73.1)	136 (70.8)	149 (78.0)	178 (72.1)
Age, mean (SD)	28.9 (10.7)	29.2 (10.3)	29.1 (10.9)	28.7 (10.1)	29.1 (10.5)	29.0 (10.5)	29.3 (10.6)
Gender, No. (%)							
Male	125 (65.4)	126 (66.3)	123 (66.1)	121 (65.1)	133 (69.3)	120 (62.8)	152 (61.5)
Female	66 (34.6)	64 (33.7)	63 (33.9)	65 (34.9)	59 (30.7)	64 (37.2)	95 (38.5)
Clinical stage, No. (%)							
Stage I	21 (11.0)	27 (14.2)	27 (14.5)	20 (10.8)	29 (15.1)	20 (10.5)	40 (16.2)
Stage II	65 (34.0)	56 (29.5)	60 (32.3)	58 (31.2)	61 (31.8)	60 (31.4)	97 (39.3)
Stage III	85 (44.5)	85 (44.7)	77 (41.4)	88 (47.3)	80 (41.7)	91 (47.6)	106 (42.9)
Stage IV	16 (8.4)	15 (7.9)	14 (7.5)	17 (9.1)	18 (9.4)	13 (6.8)	-
Unknown	4 (2.1)	7 (3.7)	8 (4.3)	3 (1.6)	4 (2.1)	7 (3.7)	4 (1.6)
Histologic grade, No. (%)							
G1-2	67 (35.1)	75 (39.5)	76 (40.9)	63 (33.9)	73 (38.0)	70 (36.6)	91 (36.8)
G3	120 (62.8)	110 (57.9)	106 (57.0)	118 (63.4)	113 (58.9)	118 (61.8)	151 (61.1)
Unknown	4 (2.1)	5 (2.6)	4 (2.2)	5 (2.7)	6 (3.1)	3 (1.6)	5 (2.0)
Radiation therapy, No. (%)							
Yes	29 (15.2)	21 (11.1)	19 (10.2)	31 (16.7)	30 (15.6)	22 (11.5)	40 (16.2)
No	127 (66.5)	123 (64.7)	129 (69.4)	120 (64.5)	119 (62.0)	131 (68.6)	187 (75.7)
Unknown	35 (18.3)	46 (24.2)	38 (20.4)	35 (18.8)	43 (22.4)	38 (19.9)	20 (8.1)
<i>H. pylori</i> infection, No. (%)							
Yes	11 (5.8)	9 (4.7)	9 (4.8)	11 (5.9)	7 (3.6)	13 (6.8)	15 (6.1)
No	84 (44.0)	80 (42.1)	79 (42.5)	80 (43.0)	90 (46.9)	75 (39.3)	93 (37.7)

Unknown	96 (50.3)	101 (53.2)	98 (52.7)	95 (51.1)	95 (49.5)	103 (53.9)	139 (56.3)
Family history of GC, No. (%)							
Yes	11 (5.8)	6 (3.2)	8 (4.3)	9 (4.8)	8 (4.2)	10 (5.2)	11 (4.5)
No	158 (82.7)	157 (82.6)	155 (83.3)	154 (82.8)	162 (84.4)	153 (80.1)	203 (82.2)
Unknown	22 (11.5)	27 (14.2)	23 (12.4)	23 (12.4)	22 (11.5)	28 (14.7)	33 (13.4)
Anatomic subdivision (%)							
Antrum/Distal	77 (40.3)	64 (33.7)	65 (34.9)	71 (38.2)	71 (37.0)	70 (36.6)	92 (37.2)
Cardia/Proximal	24 (12.6)	28 (14.7)	34 (18.3)	18 (9.7)	27 (14.1)	25 (13.1)	28 (11.3)
Fundus/Body	66 (34.6)	72 (37.9)	59 (31.7)	76 (40.9)	70 (36.5)	68 (35.6)	99 (40.1)
Gastroesophageal Junction	19 (9.9)	22 (11.6)	21 (11.3)	19 (10.2)	19 (9.9)	23 (12.0)	23 (9.3)
Unknown	5 (2.6)	4 (2.1)	7 (3.8)	2 (1.1)	5 (2.6)	5 (2.6)	5 (2.0)
Pathologic T, No. (%)							
T1-2	44 (23.0)	48 (25.3)	52 (28.0)	38 (20.4)	49 (25.5)	44 (23.0)	63 (25.5)
T3-4	147 (77.0)	142 (74.7)	134 (72.0)	148 (79.6)	143 (74.5)	147 (77.0)	184 (74.5)
Pathologic N, No. (%)							
N0-1	114 (59.7)	103 (54.2)	111 (59.7)	99 (53.2)	109 (56.8)	109 (57.1)	164 (66.4)
N2-3	74 (38.7)	82 (43.2)	70 (37.6)	84 (45.2)	81 (42.2)	76 (39.8)	78 (31.6)
Unknown	3 (1.6)	5 (2.6)	5 (2.7)	3 (1.6)	2 (1.0)	6 (3.1)	5 (2.0)
Pathologic M, No. (%)							
M0	173 (90.6)	169 (88.9)	166 (89.2)	169 (90.9)	171 (89.1)	172 (90.1)	238 (96.4)
M1	9 (4.7)	13 (6.8)	12 (6.5)	10 (5.4)	11 (5.7)	11 (5.8)	-
Unknown	9 (4.7)	8 (4.2)	8 (4.3)	7 (3.8)	10 (5.2)	8 (4.2)	9 (3.6)
Prognostic endpoints, No. (%)							
Event	77 (40.3)	72 (37.9)	55 (29.6)	44 (23.7)	77 (40.1)	60 (31.4)	46 (18.6)
Censored	114 (59.7)	118 (62.1)	131 (70.4)	142 (76.3)	115 (59.9)	131 (68.6)	201 (81.4)

¹Missing values of ethnicity were imputed using R package SeSAmE.

Abbreviations: GC, gastric cancer; OS, overall survival; DSS, disease-specific survival; DFI, disease-free interval; PFI, progression-free interval.

Table 2. The hazard ratios and *P*-values for the associations between DNA methylation of seven key CpG loci and four clinical prognostic endpoints of GC patients in TCGA

CpG sites	Genomic positions	Gene names	Genomic groups (CpG contents)	Training set			Validation set			Pooled set	
				HR (95% CI) ¹	P-value	FDR	HR (95% CI) ¹	P-value	HR (95% CI) ¹	P-value	
OS											
cg11348188	chr2-222655366	EPHA4, PAX3	Intergenic (open sea)	1.69 (1.32-2.18)	3.84×10 ⁻⁵	0.012	1.33 (1.05-1.68)	0.017	1.46 (1.23-1.72)	1.44×10 ⁻⁵	
cg25117092	chr3-150948274	MED12L	Body (open sea)	1.67 (1.29-2.16)	1.03×10 ⁻⁴	0.017	1.28 (1.01-1.62)	0.037	1.42 (1.20-1.69)	4.70×10 ⁻⁵	
cg05275153	chr4-3431837	RGS12	Body (open sea)	0.79 (0.68-0.91)	9.97×10 ⁻⁴	0.042	0.69 (0.56-0.86)	8.99×10 ⁻⁴	0.74 (0.65-0.84)	3.02×10 ⁻⁶	
cg11671115	chr10-87336486	FAM190B, GRID1	Intergenic (open sea)	1.58 (1.22-2.04)	4.85×10 ⁻⁴	0.032	1.36 (1.08-1.71)	0.010	1.42 (1.20-1.67)	3.52×10 ⁻⁵	
cg14719951	chr18-28622474	DSC3	Body (island)	0.69 (0.55-0.87)	1.33×10 ⁻³	0.048	0.74 (0.58-0.93)	0.010	0.72 (0.61-0.84)	4.22×10 ⁻⁵	
cg10399824	chr10-121048354	GRK5	Body (open sea)	0.61 (0.50-0.75)	3.67×10 ⁻⁶	0.005	0.65 (0.53-0.80)	4.01×10 ⁻⁵	0.65 (0.56-0.74)	7.00×10 ⁻¹⁰	
cg24406668	chr20-44639431	MMP9	Body (island)	0.59 (0.46-0.76)	2.88×10 ⁻⁵	0.011	0.78 (0.62-0.99)	0.037	0.70 (0.59-0.82)	2.29×10 ⁻⁵	
DSS											
cg11348188	chr2-222655366	EPHA4, PAX3	Intergenic (open sea)	1.54 (1.16-2.06)	3.10×10 ⁻³	0.028	2.05 (1.45-2.90)	4.41×10 ⁻⁵	1.73 (1.39-2.15)	9.42×10 ⁻⁷	
cg25117092	chr3-150948274	MED12L	Body (open sea)	1.49 (1.10-2.00)	8.95×10 ⁻³	0.043	1.75 (1.27-2.40)	5.66×10 ⁻⁴	1.60 (1.29-1.98)	2.17×10 ⁻⁵	
cg05275153	chr4-3431837	RGS12	Body (open sea)	0.67 (0.55-0.82)	1.04×10 ⁻⁴	0.005	0.70 (0.53-0.91)	0.009	0.69 (0.59-0.80)	1.32×10 ⁻⁶	
cg11671115	chr10-87336486	FAM190B, GRID1	Intergenic (open sea)	1.59 (1.19-2.12)	1.79×10 ⁻³	0.020	1.71 (1.26-2.31)	5.65×10 ⁻⁴	1.65 (1.34-2.04)	2.43×10 ⁻⁶	
cg14719951	chr18-28622474	DSC3	Body (island)	0.64 (0.49-0.85)	1.62×10 ⁻³	0.020	0.65 (0.49-0.88)	0.005	0.65 (0.53-0.79)	2.16×10 ⁻⁵	
cg10399824	chr10-121048354	GRK5	Body (open sea)	0.54 (0.42-0.69)	9.20×10 ⁻⁷	1.31×10 ⁻⁴	0.57 (0.44-0.73)	7.12×10 ⁻⁶	0.57 (0.48-0.67)	1.65×10 ⁻¹¹	
cg24406668	chr20-44639431	MMP9	Body (island)	0.71 (0.53-0.93)	1.46×10 ⁻²	0.050	0.54 (0.39-0.74)	1.49×10 ⁻⁴	0.62 (0.50-0.76)	6.56×10 ⁻⁶	
PFI											
cg11348188	chr2-222655366	EPHA4, PAX3	Intergenic (open sea)	1.65 (1.27-2.13)	1.53×10 ⁻⁴	0.021	1.53 (1.17-1.99)	0.002	1.58 (1.32-1.89)	7.00×10 ⁻⁷	
cg25117092	chr3-150948274	MED12L	Body (open sea)	1.47 (1.16-1.86)	1.53×10 ⁻³	0.049	1.48 (1.12-1.94)	0.005	1.49 (1.25-1.79)	1.07×10 ⁻⁵	
cg05275153	chr4-3431837	RGS12	Body (open sea)	0.71 (0.59-0.87)	6.84×10 ⁻⁴	0.043	0.54 (0.41-0.72)	2.11×10 ⁻⁵	0.70 (0.61-0.80)	2.17×10 ⁻⁷	

cg11671115	chr10-87336486	<i>FAM190B</i> <i>, GRID1</i>	Intergenic (open sea)	1.57 (1.24-1.99)	1.86×10 ⁻⁴	0.021	1.40 (1.08-1.83)	0.011	1.51 (1.27-1.80)	3.12×10 ⁻⁶
cg14719951	chr18-28622474	<i>DSC3</i>	Body (island)	0.69 (0.55-0.87)	1.52×10 ⁻³	0.049	0.77 (0.60-0.99)	0.043	0.73 (0.62-0.86)	2.48×10 ⁻⁴
cg10399824	chr10-121048354	<i>GRK5</i>	Body (open sea)	0.57 (0.46-0.7)	1.26×10 ⁻⁷	5.62× 10 ⁻⁵	0.75 (0.59-0.96)	0.020	0.63 (0.54-0.74)	5.10×10 ⁻⁹
cg24406668	chr20-44639431	<i>MMP9</i>	Body (island)	0.68 (0.54-0.86)	1.24×10 ⁻³	0.049	0.68 (0.52-0.90)	0.007	0.69 (0.58-0.83)	4.74×10 ⁻⁵
DFI										
cg11348188	chr2-222655366	<i>EPHA4</i> , <i>PAX3</i>	Intergenic (open sea)	-	-	-	-	-	1.51 (1.12-2.05)	0.007
cg25117092	chr3-150948274	<i>MED12L</i>	Body (open sea)	-	-	-	-	-	1.75 (1.27-2.42)	0.001
cg05275153	chr4-3431837	<i>RGS12</i>	Body (open sea)	-	-	-	-	-	0.77 (0.63-0.94)	0.012
cg11671115	chr10-87336486	<i>FAM190B</i> <i>, GRID1</i>	Intergenic (open sea)	-	-	-	-	-	1.59 (1.18-2.14)	0.002
cg14719951	chr18-28622474	<i>DSC3</i>	Body (island)	-	-	-	-	-	0.74 (0.56-0.99)	0.042
cg10399824	chr10-121048354	<i>GRK5</i>	Body (open sea)	-	-	-	-	-	0.78 (0.60-1.01)	0.059
cg24406668	chr20-44639431	<i>MMP9</i>	Body (island)	-	-	-	-	-	0.77 (0.57-1.04)	0.089

¹ Cox regression analyses adjusting for age at initial pathologic diagnosis (continuous) and gender (male, female).

Abbreviations: GC, gastric cancer; OS, overall survival; DSS, disease-specific survival; DFI, disease-free interval; PFI, progression-free interval; HRs, hazard ratios; CI, confidence interval; FDR: false discovery rate.

Table 3. Gene set enrichment analysis for gene body DNA methylation in relation to four clinical endpoints of GC patients in TCGA

			OS		DSS		PFI		DFI	
ID	Description	Size	<i>P</i> -value	FDR	<i>P</i> -value	FDR	<i>P</i> -value	FDR	<i>P</i> -value	FDR
GO										
35267	NuA4 histone acetyltransferase complex	31	0.002	0.066	0.007	0.197	0.047	0.477	0.002	0.042
36019	Endolysosome	21	0.012	0.219	0.002	0.078	0.006	0.152	0.017	0.220
36020	Endolysosome membrane	14	0.007	0.154	0.001	0.036	0.003	0.106	0.036	0.364
43189	H4/H2A histone acetyltransferase complex	31	0.002	0.066	0.007	0.197	0.047	0.477	0.002	0.042
44346	fibroblast apoptotic process	24	0.036	0.409	0.024	0.409	0.029	0.370	1.51×10 ⁻⁴	0.008
KEGG										
04080	Neuroactive ligand-receptor interaction	272	1.24×10 ⁻¹⁶	2.64×10 ⁻¹⁴	2.32×10 ⁻¹⁶	4.95×10 ⁻¹⁴	3.44×10 ⁻¹⁰	7.34×10 ⁻⁸	1.000	1.000
04144	Endocytosis	201	0.005	0.324	0.026	0.923	0.027	0.645	0.194	0.858
04950	Maturity onset diabetes of the young	25	0.037	0.901	0.010	0.634	0.008	0.329	1.000	1.000
05217	Basal cell carcinoma	55	0.017	0.712	0.011	0.634	0.035	0.749	1.000	1.000

Abbreviations: GC, gastric cancer; OS, overall survival; DSS, disease-specific survival; DFI, disease-free interval; PFI, progression-free interval; HRs, hazard ratios; CI, confidence interval; FDR: false discovery rate.

Figure 1. Methylation correlated blocks (MCBs) (A) MCBs based on cg11671115 and adjacent CpG sites in chromosome 10. (B) MCBs based on cg14719951 and adjacent CpG sites in chromosome 18. (C) MCBs based on cg24406668 and adjacent CpG sites in chromosome 20. (D) Forest plots indicating the association between DNA methylation of three MCBs and four clinical endpoints. For (A), (B), and (C), the circles indicate P -values of the associations between each CpG within the MCBs and OS. The red dotted lines represent the significant level ($P = 0.05$). The yellow bars represent locations of genes and the green bars represent locations of CpG islands. The blue dots represent DNA methylation level of each CpG within the MCBs. The blue lines connect mean DNA methylation level of each CpG within the MCBs.

Figure 2. Kaplan-Meier (KM) survival curves of the DNA methylation-based risk score models in TCGA. (A) The seven-CpG based risk score model in OS dataset. (B) The seven-CpG based risk score model in DSS dataset. (C) The seven-CpG based risk score model in PFI dataset. (D) The five-CpG based risk score model in DFI dataset. In each dataset, GC patients were equally categorized into three groups based on the tertile of risk scores of all patients. Blue curves represent low risk score groups; Yellow curves represent median risk score groups; Red curves represent high risk score groups. Log-rank tests were used to compare survival curves among subgroups for each prognostic outcome of GC.

Figure 3. Spearman correlations between gene body DNA methylation of three CpG loci and the mRNA expression of their host genes. (A) Spearman correlation for cg14719951 (DSC3). (B) Spearman correlation for cg10399824 (GRK5). (C) Spearman correlation for cg25117092 (MED12L). (D) Forest plots indicating the association between mRNA expression of three genes and four clinical endpoints of GC patients.

Figures

Figure 1

Methylation correlated blocks (MCBs) (A) MCBs based on cg11671115 and adjacent CpG sites in chromosome 10. (B) MCBs based on cg14719951 and adjacent CpG sites in chromosome 18. (C) MCBs based on cg24406668 and adjacent CpG sites in chromosome 20. (D) Forest plots indicating the association between DNA methylation of three MCBs and four clinical endpoints. For (A), (B), and (C), the circles indicate P-values of the associations between each CpG within the MCBs and OS. The red dotted lines represent the significant level ($P = 0.05$). The yellow bars represent locations of genes and the green bars represent locations of CpG islands. The blue dots represent DNA methylation level of each CpG within the MCBs. The blue lines connect mean DNA methylation level of each CpG within the MCBs.

Figure 2

Kaplan-Meier (KM) survival curves of the DNA methylation-based risk score models in TCGA. (A) The seven-CpG based risk score model in OS dataset. (B) The seven-CpG based risk score model in DSS dataset. (C) The seven-CpG based risk score model in PFI dataset. (D) The five-CpG based risk score model in DFI dataset. In each dataset, GC patients were equally categorized into three groups based on the tertile of risk scores of all patients. Blue curves represent low risk score groups; Yellow curves represent median risk score groups; Red curves represent high risk score groups. Log-rank tests were used to compare survival curves among subgroups for each prognostic outcome of GC.

Figure 3

Spearman correlations between gene body DNA methylation of three CpG loci and the mRNA expression of their host genes. (A) Spearman correlation for cg14719951 (DSC3). (B) Spearman correlation for cg10399824 (GRK5). (C) Spearman correlation for cg25117092 (MED12L). (D) Forest plots indicating the association between mRNA expression of three genes and four clinical endpoints of GC patients.