

Nurses' Knowledge, Attitude, and Fall Prevention Practices at South Korean Hospitals: A Cross-Sectional Survey

Mi-young Cho

Eulji University

Sun Joo Jang (✉ icedcoffee@cau.ac.kr)

Chung-Ang University

Research article

Keywords: attitude, falls, hospital, knowledge, nursing

Posted Date: April 7th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-20233/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Version of Record: A version of this preprint was published on November 24th, 2020. See the published version at <https://doi.org/10.1186/s12912-020-00507-w>.

Abstract

Background: Fall-prevention activities are nursing interventions which are designed to improve patient safety. The introduction of evaluations of medical institutions and an increase in medical litigation has led institutions to emphasize the importance of fall-prevention activities. The current situation regarding falls among patients in small and medium-sized hospitals is poorly understood. This study examined fall-prevention activities in small and medium-sized hospitals and assessed knowledge and attitudes regarding falls of nurses working in these hospitals.

Methods: Participants were 162 nurses from seven small and medium-sized hospitals. Data on participants' characteristics, education regarding patient falls, knowledge of stretcher cart use, attitudes regarding patient falls, and fall-prevention activities were collected from August 1 to September 1, 2016.

Results: Nurses' experience with patients falling was positively correlated with knowledge. Length of nursing experience and attendance at education sessions influenced attitudes regarding patient falls. Attitudes positively correlated with fall-prevention activities, but knowledge did not. Nurses' attitudes regarding patient falls were correlated with fall-prevention activities. To ensure engagement in fall-prevention activities, hospitals should develop incentive programs tailored to each hospital's specific circumstances and create attitude improvement programs based on nurses' subjective norms.

Conclusion: In short, we recommend that consistent, repeated, and custom fall-prevention education should be implemented in small and medium-sized hospitals to promote engagement in fall-prevention activities. Patient safety activities in small and medium-sized hospitals can be enhanced by creating an environment which encourages voluntary and self-directed participation in fall-prevention activities using motivation and rewards.

Background

Falls in hospitals can lead to significant secondary complications for patients [1]. Falls are considered a serious concern because they lead to financial losses and patient discomfort and affect patients' quality of life by prolonging hospitalization and incurring additional medical expenses for tests, surgery, or rehabilitation [2, 3].

According to the Korea Consumer Agency's annual report on patient safety, falls in hospitals accounted for 1,522 (49.7%) of 3,060 cases related to hospital safety management and constituted the largest proportion of safety accidents [4]. Fall rates for hospital inpatients vary considerably according to each medical institution's incident report regulations and standards [5], and the number of reported cases is lower relative to that of actual occurrences [6]. Furthermore, determining accurate fall rates for all domestic hospitals is difficult, because hospitals do not release fall reports. Patients have the right to safety and quality management to prevent falls and injuries [7], especially because falls are preventable and predictable health issues which can generally be reduced 30–50% with environmental improvements. In addition, some researchers have suggested that falls can indicate nursing quality [8]. For these reasons it behooves us to study and better understand falls and fall-prevention practices for the good of hospitals, nurses, and patients alike.

Small and medium-sized hospitals improve accessibility to healthcare, increase the speed of providing emergency care, and serve as the backbone of the medical delivery system by providing residents with appropriate and diverse services for their medical needs [9]. A total of 89.2% of all hospitals in South Korea in 2013 were small and medium-sized hospitals [10]. Although they accounted for two thirds of all medical institutions, the numbers of small and medium-sized hospitals with patient safety departments, patient safety education, and patient safety indicators were lower relative to those of medical institutions with 301 or more beds [11]. In addition, a study that compared patient safety nursing activities between university hospitals and small and medium-sized hospitals showed that the former performed more patient safety nursing activities pertaining to falls than the latter [12]. However, comparisons are difficult – research on the perception of hospital safety culture in South Korea has been conducted primarily in tertiary care medical institutions or medical institutions with 300 or more beds, and subarea classification criteria and analysis methods vary between studies [13]. The composition or

characteristics of medical staff, hospital systems, patients' illness, patients themselves, and patients' caregivers could differ between small and medium-sized hospitals and large hospitals [14].

Falls in small and medium-sized hospitals are not well understood because previous studies have been conducted mostly in tertiary care hospitals and few of these have involved small and medium-sized hospitals [15, 16]. Small- and medium-sized hospitals have challenges with implementing effective strategies and practices to reduce patient falls [17]. According to the Agency for Healthcare Research and Quality [18], successful practice strategies begin with an understanding of current practice. Therefore, an understanding of the knowledge and attitudes of nurses working at small and medium-sized hospitals, which affects engagement in fall-prevention activities, should be acquired to ensure better fall-prevention practices.

The present study examined nurses working in small and medium-sized hospitals' fall-related education and their knowledge, attitudes, and engagement in fall-prevention activities. We also explored differences in these factors and the relationships between nurses' knowledge, attitudes, and engagement in fall-prevention activities. Our results may guide the future implementation of fall-prevention activities in small and medium-sized hospitals.

Methods

Design

We distributed a cross-sectional survey to nurses working in small and medium-sized hospitals to examine their above-mentioned knowledge of, attitudes toward, and engagement in fall-prevention activities.

Participants

Our participants were nurses who worked at small and medium-sized hospitals with fewer than 300 beds located in D and G cities, who understood the purpose of the study, and who agreed to participate in the study. Nurses working at Chinese medicine clinics and long-term care hospitals were excluded from the study.

The sample size necessary for correlation analysis with an effect size of .35, a significance level of $p < .05$, and power of .80 was 142, calculated based on a previous study [19] using G*power 3.1.9 [20]. Expecting a drop-out rate of 20%, we distributed 190 questionnaires; of these, 186 questionnaires were returned. We excluded 24 questionnaires with no responses. Therefore, the final analytic sample ultimately included 162 questionnaires.

Instruments

Knowledge regarding falls

We assessed nurses' knowledge regarding falls using the Hospital Falls Knowledge Scale, which measures nurses' knowledge regarding falls [21] and was created in consultation with two nursing professors and two patient safety experts. The questionnaire consisted of 15 items pertaining to fall rates; types and severity of injuries caused by falls; intrinsic factors including high risk of falls, age group, sex, disease, state of consciousness, activity status, use of assistive devices, risk factors, and use of drugs that increased the risk of falls; and extrinsic factors including fall timing, location, and type. The questionnaire included 14 multiple-choice items and one subjective item (Item 15), and the highest possible score was 14. One point and zero points were awarded for correct and incorrect responses, respectively. The Kuder-Richardson Formula 20 reliability coefficient was .76. The content validity index for the modified instrument exceeded .80 for all items.

Attitudes regarding falls

Attitudes regarding falls refers to nurses' emotional, cognitive, and behavioral reactions to inpatient falls that occurred in hospitals. We assessed these attitudes using the 13-item Hospital Falls Attitude Scale [21], which was used in a previous

study [22]. Responses were provided using a five-point Likert scale ranging from 1 (strongly disagree) to 5 (strongly agree). Higher scores indicated higher levels of positivity in nurses' attitudes. Items 2, 8, 9, 11, and 12 were reverse scored. Cronbach's α for the scale were .75 in a previous study and .72 in the present study.

Fall-prevention activities

Nurses' engagement in fall-prevention activities was measured using a 20-item scale [15], which was based on the Hospital Nurses Association's nursing practice guidelines for falls. Responses were provided using a four-point Likert scale ranging from 1 (never use) to 4 (always use), and higher scores indicated greater engagement in fall-prevention activities. Cronbach's α for the scale was .92 in a previous study as well as in the present study.

Data collection

Data were collected from nurses working at small and medium-sized hospitals with fewer than 300 beds located in D and G cities from August 1 to September 1, 2016.

Ethical considerations

Ethical approval for the study was granted by the institutional review board of Eulji University (approval number: IRB No. EU16-28), and the study protocol conforms to the provisions of the Declaration of Helsinki. Written informed consent was obtained from all participants.

Data analysis

Participants' characteristics and fall-related education were analyzed using frequencies, percentages, means, and standard deviations. Means and standard deviations were calculated for knowledge, attitudes, and engagement in fall-prevention activities. Differences in participants' characteristics, knowledge, and attitudes regarding falls, and engagement in fall-prevention activities according to their experience of fall-related education were analyzed using Mann-Whitney U tests and Kruskal-Wallis tests. Pearson's correlation coefficients were calculated for the relationships between knowledge regarding falls, attitudes regarding falls, and engagement in fall-prevention activities. All data analyses were performed using SPSS/WIN 24.0.

Results

Differences in knowledge, attitudes, and engagement in fall-prevention by participants' experience

The participants' mean age was 32.49 years (SD 8.17). The mean duration of participants' work experience was 8 years and 5 months.

Nurses' experience with patient falls differed significantly depending on their experience. Nurses who had experience with patient falls had significantly higher levels of knowledge regarding falls than those who had no such experience. In addition, attitudes regarding falls differed significantly depending on how long our participants had been working as nurses. Nurses with 5 or more years of experience demonstrated significantly more positive attitudes toward falls than nurses with less experience. However, engagement in fall-prevention activities did not significantly differ by participants' experience (Table 1).

Table 1

Differences in Knowledge, Attitudes, and Engagement in Fall-Prevention Activities According to Participants' Characteristics (N = 162).

Characteristics	n	Knowledge				Attitude				Fall-prevention activities			
		M	SD	χ^2	p	M	SD	χ^2	p	M	SD	χ^2 (p)	p
Age													
< 30 years	70	6.96	1.48	0.72	.488	3.78	0.38	1.04	.356	3.27	0.48	0.52	.597
30–39 years	59	6.64	1.74			3.80	0.32			3.27	0.50		
≥ 40 years	33	6.94	1.43			3.88	0.30			3.37	0.52		
Sex													
Male	5	7.20	0.84	0.56 [†]	.578	3.75	0.15	0.64 [†]	.525	3.21	0.45	0.49 [†]	.628
Female	157	6.83	1.59			3.81	0.35			3.29	0.50		
Educational level													
Community college	81	6.79	1.73	0.76	.685	3.82	0.35	0.93	.630	3.30	0.49	0.68	.713
University	77	6.91	1.43			3.80	0.34			3.28	0.51		
Master's degree or higher	4		0.58			3.69	0.32			3.13	0.49		
Work experience													
< 5 years	62	6.92	1.54	0.51 [†]	.611	3.73	0.40	2.25 [†]	.026	3.25	0.51	0.75 [†]	.455
≥ 5 years	100	6.79	1.59			3.85	0.29			3.31	0.49		
Department													
Internal medicine	20	7.05	1.54	6.78	.079	3.82	0.24	0.85	.838	3.38	0.51	1.54	.674
Surgery	101	7.00	1.61			3.79	0.39			3.25	0.50		
Intensive care unit	22	6.32	1.13			3.85	0.25			3.34	0.50		
Other	14	6.21	1.58			3.82	0.29			3.31	0.52		
Experience of patient falling													
No	80	6.54	1.60	2.46 [†]	.015	3.76	0.32	1.61 [†]	.110	3.26	0.51	0.82 [†]	.415
Yes	82	7.13	1.49			3.85	0.37			3.32	0.49		

M = mean; SD = standard deviation

[†]Mann-Whitney U-test result; SD, Standard deviation

Characteristics	n	Knowledge				Attitude				Fall-prevention activities				
Number of patient falls														
1	32	7.25	1.68	5.89	.207	3.82	0.33	0.89	.926	3.30	0.54	1.21	.877	
2	22	6.55	1.10			3.82	0.40			3.39	0.48			
3	14	7.43	1.55			3.90	0.44			3.34	0.44			
4	5	7.80	1.48			3.94	0.22			3.11	0.61			
≥ 5	9	7.33	1.32			3.91	0.41			3.30	0.39			
Fall-prevention education														
No	35	6.86	1.59	0.08 [†]	.940	3.73	0.26	1.56 [†]	.122	3.14	0.57	1.95 [†]	.053	
Yes	127	6.83	1.57			3.83	0.36			3.33	0.47			
M = mean; SD = standard deviation														
†Mann-Whitney U-test result; SD, Standard deviation														

[Insert Table 1 here]

Differences in knowledge, attitudes, and engagement in fall-prevention activities by fall-related education experience

Nurses' knowledge did not differ significantly by experience with fall-related education. Nurses' attitudes regarding falls differed significantly according to the number of fall-prevention education sessions they attended. Nurses who had attended at least five fall-prevention education sessions demonstrated significantly more positive attitudes toward patient falls than nurses who had attended only one or two sessions. However, engagement in fall-prevention activities did not differ significantly by experience with fall-related education (Table 2).

Table 2

Differences in Knowledge, Attitude, and Engagement in Fall-Prevention Activities according to Experience of Fall-Related Education (N = 127)

Characteristic	n	Knowledge				Attitude				Fall-prevention activities			
		M	SD	χ^2	p	M	SD	χ^2	p	M	SD	χ^2 (p)	p
Host													
Nursing association	16	7.19	1.60	2.61	.456	3.89	0.32	6.00	.112	3.41	0.39	2.99	.394
Nursing society	3	6.33	1.15			4.18	0.25			3.43	0.55		
Hospital	106	6.77	1.58			3.82	0.37			3.32	0.48		
Other	2	8.00	1.41			3.46	0.44			2.90	0.14		
Instructor													
Nursing professor	16	7.19	1.33	3.40	.334	3.87	0.40	1.50	.683	3.26	0.52	3.34	.342
Nurse	107	6.79	1.61			3.83	0.36			3.35	0.47		
Hospital administrator	2	5.50	0.71			3.73	0.27			2.88	0.04		
Other	2	7.50	0.71			3.62	0.22			3.25	0.35		
Education method													
Lecture	38	6.66	1.55	2.82	.588	3.84	0.35	1.93	.749	3.31	0.57	1.78	.776
Audiovisual	5	7.2	1.30			3.69	0.25			3.20	0.27		
Lecture + audiovisual	79	6.91	1.63			3.82	0.37			3.35	0.44		
Lecture + discussion	3	6.00	1.00			3.92	0.20			3.17	0.47		
Other	2	7.5	0.71			4.15	0.76			3.53	0.25		
Education duration													
> 1 hour	41	6.83	1.82	5.68	.224	3.79	0.41	6.15	.188	3.31	0.54	5.59	.232
1–2 hours	60	6.93	1.38			3.82	0.36			3.28	0.45		
2–4 hours	16	6.38	1.67			3.84	0.27			3.55	0.38		
4–6 hours	3	8.67	1.53			3.92	0.08			3.53	0.48		
≥ 6 hours	7	6.29	0.76			4.12	0.27			3.27	0.41		
Number of education sessions													

† Bonferroni correction; M = mean; SD = standard deviation

Characteristic	n	Knowledge				Attitude				Fall-prevention activities			
a.1	39	6.82	1.41	0.58	.966	3.74	0.39	22.68	<.001	3.21	0.52	3.79	.435
b.2	38	6.92	1.73			3.72	0.28	e > a,b [†]		3.37	0.44		
c.3	22	6.86	2.01			3.86	0.31			3.37	0.49		
d.4	6	6.67	1.03			4.06	0.31			3.55	0.39		
e.≥5	22	6.73	1.24			4.08	0.37			3.36	0.42		
†Bonferroni correction; M = mean; SD = standard deviation													

[Insert Table 2 here]

Knowledge and attitudes regarding falls and engagement in fall-prevention activities

The mean proportion of correct answers on the items regarding knowledge about falls was 48.9% (an average of 6.84 out of 14 items). The proportion of correct answers for the items ranged from 3.7–95.1%. Item 5 (state of consciousness) had the lowest proportion of correct answers, and Item 2 (high-risk age) the highest.

The mean rate of positive attitudes regarding falls was 76.2% (Mean 3.81 out of 5, SD = 0.34; Table 3). The mean rate of engagement in fall-prevention activities was 82.3% (Mean 3.29 out of 4, SD = 0.50; Table 4).

Table 3
Average Scores for Attitudes Regarding Falls (N = 162)

Rank	Item	Items pertaining to attitudes regarding falls	M	SD
1	10	I think I should respond promptly when patients ask for help to move.	4.36	0.58
2	7	I think that patients' fall risk should be assessed upon admission.	4.21	0.59
3	11	I do not think that fall-related physical injury is severe. [†]	4.20	0.73
4	6	I think I should actively nurse patients to prevent falls.	4.14	0.62
5	1	I am interested in the occurrence of inpatient falls.	4.07	0.71
6	4	I think that fall prevention is a high priority in nursing.	4.04	0.59
7	5	I am interested in nursing interventions to prevent falls.	3.99	0.66
8	13	I would feel guilty if a patient fell.	3.80	0.80
9	3	I think that falls in hospitals are an important responsibility for nurses.	3.78	0.67
10	2	I think that inpatient falls are inevitable. [†]	3.55	0.90
11	12	I think that the hospital environment is safe with respect to falls. [†]	3.41	0.85
12	8	I think that fall-prevention education provided upon admission is sufficient. [†]	2.71	0.89
13	9	I think that falls occur because of the patient's condition. [†]	2.65	0.86
		Total	3.81	0.34
[†] Reverse scored; M = mean; SD = standard deviation				

Table 4
Participants' Engagement in Fall-Prevention Activities and Average Item Scores (N = 162)

Rank	Item	Items pertaining to engagement in fall-prevention activities	M	SD
1	20	I always raise the bed rails when moving a patient on a stretcher cart.	3.82	0.47
2	19	I always engage the lock when transferring patients to wheelchairs.	3.77	0.51
3	6	I always raise the bed rails for elderly people, children, unconscious patients, and very unstable patients.	3.73	0.53
4	11	I educate patients to ensure that they ask for help to prevent falls.	3.68	0.53
5	8	I ensure that unconscious patients, very unstable patients, or surgical patients are moved from the bed with assistance from a nurse or caregiver.	3.60	0.62
6	10	I ensure that patients at risk of falling walk with their caregivers.	3.60	0.60
7	7	I ensure that patients at risk of falling who wake up to go to bathroom are helped off the bed by a nurse or guardian	3.52	0.69
8	9	In cases of abuse of drugs that can cause falls, I monitor the occurrence of the drug's effects.	3.41	0.75
9	1	I inform all inpatients and caregivers of the possibility of falls while introducing them to hospital life.	3.38	0.75
10	13	I educate patients and caregivers in moving to the bed, chair, bathroom, and wheelchair safely.	3.36	0.76
11	16	I ensure that patients wear non-slip shoes of the correct size.	3.31	0.76
12	17	I maintain proper illumination on the bed and in the bathroom.	3.29	0.76
13	15	Paths should be cleared for easy use.	3.28	0.74
14	12	I provide patients and caregivers with instructions on fall prevention and remind them of these frequently.	3.09	0.93
15	14	I encourage high-risk patients to exercise regularly unless it is contraindicated (once per day).	3.01	0.91
16	5	I attach fall hazard signs to patient charts, patient rooms, and beds for high-risk patients.	2.94	1.11
17	4	I assess patients' levels of normal motor function.	2.93	0.88
18	2	I assess patients' fall risk factors using a fall risk assessment scale upon admission.	2.91	1.15
19	3	I regularly (e.g., twice) reassess fall risk factors in connection with changes in a patient's condition after admission.	2.77	1.06
20	18	I place a non-slip mat on the floor when taking a barrel bath or shower.	2.36	1.13
M = mean; SD = standard deviation				

[Insert Table 3 and Table 4 here]

Relationships between knowledge, attitudes, and engagement in fall-prevention activities

Attitudes about falls were positively related to engagement in fall-prevention activities ($r = .25, p = .001$), but no significant relationships were observed between knowledge of falls and engagement in fall-prevention activities ($r = -.09, p = .267$) or knowledge of falls and attitudes toward falls ($r = .09, p = .240$).

Discussion

The results of this study showed a mean score of 48.9 for the knowledge of falls, which is lower than the score of 6.84 out of 14 items reported in a previous study [22, 23] measuring knowledge regarding falls among university hospital nurses and a score of 66 reported in another study of nurses in general hospitals [24]. These findings suggest that nurses at small and medium-sized hospitals' knowledge regarding falls is lower than that of nurses at tertiary care hospitals. Perhaps this is because small and medium-sized hospitals offer fewer educational opportunities compared to tertiary care hospitals, which provide consistent patient safety education. The item with the lowest knowledge score regarding falls pertained to consciousness at the time of a fall, followed by those pertaining to frequent diseases that cause falls and bodily injury resulting from falls; these findings were similar to those of previous studies [23]. Fall-prevention methods were the most frequently reported educational content that nurses in the study experienced; however, educational content reviewing the causes and effects of falls should be increased in fall-related education.

Levels of knowledge regarding falls in participants who had experience of patients falling were significantly higher relative to those in participants who had no experience of patients falling. This finding is consistent with those of a previous study [22] that measured knowledge regarding falls in university hospital nurses and showed that participants' knowledge regarding falls increased as their interest in falls increased, once they had experienced patients falling.

Mean score for the positivity of nurses' attitudes regarding falls was higher relative to the scores reported in previous studies using the same questionnaire [19, 22, 24]. The positivity of nurses' attitudes regarding falls could have increased as the importance of patient safety increased because of small and medium-sized hospitals' voluntary participation in the medical institution accreditation system introduced through an amendment to the medical law in June 2010.

Attitudes regarding falls varied depending on the duration of work experience. The positivity of attitudes regarding falls in nurses who had worked for over 5 years was higher than reported by the nurses who had worked for less than 5 years. This finding is consistent with previous research reporting significant differences in attitudes based on length of nursing career [22, 24]. This finding might be the result of the higher level of responsibilities and sense of duty found in experienced nurses. Less experienced nurses typically are focused on general nursing activities rather than those involving patient safety.

Participants' mean score for engagement in fall-prevention activities was higher relative to the score of 81.5 previously reported in a study of nurses at a general hospital [24] and the score of 77.3 reported in a study of nurses at small and medium-sized hospitals [15], but was lower relative to the score of 89.8 reported in a study of nurses in geriatric hospitals [19]. The lowest scores were given to Item 18 (e.g., providing a non-slip mat when patients take a barrel bath or shower) and Item 3 (e.g., reassessing fall risk factors after admission). This was consistent with the findings of a study of nurses in a general hospital [24]. It could have been difficult for nurses to reassess fall risk repeatedly and perform fall-related nursing duties regularly in small and medium-sized hospitals, which have less staff performing fall-prevention duties and patient assessments, relative to those at tertiary care hospitals. Aggressive fall-prevention activities require standardized environmental assessment tools for fall risk, and the installation of preventive environmental facilities tailored to small and medium-sized hospitals is needed.

Nurses' attitudes regarding falls were more positive the greater number of fall-prevention education sessions they attended. Therefore, we suggest that regular and recurrent fall-prevention education targeted to small and medium-sized hospitals would reinforce and improve nurses' attitudes toward patient falls.

Attitudes regarding falls were significantly positively correlated with engagement in fall-prevention activities; however, we found no relationship between nurses' knowledge regarding and their attitudes regarding falls or engagement in fall-prevention activities. These findings are consistent with previous research [19, 22, 25]. For example, a multifaceted strategy, including reminders and identification systems, audits and feedback, and further education, has been found to effectively

increase fall-prevention activities [26]. Engagement in fall-prevention activities should be increased via attitude reinforcement programs for nurses.

We observed few significant differences in knowledge or attitudes regarding falls and no significant differences in engagement in fall-prevention activities according to participants' characteristics. This finding might be explained by career nurses' mannerisms resulting from dissatisfaction with their work environments— small and medium-sized hospitals see high nurse turnover and often have insufficient workforces to meet their operational needs [27]. Patient safety activities in small and medium-sized hospitals should be enhanced by creating an atmosphere to foster voluntary and self-directed engagement in fall-prevention activities through the use of motivation and rewards.

The findings described above indicate that consistent and recurrent fall-prevention education tailored to small and medium-sized hospitals should be implemented to promote engagement in fall-prevention activities. In addition, the positivity of nurses' attitudes regarding falls was an important factor affecting behavioral intention, and the promotion of changes in attitudes regarding falls via introducing attitude-reinforcement programs that consider nurses' subjective norms must enhance their sense of responsibility and duty.

Conclusions

In summary, the present study opens the possibility of increasing nurses' interest in fall-prevention activities in small and medium-sized hospitals. Nurses are considered to play a critical role in fall-prevention activities, including assessing risk factors for falls at patients' bedsides and providing interventions 24 hours a day. Therefore, in consideration of these results, small and medium-sized hospitals should implement consistent and repeated fall-prevention education tailored to their unique circumstances in order to promote engagement in fall-prevention activities. Patient safety activities in small and medium-sized hospitals can be enhanced by creating an atmosphere wherein fall-prevention activities are voluntary and self-directed, and increased by providing appropriate motivation and rewards.

This study has some limitations that should be noted. First, data were collected from only seven hospitals in a single city in South Korea; therefore, we cannot generalize too much from these results. This research should be repeated with more nurses from different cities. Second, in our study, the nurses might have understood questions differently as attitudes and engagement in fall prevention activities were evaluated using self-report data.

The present study broadens our understanding of the current situation regarding nurses' knowledge, attitude, and prevention activities regarding patient falls in small and medium-sized hospitals. Nurses' attitudes regarding falls were positively correlated with engagement in fall-prevention activities, but their knowledge regarding falls was not. This study confirmed that nurses' attitudes toward patient safety are critical; considering these findings, a multifaceted implementation program for fall prevention should be implemented to increase the positivity of nurses' attitudes.

Abbreviations

M
mean
SD
standard deviation

Declarations

Ethics approval and consent to participate: Ethical approval for the study was granted by the Institutional Review Board of Eulji University (Approval number: EU16-28). Written consent was obtained from all participants.

Consent for publication: Not applicable.

Availability of data and materials: The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

Conflicts of interest: The authors declare that they have no competing interests.

Funding: This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Authors' contributions: M-YC and SJJ both contributed to the study conception and design, the data collection, and the data analysis and interpretation. Further, both authors were involved in drafting the article, critically revising it, and approved the final version.

Acknowledgements: The authors would like to acknowledge the participation of nurses.

References

1. Gu YY, Balcaen K, Ni Y, Ampe J, Goffin J. Review on prevention of falls in hospital settings. *Chin Nurs Res.* 2016;3(1):7-10.
2. Centers for Disease Control and Prevention. Costs of falls among older adults. Atlanta 2016. <http://www.cdc.gov/HomeandRecreationalSafety/Falls/fallcost.html>. Accessed 19 Jan 2018. (in Korean).
3. Dellinger AM, Stevens JA. The injury problem among older adults: Mortality, morbidity and costs. *J Safety Res.* 2006;37(5):519-22.
4. Ministry of Health and Welfare. An alert system is implemented to prevent patient safety accidents. 2017. https://www.mohw.go.kr/react/al/sal0301vw.jsp?PAR_MENU_ID=04&MENU_ID=0403&CONT_SEQ=342830&page=1. Accessed 19 Jan 2018. (in Korean).
5. Florence CS, Bergen G, Atherly A, Burns E, Stevens J, Drake C. Medical costs of fatal and nonfatal falls in older adults. *J Am Geriatr Soc.* 2018;66(4):693-8.
6. Oliver D, Healey F, Haines TP. Preventing falls and fall-related injuries in hospitals. *Clin Geriatr Med.* 2010;26(4):645-92.
7. Gray-Miceli D, de Cordova PB, Crane GL, Quigley P, Ratcliffe SJ. Nursing home registered nurses' and licensed practical nurses' knowledge of causes of falls. *J Nurs Care Qual.* 2016;31(2):153-60.
8. Kim GS, Kim JA, Kim MS, Kim YJ, Kim ES, Park GO, et al. Development of performance measures based on the nursing process for journal prevention and management of pressure ulcers, falls and pain. *J Korean Clin Nurs Res.* 2010;16:5-23. (in Korean).
9. Han Y, Kim H, Ko J, Kim EJ. Relationship between expectations of clinical ladder system, career commitment and turnover intention of nurses in small and medium sized hospitals. *J Korean Clin Nurs Res.* 2018;24:67-73. (in Korean).
10. Sin HH. Development of roadmap for the promotion and facilitation of small and medium sized hospitals: Based on risk factor analysis. *J Hosp Manag Policy.* 2014;3:257-73. (in Korean).
11. Sin HH, Jeong SH, Yu J. A survey of the current state of patient safety activities of medical institutions and activation plan for patient safety activities. *Korean Institute Hosp Manag.* 2015;9:1-153. (in Korean).
12. Cha BK, Choi JA. A comparative study on perception of patient safety culture and safety care activities: Comparing university hospital nurses and small hospital nurses. *J Korean Nurs Adm.* 2015;21:405-16. (in Korean).
13. Kim HY, Lee ES. Effects of perceived patient safety culture on safety care activities among nurses in general hospitals. *J East-West Nurs Res.* 2013;19:46-54.
14. Lee NJ, Kim JH. Perception of patient safety culture and safety care activity among nurses in small-medium sized general hospitals. *J Korean Nurs Admin Acad Soc.* 2011;17:462-73. (in Korean).
15. Hwang IY. The knowledge, attitude, and fulfillment of practice guidelines for fall

of clinical nurses at small and medium sized hospitals (Master's thesis). Jinju, South Korea: Gyeongsang National University; 2011. p. 75. (in Korean). 16. Lee YJ, Gu MOK. Circumstances, risk factors, and the predictors of falls among patients in the small and medium-sized hospitals. *J Korean Clin Nurs Res.* 2015;21:252-65. (in Korean). 17. Park JH. Fall prevention activities of small and medium sized hospital nurses and structural analysis on related variables (Unpublished doctoral dissertation). Gyeongsan, South Korea: Catholic University of Daegu; 2016. (in Korean). 18. Agency for Healthcare Research and Quality. How will you manage change?. Rockville: MD; 2013. <https://www.ahrq.gov/professionals/systems/hospital/fallpxtoolkit/fallpxtk2.html>. Accessed 19 Jan 2018. 19. Kim SH, Seo JM. Geriatric hospital nurses' knowledge, attitude toward falls, and fall prevention activities. *J Korean Gerontol Nurs.* 2017;19:81-91. (in Korean). 20. Faul F, Erdfelder E, Buchner A, Lang AG. Statistical power analyses using G* Power 3.1: Tests for correlation and regression analyses. *Behav Res Methods.* 2009;41(4):1149-60. 21. Kim CG, Suh MJ. An analysis of fall incidence rate and its related factors of fall in inpatients. *Qual Improv Health Care.* 2002;9:210-28. (in Korean). 22. Lee IK, Choi JY. Factors associated with nurses' activities for hospital fall prevention. *Korean J Rehab Nurs.* 2013;16:55-62. (in Korean). 23. Yoo KS. Knowledge, attitude and prevention activities related to fall among geriatric hospital nurse. *J Korean Pub Health Nurs.* 2017;31:436-50. (in Korean). 24. Choi AS, Oh PJ. A study on self-leadership, fall attitude and nurses' behavior to prevent patient falls. *J Korean Acad Nurs Admin.* 2013;19:394-403. (in Korean). 25. Hiroko KT. Fall Prevention Practice Gap Analysis: aiming for targeted improvements. *MEDSURG Nurs.* 2017;26:332-5. 26. Koh SLS, Hafizah N, Lee JY, Loo YL, Muthu R. Impact of a fall prevention programme in acute hospital settings in Singapore. *Singapore Med J.* 2009;50:425-32. 27. Kim MA, Park GO, Yu SJ, Kim MJ, Kim ES. A survey of nursing activities in small and medium-size hospitals: Reasons for turnover. *J Korean Clin Nurs Res.* 2009;15:149-65. (in Korean).