Cloning and Expressions of *Chop* in Loach (*Misgurnus Anguillicaudatus*) and Its Response to Hydrogen Peroxide (H₂O₂) Stress

Hui Li

Huazhong Agricultural University: Huazhong Agriculture University

Minxin Kang

Huazhong Agricultural University: Huazhong Agriculture University

Shouxiang Sun

Huazhong Agricultural University: Huazhong Agriculture University

Jian Gao

Huazhong Agricultural University: Huazhong Agriculture University

Zhiying Jia

Heilongjiang River Fishery Research Institute of Chinese Academy of Fishery Sciences

Xiaojuan Cao (✉ caoxiaojuan@mail.hzau.edu.cn)

College of fisheries huazhong agriculture university <https://orcid.org/0000-0002-9676-5838>

Research Article

Keywords: *Misgurnus anguillicaudatus*, chop, Gene cloning, Spatio-temporal expressions, Hydrogen peroxide stress

Posted Date: July 13th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-202985/v2>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Fish Physiology and Biochemistry on April 9th, 2022. See the published version at <https://doi.org/10.1007/s10695-022-01067-3>.

Abstract

C/EBP [CCAAT/enhancer-binding protein]-homologous protein gene (*chop*) which plays an important role in endoplasmic reticulum stress-induced apoptosis, was investigated here by RACE and qPCR in an aquaculture animal for the first time. The full-length cDNA sequences of loach (*Misgurnus anguillicaudatus*) *chop* were 2533bp, encoding 266 amino acids. The expression levels of loach *chop* changed obviously during different early life stages, and the highest appeared at the 8-cell stage. Among different tissues, loach *chop* predominantly expressed in gill, spleen and gonad. Then, we performed a hydrogen peroxide (H₂O₂, a common-used disinfectant) stress trial to explore the role of loach *chop*, with three different concentrations (0 μM, 50 μM and 100 μM) of H₂O₂. Half of the loaches from the 100 μM group were dead, while all loaches from the other two treatment groups were alive. The activities of CAT, SOD and GPX in loach gill, liver and spleen decreased with extended stress time and increased H₂O₂ concentration. The expression levels of gill *chop* in loaches from the 100 μM group were significantly higher than those from the other two treatment groups between 12 and 24 hours of exposure. *atf4* and *bax*, two proapoptotic genes, were significantly up-regulated in gills of loaches from the 100 μM group compared to the other two groups 18 hours and 24 hours after treatment, while *bcl2*, an antiapoptotic gene, presented an opposite trend. These results indicated a close relationship between H₂O₂ stress and fish apoptosis, and loach *chop* played an important role in H₂O₂ stress response.

1. Introduction

C/EBP [CCAAT/enhancer-binding protein]-homologous protein gene (*chop*), also known as the growth arrest and DNA damage-inducible gene 153 (GADD153), is considered as a master proapoptotic molecule of endoplasmic reticulum (ER) stress-induced apoptosis (Li et al. 2014). Although ubiquitously expressing at very low levels under normal conditions (Ariyama et al. 2008), *chop* is a highly stress-inducible expressed gene, markedly following the ER stress (Zhang et al. 1999; Cheng et al. 2009; Pirot et al. 2007). Cells respond to ER stress by activating a series of signaling pathways, collectively termed the unfolded protein response (UPR) pathway (Walter and Ron 2011). It has been proved that three branches of the UPR pathway, namely inositol requiring enzyme 1 (IRE1), prospective evaluation of radial keratotomy (PERK) and activating transcription factor 6 (ATF6) branch, trigger apoptosis when the ER stress is severe, prolonged, or chronic unmitigated (Zhang et al. 2016; Deegan et al. 2015; Shang and Lehrman 2004). The apoptotic program is activated by the elevated expression of massive kinds of transcription factors (Rutkowski et al. 2006), and *chop* is the key one.

More recently, it is reported that *chop* is mainly up-regulated in the PERK branch when organisms are exposed to the ER stress (Cao et al. 2012). PERK is also known as eukaryotic translation initiation factor 2α (eIF2α) kinase 3 (Mounir et al. 2011). It can up-regulate the translations of some specific genes, like activation of transcription factor-4 (*atf4*) (B'chir et al. 2013), by phosphorylating its substrate eIF2α. *atf4* is a member of C/EBP family, and it elevates expression of its key downstream target gene *chop* when stress cannot be alleviated, resulting in the onset of apoptosis (Su and Kilberg 2008). In the PERK branch,

chop is suggested to facilitate apoptosis through repressing the transcriptional expression of the antiapoptotic gene B-cell lymphoma-2 (*bcl2*) in response to ER stress (McCullough et al. 2001). Another widely accepted mechanism of *chop*-induced apoptosis is the induced overexpression of proapoptotic gene BCL2-associated X (*bax*), which may function as an executioner in ER stress-mediated apoptosis (Zou et al. 2013). *chop*-mediated apoptosis enhanced by ER stress plays a significant role in the pathophysiology of many mammalian diseases (Gopalan et al. 2013; Oyadomari et al. 2002; Lindholm et al. 2006; Thorp et al. 2009; Gotoh et al. 2010; Lee et al. 2012), affecting the survival of organisms. So far, many studies on functions of *chop* have been carried out in mammals. However, no information on the cloning and roles of *chop* in aquaculture animals is available.

The water environment changes are easy to cause ER stress responses in aquaculture animals (Chen et al. 2015; Zhu et al. 2013; Ji 2014), which have many adverse effects (Cripps and Bergheim 2000), including poor growth, greater incidence of disease, increased mortality and low production (Barton and Iwama 1991). At present, due to the impact of the 2019-nCoV epidemic, hydrogen peroxide (H₂O₂) has been widely used as an environmental disinfectant (https://www.sohu.com/a/375631984_120113054). Part of the H₂O₂ residual might enter the aquaculture environment, resulting in changes of environment parameters. In order to understand the impact of H₂O₂ on aquaculture animals, loaches (*Misgurnus anguillicaudatus*) are used in this study. The loach, belonging to the family Cobitidae, is one of the most commercially momentous cultured species in several Eastern Asian countries including Korea, Japan and China. In this study, we first cloned the *chop* gene in loaches, and then monitored its expression profiles in different tissues and early life stages. In addition, a H₂O₂ stress experiment was performed to study the stress response in aquaculture fish in order to explore the role of *chop*. The results will further our understanding of the H₂O₂ stress responses and the role of *chop* in aquaculture animals.

2. Materials And Methods

2.1. Fish and sample collections

Loaches were obtained from Baishazhou fish market (Wuhan, China). All loaches were subjected to a ploidy analyzer (Partec, Germany) to determine their ploidy level. Diploid loaches were used and divided into four groups. The first group (15g ± 0.2) was used for cloning *chop*. Three female and three male adult loaches of the second group (15g ± 0.2) were used for determining expression levels of *chop* in ten tissues including liver, gill, brain, spleen, heart, intestine, muscle, gonad, kidney and skin, and each tissue served as an experimental unit (n = 3). A pair of parents from the third group (20g ± 0.2) was used for reproduction and embryos were collected at different early life stages, including oosperm, 2-cell stage, 8-cell stage, 32-cell stage, blastula stage, gastrula stage, neurula stage, tail-bud forming stage, muscle effect stage, heart-beating stage, and hatching stage. All samples were frozen at -80°C prior to RNA extraction. The fourth group (5g ± 0.2) contained 270 individuals, was used for H₂O₂ stress experiment.

2.2. Cloning the full-length cDNA of *chop* in loach

Total RNA was isolated from the liver tissues by using RNA isoPlus (TaKaRa, Japan). Quantities and qualities of isolated RNAs were measured by electrophoresis and spectrophotometry (Nanodrop 2000, Thermo Scientific, USA). Following the method of the SMART™ RACE cDNA amplification kit (Clontech, USA), the RNA obtained was completely reverse-transcribed into cDNA, which was next used to clone the gene.

A pair of universal amplified primers was designed (Table S1), depending on the multiple sequences alignments of numerous species including zebrafish (*Barchydanio rerio var*), common carp (*Cyprinus carpio*), roughskin sculpin (*Trachidermus fasciatus*), yellow catfish (*Pseudobagrus fulvidraco*), etc. The PCR program was set as follow: initial denaturing for 5 min at 95°C, 35 cycles of 30 s at 95°C, 30 s at 55°C and 18 s at 72°C, and extra elongation for 5 min at 72°C. The PCR products were collected with 2% agarose (Sangon, China) and then purified with a TaKaRa Agarose Gel DNA Purification Kit Ver.2.0 (TaKaRa). After purification, the DNA fragments were ligated into PMD19 (TaKaRa) and the randomly selected positive transformants were opted and sequenced (Invitrogen, China).

2.3. Sequence and phylogenetic analysis

Similarity analysis of all the sequences was conducted by BLAST program at the National Center of Biotechnology Information (NCBI) (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>). The sequences of loach *chop* were translated into amino acid sequences by ORF Finder (<http://www.ncbi.nlm.nih.gov/projects/gorf/>). Multiple alignments of the deduced amino acid sequences were conducted by BioEdit software. Other vertebrate *chop* amino acid sequences for alignments and constructing phylogenetic tree were obtained from NCBI and the identities of these sequences were blasted by Blastp (NCBI). The phylogenetic tree of amino acid sequence of *chop* was constructed by MEGA6.0 program using the neighbor-joining method (NJ, bootstrap method: 1000 replications, Arizona State University, USA). The evolutionary distance between loach *chop* and other *chop* sequences was calculated by using p-distance and gaps were removed by pairwise detection, using default parameters.

2.4. H₂O₂ stress experiment

270 loaches were stocked in nine 30 L white tanks (10 L water volume and 25 ± 0.5°C water temperature) at a stocking density of 30 fishes/tank. Loaches were respectively treated with 0 µM, 50 µM and 100 µM H₂O₂ with triplicate for 24 h. During the experiment, there was no feeding and human interference to prevent introducing other stressors. The number of dead individuals in each tank was recorded every 6 hours. The data were input into GraphPad Prism 6.0 and the survival rate curves were generated.

Gill, liver and spleen tissues were collected from three loaches per tank at 6-, 12-, 18-, and 24-hour H₂O₂ stress exposure. All tissues were frozen in liquid nitrogen and stored at -80°C for determining catalase (CAT), glutathione peroxidase (GPX) and superoxide dismutase (SOD) activities. In addition, gills were also used for detecting expression levels of *chop* and its related genes (*atf4*, *bcl2* and *bax*).

2.5. Real-time quantitative PCR (qPCR)

qPCR was performed using a Mini opticon real-time detector (BIO-RAD, Hercules, CA USA). Specific primers of the target genes and reference genes (*β-actin* and *gapdh* (glyceraldehyde-3-phosphate dehydrogenase)) are summarized in Table S1. The relative expressions of the target genes were calculated with the comparative Ct method ($2^{-\Delta\Delta Ct}$). All the procedures were based on the methods from our laboratory described by Cui et al. (2018).

2.6. Determination of antioxidant enzyme activities

The activities of SOD, GPX and CAT were estimated by using the commercially available kits (Nanjing Jiancheng Bioengineering Institute, China). The examinations of the three antioxidant enzyme activities were carried out according to the manuals of the kits.

2.7. Statistical analysis

The data were expressed as the means $\pm$ SD (standard deviation) or SE (standard error). One-way analysis of variance (ANOVA) was conducted by using Tukey's *post hoc* test in SPSS statistical package version 25.0 (SPSS Inc., USA). $P < 0.05$ was significantly different (* $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$).

3. Results

3.1. Gene cloning of *chop* and its sequence analysis in loach

The full-length of loach *chop* cDNA was 2533bp, containing 801bp open reading frame (ORF) encoding 266 amino acids, a 217bp 5'-untranslated region (UTR) and a 1519bp 3'-untranslated region.

Deduced amino acid sequences of loach *chop* shared high similarity with those of many other species. The putative results indicated that several completely conserved residues might be functionally important (Fig. 1a). The resulting phylogenetic tree of *chop* amino acid sequences demonstrated that the *chop* in loach was grouped into distinct clade similar to higher vertebrate species, and the deduced amino acid sequences of *chop* in loach displayed high similarity to those in *Danio rerio* and *C. carpio* (Fig. 1b).

3.2. Spatial and temporal expression profiles of *chop* in loach

The expression levels of loach *chop* in different tissues were detected (Fig. 2a). The highest expression of *chop* was found in gill, the second-highest in spleen and the third-highest in gonad. The expression levels of loach *chop* in different early life stages were observed as well (Fig. 2b). The expression level at 8-cell stage was the highest, then in sequence were muscle effect stage and blastula stage ($P < 0.05$). There were no significant differences in *chop* expression level among oosperm, 2 cell-, 32 cell-, gastrula-, neurula-, tail-bud forming-, heart-beating-, and hatching-stage.

3.3. H₂O₂ stress experiment

3.3.1. Survival

At 12-hour of H₂O₂ stress, the loaches in the 100 µM group began to die. The percentages of death at 12-, 18- and 24-hour of the 100 µM group were respectively 10%, 25%, and 15%. After 24 hour H₂O₂ stress, half of the loaches from the 100 µM group were dead, while all loaches from the other two groups (0 µM and 50 µM) were alive (Fig.S1).

3.3.2. The antioxidant enzyme activities

As shown in Fig. 3a, at each time point of H₂O₂ stress, gill CAT activities of loaches from the 100 µM group were significantly lower than those from the 0 µM and 50 µM groups. Between 12 and 24 hours of H₂O₂ stress, loaches from the 50 µM and 100 µM groups presented significantly lower hepatic CAT activities, compared with the control group (Fig. 3b). Figure 3c showed that spleen CAT activities in the 100 µM group were the lowest. In addition, the trends of SOD and GPX activities were similar to the CAT activity trend (Fig. 3d-3i).

3.3.3. Expression levels of *chop* and its related genes in gill

Expression levels of *chop* and its related genes *atf4*, *bcl2* and *bax* in gills of loaches from the 0 µM, 50 µM and 100 µM groups were analyzed (Fig. 4). The expression levels of *chop* from the 50 µM and 100 µM groups were significantly higher than those from the 0 µM group at 12-, 18- and 24- hour of H₂O₂ stress (Fig. 4a). The expression level of *chop* from the 100 µM group reached the highest at 24-hour of H₂O₂ stress, 10 times of that compared to the 0 µM group.

There were no significant differences in the expression levels of *atf4* and *bax* (two proapoptotic genes) among the three groups (0 µM, 50 µM and 100 µM) at the 6-hour and 12-hour of H₂O₂ stress (Fig. 4b,4d). Similar to *chop*, the expression levels of *atf4* and *bax* from the 100 µM group were the highest at the 24-hour of H₂O₂ stress. However, Fig. 4c showed that the expression profile of *bcl2* was opposite to those of *chop*, *atf4* and *bax*.

4. Discussion

This study cloned *chop* gene from an aquaculture animal (namely *M. anguillicaudatus*) for the first time. The multiple alignments and phylogenetic tree of the deduced amino acid sequences of *chop* genes showed that the loach *chop* kept relatively conservative sequences and structural homology compared to other vertebrates, suggesting *chop* functions are stable during vertebrate evolution (Luethy et al. 1990; Lee et al. 2011). The expression profiles of loach *chop* were determined here as well. The results showed that loach *chop* was highly expressed during certain embryogenesis periods, consistent with *Xenopus* (Iijima et al. 2003). In addition, the tissue expression analysis showed that *chop* predominantly expressed

in the gill and spleen of loach. It has been previously reported that gills and spleens are parts of the immune system, responding to the organism stress (Dautremepuits et al. 2009; Kocabas et al. 2002). The elevated loach *chop* expression in gills and spleens may indicate its immunological functions related to stress response.

At present, due to the impact of the 2019-nCoV epidemic, part of the H₂O₂ residual might enter the aquaculture environment. In order to explore the effects of H₂O₂ stress in fish, we treated the loaches with different concentrations of H₂O₂. Our results confirmed that activities of antioxidant enzymes in the loaches had a highly negative correlation with the prolongation of H₂O₂ stress and the increase of H₂O₂ concentrations, which were similar to the results of other stress studies. For example, when oxidative stress occurred, the activities of SOD and CAT in gills of zebrafish significantly dropped (Jin et al. 2015). Analogous results were also found in mice liver and spleen (Meng et al. 2004; Matsumoto et al. 2002). In addition, oxidative stress reduced GPX activities in gills and livers in black porgy (Chen et al. 2009), as well as in spleens of zebrafish (Komoike and Matsuoka 2013). In summary, H₂O₂ appears to induce stress responses in the loach due to decreased antioxidant enzymes expressions.

In the H₂O₂ stress experiment, we analyzed the expression levels of *chop* and its related genes *atf4*, *bcl2* and *bax* in gills of the loach. Our study evidenced that the expression level of *atf4* in the loach surged under high concentration H₂O₂ stress. The expression of loach *chop* significantly up-regulated with the increase of concentrations and the prolongation of H₂O₂ stress. Correspondingly, the more severe the inhibition of *bcl2* expression the more the acceleration of *bax* expression. These results were consistent with previous studies. When the apoptotic pathway is turned on, under the regulation of a series of upstream factors, the mRNA transcription of *atf4* increased (Cripps and Bergheim 2000). Afterwards, expression of its key downstream transcriptional target *chop* elevated (Barton and Iwama 1991), resulting in the repression of the antiapoptotic gene *bcl2* and the overexpression of proapoptotic gene *bax* (Luethy et al. 1990). A reasonable conclusion is that *chop* may promote apoptosis in the loach under H₂O₂ stress. However, the specific functions of *chop* warrants further studies.

In conclusion, we identified and characterized full-length cDNAs of loach *chop* for the first time. Expression profiles of *chop* in different tissues and different early life stages suggested its potential immunological functions in loach. The variety of *chop* expression levels in loaches treated with various concentrations of H₂O₂ indicated that loach *chop* expression was sensitive to H₂O₂ stress and that the stress induced apoptosis in loach. Finally, this study provides critical information for further investigation of *chop* functions in fish under H₂O₂ stress.

Declarations

Funding

This study was financially supported by National and Local Joint Engineering Laboratory for Freshwater fish Breeding Research Fund (NO. HSY201913K1-1) and Jingmen Science and Technology Project

(Project Number: 2019YFZD024).

Acknowledgements

Special thanks to Dr Wang from University of California Riverside for revising the English writing of this manuscript.

Conflicts of interest

The authors declare that they have no conflict of interest.

Ethical approval

This study was conducted in strict accordance with the recommendations in the Guide for the Care and Use of Laboratory Animals of Huazhong Agricultural University. All efforts were made to minimize suffering of the loaches.

Availability of data and material

Not applicable.

Code availability

Not applicable.

Authors' contributions

Xiaojuan Cao and Zhiying Jia designed the experiment. Hui Li, Minxin Kang, Shouxiang Sun and Jian Gao carried out the experiments. Hui Li and Minxin Kang analyzed the data. The article was written by Hui Li with input from all authors. Xiaojuan Cao contributed to the revision of the manuscript. All authors reviewed and approved the paper.

References

1. Li YM, Guo YS, Tang J, Jiang JL, Chen ZN (2014) New insights into the roles of CHOP-induced apoptosis in ER stress. *Acta Biochim Biophys Sin* 46:629–640. <https://doi.org/10.1093/abbs/gmu048>
2. Ariyama Y, Tanaka Y, Shimizu H, Shimomura K, Okada S, Saito T, Yamada E, Oyadomari S, Mori M (2008) The role of CHOP messenger RNA expression in the link between oxidative stress and apoptosis. *Metabolism* 57:1625–1635. <https://doi.org/10.1016/j.metabol.2008.06.019>
3. Zhang Z, Yang XY, Cohen DM (1999) Urea-associated oxidative stress and Gadd153/CHOP induction. *Am J Physiol-Renal Physiol* 276:F786–F793. <https://doi.org/10.1152/ajprenal.1999.276.5.F786>

4. Cheng WP, Wang BW, Shyu KG (2009) Regulation of GADD153 induced by mechanical stress in cardiomyocytes. *Eur J Clin Invest* 39:960–971. <https://doi.org/10.1111/j.1365-2362.2009.02193.x>
5. Pirot P, Ortis F, Cnop M, Ma YJ, Hendershot LM, Eizirik DL, Cardozo AK (2007) Transcriptional regulation of the endoplasmic reticulum stress gene chop in pancreatic insulin-producing cells. *Diabetes* 56:1069–1077. <https://doi.org/10.2337/db06-1253>
6. Walter P, Ron D (2011) The unfolded protein response: from stress pathway to homeostatic regulation. *Science* 334:1081–1086. <https://doi.org/10.1126/science.1209038>
7. Zhang LR, Zhang CW, Wang A (2016) Divergence and conservation of the major UPR branch IRE1-bZIP signaling pathway across eukaryotes. *Sci Rep* 6:27362–27376. <https://doi.org/10.1038/srep27362>
8. Deegan S, Koryga I, Glynn SA, Gupta S, Gorman AM, Samali A (2015) A close connection between the PERK and IRE arms of the UPR and the transcriptional regulation of autophagy. *Biochem Biophys Res Commun* 456:305–311. <https://doi.org/10.1016/j.bbrc.2014.11.076>
9. Shang J, Lehrman MA (2004) Discordance of UPR signaling by ATF6 and Ire1p-XBP1 with levels of target transcripts. *Biochem Biophys Res Commun* 317:390–396. <https://doi.org/10.1016/j.bbrc.2004.03.058>
10. Rutkowski DT, Arnold SM, Miller CN, Wu J, Li J, Gunnison KM, Mori K, Sadighi Akha AA, Raden D, Kaufman RJ (2006) Adaptation to ER stress is mediated by differential stabilities of pro-survival and pro-apoptotic mRNAs and proteins. *PLoS Biol* 4(11):e374. <https://doi.org/10.1371/journal.pbio.0040374>
11. Cao J, Dai DL, Yao L, Yu HH, Ning B, Zhang Q, Chen J, Cheng WH, Shen W, Yang ZX (2012) Saturated fatty acid induction of endoplasmic reticulum stress and apoptosis in human liver cells via the PERK/ATF4/CHOP signaling pathway. *Mol Cell Biochem* 364:115–129. <https://doi.org/10.1007/s11010-011-1211-9>
12. Mounir Z, Krishnamoorthy JL, Wang S, Papadopoulou B, Campbell S, Muller WJ, Hatzoglou M, Koromilas AE (2011) Akt determines cell fate through inhibition of the PERK-eIF2 α phosphorylation pathway. *Sci Signal* 4:ra62. <https://doi.org/10.1126/scisignal.2001630>
13. B'chir W, Maurin AC, Carraro V, Averous J, Jousse C, Muranishi YK, Parry L, Stepien G, Fafournoux P, Bruhat A (2013) The eIF2 α /ATF4 pathway is essential for stress-induced autophagy gene expression. *Nucleic Acids Res* 41:7683–7699. <https://doi.org/10.1093/nar/gkt563>
14. Su N, Kilberg MS (2008) C/EBP homology protein (CHOP) interacts with activating transcription factor 4 (ATF4) and negatively regulates the stress-dependent induction of the asparagine synthetase gene. *J Biol Chem* 283:35106–35117. <https://doi.org/10.1074/jbc.M806874200>
15. Mccullough KD, Martindale JL, Klotz LO, Aw TY, Holbrook NJ (2001) Gadd153 sensitizes cells to endoplasmic reticulum stress by down-regulating Bcl2 and perturbing the cellular redox state. *Mol Cell Biol* 21:1249–1259. <https://doi.org/10.1128/MCB.21.4.1249-1259.2001>
16. Zou XJ, Yang L, Yao SL (2012) Endoplasmic reticulum stress and C/EBP homologous protein-induced Bax translocation are involved in angiotensin II-induced apoptosis in cultured neonatal rat

- cardiomyocytes. *Exp Biol Med* 237:1341–1349. <https://doi.org/10.1258/ebm.2012.012041>
17. Gopalan A, Yu W, Sanders BG, Kline K (2013) Simvastatin inhibition of mevalonate pathway induces apoptosis in human breast cancer cells via activation of JNK/CHOP/DR5 signaling pathway. *Cancer Lett* 329:9–16. <https://doi.org/10.1016/j.canlet.2012.08.031>
 18. Oyadomari S, Koizumi A, Takeda K, Gotoh T, Akira S, Araki E, Mori M (2002) Targeted disruption of the Chop gene delays endoplasmic reticulum stress-mediated diabetes. *J Clin Investig* 109:525–532. <https://doi.org/10.1172/JCI14550>
 19. Lindholm D, Wootz H, Korhonen L (2006) ER stress and neurodegenerative diseases. *Cell Death Differ* 13:385–392. <https://doi.org/10.1038/sj.cdd.4401778>
 20. Thorp E, Li G, Seimon TA, Kuriakose G, Ron D, Tabas I (2009) Reduced apoptosis and plaque necrosis in advanced atherosclerotic lesions of Apoe^{-/-} and Ldlr^{-/-} mice lacking CHOP. *Cell Metab* 9:474–481. <https://doi.org/10.1016/j.cmet.2009.03.003>
 21. Gotoh T, Endo M, Oike Y (2010) Endoplasmic reticulum stress-induced transcriptional factor CHOP and cardiovascular diseases. *Curr Hypertens Rev* 6:55–65. <https://doi.org/10.2174/157340210790231717>
 22. Lee WK, Chakraborty PK, Roussa E, Wolff NA, Thévenod F (2012) ERK1/2-dependent bestrophin-3 expression prevents ER-stress-induced cell death in renal epithelial cells by reducing CHOP. *Biochimica et Biophysica Acta (BBA)-Molecular Cell Res* 1823:1864–1876. <https://doi.org/10.1016/j.bbamcr.2012.06.003>
 23. Chen CY, Zhang SL, Liu ZY, Tian Y, Sun Q (2015) Cadmium toxicity induces ER stress and apoptosis via impairing energy homeostasis in cardiomyocytes. *Bioscience reports* 35:e00214. <https://doi.org/10.1042/BSR20140170>
 24. Zhu GG, Zheng YC, Zhang LL, Shi YY, Li WH, Liu ZC, Peng B, Yin J, LiuWH, He XH (2013) Cocksackievirus A16 infection triggers apoptosis in RD cells by inducing ER stress. *Biochem Biophys Res Commun* 441:856–861. <https://doi.org/10.1016/j.bbrc.2013.10.142>
 25. Ji C (2014) New insights into the pathogenesis of alcohol-induced ER stress and liver diseases. *International journal of hepatology*. <https://doi.org/10.1155/2014/513787>
 26. Cripps SJ, Bergheim A (2000) Solids management and removal for intensive land-based aquaculture production systems. *Aquac Eng* 22:33–56. [https://doi.org/10.1016/S0144-8609\(00\)00031-5](https://doi.org/10.1016/S0144-8609(00)00031-5)
 27. Barton BA, Iwama GK (1991) Physiological changes in fish from stress in aquaculture with emphasis on the response and effects of corticosteroids. *Annual Review of fish diseases* 1:3–26. [https://doi.org/10.1016/0959-8030\(91\)90019-G](https://doi.org/10.1016/0959-8030(91)90019-G)
 28. Sousa DPF, Braga BB, Gondim FA, Gomes-Filho E, Martins K, Brito POBD (2016) Increased drought tolerance in maize plants induced by H₂O₂ is closely related to an enhanced enzymatic antioxidant system and higher soluble protein and organic solutes contents. *Theoretical Experimental Plant Physiology* 28:297–306. <https://doi.org/10.1007/s40626-016-0069-3>
 29. Taiebeh Z, Mohammad JA (2016) Hydrogen peroxide affects plant growth promoting effects of Azospirillum. *Journal of Crop Science Biotechnology* 19:167–175. <https://doi.org/10.1007/s12892->

30. Cui Y, Liang X, Cao XJ, Gao J (2018) Molecular characterization of peroxisome proliferator activated receptor gamma (PPAR γ) in loach *Misgurnus anguillicaudatus* and its potential roles in fatty acid metabolism in vitro. *Process Biochem* 66:205–211. <https://doi.org/10.1016/j.procbio.2018.01.008>
31. Luethy JD, Fargnoli J, Park JS, Fornace AJ, Holbrook NJ (1990) Isolation and characterization of the hamster gadd153 gene. Activation of promoter activity by agents that damage DNA. *J Biol Chem* 265:16521–16526. <https://doi.org/10.1021/bi00490a029>
32. Lee HC, Chen YJ, Liu YW, Lin KY, Chen SW, Lin CY, Lu YC, Hsu PC, Lee SC, Tsai HJ (2011) Transgenic zebrafish model to study translational control mediated by upstream open reading frame of human chop gene. *Nucleic Acids Res* 39:e139. <https://doi.org/10.1093/nar/gkr645>
33. Iijima R, Homma KJ, Natori S (2003) Participation of proteasomes in *Xenopus* embryogenesis. *J Biochem* 134:467–471. <https://doi.org/10.1093/jb/mvg165>
34. Dautremepuits C, Marcogliese DJ, Gendron AD, Fourniera M (2009) Gill and head kidney antioxidant processes and innate immune system responses of yellow perch (*Perca flavescens*) exposed to different contaminants in the St. Lawrence River, Canada, *Sci. Total Environ* 407:1055–1064. <https://doi.org/10.1016/j.scitotenv.2008.10.004>
35. Kocabas AM, Li P, Cao DF, Karsi A, He CB, Patterson A, Ju ZL, Dunham RA, Liu ZJ (2002) Expression profile of the channel catfish spleen: analysis of genes involved in immune functions. *Mar Biotechnol* 4:526–536. <https://doi.org/10.1007/s10126-002-0067-0>
36. Jin YX, Liu ZZ, Liu F, Ye Y, Peng T, Fu ZW (2015) Embryonic exposure to cadmium (II) and chromium (VI) induce behavioral alterations, oxidative stress and immunotoxicity in zebrafish (*Danio rerio*). *Neurotoxicol Teratol* 48:9–17. <https://doi.org/10.1016/j.ntt.2015.01.002>
37. Meng ZQ, Geng H, Qin G (2004) Oxidative stress on spleens of mice exposed to sulfur dioxide. *Journal of Health Toxicology* 18:1–3. [https://doi.org/10.1016/S0898-6568\(03\)00096-2](https://doi.org/10.1016/S0898-6568(03)00096-2)
38. Matsumoto K, Ui I, Mochida T, Sakamoto R, Endo K (2002) Relation of bile hydrogen peroxide level and liver super oxide dismutase activity in selenium-deficient rats. *Yakugaku Zasshi-J Pharm Soc Jpn* 122:407–411. <https://doi.org/10.1248/yakushi.122.407>
39. Chen HG, Ma SW, Lin Q, Gan JL, Cai WG, Jia XP (2009) Effects of tributyltin chloride (TBTCl) on SOD activities, MDA contents and GPx activities in gill and liver of the black porgy (*Sparus macrocephalus*). *South China Fisheries Scienc* 5:23–27. <https://doi.org/10.3969/j.issn.1673-2227.2009.02.004>
40. Komoike Y, Matsuoka M (2013) Exposure to tributyltin induces endoplasmic reticulum stress and the unfolded protein response in zebrafish. *Aquat Toxicol* 142–143:221–229. <https://doi.org/10.1016/j.aquatox.2013.08.017>

Figures

Figure 1

Alignment and phylogenetic tree of the deduced amino acid sequences of chop in *Misgurnus anguillicaudatus* and other species. (a) Alignment of the deduced amino acid sequences of chop. The same or similar amino acids are respectively highlighted by black and gray. Black is used to labeled amino acids with a similarity more than 75% and light gray is to those with a similarity merely more than 50%. (b) Phylogenetic tree of the deduced amino acid sequences of chop. The horizontal branch length is proportional to the amino acid substitution rate at each position (The scale is 0.2). The genbank accession numbers for animals involved in the figure, except *Misgurnus anguillicaudatus*, are set out as follows: *Homo sapiens* (AAB22646.1), *Mus musculus* (BAE20435.1), *Cyprinus carpio* (KTF75003.1), *Danio rerio* (XP_005166228.1), *Larimichthys crocea* (XP_010730414.2), *Lates calcarifer* (XP_018534443.1), *Nothobranchius furzeri* (XP_015814628.1), *Salmo salar* (XP_014021417.1), *Hippocampus comes* (XP_019716650.1), *Bos taurus* (NP_001071631.1), *Gallus gallus* (AJA72779.1), *Oryzias latipes* (BAL14286.1), *Rattus norvegicus* (NP_001103456.1), *Sus scrofa* (NP_001138317.1), *Xenopus laevis* (NP_001082635.1).

Figure 2

Expression levels of chop in ten different tissues (namely, liver, gill, brain, spleen, heart, intestine, muscle, gonad, kidney and skin) (a) and in eleven different early life stages (namely, oosperm, 2 cell stage, 8 cell stage, 32 cell stage, blastula stage, gsatrula stage, neurula stage, tail-bud forming stage, muscle effect stage, heart-beating stage and hatching stage) (b) of *Misgurnus anguillicaudatus*, determined by real-time quantitative PCR. Significant differences are represented by different small letters.

Figure 3

Determinations of three antioxidant enzyme activities in *Misgurnus anguillicaudatus* under three different concentrations (0μM, 50μM and 100μM) of H₂O₂ stress. Catalase (CAT) activities in gills (a), livers (b) and spleens (c) of the loaches at the 6-, 12-, 18- and 24-hour of H₂O₂ stress; Glutathione peroxidase (GPX) activities in gills (d), livers (e) and spleens (f) of the loaches at the 6-, 12-, 18- and 24-hour of H₂O₂ stress; Superoxide dismutase (SOD) activities in gills (g), livers (h) and spleens (i) of the loaches at the 6-, 12-, 18- and 24-hour of H₂O₂ stress. Significant differences for each antioxidant enzyme activity at each time point among the three different stress groups are represented by different small letters.

Figure 4

Expression levels of chop (a) and its related genes (atf4 (b), bcl2 (c) and bax (d)) in gills of *Misgurnus anguillicaudatus* under three different concentrations (0μM, 50μM and 100μM) of H₂O₂ stress. The values are expressed as the means ± SEs. *P < 0.05, **P < 0.01, ***P < 0.001. chop, C/EBP [CCAAT/enhancer-binding protein]-homologous protein gene; atf4, activation of transcription factor-4; bcl2, B-cell lymphoma-2; bax, BCL2-associated X.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [5supplementaltableandfigure.doc](#)