Bioinformatics analysis of BUB1 expression and gene regulation network in lung adenocarcinoma

Luyao Wang (✉ 1318908938@qq.com)

Qingdao University

Xue Yang

Qingdao University

Ning An

Qingdao University

Jia Liu

Qingdao University

Research article

Keywords: bioinformatics analysis; BUB1; lung adenocarcinoma

Posted Date: July 9th, 2019

DOI: <https://doi.org/10.21203/rs.2.11120/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Translational Cancer Research on August 1st, 2020. See the published version at <https://doi.org/10.21037/tcr-20-1045>.

Abstract

Lung adenocarcinoma is the most common type of lung cancer with high morbidity and mortality. Potential mechanisms and therapeutic targets of lung adenocarcinoma need further study. BUB1 (BUB1 mitotic checkpoint serine/threonine kinase) encodes a serine/threonine protein kinase which is critical in the mitosis. It is associated with poor prognosis in multiple cancer types. Oncomine database was used to determine the differential expression of BUB1 in normal and lung adenocarcinoma tissues, while UALCAN was used to perform analysis of the relative expression and survival of BUB1 between tumor and normal tissues in different tumor subgroups. We used the cBioPortal for Cancer Genomics to perform GO analysis and KEGG analysis of the top 50 altered neighbor genes of BUB1. The LinkedOmics database was used to determine differential gene expression with BUB1 and to perform functional analysis. The kinase, miRNA and transcription factor target networks correlated with BUB1 were also analysed by LinkedOmics database. The results revealed that BUB1 was highly expressed in lung adenocarcinoma patients. BUB1 involved multiple tumor-related pathways, such as cell cycle, oocyte meiosis and p53 signaling pathway. BUB1 is associated with tumor-associated kinases, microRNAs and transcription factors. Our study reveal BUB1 expression and potential gene regulation networks in lung adenocarcinoma based on bioinformatics analysis, guiding further study on the role and regulation of BUB1 in lung adenocarcinoma.

Background

Lung cancer is the most commonly diagnosed cancer and the most common mortality cause of cancer worldwide [1]. Lung cancer patients are usually in advanced stage at the time of initial diagnosis and their 5-year net survival is low (10-20%) [2]. Lung adenocarcinoma (LUAD), which is the most common form of lung cancer, is a multifactorial disease. Although significant progress has been made in the treatment of lung adenocarcinoma in recent years [3,4], the survival rate is still very low. The pathogenesis of lung adenocarcinoma is complex and may involve the interaction of multiple genes in multiple processes. As data mining has provided a new method for finding tumor biomarkers and therapeutic targets, it may be also possible to identify novel therapeutic targets for lung adenocarcinoma by establishing gene networks associated with tumorigenesis and progression.

BUB1, whose official full name is BUB1 mitotic checkpoint serine/threonine kinase, encodes a serine/threonine protein kinase which is critical in the process of mitosis [5]. The kinase BUB1 is essential in the spindle assembly checkpoint (SAC) and chromosome congression [6]. BUB1 may also plays a role in DNA damage response. Studies have shown that BUB1 is associated with the maintenance of cancer stem cells [7]. Also, BUB1 may be a promising prognostic biomarker in cancer as high expression of BUB1 has been proved to be related to poor prognosis in multiple cancer types [8,9,10].

Our previous analysis of lung adenocarcinoma data in the Gene Expression Omnibus database revealed that BUB1 may be a key protein-coding gene in lung adenocarcinoma. Our studies in patients with lung adenocarcinoma have shown that BUB1 is overexpressed in tumor tissues than in adjacent normal tissues. Survival analysis has shown that overexpression of BUB1 was associated with poor prognosis of lung adenocarcinoma patients. Therefore, BUB1 may be a new proto-oncogene closely related to lung adenocarcinoma. Thus, we further studied the expression and mutations of BUB1 in lung adenocarcinoma patients in a variety of public databases. We also analyzed the genomic changes and functional networks associated with BUB1 in lung adenocarcinoma using multidimensional analysis methods. Our study may lead to novel targets and novel approaches for diagnosis and treatment of lung adenocarcinoma.

Methods

Oncomine analysis

The expression of BUB1 between normal tissues and lung adenocarcinoma tissues was analyzed by Oncomine database (www.oncomine.org), which is the biggest oncogene chip database in the world [11]. This analysis involved Garber Lung, Hou Lung, Landi Lung and Su Lung. In lung adenocarcinoma tissues, BUB1 expression was assessed in LUAD tissues relative to its expression in normal tissues. $P \leq 0.01$ was considered significant. Oncomine database was also used to analyse survival of LUAD patients in various subgroups. $P \leq 0.01$ was considered significant.

UALCAN analysis

UALCAN (<http://ualcan.path.uab.edu>), an open web-portal for tumor subgroup gene expression and survival analyses, contains data from The Cancer Genome Atlas (TCGA) project [12]. UALCAN was used to perform analysis of the relative expression of BUB1 between tumor and normal tissues in different tumor subgroups based on gender, age, race, cancer stage and smoking habits. UALCAN was also used to analyse survival of LUAD patients in various subgroups.

c-BioPortal analysis

The cBioPortal for Cancer Genomics, which contains 225 cancer studies, provides resource for exploration and analysis of multidimensional cancer genomics data [13,14]. The cBioPortal was used to

analyse type and frequency of RBM8A alterations in LUAD. The OncoPrint shows an overview of genetic alterations each sample in BUB1. The cBioPortal is publicly available at <http://cbiportal.org>. STRING (<http://string-db.org>), a online tool for the retrieval of interacting genes and proteins, was used to establish PPI network of the top 50 altered neighbor genes of BUB1. We then performed GO analysis and KEGG analysis of these 50 genes using database for annotation, visualization and integrated discovery (DAVID; <http://david.ncifcrf.gov>), which is a free and open-access website for researchers to analyse the biological significance behind genes[15]. $P < 0.001$ was considered significant.

LinkedOmics analysis

The LinkedOmics database is a multi-omics database that contains data of 32 cancer types and 11158 patients from TCGA project [16]. We used LinkCompare module of LinkedOmics to analyse genes differentially expressed in correlation with BUB1. Pearson's correlation coefficient was used to perform statistical analysis. The tab LinkInterpreter of LinkedOmics performs enrichment analysis of differentially expressed genes using Gene Set Enrichment Analysis (GSEA).

GeneMANIA analysis

GeneMANIA (<http://www.genemania.org>) is an open-access web interface which contains hundreds of data sets from several databases [17]. GeneMANIA was used to construct protein-protein interaction (PPI) network of genes that GSEA identified as being enriched in LUAD: kinase ATR and mi-RNA 143.

Results

The gene expression level of BUB1 was analyzed by Oncomine database. The results of Oncomine analysis revealed that the expression and DNA copy number variation (CNV) of BUB1 in LUAD tissues were significantly higher than that in normal tissues ($P < 0.01$). Moreover, the fold differences were more than 2 and BUB1 ranked within the top 4% at the level of mRNA expression (Figure 1). The results of UALCAN analysis showed that BUB1 transcription level in lung adenocarcinoma patients was significantly higher than that in healthy people in subgroup analysis based on gender, age, race, cancer stage and smoking habits ($P < 0.001$) (Figure 2). Therefore, the expression of BUB1 may be a potential diagnostic indicator of LUAD.

Survival analysis of LUAD patients based on various subgroups

Survival analysis based on UALCAN showed that the expression of BUB1 and patients' race may have an impact on LUAD patients survival ($P < 0.05$) (Figure 3).

BUB1 alterations in LUAD

Then we used cBioPortal to determine the type and frequency of RBM8A alterations in LUAD according to the sequencing data of patients in TCGA database. BUB1 was altered in 59 (12%) of LUAD patients (Figure 4). The alteration frequency was shown in Figure 5. The most frequent alteration is mRNA upregulation.

Biological interaction network of BUB1 alterations in LUAD

The PPI network of the top 50 altered neighbor genes of BUB1 established by STRING online database was shown in Figure 6A. The GO analysis based on DAVID database showed that these 50 genes were enriched in cancer-associated biological pathways including protein binding, microtubule binding, ATP binding, microtubule motor activity, as well as protein serine/threonine kinase activity. For cellular component, these genes were enriched in condensed chromosome kinetochore, chromosome, centromeric region, cytosol, spindle and kinetochore (Figure 6B). GO enrichment analysis in the category biological process showed that these genes were enriched in cell division, mitotic nuclear division, sister chromatid cohesion, chromosome segregation, as well as mitotic sister chromatid segregation (Figure 6B). The results of the KEGG analysis showed that these 50 genes were enriched in cell cycle, oocyte meiosis, progesterone-mediated oocyte maturation and p53 signaling pathway (Figure 6C). Above results indicated that biological interaction network of BUB1 alterations is mainly in the chromosome and is mainly associated with the cell cycle.

Co-expression network of BUB1 in LUAD

We used LinkedOmics Functional module to analyze the sequencing mRNA data of patients in the TCGA. Genes differentially expressed in correlation with BUB1 in LUAD was shown in Figure 6. This result

suggests that BUB1 has a wide influence on the transcriptome. GO terms show that genes differentially expressed in correlation with BUB1 mainly concentrated in the condensed chromosome, chromosomal region and spindle (Figure 7A). These genes were enriched in the category biological process, including chromosome segregation, microtubule cytoskeleton organization involved in mitosis and DNA replication (Figure 7B). In addition, these genes were enriched in cancer-associated terms in the category molecular function, including histone deacetylase binding, catalytic activity acting on DNA and ATPase activity (Figure 7C). The results of the KEGG analysis showed that these genes were enriched in biological pathways including cell cycle, oocyte meiosis and DNA replication (Figure 7D, Figure 8A, Figure 8B, Figure 8C, Figure 9).

Kinase, miRNA and transcription factor target networks correlated with BUB1 in LUAD

We then analysed the kinase, miRNA and transcription factor target networks correlated with BUB1 based on GSEA. The miRNA, kinase and transcription factor-target networks of RBM8A in LUAD was shown in Figure 10. The top 5 miRNA-target network was associated with (TCATCTC) MIR-143, (GTAAGAT) MIR-200A, (GTACTGT) MIR-101, (ATGCAGT) MIR-217, (TATTATA) MIR-374 (Supplementary Table 1). The top 5 most significant kinase-target networks were cyclin dependent kinase 1 (CDK1), cyclin dependent kinase 2 (CDK2), ATR serine/threonine kinase (ATR), ATM serine/threonine kinase (ATM) and aurora kinase B (AURKB) (Supplementary Table 2). The transcription factor-target network was significantly related to E2F family, such as V\$E2F1_Q6, V\$E2F_Q6, V\$E2F_Q4, V\$E2F4DP1_01 and V\$E2F1DP1_01 (Supplementary Table 3). PPI network of genes that GSEA identified as being enriched in LUAD (kinase ATR and mi-RNA 143) was shown in Figure 11.

Discussion

BUB1, which is related to kinase function and the spindle assembly checkpoint, has been reported to be high expressed in the process of mitosis and cell proliferation [18]. Our previous study has shown that BUB1 may be a key gene in lung adenocarcinomas. Our previous study has revealed that BUB1 is over-expressed in lung adenocarcinoma tissue compared to normal tissue and high BUB1 expression is associated with poor prognosis of lung adenocarcinoma patients. In order to understand the potential function of BUB1 in lung adenocarcinoma and its regulatory network in more detail, we used multiple databases to conduct bioinformatics analysis of public sequencing data to guide future research on lung adenocarcinoma.

Lung adenocarcinoma is the most common form of lung cancer [19]. Patients with lung cancer are usually in advanced stage at the time of first diagnosis and the survival rate is very low today. New markers may be helpful in the diagnosis and treatment of lung adenocarcinoma. Analysis of sequencing data of lung adenocarcinoma from GEO database and TCGA database showed that BUB1 mRNA level and DNA copy number variation (CNV) in lung adenocarcinoma were significantly higher than that in normal tissues. The fold differences were more than 2 and BUB1 ranked between top 1–4% at the level of mRNA expression. The over-expression of BUB1 in LUAD revealed that BUB1 may be a promising biomarker of LUAD and it is worth further study. Our study also found that the most frequent change in RBM8A in LUAD was mRNA upregulation. Altered BUB1 expression may be critical for the occurrence and progress of lung adenocarcinoma since BUB1 plays an important role in DNA damage response and BUB1 is required for the maintain of cancer stem cells [7]. Alteration of BUB1 may cause changes in downstream signaling pathways. The top 50 altered neighbor genes of BUB1 were enriched in cell cycle, oocyte meiosis, progesterone-mediated oocyte maturation, as well as p53 signaling pathway. The biological interaction network of BUB1 alterations is mostly in the chromosome and is mainly associated with the cell cycle and p53 pathway. This result is consistent with the functions of BUB1 [20-22].

GSEA was used to analyse the co-expressed network of BUB1. The results showed that genes differentially expressed in correlation with BUB1 were enriched in cell cycle, oocyte meiosis and DNA replication. These genes were also enriched in tumor-associated terms in the category molecular function, biological process and cellular component, proving that BUB1 has a wide influence on the transcriptome. Genome instability and mutation are the basic characteristics of cancer cells [23,24]. Kinases and their related signaling pathways contribute to the stabilization and repair of genomic DNA [25,26]. Therefore, we use GSEA to analyse the kinase, miRNA and transcription factor target networks correlated with BUB1. The results revealed that BUB1 in LUAD is associated with tumor-associated kinases, including CDK1, CDK2, ATR, ATM and AURKB. These kinases play a role in the cell cycle, mitosis and genomic stability [27-29].

The target network of transcription factors has significant relationship with E2F family, including V \$E2F1 UQ6, V \$E2F UQ6, V \$E2F UQ4, V \$E2F4DP1 U01 and V \$E2F1DP1 U01. It has been reported that the E2F transcription factor is one of the key regulators of cell cycle progression and triggering apoptosis [30]. Also, alterations of E2F may be associated with shorter survival of cancer patients [31]. Moreover, the E2F1 member of the E2F family is associated with metastasis of cancer cells [32]. Studies have shown that up-regulated transcription of E2F1 may promote the progression of non-small cell lung cancer [33]. Thus, E2F transcription factor family may be a critical target of BUB1, and that BUB1 may regulate cell cycle through this transcription factor.

The significant miRNA-target networks of BUB1 were identified by GSEA. MicroRNAs, endogenous RNAs of about 22 nt, is an abundant class of gene regulators in multicellular organisms [34]. MicroRNAs may affect the expression of many protein-coding genes. Studies have shown that the dysregulation of microRNAs may be the cause of many cancers, and thus miRNA-targeted therapeutics has attracted much attention [35]. In our study, miR-143, miR-200A, miR-101, miR-217 and miR-374 were identified as miRNAs significantly associated with BUB1. MiR-143 has been considered to have potential as tumor markers for non-small cell lung cancer [36], while miR-200A has been indicated to play a role in the progression of lung adenocarcinoma [37]. MiR-101 may exert tumor inhibitory effect in non-small cell lung cancer [38]. MiR-217 has been reported to be highly associated with survival time of lung adenocarcinoma patients [39]. MiR-374 has been indicated to play a role in various cancers and it may be a promising prognostic biomarker and therapeutic target of cancer [40]. The relationship between these miRNAs and BUB1 and their effects on lung adenocarcinoma need further experimental verification. They may provide novel tumor biomarkers and therapeutic targets for lung adenocarcinoma.

Our study used multiple databases which contain a large number of clinical data to study BUB1 expression and gene regulatory network in lung adenocarcinoma. Our study provides evidence for the importance of BUB1 in lung adenocarcinoma and its potential as a marker of lung adenocarcinoma. The results revealed that BUB1 was highly expressed in lung adenocarcinoma patients. The over-expression of BUB1 had a profound impact on genomic stability and involved multiple tumor-related pathways, such as cell cycle, oocyte meiosis and p53 signaling pathway. BUB1 is associated with several tumor-associated kinases (such as CDK1, CDK2, ATR, ATM and AURKB) microRNAs (such as miR-143, miR-200A, miR-101, miR-217 and miR-374) and transcription factors (such as E2F family).

In conclusion, BUB1 may hopefully become a new marker and therapeutic target for lung adenocarcinoma. In future study, we will further validate the regulatory network of BUB1 in lung adenocarcinoma.

Declarations

Competing interests

The authors declare that no competing interests exist.

Funding

References

1. Ferlay J, Soerjomataram I, Dikshit R, Eser S, Mathers C, Rebelo M, Parkin DM, Forman D, Bray F. Cancer incidence and mortality worldwide: sources, methods and major patterns in GLOBOCAN 2012. *Int J Cancer*. 2015 Mar 1;136(5):E359-86. doi: 10.1002/ijc.29210. Epub 2014 Oct 9. PubMed PMID:25220842.
2. Richards TB, Henley SJ, Puckett MC, Weir HK, Huang B, Tucker TC, Allemani C. Lung cancer survival in the United States by race and stage (2001-2009): Findings from the CONCORD-2 study. 2017 Dec 15;123 Suppl 24:5079-5099. doi: 10.1002/cncr.31029. PubMed PMID:29205305;PubMed Central PMCID:PMC 5812722.
3. Reck M, Heigener DF, Mok T, Soria JC, Rabe KF. Management of non-small-cell lung cancer: recent developments. 2013 Aug 24;382(9893):709-19. doi: 10.1016/S0140-6736(13)61502-0. PubMed PMID:23972814.
4. Saito M, Suzuki H, Kono K, Takenoshita S, Kohno T. Treatment of lung adenocarcinoma by molecular-targeted therapy and immunotherapy. *Surg Today*. 2018 Jan;48(1):1-8. doi: 10.1007/s00595-017-1497-7. Epub 2017 Mar 9. PubMed PMID:28280984.
5. Baron AP, von Schubert C, Cubizolles F, Siemeister G, Hitchcock M, Mengel A, Schröder J, Fernández-Montalván A, von Nussbaum F, Mumberg D, Nigg EA. Probing the catalytic functions of Bub1 kinase using the small molecule inhibitors BAY-320 and BAY-524. 2016 Feb 17;5. pii: e12187. doi: 10.7554/eLife.12187. PubMed PMID:26885717;PubMed Central PMCID:PMC 4769170.
6. Raaijmakers JA, van Heesbeen RGHP, Blomen VA, Janssen LME, van Diemen F, Brummelkamp TR, Medema RH. BUB1 Is Essential for the Viability of Human Cells in which the Spindle Assembly Checkpoint Is Compromised. *Cell Rep*. 2018 Feb 6;22(6):1424-1438. doi: 10.1016/j.celrep.2018.01.034. PubMed PMID:29425499.
7. Han JY, Han YK, Park GY, Kim SD, Lee CG. Bub1 is required for maintaining cancer stem cells in breast cancer cell lines. *Sci Rep*. 2015 Nov 2;5:15993. doi: 10.1038/srep15993. PubMed PMID:26522589;PubMed Central PMCID:PMC
8. Piao J, Zhu L, Sun J, Li N, Dong B, Yang Y, Chen L. High expression of CDK1 and BUB1 predicts poor prognosis of pancreatic ductal adenocarcinoma. 2019 Mar 18;701:15-22. doi: 10.1016/j.gene.2019.02.081. PubMed PMID:30898709.
9. Stahl D, Braun M, Gentles AJ, Lingohr P, Walter A, Kristiansen G, Gütgemann I. Low BUB1 expression is an adverse prognostic marker in gastric adenocarcinoma. 2017 Jul 18;8(44):76329-76339. doi: 10.18632/oncotarget.19357. PubMed PMID:29100315;PubMed Central PMCID: PMC5652709.
10. Ocaña A, Pérez-Peña J, Alcaraz-Sanabria A, Sánchez-Corrales V, Nieto-Jiménez C, Templeton AJ, Seruga B, Pandiella A, Amir E. In silico analyses identify gene-sets, associated with clinical outcome in ovarian cancer: role of mitotic kinases. 2016 Apr 19;7(16):22865-72. doi: 10.18632/oncotarget.8118. PubMed PMID: 26992217;PubMed Central PMCID:PMC5008407.

11. Rhodes DR, Kalyana-Sundaram S, Mahavisno V, Varambally R, Yu J, Briggs BB, Barrette TR, Anstet MJ, Kincead-Beal C, Kulkarni P, Varambally S, Ghosh, Chinnaiyan AM. Oncomine 3.0: genes, pathways, and networks in a collection of 18,000 cancer gene expression profiles. *Neoplasia*. 2007 Feb; 9(2): 166–180. PubMed PMID: 17356713;PubMed Central PMCID:PMC1
12. Chandrashekar DS, Bashel B, Balasubramanya SAH, Creighton CJ, Rodriguez IP, Chakravarthi BVSK and Varambally S. UALCAN: A portal for facilitating tumor subgroup gene expression and survival analyses. *Neoplasia*. 2017 Aug;19(8):649-658. doi: 10.1016/j.neo.2017.05.002. PMID:28732212.
13. Gao J, Aksoy BA, Dogrusoz U, Dresdner G, Gross B, Sumer SO, Sun Y, Jacobsen A, Sinha R, Larsson E, Cerami E, Sander C, Schultz N. Integrative analysis of complex cancer genomics and clinical profiles using the cBioPortal. *Sci Signal*. 2013 Apr 2;6(269):pl1. doi: 10.1126/scisignal.2004088. PubMed PMID: 23550210;PubMed Central PMCID:PMC4
14. Cerami E, Gao J, Dogrusoz U, Gross BE, Sumer SO, Aksoy BA, Jacobsen A, Byrne CJ, Heuer ML, Larsson E, Antipin Y, Reva B, Goldberg AP, Sander C, Schultz N. The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data. *Cancer Discov*. 2012 May;2(5):401-4. doi: 10.1158/2159-8290.CD-12-0095. PubMed PMID: 22588877;PubMed Central PMCID:PMC3
15. Huang da W, Sherman BT, Lempicki RA. Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nat Protoc*.2009;4(1):44-57. doi: 10.1038/nprot.2008.211. PubMed PMID: 19131956.
16. Vasaikar SV, Straub P, Wang J, Zhang B. LinkedOmics: analyzing multi-omics data within and across 32 cancer types. *Nucleic Acids Res*. 2018; 46:D956-D63. doi: 10.1093/nar/gkx1090. PubMed PMID: 29136207;PubMed Central PMCID:PMc5753188.
17. Warde-Farley D, Donaldson SL, Comes O, Zuberi K, Badrawi R, Chao P, Franz M, Grouios C, Kazi F, Lopes CT, Maitland A, Mostafavi S, Montojo J, Shao Q, Wright G, Bader GD, Morris Q. The GeneMANIA prediction server: biological network integration for gene prioritization and predicting gene function. *Nucleic Acids Res*. 2010 Jul;38(Web Server issue):W214-20. doi: 10.1093/nar/gkq537. PubMed PMID: 20576703;PubMed Central PMCID:PMc2
18. Lara-Gonzalez P, Westhorpe FG, Taylor SS. The spindle assembly checkpoint. *Curr Biol*. 2012 Nov 20;22(22):R966-80. doi: 10.1016/j.cub.2012.10.006. PubMed PMID:23174302
19. Travis WD. Pathology of lung cancer. *Clin Chest Med*. 2011 Dec;32(4):669-92. doi: 10.1016/j.ccm.2011.08.005. PubMed PMID:22054879.
20. Kawashima SA, Yamagishi Y, Honda T, Ishiguro K, Watanabe Y. Phosphorylation of H2A by Bub1 prevents chromosomal instability through localizing shugoshin. 2010 Jan 8;327(5962):172-7. doi: 10.1126/science.1180189. Epub 2009 Nov 19. PubMed PMID: 19965387.
21. Fischer M, Quaas M, Steiner L, Engeland K. The p53-p21-DREAM-CDE/CHR pathway regulates G2/M cell cycle genes. *Nucleic Acids Res*. 2016 Jan 8;44(1):164-74. doi: 10.1093/nar/gkv927. Epub 2015 Sep 17. PubMed PMID: 26384566;PubMed Central PMCID:PMc4

22. [Gjoerup OV, Wu J, Chandler-Militello D, Williams GL, Zhao J, Schaffhausen B, Jat PS, Roberts TM.](#) Surveillance mechanism linking Bub1 loss to the p53 pathway. [Proc Natl Acad Sci U S A.](#) 2007 May 15;104(20):8334-9. Epub 2007 May 8. PubMed PMID: 17488820;PubMed Central PMCID:[PMC1](#)
23. [Hanahan D, Weinberg RA.](#) Hallmarks of cancer: the next generation. 2011 Mar 4;144(5):646-74. doi: 10.1016/j.cell.2011.02.013. PubMed PMID: 21376230.
24. [Jeggo PA, Pearl LH, Carr AM.](#) DNA repair, genome stability and cancer: a historical perspective. [Nat Rev Cancer.](#) 2016 Jan;16(1):35-42. doi: 10.1038/nrc.2015.4. Epub 2015 Dec 15. PubMed PMID: 26667849.
25. [Yogosawa S, Yoshida K.](#) Tumor suppressive role for kinases phosphorylating p53 in DNA damage-induced apoptosis. [Cancer Sci.](#) 2018 Nov;109(11):3376-3382. doi: 10.1111/cas.13792. Epub 2018 Oct 6. PubMed PMID: 30191640;PubMed Central PMCID:[PMC6](#)
26. [Zhang HH, Li SZ, Zhang ZY, Hu XM, Hou PN, Gao L, Du RL, Zhang XD.](#) Nemo-like kinase is critical for p53 stabilization and function in response to DNA damage. [Cell Death Differ.](#) 2014 Oct;21(10):1656-63. doi: 10.1038/cdd.2014.78. Epub 2014 Jun 13. PubMed PMID: 24926618;PubMed Central PMCID:[PMC4](#)
27. [Herrera MC, Chymkowitch P, Robertson JM, Eriksson J, Bøe SO, Alseth I, Enserink JM.](#) Cdk1 gates cell cycle-dependent tRNA synthesis by regulating RNA polymerase III activity. [Nucleic Acids Res.](#) 2018 Dec 14;46(22):12188-12189. doi: 10.1093/nar/gky1102. PubMed PMID: 30395271;PubMed Central PMCID:[PMC6](#)
28. [Kabeche L, Nguyen HD, Buisson R, Zou L.](#) A mitosis-specific and R loop-driven ATR pathway promotes faithful chromosome segregation. 2018 Jan 5;359(6371):108-114. doi: 10.1126/science.aan6490. Epub 2017 Nov 23. PubMed PMID: 29170278;PubMed Central PMCID:[PMC5875943](#).
29. [Fang H, Niu K, Mo D, Zhu Y, Tan Q, Wei D, Li Y, Chen Z, Yang S, Balajee AS, Zhao Y.](#) RecQL4-Aurora B kinase axis is essential for cellular proliferation, cell cycle progression, and mitotic integrity. 2018 Sep 12;7(9):68. doi: 10.1038/s41389-018-0080-4. PubMed PMID: 30206236;PubMed Central PMCID:[PMC6134139](#).
30. [Wang L, Chen H, Wang C, Hu Z, Yan S.](#) Negative regulator of E2F transcription factors links cell cycle checkpoint and DNA damage repair. [Proc Natl Acad Sci U S A.](#) 2018 Apr 17;115(16):E3837-E3845. doi: 10.1073/pnas.1720094115. Epub 2018 Apr 2. PubMed PMID: 29610335;PubMed Central PMCID:[PMC5](#)
31. [Liu ZL, Bi XW, Liu PP, Lei DX, Wang Y, Li ZM, Jiang WQ, Xia Y.](#) Expressions and prognostic values of the E2F transcription factors in human breast carcinoma. [Cancer Manag Res.](#) 2018 Sep 13;10:3521-3532. doi: 10.2147/CMAR.S172332. eCollection 2018. PubMed PMID: 30271201;PubMed Central PMCID:[PMC6](#)
32. [Chen J, Gong C, Mao H, Li Z, Fang Z, Chen Q, Lin M, Jiang X, Hu Y, Wang W, Zhang X, Chen X, Li H.](#) E2F1/SP3/STAT6 axis is required for IL-4-induced epithelial-mesenchymal transition of colorectal

- cancer cells. *Int J Oncol*. 2018 Aug;53(2):567-578. doi: 10.3892/ijo.2018.4429. Epub 2018 Jun 5. PubMed PMID: 29901191;PubMed Central PMCID:PMC6
33. Yin J, Fu W, Dai L, Jiang Z, Liao H, Chen W, Pan L, Zhao J. ANKRD22 promotes progression of non-small cell lung cancer through transcriptional up-regulation of E2F1. *Sci Rep*. 2017 Jun 30;7(1):4430. doi: 10.1038/s41598-017-04818-y. PubMed PMID: 28667340;PubMed Central PMCID:PMC5
 34. Bartel DP. MicroRNAs: genomics, biogenesis, mechanism, and function. 2004 Jan 23;116(2):281-97. PubMed PMID: 14744438.
 35. Rupaimoole R, Slack FJ. MicroRNA therapeutics: towards a new era for the management of cancer and other diseases. *Nat Rev Drug Discov*. 2017 Mar;16(3):203-222. doi: 10.1038/nrd.2016.246. Epub 2017 Feb 17. PubMed PMID: 28209991.
 36. Zeng XL, Zhang SY, Zheng JF, Yuan H, Wang Y. Altered miR-143 and miR-150 expressions in peripheral blood mononuclear cells for diagnosis of non-small cell lung cancer. *Chin Med J (Engl)*. 2013 Dec;126(23):4510-6. PubMed PMID: 24286416.
 37. Lin K, Xu T, He BS, Pan YQ, Sun HL, Peng HX, Hu XX, Wang SK. MicroRNA expression profiles predict progression and clinical outcome in lung adenocarcinoma. *Onco Targets Ther*. 2016 Sep 15;9:5679-5692. eCollection 2016. PubMed PMID: 27695346;PubMed Central PMCID:PMC5
 38. Zhang JG, Guo JF, Liu DL, Liu Q, Wang JJ. MicroRNA-101 exerts tumor-suppressive functions in non-small cell lung cancer through directly targeting enhancer of zeste homolog 2. *J Thorac Oncol*. 2011 Apr;6(4):671-8. doi: 10.1097/JTO.0b013e318208eb35. PubMed PMID: 21270667.
 39. Yerukala Sathipati S, Ho SY. Identifying the miRNA signature associated with survival time in patients with lung adenocarcinoma using miRNA expression profiles. *Sci Rep*. 2017 Aug 8;7(1):7507. doi: 10.1038/s41598-017-07739-y. PubMed PMID: 28790336;PubMed Central PMCID:PMC5
 40. Bian H, Zhou Y, Zhou D, Zhang Y, Shang D, Qi J. The latest progress on miR-374 and its functional implications in physiological and pathological processes. *J Cell Mol Med*. 2019 Feb 17. doi: 10.1111/jcmm.14219. [Epub ahead of print] PubMed PMID: 30772950.

Figures

Figure 1

The gene expression level and copy number of BUB1 in lung adenocarcinoma (Oncomine). RBM8A mRNA level and DNA copy number were significantly higher in lung adenocarcinoma than in normal tissue. (A) BUB1 expression in Garber lung, (B) BUB1 expression in Hou lung, (C) BUB1 expression in Landi lung and (D) BUB1 expression in Su lung.

Figure 2

The gene expression level of BUB1 in subgroups of patients with lung adenocarcinoma (UALCAN). BUB1 transcription level in lung adenocarcinoma patients was significantly higher than in healthy people in subgroup analysis based on gender, age, race, cancer stage and smoking habits ($P < 0.001$). (A) BUB1 expression in lung adenocarcinoma based on sample types, (B) BUB1 expression in lung adenocarcinoma based on individual cancer stages, (C) BUB1 expression in lung adenocarcinoma based on patient's race, (D) BUB1 expression in lung adenocarcinoma based on patient's gender, (E) BUB1 expression in lung adenocarcinoma based on patient's age and (F) BUB1 expression in lung adenocarcinoma based on patient's smoking habits.

Figure 3

Survival analysis of BUB1 in subgroups of patients with lung adenocarcinoma (UALCAN). The BUB1 level and race have significant influence on the survival of lung adenocarcinoma patients ($P < 0.05$), while gender and smoking habit not. (A) effect of BUB1 expression level on lung adenocarcinoma patient survival, (B) effect of BUB1 expression level and race on lung adenocarcinoma patient survival, (C) effect of BUB1 expression level and gender on lung adenocarcinoma patient survival and (D) effect of BUB1 expression level and smoking habit on lung adenocarcinoma patient survival.

Figure 4

RBM8A alterations in lung adenocarcinoma (cBioPortal). OncoPrint of BUB1 alterations in LUAD. Different types of genetic alterations are displayed in different colors.

Figure 5

Frequency of alterations in BUB1 in lung adenocarcinoma (cBioPortal). Upregulation of miRNA is the most frequent alteration.

Figure 6

Enrichment analysis of the genes altered in the BUB1 neighborhood in lung adenocarcinoma. (A) The PPI network of the top 50 altered neighbor genes of BUB1 (STRING), (B) GO analysis (DAVID) and (C) KEGG pathway (DAVID).

Figure 7

Genes differentially expressed in correlation with BUB1 in lung adenocarcinoma (LinkedOmics). (A) positively correlated genes, (B) negatively correlated genes, (C) positively correlated genes and (D) negatively correlated genes.

Figure 8

Enrichment analysis of the genes differentially expressed in correlation with BUB1 in lung adenocarcinoma (LinkedOmics; GSEA). (A) Cellular components, (B) Biological processes, (C) Molecular functions, (D) KEGG pathway.

Figure 10

The kinase, miRNA and transcription factor-target networks of BUB1 in lung adenocarcinoma (LinkedOmics). (A) miRNAs, (B) kinases, (C) transcription factors.