

Irreversible fatal contrast-induced encephalopathy: a case report

Wei Zhao

Department of neurology, Shandong Qianfoshan Hospital

Jinping Zhang

Department of neurology, Shandong Qianfoshan Hospital

Yun Song

Department of neurology, Shandong Qianfoshan Hospital

Lili Sun

Department of neurology, Shandong Qianfoshan Hospital

Meimei Zheng

Shandong Qianfoshan Hospital

Hao Yin

Shandong Qianfoshan Hospital

Jun Zhang

Shandong Qianfoshan Hospital

Wei Wang

Shandong Qianfoshan Hospital

Ju Han (✉ hanjujack@163.com)

Case report

Keywords: Digital subtraction angiography; iodinated contrast agents; complications; contrast-induced encephalopathy

Posted Date: February 25th, 2019

DOI: <https://doi.org/10.21203/rs.2.204/v2>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published on March 28th, 2019. See the published version at <https://doi.org/10.1186/s12883-019-1279-5>.

Abstract

Background: Contrast-induced encephalopathy (CIE) is a well-known complication of iodinated contrast agents during angiography and vascular interventions. It can manifest as hemiparesis, cortical blindness, speech changes, Parkinsonism, confusion, seizure, and coma. Most of the reported CIE cases have been transient and reversible. Irreversible fatal CIE cases have been rarely reported. All the fatal CIE cases reported involved the use of ionic high osmolar contrast agents. Here, we document a heretofore unreported fatal CIE after digital subtraction angiography (DSA) using iopamidol, which is a type of non-ionic monomer low osmolar contrast agent. Case presentation: A 71-year-old woman was admitted to our Department of Neurology for paroxysmal buzzing of the brain. The cerebral magnetic resonance angiography (MRA) indicated the presence of arteriosclerosis of the cerebral arteries and bilateral stenosis of the middle cerebral arteries. The patient underwent DSA for further diagnostic work-up. The total amount of iopamidol used during the procedure was 110 ml. The patient experienced headache during the procedure, followed by dizziness with nausea and vomiting. Although the patient was treated with anti-oedema drugs, her deterioration was continuous. She gradually became comatose and suffered irreversible fatal cerebral oedema, which was confirmed by cerebral computed tomography (CT). Finally, the patient died 56 days after the procedure due to irreversible fatal cerebral oedema. Conclusions: This report documents that iopamidol-induced encephalopathy may not always have a benign outcome and can result in irreversible fatal cerebral oedema.

Background

Contrast-induced encephalopathy (CIE) is a known but rare complication of angiography and endovascular interventions. The presentations may include hemiparesis, cortical blindness, speech changes, Parkinsonism, confusion, seizure, and coma [1, 2]. In most reported cases, the symptoms are reversible, and fatal encephalopathy following iodinated contrast administration has been rarely reported. Only 8 cases of autopsy-proven fatal cerebral oedema due to contrast neurotoxicity in the early stage of angiography have been reported [1, 3, 4]. All these reported fatal cases involved the use of high osmolar contrast agents. Iopamidol is a non-ionic monomer low osmolar contrast agent, which has been reported in cases of reversible contrast-induced encephalopathy [5-9]. Here, we describe a patient who suffered irreversible fatal encephalopathy after DSA using iopamidol.

Case Presentation

A 71-year-old woman with a history of hypertension, hyperlipemia, and angina was admitted to our Department of Neurology for paroxysmal buzzing of the brain. On physical examination, she had decreased hearing bilaterally. The cerebral magnetic resonance imaging (MRI) of the patient indicated multiple cerebral infarctions and bilateral demyelination in the centrum semiovale. The cerebral MRA indicated the presence of arteriosclerosis of the cerebral arteries, and bilateral stenosis of the middle cerebral arteries. For further diagnosis, the patient underwent DSA subsequently. The total amount of iopamidol (Bracco Imaging Italia S.r.l.) administered during the procedure was 110 ml. The DSA showed

that the patient had bilateral embryonic posterior cerebral arteries, 40% stenosis of the left middle cerebral artery and tortuous vertebral arteries bilaterally. There was no obvious calcification of the aortic arch; angiography of the arch using 25 ml iopamidol was performed only once. Ten minutes after the aortic arch angiography, the patient experienced mild headache. The pain was bearable, and the patient could cooperate during the remainder of the procedure. The DSA was completed 20 minutes later. No haemorrhage or vasospasm was detected during the procedure. The headache was continuous, and the patient suffered nausea and vomiting. The immediate physical examination showed no obvious abnormal sign. The patient was treated with 8 mg ramosetron and 10 mg dexamethasone. After 20 minutes of observation, the symptoms were relieved. Her cerebral CT scan at the time was normal (Figure 1A, B, C). Two hours later, the patient manifested dizziness with nausea and vomiting and was treated with 8 mg ondansetron and 20 mg diphenhydramine. Meanwhile, compound sodium chloride injection was used to facilitate the elimination of the contrast agent. The treatment alleviated her symptoms. Four hours after the procedure, the patient manifested dizziness again; 5 mg dexamethasone was administered, but the dizziness was not relieved, and she vomited again. At 11 hours, her dizziness was alleviated, but her blood pressure rose to 183/92 mmHg. The patient was drowsy but could answer questions correctly; the pupil diameters of both eyes were normal and equal, pupillary light reflex was reactive, and the movement of the limbs was normal. The patient was treated with 30 mg nimodipine tablets to alleviate the high blood pressure. Fourteen hours after the procedure, the patient fell asleep, but at 17 hours, the electrocardiograph monitor showed sudden respiratory failure, with a pulse oxygen saturation reduced to 88% and gradually declining. The patient was in a coma state, with sighing respiration; the pupil diameters of both eyes were unequal and unreactive, the left one was 4.5 mm and the right one was 3 mm, the limb drop test was positive, and Babinski sign was negative. Cerebral hernia was considered due to the sudden deterioration. The patient was treated with oxygen inhalation, 20% mannitol, nikethamide, lobeline and diprophylline. The patient was transferred to the intensive care unit for further treatment after cardio-pulmonary resuscitation, endotracheal intubation and mechanical ventilation. Two days after the procedure, cerebral CT scan indicated diffuse cerebral oedema, loss of grey-white differentiation, effacement of the cerebral sulci and decrease in cerebrospinal fluid space (Figure 1D, E, F). The patient was treated with dehydration, mechanical ventilation, and anti-infectious agent, but the diffuse cerebral oedema did not improve. Nine days after the procedure, the third cerebral CT scan showed that the cerebral oedema had become much more severe, the ventricles had disappeared and there was hyperdense signal in the subarachnoid space, which was considered to be indicative of a pseudo-subarachnoid haemorrhage due to the severe cerebral oedema [10] (Figure 1G, H, I). Fifteen days after the procedure, the cerebral CT showed unrelieved diffuse oedema of the brain, and the hyperdense signal in the subarachnoid space persisted (Figure 1J, K, L). None of these cerebral CT scans showed intracerebral haemorrhage or infarct in this patient. The patient remained in a continuous deep coma state, and the brainstem reflexes had disappeared; she died 56 days after that sudden deterioration.

Discussion And Conclusions

The prognosis of most CIE is generally reported to be good with a rapid recovery, and only rare cases with the persistent deficits have been reported [2]. Notably, there were 8 cases of autopsy-proven fatal cerebral oedema due to contrast neurotoxicity in the early stage of angiography [1, 3, 4]. The 8 deaths included 6 infants; 5 of these patients underwent cardiac angiography, and the other 3 received aortography. All fatal cerebral oedema cases reported involved the use of ionic high osmolar contrast agents, and ionic high osmolar contrast agents are no longer used in routine angiography and intervention procedures. The case which we report here may be the first fatal cerebral oedema after DSA using iopamidol. This case highlights the potential for other types of iodinated contrast agents to induce fatal encephalopathy.

The diagnosis of CIE is important, as its presentation may be similar to embolism, and haemorrhagic complications following angiography or endovascular interventions. Typical radiological findings include abnormal cortical contrast enhancement and cerebral oedema, subarachnoid contrast enhancement and striatal contrast enhancement [2, 11]. CT or MRI of the brain helps us to differentiate CIE from haemorrhage or infarct. In the case which we report here, none of the CT scans of the brain after DSA (immediately, 2 days, 9 days and 15 days after the procedure) indicated intracerebral haemorrhage or infarct. Therefore, the possibility of multiple embolisms was not considered in this case. The hyperdense signal in the subarachnoid space in the cerebral CT scans was considered to be due to the severe diffuse cerebral oedema. The hyperdense appearance results from a combination of loss of grey-white differentiation, narrowing and effacement of the subarachnoid spaces, and corresponding engorgement of superficial pial veins [10].

The mechanism of CIE is controversial. The temporary disruption of the blood-brain barrier (BBB) after injection of the iodinated contrast agent is widely accepted [2, 11-15]. Experimental studies have demonstrated that contrast agents can penetrate the altered BBB and that this is dependent on the contact time, anions and dosage [1, 12, 13, 15]. Both the hyperosmolality and chemotoxicity of the contrast agents contribute to the disruption of the BBB. Hyperosmolality of the contrast medium is hypothesised to cause shrinkage of endothelial cells and disrupt tight junctions [12]. Other studies suggest that the alteration of the BBB is due to the physical or chemical effects of the contrast medium on the BBB instead of the hyperosmolality [14]. The expression of endothelin, which can be induced by radiocontrast agents, can increase human brain endothelial cell permeability and has been implicated in the pathophysiology of disorders associated with BBB injury [2, 15].

Studies have indicated that opening of the BBB may be accompanied by brain oedema, resulting from the flux of proteins, electrolytes, and water across the abnormally permeable cerebral vessels into the extracellular space [4]. An idiosyncratic response to small doses of contrast agent, which may be related to the areas of incompleteness of their BBB, has been reported [1]. We postulate that the idiosyncratic response to contrast agents may have contributed to the patient's prolonged and progressive brain oedema. Contrast agents can produce direct neurotoxic effects on the neurons and astrocytes when they penetrate the altered BBB. Experimental studies have shown that ionised contrast agents can severely alter neuronal function when directly introduced into the nervous system [1, 12, 13, 15]. We hypothesised

that the direct neurotoxic effect of the contrast agent also contributed to the patient's progressive and fatal brain oedema.

All types of iodinated contrast agents can induce the development of neurotoxicity, but the occurrence of fatal cerebral oedema is very rare. Unfortunately, there is no currently available effective treatment for such a severe fatal CIE. In the case reported by L. Junck and W.H. Marshall [4], the post mortem tissue iodine concentrations were the highest in the urine, serum and kidney. The use of continuous renal replacement therapy and continuous blood purification may be potential treatments for cases of fatal CIE.

In summary, although CIE has typically been associated with benign outcomes in previous studies, we present a case of fatal cerebral oedema after DSA using iopamidol. This case illustrates the potential to cause severe complications, even fatal cerebral oedema, with all types of iodinated contrast agents. The doctors performing angiography and interventions should be aware of this severe potentially harmful effect. The rare occurrence of severe contrast-induced complications renders their prevention very difficult. Further studies are needed to define the risk factors and the mechanism of the iodinated contrast agent neurotoxicity, which may help minimise the occurrence of severe complications.

List Of Abbreviations

BBB: blood-brain barrier

CIE: contrast-induced encephalopathy

CT: computed tomography

DSA: digital subtraction angiography

MRA: magnetic resonance angiography

MRI: magnetic resonance imaging

Declarations

Ethics approval and consent to participate

The Institutional Review Board of Qianfoshan Hospital affiliated to Shandong University approved the study.

Consent for publication

A signed informed consent was obtained from the patient's guardian for publication of this case report and accompanying neuroimages.

Availability of data and material

All data generated and analysed during this study are included in this article.

Competing interests

The authors declare that they have no competing interests.

Funding

None.

Authors' contributions

WZ designed and wrote the manuscript. JPZ, YS, and LLS examined the patient. MMZ, HY, JZ and WW analysed the neuroimages. JH examined the patient, designed the case report and helped draft the manuscript. All authors read and approved the final manuscript.

Acknowledgements

None.

References

1. Lalli AF. Contrast media reactions: data analysis and hypothesis. *Radiology*. 1980;134 1:1-12; doi: 10.1148/radiology.134.1.6985735.
2. Leong S, Fanning NF. Persistent neurological deficit from iodinated contrast encephalopathy following intracranial aneurysm coiling. A case report and review of the literature. *Interv Neuroradiol*. 2012;18 1:33-41.
3. Shrivastava S, Mohan JC, Chopra P. Fatal cerebral edema following angiocardiology: a case report. *Int J Cardiol*. 1985;8 4:490-1.
4. Junck L, Marshall WH. Fatal brain edema after contrast-agent overdose. *AJNR Am J Neuroradiol*. 1986;7 3:522-5.
5. Parry R, Rees JR, Wilde P. Transient cortical blindness after coronary angiography. *British heart journal*. 1993;70 6:563-4.
6. Kamata J, Fukami K, Yoshida H, Mizunuma Y, Moriai N, Takino T, et al. Transient cortical blindness following bypass graft angiography. A case report. *Angiology*. 1995;46 10:937-46.
7. Uchiyama Y, Abe T, Hirohata M, Tanaka N, Kojima K, Nishimura H, et al. Blood brain-barrier disruption of nonionic iodinated contrast medium following coil embolization of a ruptured intracerebral aneurysm. *AJNR Am J Neuroradiol*. 2004;25 10:1783-6.

8. Gellen B, Remp T, Mayer T, Milz P, Franz WM. Cortical blindness: a rare but dramatic complication following coronary angiography. *Cardiology*. 2003;99 1:57-9; doi: 10.1159/000068443.
9. Merchut MP, Richie B. Transient visuospatial disorder from angiographic contrast. *Arch Neurol*. 2002;59 5:851-4.
10. Hasan TF, Duarte W, Akinduro OO, Goldstein ED, Hurst R, Haranhalli N, et al. Nonaneurysmal "Pseudo-Subarachnoid Hemorrhage" Computed Tomography Patterns: Challenges in an Acute Decision-Making Heuristics. *J Stroke Cerebrovasc Dis*. 2018;27 9:2319-26; doi: 10.1016/j.jstrokecerebrovasdis.2018.04.016.
11. Lantos G. Cortical blindness due to osmotic disruption of the blood-brain barrier by angiographic contrast material: CT and MRI studies. *Neurology*. 1989;39 4:567-71.
12. Junck L, Marshall WH. Neurotoxicity of radiological contrast agents. *Ann Neurol*. 1983;13 5:469-84; doi: 10.1002/ana.410130502.
13. Torvik A, Walday P. Neurotoxicity of water-soluble contrast media. *Acta Radiol Suppl*. 1995;399:221-9.
14. Wilson AJ, Evill CA, Sage MR. Effects of nonionic contrast media on the blood-brain barrier. Osmolality versus chemotoxicity. *Invest Radiol*. 1991;26 12:1091-4.
15. Heyman SN, Clark BA, Kaiser N, Spokes K, Rosen S, Brezis M, et al. Radiocontrast agents induce endothelin release in vivo and in vitro. *J Am Soc Nephrol*. 1992;3 1:58-65.

Figures




Figure 1

Cerebral CT immediately after DSA did not indicate any obvious abnormal sign (A, B, C); 2 days after the procedure, a repeat cerebral CT revealed diffuse cerebral oedema, loss of grey-white differentiation, effacement of the cerebral sulci and decrease in cerebrospinal fluid space (D, E, F); 9 days after the procedure, the cerebral CT showed more severe diffuse oedema of the brain, loss of grey-white differentiation, effacement of the cerebral sulci and subarachnoid space, disappearance of the cerebral

ventricles, enhancement of the subarachnoid space, and darkened brain tissue in Hounsfield units (G, H, I); 15 days after the procedure, the cerebral CT showed persistent diffuse oedema of the brain with effacement of the cerebral ventricles and sulci, darkened brain tissue in Hounsfield units, and enhancement of the subarachnoid space (J, K, L).

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [supplement1.jpg](#)