Nest Size Matters: Common Cuckoos Prefer to Parasitize Larger Nests of Oriental Reed Warblers

Longwu Wang

Guizhou Normal University School of Life Sciences <https://orcid.org/0000-0002-7020-4397>

Gangbin He

Guizhou Normal University

Canchao Yang

Hainan Normal University

Anders Pape Møller

Universite Paris-Saclay

Wei Liang (✉ liangwei@hainnu.edu.cn)

Hainan Normal University

Research

Keywords: brood parasitism, common cuckoo, nest preference, nest size, Oriental reed warbler

Posted Date: February 18th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-204677/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Animal Cognition on November 12th, 2021.
See the published version at <https://doi.org/10.1007/s10071-021-01574-5>.

Abstract

Background

Avian brood parasites leave parental care of their offspring to foster parents. Theory predicts that parasites should select for large host nests when they have sufficient available host nests at a given time. We developed an empirical experimental design to address this hypothesis by studying nest choice of common cuckoos (*Cuculus canorus*) among nests of its Oriental reed warbler (*Acrocephalus orientalis*) hosts.

Results

We presented two groups of experimental nests: 1) nest dyads comprise one large and one small artificial nest from reed leaves, and 2) nest triads tied together use the modified old own warbler nests including enlarged, reduced and medium sized nests to elicit parasitism by common cuckoos. We predicted that cuckoos prefer larger nests over medium sized ones, and over the smallest nest. Our experimental findings show that common cuckoo females generally prefer large nests over medium or small sized nests. Furthermore, experiments showed that cuckoo parasitism was significantly more common than in previous studies of the same warbler population.

Conclusions

Our results implying that larger, taller and more exposed host nests effectively increased the probability of cuckoo parasitism.

Background

Avian brood parasites increase their fitness through evolved egg-laying behaviour in the nests of hosts while leaving parental care of their offspring to the hosts (Davies 2000). Brood parasites always depend on parasitized hosts for the care of their offspring. However, foster parents differ in their food provisioning ability of cuckoo nestlings caused by variation in parental quality (Brooke and Davies 1989; Grim et al. 2017). Therefore, if altricial obligate brood parasites maximize the benefits from the parental effort of their hosts, cuckoos should search for and select host nests with better quality during parasitism (de Neve et al. 2004; Álvarez and Barba 2008; Jelínek et al. 2016).

Birds' nests are used for containing the eggs and/or offspring during breeding (Hansell 2000, 2007), the potential signaling function of a high-quality nest being important for successful fledging to avoid hosts from suffering from predation or from falling out of the nest (Álvarez and Barba 2008; Heenan and Seymour 2011; Møller et al. 2014; Møller 2017). Thus, parasites should search for the most suitable host nest (Davies 2000). How cuckoos find and choose host nests for parasitism is still poorly understood, and previous studies suggested that egg-laying by cuckoos should match the phenotypes or appearance of their host eggs (Avilés et al. 2006; Cherry et al. 2007; Honza et al. 2014; Li et al. 2016), while other

studies challenged these ideas and suggested that cuckoos randomly searched host nests for parasitism (Antonov et al. 2012; Yang et al. 2016b). Recent empirical work showed that when experimental nests in which some have different model eggs were given a similar opportunity for cuckoo parasitism, cuckoos selected these nests for parasitism without use of the content of nests including egg color, shape and size as search cues (Yang et al. 2016a). They randomly selected any one nest for parasitism and neglected the egg categories in the nest. Furthermore, the cuckoo parasitism frequency in their population associated with host nest size tended to be higher in large compared to small nests, they found that cuckoo females more often selected larger Oriental reed warbler nests (*Acrocephalus orientalis*) over smaller ones of black-browed reed warbler (*Acrocephalus bistrigiceps*) when these two types of nests were available for cuckoo choice (Yang et al. 2017).

The optimal egg-laying strategy proposes that when there are sufficiently many available nests at a given time, cuckoo females should prefer the nest made by the best quality hosts (Hauber 2001; Parejo and Avilés 2007; Jelínek et al. 2014). In this scenario, cuckoos may select larger nests associated with increased fitness because larger nests may indicate superior parenting ability (Soler et al. 1998; de Neve et al. 2004; Avilés et al. 2009; Jelínek et al. 2015). A study spanning eight consecutive years showed that large magpie (*Pica pica*) nests were more likely to be parasitized by great spotted cuckoos (*Clamator glandarius*) than smaller nests when cuckoos have a larger availability of host nests among which to choose. Therefore, selection of active host nests may arise from nest choice by the great spotted cuckoo (Molina-Morales et al. 2016), probably because nest size correlates positively with parental abilities of magpies (Soler et al. 1995).

Here, we developed a novel field experimental method to test whether nest-size manipulation affects parasite nest preference when there are sufficiently many available host nests. The common cuckoo (*Cuculus canorus*) is the most common brood parasite using the Oriental reed warbler as its major host in our study area (Yang et al. 2016a, 2017; Wang et al. 2020a). Different types of artificial experimental nests can be categorized as large, medium or small size, and we divided these nests into two groups (nest dyads and triads) to test our hypotheses. We predicted that cuckoos would lay their eggs in larger host nests in both the two groups if they had evolved the ability to discriminate among different nest sizes.

Materials And Methods

Study site and study species

This study was performed in Zhalong National Nature Reserve (46°48'–47°31' N, 123°51'–124°37' E), which is located in Heilongjiang, northeast China. Fieldwork was carried out during the breeding season (June to August) in years 2015–2019. Habitats were primarily reed swamps with several small villages (Yang et al. 2017; Wang et al. 2020b). The common cuckoo is the main parasitic cuckoo in this study area, and the Oriental reed warbler is its predominant host with a high parasitism rate ranging from 34.3–65.5% among years (Yang et al. 2017), cuckoos often remove one or two eggs before they parasitize the

host nest (Wang et al. 2020a). Furthermore, the appearance of cuckoo eggs resembled the eggs of their hosts well (Yang et al. 2015; Li et al. 2016a).

Experimental Nest Manufacture

We designed two experiments to test the nest size hypothesis. For group 1, one large nest (internal diameter, mean $\pm$ SD: 86.42 ± 0.91 mm, $n = 36$) and one small nest (internal diameter, 67.57 ± 1.05 mm, $n = 36$) were tied together as a combination of nests (Fig. 1, group 1). We tied two randomly chosen and different sized nests together to constrain the choice of host nests to the experimental variable of nest size. This constrained the choice between the nest dyads, avoiding an absence of selection caused by distance between nests. If these two experimental nests were located at a distance, the cuckoo may not find and compare the two different-sized nests, especially when nests were located in dense reeds. These nests were weaved together with a mixture of dry reed leaves and sticks, sewing them with a needle and string. The purpose of this experiment was to control the nest structure, including reed material, shape, duration of nest construction, except for nest size, which was a unique character dividing nests into two categories. This helped us determine the relevance of nest size choice by cuckoos.

For group 2, 'triple nests' were installed in the field including large, medium and small sized old nests as described above. All nests used for this study were modified from natural old Oriental reed warbler nests. We used natural warbler nests as the medium category (internal diameter: 57.88 ± 5.00 mm, $n = 33$), while large nests were enlarged by 20 mm (internal diameter: 78.74 ± 4.34 mm, $n = 33$). Similarly, small nests were reduced by approximately 20 mm (internal diameter: 39.18 ± 3.42 mm, $n = 33$).

We collected natural old nests that had fledged young or were deserted the previous year. Nests were collected and preserved in cardboard boxes. For the large nest treatment, we selected two old nests that were most similar in appearance (size, shape and nest materials) for our experiment. First, we randomly selected one nest and made a cut in the rim of the nest with scissors. Second, we put one third of the other nest piece together with the rim cut and sewed it up. Similarly, for small nests, we cut nest material and re-sewed them to keep the internal diameter at approximately 40 mm. For the medium nests, we did not cut them, although we also sewed them with the same color string. We sewed these nests using needle and string to make them look similar in appearance to natural nests. We enlarged and reduced 36 nests in total for our nest experiments. We randomly selected one of three types of nests and sewed them together with string (Fig. 1, group 2).

Experiments For Attracting Cuckoo Parasitism

We searched systematically for naturally active Oriental reed warbler host nests and monitored reproductive activities of hosts daily. Observed nests were checked every day to confirm the first date of laying an egg. If these target nests were found to have one host egg, one set of the prepared combination of nests were tied to the reeds one meter away from this neighboring active nest, and two common quail (*Coturnix coturnix*) eggs were inserted into each of the three combinations of nests, respectively (Fig. 1). Here, we controlled that the two quail eggs stayed the same throughout the test period, avoiding that cuckoo females showed a preference bias towards variation in the number of eggs, and the large quail

eggs had the advantage that warbler hosts cannot remove them, but the cuckoos can easily remove them.

To increase the risk of brood parasitism, we specifically tied the combination of nests taller (approximately 0.5 m) in the vegetation (the nest height hypothesis, Budnik et al. 2002; Patten et al. 2011) than the position of the active host nest. We made them easier to see from the angle of view (nest exposure hypothesis, Moskát and Honza 2000; Clarke et al. 2001), which allowed cuckoos to more easily choose a specific type of nest during parasitism. Cuckoos generally selected to lay their eggs in Oriental reed warbler nests (Yang et al. 2016a, 2017). Thus, we did not design control nests that simultaneously set up another combination of nests in a random direction and the same nest content and position, as there was little probability for cuckoo parasitism in nests without host activity (Yang et al. 2017).

All experimental and naturally active nests were monitored for up to 6 days to confirm cuckoo parasitism until completion of the clutch and start of incubation. All experimental nests were video recorded and checked once per day. Video devices were retrieved at dusk. For active warbler nests, we did not manipulate, but checked and monitored clutch size and cuckoo parasitism.

Statistical Analyses

Binomial tests (two-tailed test) were used to test for a preference of nests during choice by cuckoos. For combinations of nests, when more than one nest was parasitized, the cuckoo egg was removed, and the original experimental status was re-established to ensure that all combinations of nests were parasitized with similar probability. Differences were considered significant at $P < 0.05$. Statistical analysis was conducted using IBM SPSS Version 22.0 (IBM Corp., Armonk, NY, USA), and unless otherwise specified, all results are presented as mean $\pm$ SD.

Results

The rate of parasitism was 58% (21/36) and 70% (23/33) for groups 1 and group 2, respectively. For group 1, the frequency of selection of the larger nests was 81% (17/21), and the preference for large nests was significant compared to small nests (Binomial test, $p = 0.007$, Fig. 2). For group 2, the frequency of selection of large nests was 78% (18/23) and the medium sized nests was 22% (5/23), respectively (Fig. 2). The smaller nest was never parasitized (0%, 0/23). The parasitism rate of the larger nest was significantly higher than that of the medium sized nest (Binomial test, $p = 0.011$; Fig. 2).

There was no case of parasitism among naturally active warbler nests in either group (0/36, 0/33). Four cases of egg laying in large, medium and small nests are shown in Electronic Supplementary Materials Videos S1-S2 in group 1 and Videos S3-S4 in group 2.

Discussion

The main finding of this study was that common cuckoos prefer large host nests for parasitism when they have sufficiently many synchronous available nests. Our findings support the active selection hypothesis suggesting that cuckoos prefer to lay eggs in larger nests providing evidence for the ability to distinguish among host nests varying in size when monitoring available nests. In addition, the frequency of parasitism was higher than in previous studies of the same warbler population, which implies that host nests with a taller and exposed position would effectively increase the possibility of cuckoo parasitism (Yang et al. 2016a, 2017).

Nest size is an indicator of parental care, and it is a likely reason why cuckoos choose large host nests for parasitism. For example, magpie pairs with large nests provide higher quality of food for their nestlings compared to pairs with smaller cuckoo nests (de Neve et al. 2004). The great spotted cuckoo prefers to parasitize large magpie nests because nest owners of such nests can rear nestlings more successfully than control nests (Soler et al. 1995). Nest size in the great reed warbler (*Acrocephalus arundinaceus*) serves as a signal of female quality for investment in reproduction, because enlarged great reed warbler nests significantly increase male feeding effort (Trnka and Prokop 2011; Jelínek et al. 2016). However, no evidence supports the hypothesis that nest size affects the frequency of parasitism in the great reed warbler (Avilés et al. 2009; Jelínek et al. 2015). We showed that cuckoos laid their eggs in larger host nests suggesting that cuckoos have evolved the ability to distinguish among different nest sizes to increase nestling provisioning. This was further revealed by clutch size in birds generally being larger in larger nests (Møller et al. 2014). Clutch size remained the same in experimental nests receiving all treatments, suggesting that nest size rather than clutch size is an important cue for parasitism selection, even if the benefit may derive from the clutch size-food provisioning relationship. In addition, the combination of different types of experimental nests presented next to each other makes it feasible for cuckoos to choose a nest without consideration of distance, allowing a similar preference for different sized nests.

When cuckoos approach a combination of nests with a similar angle of view or height, they may be subject to physical constraints, such that it is easier to perch on and deposit eggs in larger rather than smaller nests. Therefore, large nests should be easy to find and facilitate egg laying. Of course, this scenario depends on sufficiently many nests being available in the neighborhood. Ma et al. (2018) showed that isolated Oriental reed warbler host nests were more vulnerable to cuckoo parasitism far away from neighbors, cuckoo females having no other choice being forced to select the only isolated nest for parasitism. Our results show that parasitism frequency is higher than in the previous studies (Li et al. 2016a, 2016b; Ma et al. 2018), the conspicuousness of host nests for the likelihood of being parasitized by the cuckoo are vulnerable (Patten et al. 2011; Clarke et al. 2001), that is, the taller and more exposed are host nests, the more effectively the increase in the probability of cuckoo parasitism, consistent with the nest height hypothesis (Budnik et al. 2002) and the nest exposure hypothesis (Moskát and Honza 2000; Patten et al. 2011).

There may be other reasons for why cuckoos prefer larger nests. For example, there might be a physical body constraint, because many small hosts may have evolved a small sized nest allowing for parasitism

by small cuckoo parasites (i.e. Bianchi's warbler *Seicercus valentine* vs. Asian emerald cuckoo *Chrysococcyx maculatus*) (for more examples, see Yang et al. 2012). The only evidence is that larger nests in many different species of birds lay larger clutches (Møller et al. 2014), and larger clutches being associated with higher quality nest owners. In this study, two large quail eggs in large nests may also better predict larger final clutch size, two large quail eggs in small nests where there is no free space for other eggs may indicate small size of a complete clutch. This could also result in a higher preference of cuckoos for larger experimental nests. Moreover, Wang et al. (2020b) revealed that cuckoos prefer to select the natural-active host nests with small clutch sizes for parasitism [e.g., 1–2 host eggs, 75% (183/245)], and the empirical study also supporting this result (78%, 25/33) (Wang et al. 2020b). In addition, a large clutch size (e.g., 5–6 eggs) also implies more space for cuckoo egg-laying. Therefore, nest space may act as an important cue for cuckoo egg-laying resulting in strong nest preferences.

In conclusion, the present study provided experimental evidence suggesting that cuckoos preferred to lay their eggs in larger host nests when they have more available nests for use at any given time. Furthermore, the conspicuousness of host nests caused by higher location and full exposure in the field resulting in more cuckoo parasitism, suggesting that host nest exposure could increase the risk of cuckoo parasitism.

Declarations

Acknowledgements

We are grateful to Wenfeng Wang and Jianhua Ma from Zhalong National Nature Reserve for their help and cooperation. We thank Tinggao Yang and Hailin Lu for their assistance with fieldwork.

Authors' contributions

WL conceived and designed this project. LW and GH carried out field experiments. LW and CY performed data analyses. LW drafted the manuscript, and WL and APM were involved in discussion and revision of the manuscript. All authors read and approved the final manuscript.

Funding

This work was supported by the National Natural Science Foundation of China (Nos. 31960105 to LW, 31772453 and 31970427 to WL). LW was funded by the Joint Fund of the National Natural Science Foundation of China and the Karst Science Research Center of Guizhou province (No. U1812401).

Ethics approval and consent to participate

The experiments comply with the current laws of China, where they were performed. Fieldwork was carried out with permission (no. ZL-GZNU-2019-06) from Zhalong National Nature Reserve, Heilongjiang, China. Experimental procedures were in agreement with the Animal Research Ethics Committee of Hainan

Provincial Education Centre for Ecology and Environment, Hainan Normal University (no. HNECEE-2012-003).

Competing interests

The authors declare that they have no competing interest.

Consent for publication

Not applicable.

Availability of data and materials

Data used in this study are available in the electronic supplementary material.

References

- Álvarez E, Barba E. Nest quality in relation to adult bird condition and its impact on reproduction in great tits *Parus major*. *Acta Ornithol.* 2008;43:3-9.
- Antonov A, Stokke BG, Fossøy F, Ranke PS, Liang W, Yang C, Moksnes A, Shykof J, Røskift E. Are cuckoos maximizing egg mimicry by selecting host individuals with better matching egg phenotypes? *PLoS One.* 2012;7:e31704.
- Avilés JM, Stokke BG, Moksnes A, Røskift E, Asmul M, Møller AP. Rapid increase in cuckoo egg matching in a recently parasitized reed warbler population. *J Evol Biol.* 2006;19:1901-1910.
- Avilés JM, Moskát C, Bán M, Hargitai R, Parejo D. Common cuckoos (*Cuculus canorus*) do not rely on indicators of parental abilities when searching for host nests: The importance of host defenses. *Auk.* 2009;126:431-438.
- Budnik JM, Thompson III FR, Ryan MR. Effect of habitat characteristics on the probability of parasitism and predation of Bell's vireo nests. *J Wildl Manage.* 2002;66:232-239.
- Cherry MI, Bennett AT, Moskát C. Do cuckoos choose nests of great reed warblers on the basis of host egg appearance? *J Evol Biol.* 2007;20:1218-1222.
- Clarke AL, Øien IJ, Honza M, Moksnes A, Røskift E. Factors affecting reed warbler risk of brood parasitism by the common cuckoo. *Auk.* 2001;118:534-538.
- Brooke M de L, Davies NB. Provisioning of nestling cuckoos *Cuculus canorus* by reed warbler *Acrocephalus scirpaceus* hosts. *Ibis.* 1989;131:250-256.

Davies NB. Cuckoos, cowbirds and other cheats. T & AD Poyser, London. 2000.

de Neve L, Soler JJ, Soler M, Pérez-Contreras T. Nest size predicts the effect of food supplementation to magpie nestlings on their immunocompetence: An experimental test of nest size indicating parental ability. *Behav Ecol*. 2004;15:1031-1036.

Grim T, Tyller Z, Samaš P. Unusual diet of brood parasitic nestlings and its fitness consequences. *Auk*. 2017;134:732-750.

Hansell MH. Bird nests and construction behaviour. Cambridge University Press, Cambridge. 2000.

Hansell MH. Built by animals. Oxford University Press, Oxford. 2007.

Hauber ME. Site selection and repeatability in brown-headed cowbird (*Molothrus ater*) parasitism of eastern phoebe (*Sayornis phoebe*) nests. *Can J Zool*. 2007;79:1518-1523.

Heenan CB, Seymour RS. Structural support, not insulation, is the primary driver for avian cup-shaped nest design. *Proc R Soc B Biol Sci*. 2011;278:2924-2929.

Honza M, Šulc M, Jelínek V, Požgayová M, Procházka P. Brood parasites lay eggs matching the appearance of host clutches. *Proc R Soc B Biol Sci*. 2014;281:20132665.

Jelínek V, Procházka P, Požgayová M, Honza M. Common cuckoos *Cuculus canorus* change their nest-searching strategy according to the number of available host nests. *Ibis*. 2014;156:189-197.

Jelínek V, Procházka P, Honza M. Experimental enlargement of nest size does not increase risk of predation or brood parasitism in the great reed warbler *Acrocephalus arundinaceus*. *Ibis*. 2015;157:396-400.

Jelínek V, Požgayová M, Honza M, Procházka P. Nest as an extended phenotype signal of female quality in the great reed warbler. *J Avian Biol*. 2016;47:428-437.

Li D, Ruan Y, Wang Y, Chang AK, Wan D, Zhang Z. Egg-spot matching in common cuckoo parasitism of the Oriental reed warbler: Effects of host nest availability and egg rejection. *Avian Res*. 2016a;7:21.

Li D, Zhang Z, Grim T, Liang W, Stokke BG. Explaining variation in brood parasitism rates between potential host species with similar habitat requirements. *Evol Ecol*. 2016b;30:905-923.

Ma L, Yang C, Liu J, Zhang J, Liang W, Møller AP. Costs of breeding far away from neighbors: Isolated host nests are more vulnerable to cuckoo parasitism. *Behav Proc*. 2018;157:327-332.

Molina-Morales M, Martínez JG, Avilés JM. Criteria for host selection in a brood parasite vary depending on parasitism rate. *Behav Ecol*. 2016;27:1441-1448.

- Møller AP. Fashion and out of fashion: Appearance and disappearance of a novel nest building innovation. *Avian Res.* 2017;8:14.
- Møller AP, Adriaensen F, Artemyev A, Banbura J, Barba E, Biard C, et al. Variation in clutch size in relation to nest size in birds. *Ecol Evol.* 2014;4:3583-3595.
- Moskát C, Honza M. Effect of nest and nest site characteristics on the risk of cuckoo *Cuculus canorus* parasitism in the great reed warbler *Acrocephalus arundinaceus*. *Ecography.* 2000;23:335-341.
- Parejo D, Avilés JM. Do avian brood parasites eavesdrop on heterospecific sexual signals revealing host quality? A review of the evidence. *Anim Cogn.* 2007;10:81-88.
- Patten MA, Reinking DL, Wolfe DH. Hierarchical cues in brood parasite nest selection. *J Ornithol.* 2011;152:521-532.
- Soler JJ, Cuervo JJ, Møller AP, de Lope F. Nest building is a sexually selected behaviour in the barn swallow. *Anim Behav.* 1998;56:1435-1442.
- Soler JJ, Soler M, Møller AP, Martínez JG. Does the great spotted cuckoo choose magpie hosts according to their parenting ability? *Behav Ecol Sociobiol.* 1995;36:201-206.
- Soler M. Long-term coevolution between avian brood parasites and their hosts. *Biol Rev.* 2014;89:688-704.
- Stoddard MC, Hauber ME. Colour, vision and coevolution in avian brood parasitism. *Phil Trans R Soc B Biol Sci.* 2017;372:20160339.
- Trnka A, Prokop P. The use and function of snake skins in the nests of great reed warblers *Acrocephalus arundinaceus*. *Ibis.* 2011;153:627-630.
- Wang L, Yang C, He G, Liang W, Møller AP. Cuckoos use host egg number to choose host nests for parasitism. *Proc R Soc B Biol Sci.* 2020b;287:20200343.
- Wang L, Zhong G, He G, Zhang Y, Liang W. Egg laying behavior of the common cuckoo (*Cuculus canorus*): Data based on field video-recordings. *Zool Res.* 2020a;41:458-464.
- Yang C, Liang W, Antonov A, Cai Y, Stokke BG, Fossøy F, Moksnes A, Røskft E. Diversity of parasitic cuckoos and their hosts in China. *Chin Birds.* 2012;3:9.
- Yang C, Wang L, Liang W, Møller AP. Egg recognition as antiparasitism defence in hosts does not select for laying of matching eggs in parasitic cuckoos. *Anim Behav.* 2016a;122:177-181.
- Yang C, Wang L, Liang W, Møller AP. Do Common cuckoos (*Cuculus canorus*) possess an optimal laying behaviour to match their own egg phenotype to that of their Oriental reed warbler (*Acrocephalus orientalis*) hosts? *Biol J Linn Soc.* 2016b;117:422-427.

Figures

Figure 1

Example of combination of host nests for eliciting cuckoo parasitism. In group 1, a) large and b) small nests are shown. In group 2, A) large, B) medium, and C) small nests are shown. Arrows shown in red refer to eggs of the common cuckoo while the remainder are quail eggs.

Figure 2

Nest parasitism frequency and sample size of experimental nest types (large and small) in group 1 and (large, medium and small) in group 2 exploited by common cuckoos in the field. Probabilities are from statistical tests of the null hypothesis of similar rates of parasitism. Numbers above the bars refer to test sample.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [VideoS1.mp4](#)
- [VideoS2.mp4](#)
- [VideoS3.mp4](#)
- [VideoS4.mp4](#)

- [datafigure1.xlsx](#)
- [datafigure2.xls](#)