

Aldosterone (dietary Na⁺ depletion) up-regulates NKCC1 expression and enhances electrogenic Cl⁻ secretion in rat proximal colon

Andrew J Nickerson

University of Pittsburgh School of Health and Rehabilitation Sciences

Vazhaikkurichi M Rajendran (✉ vrajendran@hsc.wvu.edu)

West Virginia University - Health Sciences Campus: West Virginia University Health Sciences Center

<https://orcid.org/0000-0002-8999-8721>

Research Article

Keywords: Na⁺ depletion, Aldosterone, CFTR, Cl⁻ secretion, Ussing chamber

Posted Date: September 22nd, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-2051661/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Cellular and Molecular Life Sciences on July 17th, 2023. See the published version at <https://doi.org/10.1007/s00018-023-04857-x>.

Abstract

Aldosterone markedly enhances K^+ secretion throughout the colon, a mechanism critical to its role in maintaining overall K^+ balance. Previous studies demonstrated that basolateral NKCC1 was up-regulated by aldosterone in the distal colon specifically to support K^+ secretion – distinct from the more well-established role of NKCC1 in supporting luminal Cl^- secretion. However, considerable segmental variability exists between proximal and distal ion transport processes, especially concerning their regulation by aldosterone. Furthermore, delineating such region-specific effects has important implications for the management of various gastrointestinal pathologies. Experiments were therefore designed to determine whether aldosterone similarly up-regulates NKCC1 in the proximal colon to support K^+ secretion. Using dietary Na^+ depletion as a model of secondary hyperaldosteronism in rats, we found that proximal colon NKCC1 expression was indeed enhanced by aldosterone (i.e., Na^+ depletion). Surprisingly, electrogenic K^+ secretion was not detectable by short-circuit current (I_{SC}) measurements, in response to either basolateral bumetanide (NKCC1 inhibitor) or luminal Ba^{2+} (non-selective K^+ channel blocker), despite enhanced K^+ secretion in Na^+ -depleted rats, as measured by $^{86}Rb^+$ fluxes. Expression of BK and IK channels was also found to be unaltered by dietary Na^+ depletion. However, bumetanide-sensitive basal and agonist-stimulated Cl^- secretion (I_{SC}) were significantly enhanced by Na^+ depletion, as was CFTR Cl^- channel expression. These data suggest that NKCC1-dependent secretory pathways are differentially regulated by aldosterone in proximal and distal colon. Development of therapeutic strategies in treating pathologies related to aberrant colonic K^+/Cl^- transport – such as pseudo-obstruction or ulcerative colitis – may benefit from these findings.

INTRODUCTION

Aldosterone is a primary regulator of colonic epithelial ion transport, serving to enhance Na^+ and water absorption, and simultaneously induce K^+ secretion – two functions which are critical to overall fluid and electrolyte homeostasis (1–3). A key feature of aldosterone is that its effects on colonic transport processes are known to vary substantially by anatomical region within the colon. For example, aldosterone inhibits electroneutral NaCl absorption and stimulates electrogenic Na^+ absorption through the epithelial Na^+ channel (ENaC) in the distal colon (4–6), while enhancing electroneutral (Na^+/H^+ exchanger 3; NHE-3)-dependent Na^+ absorption in the proximal colon (7, 8). With respect to K^+ transport, aldosterone enhances K^+ secretion in both distal and proximal regions of the colon (1, 9, 10), although there may be differences in the secretory pathways and channels/transporters involved (11).

In the distal colon, expression of the apical membrane large conductance K^+ (BK) channel is increased by aldosterone (15–17), correlating to enhanced BK-mediated K^+ secretion as measured by isotopic flux assays (18) or electrophysiological recordings (16, 19). Aldosterone-induced apical K^+ secretion is supported by basolateral K^+ uptake via Na^+/K^+ ATPase and $Na^+/K^+/2Cl^-$ co-transporter 1 (NKCC1) (17). $^{86}Rb^+$ (K^+ surrogate) flux experiments have established that net K^+ secretion is also enhanced by

aldosterone in the proximal colon (9, 20). Still, the mechanism of aldosterone-induced K^+ secretion in the proximal colon is not known, apart from recent functional and immunohistochemical evidence against a role for aldosterone in the regulation of BK channels or BK-mediated K^+ currents in rat proximal colon (11).

Recently, NKCC1, which facilitates basolateral K^+ uptake, was shown to be functionally up-regulated by aldosterone via the prevention of its proteasomal degradation in colonic epithelial cells *in vitro* (21, 22). Beyond this, studies from our group have demonstrated that aldosterone transcriptionally enhances NKCC1 expression both *in vivo* and *ex vivo*, specifically to support K^+ secretion through the apical membrane BK channels in rat distal colon (23). However, it is still unknown whether aldosterone controls NKCC1 expression and/or function within the proximal region of the tissue. The present study was therefore designed to test the hypothesis that aldosterone similarly up-regulates NKCC1 to support enhanced K^+ secretion in rat proximal colon. Our results indicate that NKCC1 expression is indeed upregulated by aldosterone (using the model of dietary Na^+ depletion), but that this does not support electrogenic K^+ secretion. Rather, basal and agonist-stimulated Cl^- secretion is enhanced by dietary Na^+ depletion, concurrent with an increase in NKCC1, as well as cystic fibrosis transmembrane conductance regulator (CFTR) protein abundance. Thus, the involvement of NKCC1 in aldosterone-regulated K^+ and Cl^- secretory pathways appears to be specific to the distal and proximal colon, respectively.

MATERIALS and METHODS

Animals

Male Sprague-Dawley rats (126–150 g; Charles River, Wilmington, MA) were housed 2 per cage and maintained in 12-hour light/dark cycles in temperature-controlled housing. Rats were randomly assigned to groups receiving either standard rodent chow or Na^+ -deficient diet (MP Biomedicals #02960364) for 7 days to induce secondary hyperaldosteronism, as has been described previously (2, 10, 24, 25). All rats were allowed free access to water throughout. On the day of experiments, rats were rapidly anesthetized with 5% isoflurane and maintained under a deep plane of anesthesia with 3% isoflurane delivered via nose cone. Colons were removed by cutting away mesenteric attachments and excising tissue from the caecum to the rectum and were subsequently flushed with ice cold saline prior to use in downstream experiments. After tissue removal, rats were euthanized by cutting the diaphragm. All procedures used in this study were approved by the West Virginia University Institutional Animal Care and Use Committee prior to the start of the project.

Plasma aldosterone measurements

Plasma aldosterone levels from normal and Na^+ -depleted rats were determined using an ELISA kit (IBL-America #IB79134). Aldosterone concentrations for each sample were calculated by fitting sample absorbance values to a standard curve using non-linear regression in GraphPad Prism 9.0 software, as outlined in the kit's instructions.

Ussing chamber experiments: After flushing, the colon was placed on a dissection stage and submerged in ice cold HCO_3^- Ringer's solution containing the following, in mM: 140 Na^+ , 119.8 Cl^- , 25 HCO_3^- , 5.2 K^+ , 1.2 Ca^{2+} , 1.2 Mg^{2+} , 2.4 HPO_4^{2-} , 0.4 H_2PO_4^- , and 10 glucose (pH: 7.41). The solution was continuously bubbled with 5% CO_2 balanced with O_2 to maintain pH at 7.41. The colon was opened with surgical scissors along the mesenteric border and placed mucosal side facing downward. A lateral incision was made with a razor blade just beneath the longitudinal and circular smooth muscle layers, which were peeled away to leave a submucosa/muscularis mucosae/mucosa preparation.

Segments of proximal colon, identified by striation patterns in the tissue, were then mounted on 1.1 cm^2 plastic sliders for use in an Ussing-style recording chamber (Physiologic Instruments, San Diego, CA). The recording chambers were equipped with one pair of voltage-sensing AgCl pellet electrodes and one pair of current-injecting AgCl wire electrodes, each connected to the chamber bath via agar salt bridge (3.5% agar in 3M KCl). The chamber bath itself was temperature controlled and contained 5 mL of bubbled HCO_3^- Ringer's solution in each chamber half. A multichannel voltage clamp (VCC MC8, Physiologic Instruments, San Diego, CA) was operated automatically by computer software to measure short-circuit current (I_{SC}), a function of electrogenic ion transport. By convention, positive I_{SC} values correspond to anion secretion/cation absorption and negative values correspond to cation secretion/anion absorption. Intermittent bi-polar 200 ms, 5mV pulses were applied to monitor trans-epithelial conductance (G_{TE}) at 30 second intervals.

For $^{86}\text{Rb}^+$ (PerkinElmer, Billerica, MA) flux experiments, 2 μCi of isotope was added to either the mucosa or serosal chamber bath in tissue pairs matched by conductance values within 20% of each other (26). Following a 20-minute equilibration period, 500 μL samples were taken from the chamber bath opposite to where the isotope was added (i.e., the "sink" side), as well as 10 μL from the isotope bath ("hot" side) to be used in the calculation of basal K^+ transport. Samples were mixed with scintillation cocktail solution (Sigma Aldrich, #03999-5L) and counted using a Tri-carb 4910TR liquid scintillation counter (Perkin Elmer, Waltham MA). Unidirectional $^{86}\text{Rb}^+$ flux rates were calculated from the measured counts per minute (cpm) values using the following equation:

$$\text{K}^+ \text{ flux (J) } (\mu\text{Eq}/\text{cm}^2 \cdot \text{hr}) = (5 \times \text{B} - [\text{A} \times 0.9]) / ([\text{S}/5.2] \times 1.1 \times 0.25)$$

"A" and "B" represent cpm/mL values measured from the beginning and end of the flux period, respectively. 5 is the total volume (in mL) of the chamber bath and 0.9 is dilution factor accounting for removal of 500 μL of bath volume upon sample collection. "S" represents the cpm/ μEq (in 1 mL) measured from the standard obtained from the "hot" bath of the chamber. 5.2 is the K^+ concentration in the bath (mM = mMol/L = $\mu\text{Eq}/\text{mL}$), 1.1 corresponds to the slider aperture (in cm^2) and 0.25 is the time (in hours) for the flux period (26). Net K^+ fluxes were calculated by subtracting the serosal to mucosal (S-M) from the mucosal to serosal (M-S) fluxes from paired tissues.

Mucosal Scraping: Proximal colon sacs were made by filling with ice cold buffer containing (in mM): 15.4 NaCl, 8 HEPES/Tris (pH 7.5), 5 ethylenediaminetetraacetic acid (EDTA), 0.5 dithiothreitol (DTT), and 0.5 phenylmethylsulphonyl fluoride (PMSF) and incubated in the same solution on ice for 45 minutes. The drained sacs were opened along the mesenteric border. The proximal mucosal layer was scraped off using glass slides and frozen immediately in liquid nitrogen until use. Scrapped mucosae were used for RNA isolation and protein extraction.

RNA isolation and qRT-PCR analysis

Total RNA was isolated from frozen colonic mucosa (50 mg) using the Trizol method and quantitated with NanoDrop spectrometer (Thermo Sci. ND2000C). Quantitative reverse transcriptase polymerase chain reaction (qRT-PCR) was performed using a One-Step reaction kit (NEB #E3005) per the manufacturer's protocol. Custom gene-specific primers were designed using published DNA sequences (NCBI database) and synthesized by Thermo Scientific (sequences listed in Table 1). qRT-PCR reactions performed on a 96-well plate were read using a CFX96 real time PCR machine (BioRad). mRNA abundance was determined using the $\Delta\Delta C_t$ method, with β -actin as a reference housekeeping gene and represented in the figures as fold change in mRNA transcript expression in the experimental (Na^+ -depleted) relative to the control (normal diet) animals.

Table 1
Primer sequences used in qRT-PCR analysis.

Gene Target and Accession number	Primer Sequence	Amplicon length
NKCC1 (Slc12a2) NM_001270617.1	F: 5'- TGGCAAGACTTCAACTCAGC - 3' R: 5'- GGTATCAACAAGGTCAAACCTCC - 3'	177
BK α (Kcnma1) NM_001393699.1	F: 5'- ATGTCTACAGTGGGTTACGG - 3' R: 5'- TGGGTGGTAGTTCTTTATGG - 3'	464
IK (Kcnn4) NM_001270701.1	F: 5'- TGGCTGAGCACCAAGAGC - 3' R: 5'- TACAGCACCCACTTGCAACC - 3'	197
CFTR (Cftr) NM_031506.1	F: 5'- TTCTTCAGCTGGACCACACC - 3' R: 5'- TGGAAGCTTGTTCCCTGTCC - 3'	106
TMEM16A (Ano1) NM_001107564.1	F: 5'- TTATGGCCCTCTGGGCTCG - 3' R: 5'- CACCTCCTCTTCCTCGAAGC - 3'	102
β -actin (Actb) NM_031144.3	F: 5'- AGATCAAGATCATTGCTCCTCC - 3' R: 5'- AGTAACAGTCCGCCTAGAAGC - 3'	165

Western blot analysis: Proteins were extracted from frozen colonic mucosae (50 mg) by homogenizing in ice-cold lysis buffer (150 mM NaCl, 50 mM Tris base, 0.5% sodium deoxycholate, 0.1% SDS, 1.0% Triton-X100) plus protease inhibitor cocktail (Thermo #A32965), followed by a brief sonication (Fisher Sonic Dismembrator Model 100). Samples were centrifuged for 20 minutes at 12,000 x g and the supernatant was mixed with 4 x Laemmli buffer (Life Technologies #NP0007) and heated to 90°C for 5 minutes. Samples were then chilled on ice before adding β -mercaptoethanol (5% w/v), and the aliquoted samples were flash frozen and stored at -80°C. Proteins (~ 20 μ g) were resolved on 8% Tris-glycine polyacrylamide gels and transferred on to PVDF membranes. Membranes were incubated in blocking buffer (3% BSA in tris-buffered saline plus 0.1% Tween-20 (TBS-T) for 1 hour at room temperature, followed by overnight incubation at 4°C in primary antibody solution. Table 2 provides the source and dilutions used for all primary antibodies. The membranes were then washed in TBST (5 minutes x 5) and incubated in TBST + 3% BSA containing goat anti-rabbit HRP-conjugated secondary antibody (Thermo #31462; 1:20,000 dilution) for 1 hour at room temperature before being washed again in TBST (5 min x 5). Immune complexes were detected using enhanced chemiluminescence (West Dura, Thermo #34075). Images were captured using a G:BOX (Syngene) and band intensities were quantified by densitometry using Image J software and normalized to β -actin as a loading control.

Table 2
Primary antibody product and usage information.

Target	Host Species	Supplier	Dilution (Application)
NKCC1	Rabbit	Cell Signaling	1:500 (WB)
			1:500 (IF)
BK α	Rabbit	Alomone	1:250 (WB)
IK	Rabbit	Gift from Dr. Michael Kashgarian	1:3,000 (WB)
CFTR	Rabbit	Alomone	1:250 (WB)
TMEM16A	Rabbit	Alomone	1:250 (WB)
β -actin	Mouse	SCBT	1: 1,000 (WB)
	(HRP-conjugated)		

Immunofluorescence microscopy

Segments of whole proximal colon were fixed overnight at 4°C in 4% paraformaldehyde. The next day, tissues were washed in PBS, flash frozen in isopentane cooled with liquid nitrogen and embedded in tissue freezing medium. 10 μ m sections were cut with a cryostat and mounted on charged glass slides. 40 minutes of heated antigen retrieval was performed by placing the slides in 10 mM citrate + 0.05% Tween-20 buffer and microwaving on very low power for 45 minutes before allowing gradual cooling to room temperature. Sections were then permeabilized in PBS + 5% goat serum + 0.1% Triton-X for 15 minutes. Afterward, primary antibody solution containing PBS + 0.01% Triton-X (PBST) + 5% goat serum

with a 1:500 dilution of NKCC1 primary antibody (Cell Signaling #14581) (23, 27) was placed on the sections overnight at 4°C. The next day, tissues were washed 5 times for 5 minutes each in PBST and subsequently incubated in PBST + 5% goat serum containing goat anti-rabbit AlexaFluor 488 secondary antibody (Thermo #11008; 1:2,00 dilution) for 1 hour at room temperature. After washing an additional 5 times for 5 minutes each, slides were mounted with SlowFade Diamond antifade mountant with DAPI (Thermo #36971) and sealed with clear fingernail polish. IF images were captured using a Zeiss 710 confocal microscope at 20x magnification (Carl Zeiss, Oberkochen, Germany).

RESULTS

Dietary Na⁺ depletion induces secondary hyperaldosteronism: Dietary Na⁺ depletion is a well-established model of secondary hyperaldosteronism that has been extensively characterized (24). Previous experiments from our lab using this model have demonstrated that plasma aldosterone levels increase ~10-fold after 7 days of Na⁺-deficient diet administration (23). To confirm that dietary Na⁺ depletion induced secondary hyperaldosteronism in the present study, plasma aldosterone levels were measured via ELISA. Na⁺ depleted rats exhibited a robust increase in aldosterone compared to normal diet-fed controls ($4,154 \pm 966$ vs 497 ± 151 pg/mL; $p < 0.05$) (Supplementary Fig. S1).

Dietary Na⁺ depletion enhances NKCC1 protein expression in proximal colon: Initial experiments were aimed at evaluating the effect of aldosterone (dietary Na⁺ depletion) on the expression of NKCC1 in the proximal colon. Western blot analysis of normal and Na⁺-depleted rat proximal colonic mucosae revealed a ~2-fold increase in NKCC1 protein abundance (Fig. 1A-B) ($p < 0.05$). However, no significant change in mRNA expression was detected by qRT-PCR (Fig. 1C) ($p = 0.32$). Immunofluorescent labeling of NKCC1 congruent with our western blot analysis, as the signal was more prominent in Na⁺-depleted tissues, particularly in the crypt region (Fig. 2). The observed ~2-fold increase in NKCC1 protein abundance mirrors the effect previously described in the distal colon using the same dietary model system (23). However, the present discrepancy between changes in NKCC1 mRNA transcript and protein abundance in the proximal colon suggests that the upregulation of NKCC1 in proximal colon may not be the result of a transcriptional mechanism, as was previously demonstrated in the distal colon (23). While the nature of this phenomenon was not further explored here, a similar pattern has been observed in cultured epithelial cells in response to aldosterone (21, 22). This was attributable to changes in protein turnover (e.g., the proteasomal pathway), providing a possible explanation for our current findings.

NKCC1 does not support electrogenic bumetanide- or Ba²⁺-sensitive K⁺ secretion in normal or Na⁺-depleted rat proximal colon: Experiments were next designed to assess the role of NKCC1 in supporting electrogenic K⁺ secretion in normal and Na⁺ depleted rat proximal colon, measured as short-circuit current (I_{SC}). Recording chambers were configured such that luminal cation and anion secretion are represented by negative and positive I_{SC} values, respectively (16, 28). Baseline I_{SC} was higher in proximal colon segments from Na⁺ depleted rats compared to controls (128.6 ± 16.8 vs. 89.2 ± 8.7 $\mu\text{A}/\text{cm}^2$; $p < 0.05$) (Fig. 3C), as previously shown (7). Surprisingly, we were unable to detect electrogenic K⁺ secretion that

was sensitive to serosal bumetanide (NKCC1 inhibitor) or mucosal Ba^{2+} (non-selective K^+ channel blocker). In fact, bumetanide substantially inhibited I_{SC} , likely corresponding to a reduction in NKCC1-dependent Cl^- secretion (Fig. 3B, D). However, we could not determine that bumetanide-sensitive I_{SC} was greater in Na^+ -depleted animals as compared to control ($\Delta I_{\text{SC}(\text{BUMET})} = -51.4 \pm 8.3$ vs. $-28.4 \pm 7.8 \mu\text{A}/\text{cm}^2$; $p = 0.07$) (Fig. 3D), likely due to large variability in starting I_{SC} values. Although a significant difference was not observed here, the data generally suggest that Na^+ -depletion (aldosterone) enhances NKCC1 expression to support Cl^- secretion, rather than K^+ secretion, in proximal colon.

Mucosal Ba^{2+} also induced a slight negative deflection in I_{SC} in Na^+ -depleted, but not in normal proximal colon ($\Delta I_{\text{SC}(\text{Ba}^{2+})} = -12.3 \pm 3.1$ vs. $-2.9 \pm 2.6 \mu\text{A}/\text{cm}^2$; $p < 0.05$) (Fig. 3E), although this effect was subtle. The absence of measurable Ba^{2+} -sensitive K^+ secretion was unexpected because $^{86}\text{Rb}^+$ flux studies (K^+ tracer) have established that net luminal K^+ secretion is present in the proximal colon and is markedly enhanced by dietary Na^+ depletion (Supplementary Fig. S2) (10) or by aldosterone directly (20). Our electrophysiological data therefore imply that aldosterone-induced K^+ secretion in the proximal colon is either not electrogenic or may proceed via some Ba^{2+} -insensitive pathway. Unfortunately, $^{86}\text{Rb}^+$ was taken off market during this study, and thus a thorough interrogation of a potentially yet unexplored K^+ secretory pathway could not be conducted. Nonetheless, if NKCC1 does support luminal K^+ secretion in the proximal colon to some extent, it appears to be overshadowed by a more prominent role in supporting Cl^- secretion.

Dietary Na^+ depletion does not enhance mucosal BK and IK channel expression in proximal colon: We next sought to determine whether mucosal membrane K^+ channels were detectable in proximal colon tissue lysates and whether their expression was altered by Na^+ -depletion. Perry et. al. reported the presence of large conductance K^+ (BK) channels in patch clamp recording from proximal colonocytes (11). In their study, dietary K^+ loading – another maneuver that induces secondary hyperaldosteronism – did not increase the number of functional K^+ channels in the proximal colon detectable by patch clamp, Western blot analysis or immunohistochemistry. This was in sharp contrast to effects observed in the distal colon (11).

Expression of intermediate conductance K^+ (IK) channels, which are also present in the proximal colon in rat, was not quantified in that study. Further, because the models of dietary Na^+ depletion and K^+ loading are not equivalent, quantification of both BK and IK channel expression was performed here. Neither BK nor IK abundance was found to be altered in Na^+ -depleted animals, as detected by Western blot (Fig. 4A-C) or qRT-PCR (Fig. 4D,E), corroborating the findings of Perry et. al. Interestingly, although electrogenic, Ba^{2+} -sensitive K^+ secretion was not observed in either normal or Na^+ -depleted rat proximal colon via I_{SC} recordings, both BK and IK channels were detected at both the mRNA and protein level. Thus, it is possible that BK and IK channels may not facilitate luminal K^+ secretion in the proximal colon, but rather

serve some other non-vectorial function(s), such as participating in K^+ recycling at the basolateral membrane.

Dietary Na^+ depletion enhances NKCC1-dependent, Ca^{2+} - and cAMP-stimulated Cl^- secretion in proximal colon: NKCC1 is known to support aldosterone-induced K^+ secretion in the distal colon (17, 23, 29) and other tissues (22, 30). In the proximal colon, basal I_{SC} is elevated by aldosterone (7) (Fig. 3), suggesting enhanced Cl^- secretion. Indeed, the most well-defined role of NKCC1 throughout the gastrointestinal tract is that of supporting basal and stimulated Cl^- secretion to the lumen in response to various stimuli (31, 32–34). We therefore aimed to expand upon our earlier observations by assessing whether pharmacologically stimulated Cl^- secretion, in addition to basal secretion, was also augmented in Na^+ -depleted animals.

Carbamoylcholine (carbachol; muscarinic agonist) and forskolin (adenylate cyclase activator) were chosen because they induce Ca^{2+} - and cAMP-dependent Cl^- secretion, respectively, representing the two predominant secretory pathways in intestinal epithelia. The responses to both carbachol [ΔI_{SC} (CCH): 239.6 ± 5.6 vs 105.7 ± 33.5 $\mu A/cm^2$; $p < 0.05$] and forskolin [ΔI_{SC} (FSK): 154.3 ± 8.7 vs. 115.3 ± 8.7 $\mu A/cm^2$; $p < 0.05$] were enhanced in Na^+ -depleted rat proximal colon compared to controls (Fig. 5A,5B). Responses to both carbachol and forskolin were also attenuated by the presence of bumetanide in the serosal bath. (Fig. 5A,B). These data suggest that increased NKCC1 expression correlates to an increased capacity for both Ca^{2+} - and cAMP-stimulated Cl^- secretion. In addition, cAMP-induced Cl^- secretion was significantly blunted by bumetanide in normal animals (ΔI_{SC} (FSK): 154.3 ± 8.7 vs. 61.8 ± 2.5 $\mu A/cm^2$; $p < 0.01$) (Fig. 5B), whereas Ca^{2+} -induced Cl^- secretion was not (ΔI_{SC} (CCH): 105.7 ± 33.5 vs. vs 71.0 ± 21.9 $\mu A/cm^2$; $p = 0.81$) (Fig. 5A). This discrepancy may reflect a greater contribution of HCO_3^- secretion in the response to Ca^{2+} versus cAMP signaling.

Dietary Na^+ depletion enhances CFTR channel protein expression in proximal colon: Luminal Cl^- secretion in the colon is primarily mediated by apical membrane-localized CFTR Cl^- channels (35–38) with evidence of some involvement of TMEM16A as well (38–40). Because basolateral NKCC1 protein expression and basal/stimulated Cl^- secretion were all enhanced by Na^+ depletion, we sought to determine if there was a parallel increase in the expression of either of these two Cl^- channel proteins. Western blot analysis revealed that CFTR protein, and in particular the mature, glycosylated form (“c” band) (41) was increased ~ 1.5 fold in Na^+ -depleted rat proximal colonic mucosae (Fig. 6A,D; $p < 0.05$). CFTR mRNA transcript abundance was not significantly increased (Fig. 6E). Interestingly, TMEM16A transcript (Fig. 6E) but not protein (Fig. 6B,D) was detected in these samples. The TMEM16A antibody used here has been validated previously by our lab and others (42). As a positive control, distal colonic mucosae tissue lysates were probed alongside proximal colon samples. TMEM16A was detected only in the distal colon (Fig. 6C), suggesting possible regional variability in the expression and/or functional role of TMEM16A in colonic epithelia. Therefore, CFTR is likely the apical membrane channel that mediates

the basal and agonist-stimulated Cl^- secretion that is enhanced by dietary Na^+ depletion in the proximal colon.

DISCUSSION

Our findings indicate that the expression of basolateral NKCC1 in rat proximal colon was enhanced by dietary Na^+ depletion, a model of hyperaldosteronism. Further, the increase in NKCC1 protein abundance was correlated with an augmented capacity for electrogenic Cl^- secretion, rather than K^+ secretion. This is in stark contrast to the effects of aldosterone on K^+ and Cl^- transport processes in the distal colon – where K^+ , but not Cl^- secretion is enhanced – despite a similar increase in NKCC1 expression. A parallel increase in the apical membrane CFTR Cl^- channel protein was observed, whereas changes in BK and IK channel expression were not found. This also differs from the effects of aldosterone on the distal colon. Segmental variability within the two regions of the colon has been well-documented in terms of the effects of corticosteroid hormones, including their influence of Na^+ , K^+ , and Cl^- transport. Here, we report a novel example of differential regulation of proximal and distal colonic ion transport systems by aldosterone.

The results of this study have clinical relevance because colonic K^+ and Cl^- transport is a critical aspect of both gastrointestinal (G.I.) and non-G.I. pathologies. For example, in patients with end-stage renal disease (ESRD), the colon provides an alternative outlet for K^+ which can prevent life-threatening hyperkalemia (43,44). On the other hand, colonic pseudo-obstruction (45) and ulcerative colitis (UC) (12, 46, 47) are characterized by excessive fecal K^+ losses that contribute to the manifestation of diarrhea. A mineralocorticoid receptor antagonist has even been used clinically in at least one case of pseudo-obstruction (Ogilvie's syndrome) to relieve diarrheal symptoms associated with excessive stool K^+ output (48). Other diarrheal pathologies such as enteroviral or enterobacterial infection primarily affect Cl^- secretion independently of K^+ secretion (49, 50). Given the high degree of variability between colonic regions in terms of aldosterone-regulated K^+ and Cl^- transport, results from this and related studies may help with management of such pathologies.

The overall function of aldosterone signaling in the colon is to promote salt and fluid retention, while also driving the removal of excess K^+ from the body (51). Increased NKCC1 expression that may be tied only to Cl^- secretion appears counterproductive because up-regulation of this pathway would hinder fluid absorption, in theory. However, Cl^- absorption in the proximal colon is known to be electroneutral (mediated primarily by DRA) and occurs mostly in surface absorptive cells (7). On the other hand, NKCC1 and CFTR are known to be expressed primarily within the crypt region (34). $^{36}\text{Cl}^-$ flux studies in the past have shown an increase in net Cl^- absorption within the proximal colon of Na^+ -depleted rats compared to controls. However, the magnitude of Cl^- flux did not match that of Na^+ (7, 9), even though the transport of these two ions is functionally coupled through Na^+/H^+ and $\text{Cl}^-/\text{HCO}_3^-$ exchangers. The increase in basal I_{SC} measured in those studies as well as ours, along with the augmented response to secretagogues (e.g.,

those derived from basal neurogenic input), may account for the discrepancy in net Na^+ versus Cl^- fluxes. In other words, a portion of the Cl^- absorbed through surface epithelial cells may ultimately be “recycled” back to the lumen via secretion occurring within the crypt. Despite this possible explanation, the physiological significance of having an enhanced capacity for Cl^- secretion under hyper-aldosteronaemic conditions remains unclear.

Another puzzling observation is that although the net secretion of K^+ is enhanced by aldosterone in the proximal colon, as indicated by $^{86}\text{Rb}^+$ flux experiments (Supplementary Fig. 2) (20), the nature of this lumenally-directed K^+ transit pathway has yet to be identified. Our data indicate that at least electrogenic K^+ secretion in both normal and Na^+ -depleted rat proximal colon is insensitive to Ba^{2+} (a non-selective K^+ channel blocker) because no effect on I_{SC} was observed. Other experiments from our lab indicate that basal and cAMP-stimulated K^+ secretion in normal proximal colon is also unaffected by other K^+ channel inhibitors (e.g., iberiotoxin, charybdotoxin, Tram-34, paxilline) as measured by I_{SC} and $^{86}\text{Rb}^+$ fluxes (unpublished observations). The apparent insensitivity to K^+ channel inhibitors/blockers, especially Ba^{2+} , raises the possibility that luminal K^+ secretion in the proximal colon is not facilitated an electrogenic process, but rather through some other secretory mechanism.

Perry et. al. examined BK channel activity in normal and hyper-aldosteronaemic (K^+ -loaded) rat proximal colonocytes by whole-cell and single channel patch clamp recording (11). Although this approach allows for fine resolution of the biophysical properties of K^+ channels for the purpose of identifying specific channels, vectorial transport functions cannot be measured using this technique. Nonetheless, our data are congruent with the findings from that study, as BK (and IK) channels were detectable, but aldosterone had no apparent effect on the expression of either. It is possible that the function of these K^+ channels in the proximal colon is unrelated to luminal electrogenic K^+ secretion.

In conclusion, here we have presented novel evidence of the up-regulation of NKCC1 and Cl^- secretory capacity by aldosterone in the rat proximal colon. This effect appears to be unrelated to electrogenic K^+ secretion, which differs from the effects of aldosterone on K^+ and Cl^- transport in the distal colon. Segmental differences should thus be accounted for when investigating or managing aldosterone-induced changes to colonic K^+ and Cl^- transport processes in research or clinical settings.

Abbreviations

ENaC - epithelial Na^+ channel

NKCC1 - $\text{Na}^+/\text{K}^+/\text{2Cl}^-$ co-transporter 1

BK - large conductance K^+ channel

IK - intermediate conductance K^+ channel

CFTR - cystic fibrosis transmembrane conductance regulator

EDTA - 5 ethylenediaminetetraacetic acid

DTT - dithiothreitol

PMSF - phenylmethylsulphonyl fluoride

CCH - carbachol

FSK - forskolin

I_{SC} – short circuit current

G.I - gastrointestinal

Declarations

Ethics approval and consent to participate: All animal experimental procedures used in this study were approved by the West Virginia University Institutional Animal Care and Use Committee prior to the start of the project.

Consent for publication: N/A

Availability of data and material: Data will be available on reasonable request

Competing interests: The authors declare no conflict of interest

Funding: This study was supported by the National Institute of Health NIDDK grants DK104791 and DK112085 grants to VMR

Authors' contributions: AJN conceived the idea, performed experiments, wrote manuscript. VMR conceived the idea and got funding. Both AJN and VMR discussed and edited the manuscript.

Acknowledgements: We acknowledge the technical assistance rendered by Avinash Elangovan.

Authors' information (optional): Andrew J. Nikerson, PhD., Postdoctoral Fellow, Kleyman lab, Department of Medicine – Renal Section, University of Pittsburgh, Pittsburgh, Pennsylvania, USA.

Vazhaikurichi M. Rajendran, PhD. Professor of Biochemistry and Molecular Medicine, and Professor of Medicine – Section of Gastroenterology and Hepatology, West Virginia University School of Medicine, Morgantown, West Virginia, USA.

References

1. Foster ES, Jones WJ, Hayslett JP, Binder HJ. Role of Aldosterone and Dietary Potassium in Potassium Adaptation in the Distal Colon of the Rat. *Gastroenterology*. 1985;88(1):41-46. doi:10.1016/S0016-5085(85)80130-X
2. Halevy J, Budinger ME, Hayslett JP, Binder HJ. Role of Aldosterone in the Regulation of Sodium and Chloride Transport in the Distal Colon of Sodium-Depleted Rats. *Gastroenterology*. 1986;91(5):1227-1233. doi:https://doi.org/10.1016/S0016-5085(86)80021-X
3. Rechkemmer G, Halm DR. Aldosterone stimulates K secretion across mammalian colon independent of Na absorption. *Proc Natl Acad Sci*. 1989;86(January):397-401.
4. Amasheh S, Epple HJ, Mankertz J, et al. Differential Regulation of ENaC by Aldosterone in Rat Early and Late Distal Colon. *Ann N Y Acad Sci*. 2006;915(1):92-94. doi:https://doi-org.wvu.idm.oclc.org/10.1111/j.1749-6632.2000.tb05227.x
5. Greig ER, Baker EH, Mathialahan T, Boot-Handford RP, Sandle GI. Segmental variability of ENaC subunit expression in rat colon during dietary sodium depletion. *Pflugers Arch Eur J Physiol*. 2002;444(4):476-483. doi:10.1007/s00424-002-0828-7
6. Binder HJ, Foster ES, Budinger ME, Hayslett JP. Mechanism of electroneutral sodium chloride absorption in distal colon of the rat. *Gastroenterology*. 1987;93(3):449-455. doi:10.1016/0016-5085(87)90905-X
7. Foster ES, Budinger ME, Hayslett JP, Binder HJ. Ion transport in proximal colon of the rat. Sodium depletion stimulates neutral sodium chloride absorption. *J Clin Invest*. 1986;77(1):228-235. doi:10.1172/JCI112281
8. Ikuma M, Kashgarian M, Binder HJ, Rajendran VM. Differential regulation of NHE isoforms by sodium depletion in proximal and distal segments of rat colon. *Am J Physiol - Gastrointest Liver Physiol*. 1999;276(2 39-2):539-549. doi:10.1152/ajpgi.1999.276.2.g539
9. Tumamian SG, Binder HJ. Regulation of active sodium and potassium transport in the distal colon of the rat. *J Clin Invest*. 1989;84(6):1924-1929.
10. Foster S, Hayslett JP, Binder J, Emily S, Hayslew JP, Henry J. Mechanism of active potassium absorption and secretion in the rat colon. 1984.
11. Perry MD, Rajendran VM, MacLennan KA, Sandle GI. Segmental differences in upregulated apical potassium channels in Mammalian colon during potassium adaptation. *Am J Physiol - Gastrointest Liver Physiol*. 2016;311(5):G785-G793. doi:10.1152/ajpgi.00181.2015
12. Rajendran VM, Sandle GI. Colonic Potassium Absorption and Secretion in Health and Disease. *Compr Physiol*. 2018;8(4):1513-1536. doi:10.1002/cphy.c170030
13. Rajendran VM, Singh SK, Geibel J, Binder HJ. Differential localization of colonic H⁺-K⁺-ATPase isoforms in surface and crypt cells. *Am J Physiol - Gastrointest Liver Physiol*. 1998;274(2 37-2):424-429. doi:10.1152/ajpgi.1998.274.2.g424
14. Spicer Z, Clarke LL, Gawenis LR, Shull GE. Colonic H⁽⁺⁾-K⁽⁺⁾-ATPase in K⁽⁺⁾ conservation and electrogenic Na⁽⁺⁾ absorption during Na⁽⁺⁾ restriction. *Am J Physiol Gastrointest Liver Physiol*. 2001;281(6):G1369-77. doi:10.1152/ajpgi.2001.281.6.G1369

15. Sausbier M, Matos JE, Sausbier U, et al. Distal colonic K⁺ secretion occurs via BK channels. *J Am Soc Nephrol*. 2006;17(5):1275-1282. doi:10.1681/ASN.2005101111
16. Sørensen M V, Matos JE, Sausbier M, et al. Aldosterone increases KCa1.1 (BK) channel-mediated colonic K⁺ secretion. *J Physiol*. 2008;586(Pt 17):4251-4264. doi:10.1113/jphysiol.2008.156968
17. Sweiry JH, Binder HJ. Characterization of aldosterone-induced potassium secretion in rat distal colon. *J Clin Invest*. 1989;83(3):844-851. doi:10.1172/JCI113967
18. Singh SK, O'Hara B, Talukder JR, Rajendran VM. Aldosterone induces active K⁺ secretion by enhancing mucosal expression of Kcnn4c and Kcnma1 channels in rat distal colon. *Am J Physiol - Cell Physiol*. 2012;302(9):1353-1360. doi:10.1152/ajpcell.00216.2011
19. Lomax B, McNicholas M, Lombes M, Sandle I, McNicholas M. Aldosterone-induced apical Na⁺ and K⁺ conductances are located primarily in surface cells in rat distal colon. 1994;29:G71-G82.
20. Turnamian SG, Binder HJ. Aldosterone and glucocorticoid receptor-specific agonists regulate ion transport in rat proximal colon. *Am J Physiol*. 1990;258(3 Pt 1):G492-8. doi:10.1152/ajpgi.1990.258.3.G492
21. Bazard P, Ding B, Chittam HK, et al. Aldosterone up-regulates voltage-gated potassium currents and NKCC1 protein membrane fractions. *Sci Rep*. 2020;10(1):1-14. doi:10.1038/s41598-020-72450-4
22. Ding B, Frisina RD, Zhu X, Sakai Y, Sokolowski B, Walton JP. Direct control of Na⁽⁺⁾-K⁽⁺⁾-2Cl⁽⁻⁾-cotransport protein (NKCC1) expression with aldosterone. *Am J Physiol Cell Physiol*. 2014;306(1):C66-75. doi:10.1152/ajpcell.00096.2013
23. Nickerson AJ, Rajendran VM. Aldosterone up-regulates basolateral Na⁽⁺⁾ -K⁽⁺⁾ -2Cl⁽⁻⁾ cotransporter-1 to support enhanced large-conductance K⁽⁺⁾ channel-mediated K⁽⁺⁾ secretion in rat distal colon. *FASEB J Off Publ Fed Am Soc Exp Biol*. 2021;35(5):e21606. doi:10.1096/fj.202100203R
24. Martin RS, Jones WJ, Hayslett JP. Animal model to study the effect of adrenal hormones on epithelial function. *Kidney Int*. 1983;24(3):386-391. doi:10.1038/ki.1983.171
25. Edmonds CJ, C.J. Edmonds C.L. Willis. THE EFFECT OF DIETARY SODIUM AND POTASSIUM INTAKE ON POTASSIUM SECRETION AND KINETICS IN RAT DISTAL COLON. *J Physiol*. 2019;53(9):1689-1699.
26. Clarke LL. A guide to Ussing chamber studies of mouse intestine. *AJP Gastrointest Liver Physiol*. 2009;296(6):G1151-G1166. doi:10.1152/ajpgi.90649.2008
27. Hu D, Yu Z-L, Zhang Y, et al. Bumetanide treatment during early development rescues maternal separation-induced susceptibility to stress. *Sci Rep*. 2017;7(1):11878. doi:10.1038/s41598-017-12183-z
28. Zhang J, Halm ST, Halm DR. Role of the BK channel (KCa1.1) during activation of electrogenic K⁺ secretion in guinea pig distal colon. *Am J Physiol - Gastrointest Liver Physiol*. 2012;303(12):1322-1334. doi:10.1152/ajpgi.00325.2012
29. Jiang C, Kawabe H, Rotin D. The ubiquitin ligase Nedd4L regulates the Na/K/2Cl co-transporter NKCC1/SLC12A2 in the colon. *J Biol Chem*. 2017;292(8):3137-3145. doi:10.1074/jbc.M116.770065

30. Jiang G, Cobbs S, Klein JD, O'Neill WC. Aldosterone regulates the Na-K-2Cl cotransporter in vascular smooth muscle. *Hypertens (Dallas, Tex 1979)*. 2003;41(5):1131-1135. doi:10.1161/01.HYP.0000066128.04083.CA
31. Field M. Intestinal ion transport and the pathophysiology of diarrhea. *J Clin Invest*. 2003;111(7):931-943. doi:10.1172/JCI200318326
32. Reynolds A, Parris A, Evans LA, et al. Dynamic and differential regulation of NKCC1 by calcium and cAMP in the native human colonic epithelium. *J Physiol*. 2007;582(2):507-524. doi:10.1113/jphysiol.2007.129718
33. Bachmann O, Wüchner K, Rossmann H, et al. Expression and regulation of the Na⁺-K⁺-2Cl⁻ cotransporter NKCC1 in the normal and CFTR-deficient murine colon. *J Physiol*. 2003;549(2):525-536. doi:10.1113/jphysiol.2002.030205
34. Jakab RL, Collaco AM, Ameen NA. Physiological relevance of cell-specific distribution patterns of CFTR, NKCC1, NBCe1, and NHE3 along the crypt-villus axis in the intestine. *Am J Physiol Gastrointest Liver Physiol*. 2011;300(1):G82-98. doi:10.1152/ajpgi.00245.2010
35. Greger R. Role of Cftr in the Colon. *AnnuRevPhysiol*. 2000;62:467-491. doi:10.1146/annurev.physiol.62.1.467
36. Kunzelmann K, Mehta A. CFTR: A hub for kinases and crosstalk of cAMP and Ca²⁺. *FEBS J*. 2013;280(18):4417-4429. doi:10.1111/febs.12457
37. Seidler U, Blumenstein I, Kretz A, et al. A functional CFTR protein is required for mouse intestinal cAMP-, cGMP- and CA²⁺-dependent HCO₃⁻ secretion. *J Physiol*. 1997;505(2):411-423. doi:10.1111/j.1469-7793.1997.411bb.x
38. Benedetto R, Ousingasawat J, Wanitchakool P, et al. Epithelial Chloride Transport by CFTR Requires TMEM16A. *Sci Rep*. 2017;7(1):1-13. doi:10.1038/s41598-017-10910-0
39. Rottgen TS, Nickerson AJ, Minor EA, Stewart AB, Harold AD, Rajendran VM. Dextran sulfate sodium-induced chronic colitis attenuates Ca(2+)-activated Cl(-) secretion in murine colon by downregulating TMEM16A. *Am J Physiol Cell Physiol*. 2018;315(1):C10-C20. doi:10.1152/ajpcell.00328.2017
40. Kunzelmann K, Centeio R, Wanitchakool P, et al. Control of Ion Transport by Tmem16a Expressed in Murine Intestine. *Front Physiol*. 2019;10:1262. doi:10.3389/fphys.2019.01262
41. Ramalho AS, Lewandowska MA, Farinha CM, et al. Deletion of CFTR translation start site reveals functional isoforms of the protein in CF patients. *Cell Physiol Biochem Int J Exp Cell Physiol Biochem Pharmacol*. 2009;24(5-6):335-346. doi:10.1159/000257426
42. Pineda-Farias JB, Barragán-Iglesias P, Loeza-Alcocer E, et al. Role of anoctamin-1 and bestrophin-1 in spinal nerve ligation-induced neuropathic pain in rats. *Mol Pain*. 2015;11:41. doi:10.1186/s12990-015-0042-1
43. Mathialahan T, Maclennan KA, Sandle LN, Verbeke C, Sandle GI. Enhanced large intestinal potassium permeability in end-stage renal disease. *J Pathol*. 2005;206(1):46-51. doi:10.1002/path.1750
44. Mathialahan T, Sandle GI. Dietary potassium and laxatives as regulators of colonic potassium secretion in end-stage renal disease. *Nephrol Dial Transplant*. 2003;18(2):341-347.

doi:10.1093/ndt/18.2.341

45. Van Dinter TG, Fuerst FC, Richardson CT, et al. Stimulated active potassium secretion in a patient with colonic pseudo-obstruction: A new mechanism of secretory diarrhea. *Gastroenterology*. 2005;129(4):1268-1273. doi:10.1053/j.gastro.2005.07.029
46. Sandle GI, Perry MD, Mathialahan T, et al. Altered cryptal expression of luminal potassium (BK) channels in ulcerative colitis. *J Pathol*. 2007;212(1):66-73. doi:10.1002/path.2159
47. Sandle GI. Pathogenesis of diarrhea in ulcerative colitis: new views on an old problem. *J Clin Gastroenterol*. 2005;39(4 Suppl 2):S49-52. doi:00004836-200504002-00004 [pii]
48. Ram P, Goyal A, Lu M, Sloan J, McElhaugh W. Use of Aldosterone Antagonist to Treat Diarrhea and Hypokalemia of Ogilvie's Syndrome. *Case Rep Gastrointest Med*. 2016;2016:1-3. doi:10.1155/2016/1207240
49. Lundgren O, Peregrin AT, Persson K, Kordasti S, Uhnöo I, Svensson L. Role of the enteric nervous system in the fluid and electrolyte secretion of rotavirus diarrhea. *Science*. 2000;287(5452):491-495. doi:10.1126/science.287.5452.491
50. Thiagarajah JR, Donowitz M, Verkman AS. Secretory diarrhoea: Mechanisms and emerging therapies. *Nat Rev Gastroenterol Hepatol*. 2015;12(8):446-457. doi:10.1038/nrgastro.2015.111
51. Sandle GI. Salt and water absorption in the human colon: A modern appraisal. *Gut*. 1998;43(2):294-299. doi:10.1136/gut.43.2.294

Figures






Figure 1

Dietary Na⁺ depletion enhances NKCC1 protein, but not mRNA expression in rat proximal colon. A: Western blot performed using mucosal homogenates from normal and Na⁺-depleted rat proximal colon. Standard molecular weights and primary antibody targets are indicated to the left and right of the blot images, respectively. B: NKCC1 band intensity was quantified using densitometry analysis in ImageJ and normalized to actin band intensity as a loading control for each sample. C: qRT-PCR analysis of NKCC1-

specific mRNA transcript abundance between normal and Na⁺-depleted rat proximal colon. Data are represented as fold change relative to control (Normal). Lines and error bars represent mean ± SEM. *p<0.05 compared with control (Normal) using unpaired Student's t-test.




Figure 2

Immunofluorescent labelling of NKCC1 is enhanced by dietary Na⁺ depletion in rat proximal colon. NKCC1-specific labelling (green) in normal (top) and Na⁺-depleted (bottom) rat proximal colon sections. Nuclei are labelled with DAPI (blue). Merged images are shown in the far-right column. All images were captured at 20x magnification using a laser-powered confocal microscope. Scale bar = 50 mm.




Figure 3

Dietary Na⁺ depletion does not enhance Ba²⁺- or bumetanide-sensitive electrogenic K⁺ secretion in rat proximal colon. A-B: Short-circuit current (ISC) recordings from normal (open circles) and Na⁺-depleted (closed circles) proximal colon treated with serosal 200 mM bumetanide (A), or mucosal 3 mM Ba²⁺ (B). C-E: Basal (C), bumetanide-sensitive (D) and Ba²⁺ sensitive (E) ISC from normal and Na⁺-depleted proximal colon. Ba²⁺-sensitive and bumetanide-sensitive ISC values were calculated as the change in ISC (Δ ISC) from before and after the addition of drug to the chamber bath. Lines and error bars represent mean \pm SEM. * $p < 0.05$ compared with control (Normal) using unpaired Student's t-test.




Figure 4

Proximal colon BK and IK channel expression is unaffected by dietary Na⁺ depletion. A: Western blots performed using mucosal homogenates from normal and Na⁺-depleted rat proximal colon. Standard molecular weights and primary antibody targets are indicated to the left and right of the blot images, respectively. B-C: BK α (B) and IK (C) band intensities were quantified using densitometry analysis in ImageJ and normalized to actin band intensity as a loading control for each sample. D-E: qRT-PCR analysis of BK-specific (D) and IK-specific (E) mRNA transcript abundance between normal and Na⁺-depleted rat proximal colon. Data are represented as fold change relative to control (Normal). Lines and error bars represent mean \pm SEM.




Figure 5

Dietary Na⁺ depletion enhances NKCC1-dependent, Ca²⁺- and cAMP-stimulated Cl⁻ secretion in rat proximal colon. Short-circuit current responses to serosal 100 mM carbachol (CCH; A) or serosal 10 mM forskolin (FSK; B) from normal and Na⁺-depleted rat proximal colon in the presence (open circles) or absence (closed circles) of 200 mM serosal bumetanide. ΔI_{SC} values were calculated as the change from baseline to the maximal response after carbachol or forskolin application. Lines and error bars represent mean \pm SEM. * $p < 0.05$ using an ordinary two-way ANOVA with Šidák post-hoc.




Figure 6

Dietary Na^+ depletion enhances CFTR Cl^- channel expression in rat proximal colon. A-B: Western blots performed using mucosal homogenates from normal and Na^+ -depleted rat proximal colon. Standard molecular weights and primary antibody targets are indicated to the left and right of the blot images, respectively. C: Representative blot showing TMEM16A labelling in normal distal, but not proximal colon homogenates as a positive control for antibody specificity. D: Band intensities were quantified using

densitometry analysis in ImageJ and normalized to actin band intensity as a loading control for each sample. E: qRT-PCR analysis of CFTR- and TMEM16A-specific mRNA transcript abundance between normal and Na⁺-depleted rat proximal colon. Data are represented as fold change relative to control (Normal). Lines and error bars represent mean \pm SEM. *p<0.05 compared with control (Normal) using unpaired Student's t-test.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [s1.png](#)
- [s2.png](#)