

Diet and exercise as a management strategy for symptoms of polycystic ovary syndrome (PCOS): the views of women in the community with PCOS.

Susan Arentz (✉ s.arentz@westernsydney.edu.au)

NICM Health Research Institute, Western Sydney University <https://orcid.org/0000-0002-6721-472X>

Caroline A Smith

Western Sydney University

Jason Abbott

University of New South Wales

Alan Bensoussan

Western Sydney University

Research article

Keywords: Polycystic Ovary Syndrome, lifestyle interventions, women's experiences

Posted Date: April 24th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-20598/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published on March 17th, 2021. See the published version at <https://doi.org/10.1186/s12905-021-01252-1>.

Abstract

Background The international clinical practice guidelines for PCOS emphasize diet and exercise as first-line management of clinical signs and symptoms. This study aimed to describe the patterns, perceptions and experiences of lifestyle interventions for women in the community with PCOS.

Method An electronic survey of 493 members of two PCOS consumer support groups described women's types and patterns of diet and exercise, experiences and perceptions of effectiveness. Women were recruited from the Polycystic Ovary Association of Australia (POSAA) and from the Facebook group, PCOS University Research Group. Associations between participants perceptions of effectiveness, and diet types and exercise patterns were assessed using logistic regression. Response bias for the POSAA group was assessed with a continuum of resistance model.

Results 91% of POSAA members aged 16-50 years responded to the survey. Nearly all women reported adjusting their dietary and exercise practices with the aim to improve their health and/or PCOS (82% and 73% respectively), however less than 13% reported achievement of health goals (12.2% and 8.1% respectively). Low carbohydrate, high protein diets, and vigorous activity were associated with self-perceived effectiveness ($r=0.16$, $p<0.01$; $r=0.15$ $p<0.01$ and $r=0.2$ $p<0.01$ respectively). Barriers for lifestyle interventions included psychosocial factors. Response bias was not assessed for the Facebook group, however self-reported PCOS aligned with prevalence of clinical phenotypes and suggests results are generalizable to clinical populations of women with PCOS, who are responsible for self-directing and administering lifestyle interventions to manage their PCOS.

Conclusions Perceptions of effectiveness for lifestyle interventions by women with PCOS may be complicated by a lack of rigorous evidence. The strength of recommendations in clinical practice guidelines may be enhanced by clinical trials investigating flexible and feasible lifestyle interventions for women in the community with PCOS.

Background

Polycystic ovary syndrome (PCOS) is the most common reproductive endocrinopathy affecting up to one in five women of reproductive age.^{1,2} The cause of PCOS remains unknown however increased prevalence coincides with a global increased incidence of metabolic syndrome. Metabolic disturbance is associated with increased adiposity and clinically worsens menstrual cyclicity, hyperandrogenism, ovulation rates, fertility, maternal and neonatal outcomes and increases women's risks for diabetes, cardiovascular disease and cancer.^{2,3}

The international evidence based guidelines (EBG) for the management of PCOS highlights personal lifestyle behaviours to reduce and prevent risk factors significantly associated with increased body weight in women with PCOS.⁴ Diet and exercise interventions aimed at weight reduction and prevention of weight gain are first-line management strategies due to their significant impact on clinical outcomes,

including reproductive endocrinology and clinical signs and symptoms.³⁻⁵ However, current evidence for lifestyle intervention for women with PCOS is not well established. Many studies informing the evidence base are underpowered and report high risk of bias, particularly from high attrition.⁵⁻⁸ The effectiveness of specific dietary and exercise practices for women centred outcomes has not been established. It has not been shown that if, and to what degree diet and exercise performed outside the confines of study intervention environments correlate with improved health status for women with PCOS in the community.⁹

Dietary and exercise practices of women may be influenced by a wide range of ecological factors including inter and intrapersonal, environmental and psychosocial characteristics of individuals plus levels of health literacy and potential for self-efficacy.¹⁰⁻¹⁴ Health promoting behaviours by women with PCOS may be complicated by a lack of evidence informing women of the benefits of lifestyle intervention,⁴ and may exacerbate psychological morbidity, which has a high prevalence in women with PCOS,¹⁵ and frustration with current medical treatments.^{16,17} To date there are few data describing the views of women with PCOS for engagement and optimal effectiveness of dietary and exercise strategies in community-based environments.

This study has two aims; to describe the symptoms of concern to women with PCOS living in the community, and to explore perceptions about effectiveness for dietary and exercise interventions for PCOS.

Method

The study was approved by the University of Western Sydney Human Research and Ethics Committee H9341. A survey was undertaken to examine women's perceptions of PCOS and the role of diet and exercise and other interventions which may contribute to the management of their PCOS symptoms and wellbeing. Women with PCOS seeking health support as members of internet based social networks were approached through two on-line consumer support networks, the Polycystic Ovary Syndrome Association of Australia (POSAA) and the social network site (SNS), Facebook. Participants were self-selected from POSAA following direct email to 258 financial members between November 2011 and September 2013, and secondly a Facebook sample responded to an invitation to participate in the survey and a link placed on the 'wall' of the Facebook group page www.uws.edu.au/pcosfacebook. Wall postings occurred bi-monthly between February 2013 and May 2013.

Questionnaire design

A 37-item anonymous questionnaire was designed to describe the signs and symptoms of women with PCOS and women's use of medical treatment and diet and exercise to manage PCOS. (Supplementary File 1). Fifteen items sought information regarding dietary and exercise practices and included types of diets and exercise, types and patterns of exercise practice in terms of frequency, duration and intensity. Moderate intensity exercise was described as causing an increased breath and heart rate but still able to

talk. Vigorous exercise was described as causing markedly increased breath and heart rate and not being able to talk during activity. Additional questions examined barriers to engagement including reasons for not partaking or disengaging with therapeutic diet and exercise practices and questions about self-perceived achievement of goals and satisfaction with diets and exercise. Socio-demographic characteristics were also collected. Multiple response options were available for participants to indicate which signs and symptoms of PCOS they were experiencing, and the types of dietary and exercise practices used. A cover letter informed participants' that consent was implied on completion of the questionnaire. The questionnaire took 15–20 minutes to complete. The stability of the questionnaire was tested on a pilot sample of eight women including four with PCOS to review the applicability and accuracy of the questionnaire.

The electronic survey was loaded onto Survey Monkey¹⁸ and made available to both members of POSAA and members of the Facebook group. A single response function was applied to each member and to devices.

Data analyses

Responses were analysed using the statistical package SPSS version 21.0.¹⁹ Categorical responses were reported as percentages and proportions with 95% confidence intervals. Relationships between variables were explored using Pearson's correlation (two tail) for parametric data and Spearman's Rho correlations or Gamma co-efficient (collapsed less than six variables) for non-parametric measurements. Associations between perceptions of effectiveness (dependent variable) and type and pattern of diet and exercise (covariates) were assessed using logistic regression. A p value of 0.05 was considered statistically significant.

Non-response bias was assessed using a continuum of resistance model in which late responders were viewed as proxy non-responders.²⁰ Participants who responded to the survey four months or more after the final reminder, were classified as late responders. This method was chosen due to the likelihood of early response by women who found the topic interesting.²¹ The impact of non-response bias was assessed by comparing early and late responders.

Results

Four hundred and ninety-three participants aged between 16–50 years responded to the survey; 235 of the 258 invited members of the POSAA group responded (91.1%) including 53 late responders, and 311 women replied to snowballing using the University Research Facebook group page. No differences in demographics or PCOS signs and symptoms were found between early and late responders or the two consumer support groups (POSAA and Facebook) (Supplementary file 2. Tables 1 and 2).

Four hundred and thirty-four (88.0%) subjects completed all parts of the survey, with the remainder using the skip patterns inserted into the electronic survey or completing only part of the survey.

The demographics of participants are presented in Table 1. Most women were aged between 25 and 34 years (n = 237, 48.1%, ± 4.41 , 95%CI), were tertiary educated (n = 323, 76.2%, ± 4.05), and employed, (335, 67.9%, ± 4.12).

	Frequency N=493	Valid %	95% Confidence Interval	Table 1
				Demographic
<i>Age</i>				characteristics of Survey participants.
16-17	3	.7	± 0.79	
18-24	72	17.0	± 3.58	
25-29	126	29.7	± 4.35	
30-34	111	26.2	± 4.19	
35-40	71	16.7	± 3.55	
41-44	25	5.9	± 2.24	
45-50	16	3.8	± 1.82	
Missing data	69	14		
<i>Education</i>				
Complete high School				
Yes	371	87.5	± 3.15	
No	48	11.3	± 3.01	
Currently enrolled	5	1.2	± 1.04	
Missing data	69	14		
<u>Completed tertiary education</u>				
Yes	323	76.2	± 4.05	
No	55	13.0	± 3.0	
Currently enrolled at a tertiary inst.	46	10.8	± 2.95	
Missing data	69	14		
Total	493			
Qualification from TAFE	103	20.9	± 3.59	
Private institution Diploma	70	14.2	± 3.08	
University degree	221	44.8	± 4.39	
Missing data	69	14		
<i>Ethnicity- country of birth</i>				
Australia	302	61.3	± 4.3	
Other	121	24.5	± 2.08	
Missing data	70	14.2		
<i>Employment</i>				
Home duties	117	23.7	± 3.75	
Self-employed	38	7.7	± 2.35	
Student	80	16.2	± 3.25	
Employed part time	103	20.9	± 3.59	
Employed full-time	194	39.4	± 4.31	
Other	12		± 1.35	
<i>Private Health Insurance</i>				
Yes	275	65	± 4.55	
No	148	35	± 4.55	
Missing data	70	14.2		

($r=0.08$, $p = 0.10$). Women reporting hirsutism were more likely to report depression ($r=0.24$, $p < 0.01$).
(Table 2)

Table 2
Self-reported correlation of PCOS signs and symptoms

Signs and symptoms		Depression	Waist more cm's than hip	Infertility	Cramps
Late period	Pearson Correlation	.079	.273**	.450**	.193**
	Sig. (2-tailed)	.097	< .01	< 0.01	< 0.01
	N	443	440	419	398
Hirsutism	Pearson Correlation	.241**	.301**	.082	.003
	Sig. (2-tailed)	< 0.01	< 0.01	.092	.957
	N	453	448	426	400
Overweight	Pearson Correlation	-.013	.222**	.225**	.079
	Sig. (2-tailed)	.781	< 0.01	< 0.01	.112
	N	458	455	433	410
Very overweight	Pearson Correlation	.019	.228**	.408**	.237**
	Sig. (2-tailed)	.690	< 0.01	< 0.01	< 0.01
	N	443	440	422	395
Blood sugar/insulin imbalance	Pearson Correlation	.112*	.225**	.691**	.222**
	Sig. (2-tailed)	.018	< 0.01	< 0.01	< 0.01
	N	449	448	426	402
** Correlation was significant at the 0.01 level (2-tailed).					
* Correlation was significant at the 0.05 level (2-tailed).					
cm's: centimetres					
Sig: significance					
N: frequency of valid cases					

Management of PCOS

The respondents used various medical therapies to manage PCOS (Table 3). Over sixty five percent (± 4.21) had used the oral contraceptive pill, 62.7% (± 4.26) had used other pharmaceuticals including ovulation induction, hypoglycaemic and anti-androgen drugs.

Table 3
Medical interventions used by participants.

	Frequency	%	Missing	N
Oral contraceptive pill	322	65.3%	171	493
Pharmaceuticals: ovulation induction, anti-androgens hypoglycaemics etc	309	62.7%	184	493
Ovarian stimulation as part of an IVF cycle	58	11.8%	435	493
Surgery	49	9.9%	444	493
No medical treatment	10	2%	483	493

Lifestyle interventions

Diet

Over 82% (± 3.36 , $n = 406$) of respondents reported altering their dietary intake during the previous five years for health reasons. Over 57% (± 4.37 , $n = 282$) reported consuming a diet specifically to manage symptoms associated with PCOS and had tried various types of diets. The most frequently used was the low glycaemic (GI) index diet (50.1%, ± 4.41 , $n = 247$), followed by the low calorie diet (36.9%, ± 4.26 , $n = 181$), low carbohydrate diet (33.7%, ± 4.17 , $n = 166$) and low fat diet (32.9%, ± 4.15 , $n = 162$).

Women were asked to indicate their reason for dieting. Most respondents (72.7%, ± 3.93) modified their diet with the aim to lose weight. Other reasons included maintaining general health, 59% (± 4.64), to manage blood sugar levels 37.3% (± 4.56), to improve body composition 15% (± 3.37) and for social or religious reasons 7.4% (± 2.47).

The majority of women reported they did not fully achieve their health goals using dietary changes. Over thirty three percent (± 4.5 , $n = 140$) reported they did not achieve their weight loss goals or any positive health changes from dietary practices. Forty six percent (± 4.78 , $n = 195$) reported that dietary practices had contributed partly to positive health changes and a small number of women ($n = 51$) reported fully achieving their health goals using dietary practices (12.2%, ± 3.13).

Positive associations between types of diets and achievement of health goals were found between the low carbohydrate diet ($r = 0.16$, $p < 0.01$, $N = 419$) and the high protein diet ($r = 0.15$, $p < 0.01$, $N = 419$).

Exercise

Most women, (73.1%, \pm 4.2, n = 313) reported regular participation in moderate or vigorous exercise. Women were engaged in a variety of different forms of exercise, most exercisers reported practicing two or more types (97%, \pm 4.57, n = 305). Over half of participants (61.3%, \pm 3.13) were engaged in formal exercise including supervision provided by a personal trainer, participation in a team sport or attending a gymnasium and lifting weights, running, swimming, cycling or supervised classes. Over one third (37.5%, \pm 4.27) of women reported incidental activity as exercise. Other exercise included Pilates, dance, personal trainer, tennis and team sports.

Most respondents reported multiple reasons for exercising, with 80% (\pm 4.45) using exercise to feel better in themselves, 75% (\pm 4.79) to induce weight loss and 60% (\pm 5.45) to prevent weight gain. Sixty percent of responders (\pm 5.45) were exercising specifically to manage their PCOS.

One hundred and fifteen participants did not exercise (26.9%, \pm 4.2). Reasons for not exercising included lack of time (n = 109), felt embarrassed (n = 93), and financial costs (n = 52). Fifty-one cited other reasons such as feeling unmotivated, depressed, and co-morbid overweight conditions including back pain and arthritis.

Most women reported participation in both moderate and vigorous exercise. Over 88% (\pm 3.5, n = 277) of exercising women, regularly engaged in moderate exercise, and nearly half (48.7% \pm 6.04, n = 128) in vigorous exercise. Seven women reported not doing any moderate exercise at least once per month, and 88 women (28.3%, \pm 4.67) were not engaged in any vigorous exercise at least once per month and 87 women, no vigorous exercise at all.

Most women exercised two to four times per week (42.0%, \pm 5.46, n = 132). Most responders exercised moderately for 15–30 minutes per session (46.1%, \pm 5.49, n = 146,) and just under 40% exercised vigorously for 30–60 minutes (39.9%, \pm 5.44, n = 134).

Nearly forty percent (39.4%, \pm 5.35, n = 127) reported non achievement of health goals with exercise. The majority reported part achievement using exercise (51.8%, \pm 5.46, n = 166), and only eight percent (\pm 2.93, n = 25) of respondents felt they had achieved their health goals with exercise.

We found that increased frequency and duration of exercise was significantly associated with the achievement of health goals and an inverse association between the duration of moderate exercise and the achievement of health goals. (Table 4).

Table 4
Duration of exercise and self-perceptions of the achievement of health goals.

		Vigorous exercise duration	Moderate exercise less than 60 minutes	Moderate exercise more than 60 minutes
Self-perceived achievement of goals	Pearson Correlation	.196**	.247**	-.006
	Sig. (2-tailed)	.000	.000	.919
	N	320	320	320
** Correlation was significant at the 0.01 level (2-tailed).				
* Correlation was significant at the 0.05 level (2-tailed).				
Sig: significance				
N: frequency of valid cases				

Discussion

This study highlights the views of women in the community with PCOS and describes their self-reported dietary and exercise lifestyle behaviours and perceptions of effectiveness. Nearly all women reported adjusting their dietary and exercise practices with the aim to improve their health and/or PCOS, however less than 13% reported achievement of health goals. Participants reported a range of PCOS symptoms including being overweight, reproductive disorders and negative emotional and mental health and frequent engagement with positive health behaviours as part of self-care management, with the primary aims to lose weight and manage PCOS.

Women with PCOS in the community have been found to lack support and information for self-managing PCOS symptoms using lifestyle interventions.^{9,17} In a survey of 1385 women with PCOS, only 11.9% reported satisfaction with information provided about beneficial diet and exercise.¹⁷ In the same study, less than 4% of participants were satisfied with information provided about the emotional features of PCOS, with no information or support being offered in most cases.¹⁷ Lack of information and support for community-based women with PCOS may have influenced the low perceptions of efficacy for lifestyle interventions found in the present study. Women's information needs include details about types and patterns of safe and effective lifestyle interventions, and mediators for success and goal achievement including the impact of social support and influence of expectations and experiences of lifestyle interventions.^{15,22}

Views of community-based women with PCOS presented here contrast with findings of a randomized control trial (RCT) evaluating diet and exercise on the motivators and barriers of exercise for women with PCOS.⁸ This clinical investigation found women's perceptions of exercise were significantly improved

over 20 weeks in all three study arms (diet only, diet plus aerobic exercise and diet plus aerobic and progressive resistance exercises), and correlated with improved anthropometry, health related quality of life and less depression.⁸ Greatest perceived benefits were on the psychological outlook and social interaction subscales of the Exercise Benefits/Barriers Scale (EBBS) ($p \leq 0.001$). Other significant findings were reduced barriers on subscales of exercise milieu (atmosphere), time expenditure and physical exertion ($p \leq 0.003$). The contrasting findings found in the present study may be explained by the recruitment strategy of women with PCOS from different populations and by differences in the nature and structure of lifestyle interventions.

The present study recruited participants from consumer support groups in order to assess the views of women in the community, whereas recruitment to the clinical investigation was conducted via public advertisements via medical speciality clinics. Women with PCOS seeking medical support have been shown to display different characteristics of PCOS compared to community-based women² which may explain variable perceptions of exercise interventions found in the present study. In addition, the nature of the interventions provided in the clinical trial were structured with weekly contacts, compared to self-informed, self-initiated, self-administered and self-accounted lifestyle interventions typically used by women in the community. Although nearly two thirds of participants reported engagement with supervised/structured exercise, a lack of specific information about evidence-based lifestyle recommendations for PCOS, may have limited guidance of exercise supervisors, with respect to the types and patterns of exercise provided, and a lack of information may have mediated women's expectations and experiences of efficacy.²³

It is unclear which dietary and exercise strategies are optimal for women with PCOS. Current recommendations based on reduced caloric intake, combined with moderate energy expenditure, are described as being most likely to produce sustained weight loss and favourable endocrine and reproductive outcomes.^{5,24-27} Women in this study reported similar optimal dietary and exercise practices on symptoms of PCOS, with greatest effectiveness reported for low carbohydrate and high protein diets compared to low calorie diets. Women's perceptions of effective exercise practices highlighted the importance of including vigorous activity, however many respondents reported no engagement in any vigorous activity at all. Various reasons were cited, including time limitations, feelings of embarrassment and personal injury. Additionally, exercise preferences of women with PCOS may favour non-vigorous activity. A study into the comparative exercise practices of women with and without PCOS, controlled for BMI, ($n = 163$) showed that women with PCOS were less likely to engage in vigorous activity²⁸ and was strongly determined by personal circumstances and characteristics.

Despite the present study finding most women's exercise activity aligned with recommendations in clinical practice guidelines,²⁹ the respondents perceptions of low efficacy elucidates a gap in information¹⁷ and a need for guidance despite the absence of high quality evidence.⁵ Women in the community still need to make decisions about patterns and types of lifestyle to manage PCOS. Their perceived low efficacy may be complicated by the absence of rigorous information and reflect the trial

and error process of self-directed and self-funded searches for the most effective and feasible lifestyle interventions for their case and circumstances.

This study contributes to our understanding of women with PCOS living in the community, and their experiences using lifestyle interventions to manage symptoms and health. This research gives voice to community-based women with PCOS and identifies important modifying factors for consideration in future clinical practice recommendations.³⁰ It may also guide pragmatic clinical trial investigations of flexible lifestyle interventions, conducted in community settings with real-world characteristics³¹ in order to improve the quality of evidence and strengthen recommendations of clinical practice guidelines.⁴

Limitations and strengths

There are some methodological limitations of this study including the non-response and selection bias of recruitment from Facebook support groups. We were not able to assess unsolicited invitations from group administrators and response bias analyses was not possible. Women volunteered to participate, which may have introduced response bias with responders potentially reporting more positive lifestyle behaviours compared to non-responders.³² Generalizations are limited to computer literate women with English language skills, other groups of women with PCOS may be underrepresented. The questionnaire was not validated against medical records for PCOS lifestyle interventions; however, it was designed to investigate women with PCOS in the community and their perceptions, usage and experiences of lifestyle interventions, which may not coincide with the views of clinical groups. PCOS was self-reported however a similar symptom profile to clinical populations^{1,15,33} was described, and these results may be generalisable to women with medically diagnosed PCOS, living in the community. Genetic analyses of PCOS susceptibility has demonstrated consistency between woman's self-reports of the condition and rigorous diagnoses.^{34,35}

Conclusion

Women in the community with PCOS report frequent engagement in a variety of dietary and exercise practices according to the clinical practice guidelines, and they perceive their health goals are partially met. Self-perceived lack of efficacy for dietary and exercise interventions may be due to the lack of high-quality evidence of effectiveness for specific lifestyle interventions. The quality of evidence might be improved by pragmatic RCTs of community-based samples, comparing lifestyle interventions on outcomes of value to women with PCOS.

Abbreviations

BMI Body Mass Index

EBBS Exercise Benefits/Barriers Scale

EBG Evidence Based Guidelines

GI Glycaemic Index

PCOS Polycystic Ovary Syndrome

POSAA Polycystic Ovary Association of Australia

RCT Randomized Control Trial

SNS Social Network Site

Declarations

Ethics approval and consent to participate

The study, including the procedure for obtaining informed consent to participate, was approved by the Western Sydney University Human Research Ethics Committee (EC00314) on 28th October 2011. An amendment to the recruitment protocol was approved on 19th August 2013. The reference number was H9341.

Consent to publish

The Participant Information Form informed of potential risks from participation in the survey, and that consent to publish the results was implied through participation and submission of responses. It also explained that that responses could not be withdrawn due to anonymity of the survey.

Availability of data and materials

The datasets and materials used and analysed during the current study are available from the corresponding author on reasonable request.

Competing interests

The authors have no competing interests to declare. SA was a doctoral student at NICM Health Research Institute of Western Sydney University, Australia. CAS, JA and AB formed her supervisory panel and provided critical guidance and feedback throughout the whole research project. As a medical research institute, NICM Health Research Institute receives research grants and donations from foundations, universities, government agencies, individuals and industry. Sponsors and donors also provide untied funding for work to advance the vision and mission of the Institute. The authors declare no competing financial interests. SA and JA are also in clinical practice.

Funding

This study was funded through NICM Health Research Institute doctoral student funds. SA received an Australian Postgraduate Award (APA) scholarship during the study period.

Authors contributions

All authors SA, CAS, JA and AB conceived of the study and contributed to its design, coordination and administration. SA and CAS designed the sampling strategy, recruitment, data collection and data analyses and liaised with community groups. JA and AB reviewed the quality of data and data analyses. SA, CAS, JA and AB designed and edited the tables. All authors read, edited and approved the final manuscript.

Acknowledgements

The authors would like to acknowledge the participants of this study and the valuable support of the president of POSAA Veryan McAlister, the volunteers and the assistance of the Facebook group administrators Jamie Russell and Kristina Judd. The authors would like to acknowledge the statistical guidance of Paul Fahey of the School of Science and Health, Western Sydney University.

Authors information

SA is an adjunct research fellow at NICM Health Research Institute, Western Sydney University and in clinical practice.

References

1. March WA, Moore VM, Willson KJ, Phillips DIW, Norman RJ, Davies MJ. The prevalence of polycystic ovary syndrome in a community sample assessed under contrasting diagnostic criteria. *Human Reproduction* 2010;25:544-51.
2. Azziz R, Carmina E, Chen Z, et al. Polycystic ovary syndrome. *Nature reviews Disease primers* 2016;2:1-18.
3. Teede H, Deeks A, Moran L. Polycystic ovary syndrome: a complex condition with psychological, reproductive and metabolic manifestations that impacts on health across the lifespan. *BMC medicine* 2010;8:41.
4. Teede H, Misso, M., Costello, M., Dokras, A., Laven, J., Moran, L., Piltonen, T., Norman, R. International evidence-based guideline for the assessment and management of polycystic ovary syndrome. In: Council NHaMR, ed. Australia: NHMRC; 2018.
5. Lim SS, Hutchison SK, Van Ryswyk E, Norman RJ, Teede HJ, Moran LJ. Lifestyle changes in women with polycystic ovary syndrome. *Cochrane Database of Systematic Reviews* 2019.
6. Marsh KA, Steinbeck KS, Atkinson FS, Petocz P, Brand-Miller JC. Effect of a low glycemic index compared with a conventional healthy diet on polycystic ovary syndrome. *The American journal of clinical nutrition* 2010;92:83-92.

7. Moran LJ, Noakes M, Clifton PM, Tomlinson L, Norman RJ. Dietary composition in restoring reproductive and metabolic physiology in overweight women with polycystic ovary syndrome. *Journal of Clinical Endocrinology & Metabolism* 2003;88:812-9.
8. Thomson RL, Buckley JD, Brinkworth GD. Perceived exercise barriers are reduced and benefits are improved with lifestyle modification in overweight and obese women with polycystic ovary syndrome: a randomised controlled trial. *BMC women's health* 2016;16:14.
9. Humphreys L, Costarelli V. Implementation of dietary and general lifestyle advice among women with polycystic ovarian syndrome. *The journal of the Royal Society for the Promotion of Health* 2008;128:190.
10. Glanz K, Basil M, Maibach E, Goldberg J, Snyder D. Why Americans eat what they do: taste, nutrition, cost, convenience, and weight control concerns as influences on food consumption. *Journal of the American Dietetic Association* 1998;98:1118-26.
11. Clark NM, Dodge JA. Exploring self-efficacy as a predictor of disease management. *Health Education & Behavior* 1999;26:72-89.
12. Skelly AH, Marshall JR, Haughey BP, Davis PJ, Dunford RG. Self-efficacy and confidence in outcomes as determinants of self-care practices in inner-city, African-American women with non-insulin-dependent diabetes. *The Diabetes Educator* 1995;21:38-46.
13. Nutbeam D. Health literacy as a public health goal: a challenge for contemporary health education and communication strategies into the 21st century. *Health promotion international* 2000;15:259-67.
14. Booth SL, Sallis JF, Ritenbaugh C, et al. Environmental and societal factors affect food choice and physical activity: rationale, influences, and leverage points. *Nutrition reviews* 2001;59:S21-S36.
15. Ching H, Burke V, Stuckey B. Quality of life and psychological morbidity in women with polycystic ovary syndrome: body mass index, age and the provision of patient information are significant modifiers. *Clinical Endocrinology* 2007;66:373-9.
16. Sills ES, Perloe M, Tucker MJ, Kaplan CR, Genton MG, Schattman GL. Diagnostic and treatment characteristics of polycystic ovary syndrome: descriptive measurements of patient perception and awareness from 657 confidential self-reports. *BMC women's health* 2001;1:3.
17. Gibson-Helm M, Teede H, Dunaif A, Dokras A. Delayed diagnosis and a lack of information associated with dissatisfaction in women with polycystic ovary syndrome. *The Journal of Clinical Endocrinology & Metabolism* 2017;102:604-12.
18. Survey Monkey web based software Listserv, 2013. 2012-4, at www.surveymonkey.com.)
19. IBM. SPSS version 21. 2013:Statistics software
20. Boshuizen HC, Viet A, Picavet HSJ, Botterweck A, Van Loon A. Non-response in a survey of cardiovascular risk factors in the Dutch population: determinants and resulting biases. *Public health* 2006;120:297-308.
21. Porter SR, Whitcomb ME. Non-response in student surveys: The role of demographics, engagement and personality. *Research in higher education* 2005;46:127-52.

22. Brennan L, Teede H, Skouteris H, Linardon J, Hill B, Moran L. Lifestyle and behavioral management of polycystic ovary syndrome. *Journal of Women's Health* 2017;26:836-48.
23. Gibson-Helm M, Tassone EC, Teede HJ, Dokras A, Garad R. The needs of women and healthcare providers regarding polycystic ovary syndrome information, resources, and education: A systematic search and narrative review. *Seminars in reproductive medicine*; 2018: Thieme Medical Publishers. p. 035-41.
24. Clark A, Ledger W, Galletly C, et al. Weight loss results in significant improvement in pregnancy and ovulation rates in anovulatory obese women. *Human Reproduction* 1995;10:2705-12.
25. Huber-Buchholz M-M, Carey D, Norman RJ. Restoration of reproductive potential by lifestyle modification in obese polycystic ovary syndrome: role of insulin sensitivity and luteinizing hormone. *Journal of Clinical Endocrinology & Metabolism* 1999;84:1470-4.
26. Harrison CL, Lombard CB, Moran LJ, Teede HJ. Exercise therapy in polycystic ovary syndrome: a systematic review. *Human Reproduction Update* 2011;17:171-83.
27. Lamb JD, Johnstone EB, Rousseau J-A, et al. Physical activity in women with polycystic ovary syndrome: prevalence, predictors, and positive health associations. *American Journal of Obstetrics and Gynecology* 2011;204:352. e1-. e6.
28. Wright C, Zborowski J, Talbott E, McHugh-Pemu K, Youk A. Dietary intake, physical activity, and obesity in women with polycystic ovary syndrome. *International journal of obesity* 2004;28:1026-32.
29. Teede HJ, Misso ML, Costello MF, et al. Recommendations from the international evidence-based guideline for the assessment and management of polycystic ovary syndrome. *Fertility and sterility* 2018;110:364-79.
30. Andrews JC, Schunemann HJ, Oxman AD, et al. GRADE guidelines: 15. Going from evidence to recommendation-determinants of a recommendation's direction and strength. *J Clin Epidemiol* 2013;66:726-35.
31. Zwarenstein M, Treweek S, Gagnier JJ, et al. Improving the reporting of pragmatic trials: an extension of the CONSORT statement. *Bmj* 2008;337:a2390.
32. Hill A, Roberts J, Ewings P, Gunnell D. Non-response bias in a lifestyle survey. *Journal of Public Health* 1997;19:203-7.
33. Balen AH, Conway GS, Kaltsas G, et al. Andrology: Polycystic ovary syndrome: the spectrum of the disorder in 1741 patients. *Human reproduction* 1995;10:2107-11.
34. Hayes MG, Urbanek M, Ehrmann DA, et al. Genome-wide association of polycystic ovary syndrome implicates alterations in gonadotropin secretion in European ancestry populations. *Nature Communications* 2015;6.
35. Day FR, Hinds DA, Tung JY, et al. Causal mechanisms and balancing selection inferred from genetic associations with polycystic ovary syndrome. *Nature communications* 2015;6:1-7.