

Implementation of Targeted Screening for Poverty in a Large Primary Care Team in Toronto, Canada: a feasibility study

Kimberly Wintemute

University of Toronto Faculty of Medicine: University of Toronto Temerty Faculty of Medicine

Meh Noor (✉ m.noor@mail.utoronto.ca)

University of Toronto <https://orcid.org/0000-0003-2483-4360>

Aashka Bhatt

University of Toronto Faculty of Medicine: University of Toronto Temerty Faculty of Medicine

Gary Bloch

University of Toronto Faculty of Medicine: University of Toronto Temerty Faculty of Medicine

Suja Arackal

University of Toronto Faculty of Medicine: University of Toronto Temerty Faculty of Medicine

Sumeet Kalia

University of Toronto Faculty of Medicine: University of Toronto Temerty Faculty of Medicine

Babak Aliarzadeh

University of Toronto Faculty of Medicine: University of Toronto Temerty Faculty of Medicine

Sabrina La Tona

University of Toronto Faculty of Medicine: University of Toronto Temerty Faculty of Medicine

Joyce Lo

University of Toronto Faculty of Medicine: University of Toronto Temerty Faculty of Medicine

Andrew Pinto

University of Toronto Faculty of Medicine: University of Toronto Temerty Faculty of Medicine

Michelle Greiver

University of Toronto Faculty of Medicine: University of Toronto Temerty Faculty of Medicine

Research article

Keywords: Family Practice, Primary Health Care, Mass Screening / organization and administration, Social Determinants of Health, Poverty, Poverty Areas, Feasibility Studies, Electronic Health Records

Posted Date: March 19th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-206360/v2>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background Poverty has a significant influence on health. Efforts to optimize income and reduce poverty could make a difference to the lives of patients and their families. Routine screening for poverty in primary care is an important first step but rarely occurs in Canada. We aimed to implement a targeted screening and referral process in a large, distributed primary care team in Toronto, Ontario, Canada.

Methods This implementation evaluation was conducted with a large community-based primary care team in north Toronto. The primary care team serves relatively wealthy neighborhoods that contain pockets of poverty. Physicians were invited to participate. We implemented targeted screening by combining census information on neighborhood-level deprivation with postal codes in patient records. For physicians agreeing to participate, we added prompts to screen for poverty to the charts of adult patients living in the most deprived areas. Standardized electronic medical record templates recommended a referral to a team case worker for income optimization, for those patients screening positive. We recorded the number and percentages of participants at each stage, from screening to receiving advice on income optimization.

Results The primary care team included 86 physicians distributed across 19 clinical locations. Thirty-four physicians (39%) participated. Their practices provided care for 27,290 patients aged 18 or older; 852 patients (3%) were found to be living in the most deprived neighborhoods. 509 (60%) had at least one office visit over the 6 months of follow up, and 128 (25%) of those seen were screened. Only 25 (20%) of these patients screened positive for poverty, and 13 (52%) were referred. Eight patients (62% of those referred) were ultimately seen by a caseworker for income optimization.

Discussion & Conclusions We implemented a targeted poverty screening program combined with resources to optimize income for patients in a large, distributed community-based primary care team. However, only a small number of patients were identified and linked to the intervention. Further efforts to scale and spread screening and mitigation of poverty are warranted; these should include broadening the targeted population beyond those living in the most deprived areas.

Contributions To Literature

- Poverty affects health; screening and addressing poverty in primary care could lead to higher income and better lives
- This is rarely done in Canadian primary care
- We implemented a screening and income optimization program in a large “real world” community-based primary care team and evaluated its feasibility. Screening was targeted towards those living in the most deprived neighborhoods
- A quarter of patients living in deprived neighborhoods that visited the practices were screened and 20% of those screened positive, half of whom were referred for income optimization.
- Targeted screening may be a useful entry point towards universal screening

Background

Socioeconomic status (SES) is one of the most important determinants of health (1). Poverty adversely influences health outcomes for individuals (2-4). Persons living with income insecurity have greater rates of chronic health conditions and a higher risk of reduced lifespan (3-5). The COVID-19 pandemic has recently highlighted the effects of income disparities for communicable diseases: those living in economically deprived neighborhoods have disproportionately greater odds of contracting the infection, of being hospitalized and of dying from COVID-19 complications (6-8).

The pandemic has also highlighted gaps in socio-demographic data collection. Compared to international settings (7, 9), Canada lags in the collection and reporting of social determinants of health; these are associated with differences in COVID-19 infection rates and outcomes (10). Some gaps have been partially addressed, leading to attention and efforts to direct appropriate resources to communities at greater risk (11, 12).

Primary care represents an ideal setting for the collection of information on social determinants of health, and for taking action on these (13-16). Family physicians and their teams provide community-based longitudinal care for patients and families, building relationships based on trust, and generating knowledge about the context patients live in (13, 17). The pandemic has led to increased attention to social context; family physicians have been provided with tools to address poverty and other social determinants of health during and after the crisis (18).

As with other major risk factors such as tobacco use, a critical first step involves clinicians asking their patients so that risk status can be identified and documented (19, 20). An evidence-based tool for poverty screening and intervention in primary care has been developed and studied by Bloch et.al (21, 22). It is central to current screening recommendations in Ontario and Canada (22). The screening questions are: *“Do you ever have difficulty making ends meet at the end of the month?”* (sensitivity 98%, specificity 40% for living below the poverty line) (2) and *“Have you filled out and sent in your tax forms?”*

However, in Canada, screening for poverty is not currently a routine part of family practice, with some exceptions (23). Barriers include lack of provider training, lack of time, lack of knowledge and expertise, and difficulty changing workflows (24, 25). There may be multiple competing priorities as family physicians look after many issues and conditions that require attention. Pilot work in several sites in Toronto determined that poverty screening and intervention were feasible in the practices studied (14, 23). However, these sites had a mandate to address social inequities as a priority, practice populations with high levels of poverty, dedicated champions and resources devoted to the intervention. An exploratory study in a variety of “real world” primary care settings found that only 9% of patients were screened, despite training and the presence of motivated healthcare providers (24).

Implementing poverty screening requires addressing barriers. This is not a trivial activity in practices already overwhelmed by current care demands. Lack of time is one of the most significant barriers to change in primary care (26). In one highly cited study, Yarnall noted in order “to fully satisfy the US

Preventive Services Task Force recommendations, 1773 hours of a physician's annual time, or 7.4 hours per working day, is needed for the provision of preventive services" (27).

We used Diffusion of Innovations theory to plan the intervention (28). Implementation may be more successful if the intervention is perceived as not being complex, as taking little time or practice resources, and as providing high value compared with usual care (28-30). Our team had previous experience implementing screening interventions and integrating those in primary care Electronic Medical Records (EMRs) (31, 32). We used this knowledge for the design of the poverty screen. We adapted approaches (EMR prompts, templates) found as part of other screening workflows that physicians were already using and were familiar with (33) to provide an intervention that was simple and easy to use.

We then devised an implementation strategy consistent with good design principles, considering inner and outer settings (30). Briefly, a pilot was implemented and evaluated in a single office, with four motivated family physicians; this demonstrated initial feasibility in a community-based setting similar to ours. We scaled up using strategies known to be effective (presence of champion, building consensus on the importance of the issue, clinician education, leadership endorsement, availability of resources, adapting and tailoring strategies to local context) (30). We implemented and evaluated the poverty screening strategy in a large community primary care inter-professional team in Toronto, Ontario, Canada, the North York Family Team (NYFHT, <http://nyfht.com>).

An innovation in our approach was targeted screening for poverty. Using postal codes, we identified patients living in the most deprived neighborhoods. We expected that those areas included more patients living with poverty, and therefore would increase screening efficiency. The total number of patients requiring screening and the workload for physicians would be much lower than that required for universal screening. This would enhance trialability (28) through a limited initial implementation.

While universal screening for poverty is the recommended approach (16, 21, 22), targeted screening may present a useful initial step in the face of continuing limited uptake of screening. A similar approach (targeted vs universal screening) is currently being tested as part of a comparative effectiveness trial for major depression screening in adolescents, a screen with limited uptake (34).

If successful, further phases are being planned. A logic model is presented in Additional File 1. Phase One, reported here, consisted of assessing the feasibility of carrying out targeted screening for poverty and providing liaison to income security supports, and then assessing the feasibility of having a Case Worker collect income information sufficient to enable the calculation of sensitivity and specificity of the screening questions in this targeted study population. In Phase Two, we will determine sensitivity and specificity of the screening questions, when applied to a targeted population. In Phase 3, we will evaluate the effect of the intervention on patient household income.

Objectives

Our objectives were to evaluate the feasibility and extent of adoption of targeted poverty screening; to determine the proportion of patients referred for income optimization; and to evaluate the feasibility of capturing income information for patients seen after referral in a large, community-based interprofessional primary care team.

Methods

Study design

This was an implementation study with a progress-focused evaluation. We used the Standards for Reporting Implementation studies (35) to report the findings and the RE-AIM framework to report program elements (36).

Setting

At the time of the project, NYFHT provided services at 19 clinical locations; the team included 86 family physicians practicing in office groups ranging from one to six physicians, and 40 Allied Health Providers. The NYFHT served 89,000 patients, most living in the north part of Toronto. The geographic area included neighborhoods with concentrations of high-income earners as well as pockets of poverty, including the provincial health region with the greatest absolute number and the second highest proportion of residents living below the low-income cut-off (37). Prior to the study, most providers at the NYFHT did not routinely screen for poverty and there were no policies or formal educational efforts to address poverty.

Participants

All 86 NYFHT physicians were invited to participate by email. We included patients of participating physicians that were 18 years of age or older as of April 1st 2017 and had at least one primary care visit over the three previous years.

NYFHT participates in the University of Toronto Practice Based Research Network, UTOPIAN (38). UTOPIAN has expertise in the collection, management and linkage of EMR data. Through a collaborative effort, external census information using the Statistics Canada postal code conversion file (39), which contains SES quintile data for small geographic areas by postal codes, was used to determine neighborhood-level deprivation (40, 41), and this data was linked with the NYFHT's EMR data. We then proceeded to identify areas with quintiles indicating lowest income and greatest degree of material deprivation. In summary, material deprivation reflects neighborhood indicators for 1) proportion of people who have not graduated from high school, 2) ratio of employment to population, 3) mean income (40). A similar approach had been used in previous UTOPIAN projects (42, 43).

Intervention

NYFHT's Leadership endorsed the approach and provided resources, including a data manager and a project manager. The intervention lasted six months, from June 1st 2017 to November 30th 2017.

Interested physicians and their teams were offered a one-hour didactic education session prior to screening alerts being installed in their EMRs. During the session, clinicians were oriented to the importance of poverty screening; the clinical pathway for the study; and the availability of supports such as a case worker within the team. Following the education session, a referral to a case manager was added to the standard NYFHT referral form.

The medical records of adult patients with postal codes associated with the combination of lowest income and greatest degree of material deprivation were then flagged with an EMR alert. This prompted physicians or any other clinician accessing the chart to screen patients for poverty and load a standard template to record replies. To minimize cognitive load (44), the template also suggested standard workflows, including a referral to the case worker for positive screens (Additional file 2). The addition of the template to a record automatically turned off the alert.

Clinicians were prompted to ask the standard screening questions (2) during a visit for any reason.

Patients with a positive screen had the opportunity to be referred to a trained FHT caseworker that assisted with income optimization and linkage to community support, at no cost to the patient.

Data and Process measures

We recorded the following process measures: proportion of physicians participating, number of charts flagged, number of patients screened, number of patients responding 'yes' to the first screening question, of those, number of patients with up-to-date income tax filing, number of patients referred to the case worker and number of patients who saw a caseworker.

We recorded the number of physicians with a positive reply to the email invitation. UTOPIAN data was used to determine the number of patients in deprived neighborhoods. A data clerk entered the alerts in each chart.

We used EMR data for age and gender of patients; counts of patients screened; presence of the standard template in the chart (this indicated that the patient had been screened and that the screen was recorded); number of referrals and number of patients seen by the case worker. Data in the templates were used to determine answers to the screening questions. Following signed, informed consent, we surveyed patients seen by the case worker to record household income, number of financial dependents (21) and receipt of social benefits.

We used descriptive analyses (counts, proportions) for our results.

Results

Thirty-four physicians (39% of the 86 team physicians) agreed to participate in the study, ten of whom (29%) were male; 29% of NYFHT physicians (25 / 86) were male. There were 27,290 patients aged 18 or older, with at least one encounter in the previous three years, in the practices of participating physicians.

852 patients (3%) were found to be living in the most deprived neighborhoods; an alert to screen for poverty was added to their electronic chart.

Table 1 provides patient characteristics by age range and sex. Ages ranged from 18 to 95 years old. Patient age ranges and sex were generally similar for those living or not living in deprived areas.

Figure 1 provides a flow diagram and process measures for patients whose charts were flagged for poverty screening. 509 (60%) of the 852 patients living in deprived areas had at least one visit over the 6 months of follow up, and 25% of those with a visit had screening information (the standard template) entered in their record. 20% of those screened reported having trouble making ends meet, and all those had also filed their income tax reports for the previous year. 52% of patients screening positive were referred to a caseworker. Eight patients (62% of those referred) were seen by a caseworker during the study period. Patients seen by the caseworker represent less than 1% (8/852) of all flagged patients.

Table 2 summarizes the characteristics of study patients seen by a caseworker. The mean annual income was \$21,989.

During the study period, 85 additional patients (whose charts had not been flagged) were seen by the case worker for income optimization.

Discussion

Within the “routine” context of a large community-based primary care setting, we successfully identified people living in the most deprived neighborhoods and added clinical prompts to screen for poverty to their charts. A quarter of all patients flagged and with at least one visit were screened during a six-month period; 20% of those screened positive. A third of those patients were evaluated for income optimization by a caseworker, representing less than 1% of all patients flagged.

Our results demonstrate the challenges of implementing screening for poverty under usual circumstances in community-based primary care. We also demonstrate the feasibility of adding external census data to EMRs for clinical purposes, and of using targeted screening as a potentially useful method to initiate poverty screening for practice populations.

In our study, 25% of patients at risk were screened. This was in the context of limited health care provider training and multiple other priorities competing for clinician attention. This may be more representative of “routine” primary care in Canada than settings with a mandate to address poverty and other social determinants of health (14). A similar study in a Community Health Centre, with such a mandate, found that 20% of patients were screened (14). Although physicians welcomed screening tools, the authors noted that screening was likely not feasible if the physician was solely responsible, and recommended a team based approach (14). A recent study of universal screening in “real world” primary care settings reported that 9% of patients had been screened (24). Participating physicians were self-selected early adopters, committed to the process; 28% of patients screened positive, possibly indicating some targeting

of patients. In our study, 39% of physicians participated, representing a less selected group (early majority) (28). The EMR prompts could be seen by any clinician in the practice upon loading the chart; it is possible that a non-physician, such as a practice nurse, provided the screen, although we did not collect these data.

Following the RE-AIM framework (36), we report decreases at each step of the process. The largest decrease was at the initial, screening step. However, gaps developed at each following step and about 40% of patients referred did not go for an appointment. While we had taken steps to address known barriers for different stages, further exploration of specific barriers at each step, followed by specific strategies to address these are needed.

The proportion of the targeted population that received assistance with income optimization was small (1%). We do not know whether the assistance will result in increased income for those patients, whether the screening can be extended to those not living in deprived areas or could potentially become universal, or whether scaling up to other primary care settings, including non-interprofessional groups, is feasible. A large Canadian randomized controlled trial addressing several of these issues is currently under way (45).

In person primary care visits have plummeted during the Pandemic (46), limiting the ability to screen during these encounters. We do not know to what degree virtual care (phone, video visits) is associated with screening opportunities, nor what the mix of in person versus virtual visits will be, post-pandemic. It may be possible to screen for poverty outside of encounters; as an example, automated questionnaires can be securely emailed by primary care teams to their practice populations, with replies incorporated in EMRs (47).

Unexpectedly, we noted that the number of patients seen by the case worker for income optimization (85 patients) greatly outnumbered those seen through targeted screening (8 patients), possibly indicating knowledge gained or increased comfort with referrals to the case worker. Following the project, the two screening questions were added to the templates used by NYFHT physicians for routine periodic preventive health checks.

Societal inequities, including poverty, are a major public health concern. The goal should be to alleviate poverty for all those impacted, to improve their health and lives. While this intervention does not meet that goal, it provides information on a new pathway, through the identification of groups of patients at risk at the point of care. Calls for action to support those affected include social prescribing, enhanced preventive and chronic disease management visits in health care settings, and better access to psychological therapies (48). A key point of entry is the identification of persons living with income insecurity; measures to enhance this include appropriately funding and supporting screening and making it a priority in healthcare settings.

Strengths and limitations

A strength of this study is its context: it is situated within a large, multi-site community-based primary care team not specifically focused on social determinants of health. This improved the generalizability of our results.

Insights into real-world implementation challenges are highly valued by decision-makers (49); this study provides information on these challenges and we were able to address several barriers to implementation. Physicians had reported not being able to remember to screen and not remembering what questions to ask (14); we added alerts and templates that included the questions. Documentation in the EMR can be challenging (14); standardized templates made this task easier. A lack of resources for those screening positive could pose challenges (14); a referral to a NYFHT case worker was easily available for patients identified as screening positive and this was suggested in the template.

The study had several weaknesses. While a large substantial proportion of physicians volunteered, this was a self-selected convenience sample, and we did not address barriers or explore reasons for not participating. We could not capture income information on all patients that were screened; income data reflects only the sample seen by the case worker which would preclude calculations of sensitivity and specificity of screening for the targeted population. NYFHT had resources, such as a case worker and data manager, that may not be available to non-interprofessional teams.

The study duration was six months; it is possible that a longer period may have resulted in more patients being referred to a caseworker. Neighbourhood-level SES does not necessarily reflect individual SES; our targeted approach will miss “hidden poverty”.

Conclusions

A targeted poverty screening program combined with resources to optimize income for patients can be implemented in a large, distributed primary care team. Efforts to scale and spread screening for poverty, broaden the targeted population and optimize income for many more of those living in poverty, are warranted.

List Of Abbreviations

EMR: Electronic Medical Record

NYFHT: North York Family Health Team

SES: Socio-economic status

UTOPIAN: University of Toronto Practice Based Research Network

Declarations

Ethics Approval and consent to participate

The study was reviewed and approved by the North York General Hospital Research Ethics Board. Physicians agreed to participate, and patients seen by the case worker provided signed, informed consent for the survey and collection of their information.

Consent for publication

Not applicable.

Availability of data and materials

Data generated during this study area included in the published article. Data from UTOPIAN to support findings are available from UTOPIAN <https://www.dfcm.utoronto.ca/utopian-data-safe-haven> , but restrictions apply to the availability of these data.

Competing interests

The authors declare that they have no competing interests

Funding

This project was funded through a Patient's Medical Home Caring and Compassion Grant from the College of Family Physicians of Canada. Dr Greiver is supported through the Gordon F. Cheesbrough Research Chair at North York General Hospital. None of the funding sources had any role in the writing of the manuscript or the decision to submit it for publication. None of the authors received payment to write this article by a pharmaceutical company or other agency.

Authors' contributions

KW conceived the study, and was a major contributor to its design; she interpreted the data and participated in manuscript writing. MG contributed to the design, interpreted the data and wrote the first draft with MN and was a major contributor to manuscript writing. MN wrote the first draft with MG, and interpreted the data. SA was responsible for the acquisition of the data and for alerts and template design in the EMRs. BA was responsible for the generation of SES quintiles and linkage of data. SK contributed substantially to the analysis of the data. SLT contributed to the design and implementation of the project. SLT also provided caseworker support to participants. JL oversaw the implementation of the project and was involved in acquiring grant funds and training. GB participated in the design and contributed to the manuscript. ADP contributed to the manuscript. All authors reviewed the article and gave final approval for the version to be submitted for publication.

Acknowledgements

We thank NYFHT's Executive Director, Susan Griffis, for her support and enthusiasm for this project. We are grateful to the physicians and Allied Health Providers at NYFHT for their participation.

References

1. Marmot M. Social determinants of health inequalities. *Lancet*. 2005;365(9464):1099-104.
2. Brcic V, Eberdt C, Kaczorowski J. Corrigendum to "Development of a Tool to Identify Poverty in a Family Practice Setting: A Pilot Study". *International journal of family medicine*. 2015;2015:418125-.
3. Rosella LC, Fitzpatrick T, Wodchis WP, Calzavara A, Manson H, Goel V. High-cost health care users in Ontario, Canada: demographic, socio-economic, and health status characteristics. *BMC Health Serv Res*. 2014;14:532.
4. Fitzpatrick T, Rosella LC, Calzavara A, Petch J, Pinto AD, Manson H, et al. Looking Beyond Income and Education: Socioeconomic Status Gradients Among Future High-Cost Users of Health Care. *Am J Prev Med*. 2015;49(2):161-71.
5. Wilkinson RG, Pickett KE. Income inequality and population health: a review and explanation of the evidence. *Soc Sci Med*. 2006;62(7):1768-84.
6. Abedi V, Olulana O, Avula V, Chaudhary D, Khan A, Shahjouei S, et al. Racial, Economic, and Health Inequality and COVID-19 Infection in the United States. *Journal of racial and ethnic health disparities*. 2020:1-11.
7. Price-Haywood EG, Burton J, Fort D, Seoane L. Hospitalization and Mortality among Black Patients and White Patients with Covid-19. *New England Journal of Medicine*. 2020;382(26):2534-43.
8. Figueroa JF, Wadhwa RK, Mehtsun WT, Riley K, Phelan J, Jha AK. Association of race, ethnicity, and community-level factors with COVID-19 cases and deaths across U.S. counties. *Healthcare*. 2021;9(1):100495.
9. de Lusignan S, Lopez Bernal J, Zambon M, Akinyemi O, Amirthalingam G, Andrews N, et al. Emergence of a Novel Coronavirus (COVID-19): A protocol for Extending Surveillance Used by the Royal College of General Practitioners (RCGP) Research and Surveillance Centre (RSC) and Public Health England (PHE). *JMIR*. 2020.
10. Boyd A. Race-based coronavirus data not needed in Canada yet, health officials say. *The Toronto Star*. 2020.
11. Subedi R, Greenberg L, Turcotte M. StatCan: COVID-19 mortality rates in Canada's ethno-cultural neighborhoods. Ottawa, Ontario, Canada; 2020.
12. Grant K. Data show poverty, overcrowded housing connected to COVID-19 rates among racial minorities in Toronto. *The Globe and Mail*. 2020.
13. Starfield B, Shi L, Macinko J. Contribution of primary care to health systems and health. *Milbank Q*. 2005;83(3):457-502.
14. Pinto AD, Bondy M, Rucchetto A, Ihnat J, Kaufman A. Screening for poverty and intervening in a primary care setting: an acceptability and feasibility study. *Family practice*. 2019;36(5):634-8.
15. Kiran T, Pinto AD. Swimming 'upstream' to tackle the social determinants of health. *BMJ Qual Saf*. 25. England2016. p. 138-40.

16. Pinto AD, Bloch G. Framework for building primary care capacity to address the social determinants of health. *Canadian Family Physician*. 2017;63(11):e476.
17. Buchman S. Screening for poverty in family practice. *Canadian family physician Medecin de famille canadien*. 2012;58(6):709-10.
18. Social Care Guidance in the COVID-19 context Toronto, Ontario: Centre for Effective Practice; 2020 [Available from: <https://tools.cep.health/tool/covid-19-social-care-guidance/>].
19. Systems Change: Treating Tobacco Use and Dependence: Based on the Public Health Service (PHS) Clinical Practice Guideline—2008 Update. December 2012 Rockville, MD: Agency for Healthcare Research and Quality; 2012 [Available from: <http://www.ahrq.gov/professionals/clinicians-providers/guidelines-recommendations/tobacco/decisionmakers/systems/index.html>].
20. Greiver M, Aliarzadeh B, Meaney C, Moineddin R, Southgate CA, Barber DTS, et al. Are We Asking Patients if They Smoke? Missing Information on Tobacco Use in Canadian Electronic Medical Records. *American Journal of Preventive Medicine*. 2015;49(2):264-8.
21. Pinto AD, Glattstein-Young G, Mohamed A, Bloch G, Leung F-H, Glazier RH. Building a Foundation to Reduce Health Inequities: Routine Collection of Sociodemographic Data in Primary Care. *The Journal of the American Board of Family Medicine*. 2016;29(3):348.
22. Centre for Effective Practice: Poverty, a clinical tool for primary care providers Toronto2020 [Available from: <https://cep.health/clinical-products/poverty-a-clinical-tool-for-primary-care-providers/>].
23. Tu K, Sodhi S, Kidd M, Grunfeld E, Ji C, Greiver M, et al. The University of Toronto Family Medicine Report: Caring for our Diverse Populations. Toronto, Ontario: Department of Family and Community Medicine, University of Toronto; 2020.
24. Purkey E, Bayoumi I, Coe H, Maier A, Pinto AD, Olomola B, et al. Exploratory study of "real world" implementation of a clinical poverty tool in diverse family medicine and pediatric care settings. *International journal for equity in health*. 2019;18(1):200-.
25. Bloch G, Rozmovits L, Giambrone B. Barriers to primary care responsiveness to poverty as a risk factor for health. *BMC Fam Pract*. 2011;12:62.
26. Levesque JF, Haggerty JL, Hogg W, Burge F, Wong ST, Katz A, et al. Barriers and Facilitators for Primary Care Reform in Canada: Results from a Deliberative Synthesis across Five Provinces. *Health Policy*. 2015;11(2):44-57.
27. Yarnall KS, Pollak KI, Ostbye T, Krause KM, Michener JL. Primary care: is there enough time for prevention? *Am J Public Health*. 2003;93(4):635-41.
28. Rogers E. *Diffusion of Innovations*. 4th Edition ed. New York: The Free Press; 1995.
29. Tomlin Z, Humphrey C, Rogers S. General practitioners' perceptions of effective health care. *BMJ*. 1999;318(7197):1532-5.
30. Kirk MA, Kelley C, Yankey N, Birken SA, Abadie B, Damschroder L. A systematic review of the use of the Consolidated Framework for Implementation Research. *Implement Sci*. 2016;11:72.

31. Greiver M, Barnsley J, Glazier RH, Moineddin R, Harvey BJ. Implementation of electronic medical records: effect on the provision of preventive services in a pay-for-performance environment. *Can Fam Physician*. 2011;57(10):e381-9.
32. Greiver M, Barnsley J, Glazier RH, Moineddin R, Harvey BJ. Implementation of electronic medical records: theory-informed qualitative study. *Can Fam Physician*. 2011;57(10):e390-7.
33. Greiver M, Wintemute K, Griffis S, Moineddin M. Using evidence for the care of practice team populations. *Can Fam Physician*. 2014;60(3):223.
34. Sekhar DL, Pattison KL, Confair A, Molinari A, Schaefer EW, Waxmonsky JG, et al. Effectiveness of Universal School-Based Screening vs Targeted Screening for Major Depressive Disorder Among Adolescents: A Trial Protocol for the Screening in High Schools to Identify, Evaluate, and Lower Depression (SHIELD) Randomized Clinical Trial. *JAMA Network Open*. 2019;2(11):e1914427-e.
35. Pinnock H, Barwick M, Carpenter CR, Eldridge S, Grandes G, Griffiths CJ, et al. Standards for Reporting Implementation Studies (StaRI): explanation and elaboration document. *BMJ Open*. 2017;7(4):e013318.
36. Dzewaltowski DA, Glasgow RE, Klesges LM, Estabrooks PA, Brock E. RE-AIM: Evidence-based standards and a web resource to improve translation of research into practice. *Annals of Behavioral Medicine*. 2004;28(2):75-80.
37. Ontario Community Health Profiles Partnership (OCHPP) Toronto2020 [Available from: <http://www.ontariohealthprofiles.ca/>].
38. UTOPIAN Toronto: Department of Family and Community Medicine, University of Toronto; [Available from: <http://www.dfcm.utoronto.ca/research/UTOPIAN.htm>].
39. Statistics Canada Postal Code Conversion File (PCCF), Reference Guide, 2017 Ottawa, Ontario: Statistics Canada; 2020 [Available from: <https://www150.statcan.gc.ca/n1/pub/92-154-g/92-154-g2017001-eng.htm>].
40. Pampalon R, Hamel D, Gamache P, Raymond G. A deprivation index for health planning in Canada. *Chronic Dis Can*. 2009;29(4):178-91.
41. Wilkins R. Use of postal codes and addresses in the analysis of health data. *Health Rep*. 1993;5(2):157-77.
42. Aliarzadeh B, Greiver M, Moineddin R, Meaney C, White D, Moazzam A, et al. Association between socio-economic status and hemoglobin A1c levels in a Canadian primary care adult population without diabetes. *BMC Fam Pract*. 2014;15(1):7.
43. Loo CK, Greiver M, Aliarzadeh B, Lewis D. Association between neighbourhood walkability and metabolic risk factors influenced by physical activity: a cross-sectional study of adults in Toronto, Canada. *BMJ Open*. 2017;7(4):e013889.
44. Brehaut JC, Colquhoun HL, Eva KW, Carroll K, Sales A, Michie S, et al. Practice Feedback Interventions: 15 Suggestions for Optimizing Effectiveness. *Ann Intern Med*. 2016.

45. SPARK Study: Screening for Poverty And Related social determinants and intervening to improve Knowledge of and links to resources 2020 [Available from: <https://upstreamlab.org/spark-study/>].
46. Joy M, McGagh D, Jones N, Liyanage H, Sherlock J, Parimalanathan V, et al. Reorganisation of primary care for older adults during COVID-19: a cross-sectional database study in the UK. *Br J Gen Pract.* 2020;70(697):e540-e7.
47. Lofters AK, Telner D, Kalia S, Slater M. Association Between Adherence to Cancer Screening and Knowledge of Screening Guidelines: Feasibility Study Linking Self-Reported Survey Data With Medical Records. *JMIR Cancer.* 2018;4(2):e10529.
48. Moscrop A, Ziebland S, Bloch G, Iraola JR. If social determinants of health are so important, shouldn't we ask patients about them? *Bmj.* 2020;371:m4150.
49. Urquhart R, Kendell C, Geldenhuys L, Ross A, Rajaraman M, Folkes A, et al. The role of scientific evidence in decisions to adopt complex innovations in cancer care settings: a multiple case study in Nova Scotia, Canada. *Implementation Science.* 2019;14(1):14.

Tables

Table 1. Patient characteristics by age ranges and sex

| | | Patients not living in the most deprived areas, N (%)* | Patients living in the most deprived areas, N (%)* |
|-------------------|-------|--|--|
| TOTAL | | 26438 | 852 |
| Age range (years) | 18-24 | 3331 (13) | 93 (11) |
| | 25-34 | 4188 (16) | 177 (21) |
| | 35-44 | 4676 (18) | 165 (19) |
| | 45-54 | 4518 (17) | 151 (18) |
| | 55-64 | 4057 (15) | 111 (13) |
| | 65-74 | 2920 (11) | 90 (11) |
| | 75-95 | 2748 (10) | 65 (8) |
| Sex | M | 9557 (36) | 305 (36) |
| | F | 16881 (64) | 547 (64) |

*percentage may not add up to 100% due to rounding

Table 2. Characteristics of study patients who were seen by caseworker

| Total number, N | Number female, N | Mean age (range) | Number already receiving social benefits, N | Mean annual income | Number with financial dependents, N |
|-----------------|------------------|------------------|---|--------------------|-------------------------------------|
| 8 | 4 | 48 (29-63) | 4 | \$21,989* | 5 |

**Income was self-reported for seven patients and was confirmed through documentation for one patient.*

Figures




Figure 1

Flow diagram for patients flagged as living in deprived neighborhoods

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Appendices.docx](#)
- [StarIchecklistforauthorcompletion.docx](#)