

The Best Choices: the Diversity and Functions of the Plants in the Home Gardens of the Tsang-la (Motuo Menba) Communities in Yarlung Tsangpo Grand Canyon, Southwest China

Yu Zhang^{1,2}, Li-Xin Yang^{1,2}, Ming-Xiang Li^{3,6}, Yong-Jie Guo^{3,4}, Shan Li^{1,2,5}, Yu-Hua Wang^{1,2**}

1 Yunnan Key Laboratory for Wild Plant Resources, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming, China

2 Key Laboratory of Economic Plants and Biotechnology, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming, China

3 University of Chinese Academy of Sciences, Beijing, China

4 Germplasm Bank of Wild Species of China, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming, 650201, Yunnan, China

5 Key Laboratory for Microbial Resources of the Ministry of Education, Yunnan Institute of Microbiology, School of Life Sciences, Yunnan University, Kunming, 650091, China

6 State Key Laboratory of Photochemistry and Plant Resources in West China, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming, China

Abstract:

Background: Home garden is identified as a kind of small-scale land-use system which is used to manage and cultivate useful plants by local people, and home gardens can provide various plant products and services. Investigating home gardens was regarded as an effective way to understand the biodiversity related local knowledge and culture of native people in Ethnobiology and Ethnoecology. Home garden is important in less developed and remote areas. The grand canyon of Yarlung Tsangpo is designed as one of the biodiversity hotspots of China, and it is one of the most remote region of China, because of the rough traffic conditions. The aim of the present study is to collect, record and document the plants and their local knowledge and functions in the local home gardens, then attempt to answer the question: “why local people selected these plants?”

Material and Methods: The study area was in Beibeng Township of Motuo County in the grand canyon of Yarlung Tsangpo. Observation and semi-structure interviews with informed consent were used to collect data in field study. All information was collected and organized, then documented based on “ethno-species” as a fundamental unit. All of the information of local use and knowledge were organized as the list of “use-report” for quantitative analysis, and the local uses of plants were merged into 14 use categories. Frequency of citation (FC), relative frequency of citation (RFC),

** Corresponding author: Yunnan Key Laboratory for Wild Plant Resources, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming, China. E-mail: wangyuhua@mail.kib.ac.cn
Yu Zhang and Li-Xin Yang contribute equally to the article.

Cultural importance index (CI) and Cultural value index (CV) were used in quantitative analysis. Besides, the Jaccard Index was used to compare the similarity in plant species selection among different communities.

Results: A total of 78 home gardens in the nine communities of Beibeng Township were visited, and 196 ethno-species were collected. These ethno-species were identified into 188 Botanical taxa. A total of 87 home garden owners as informants were interviewed in the present study, and they provided 625 use-reports to us. The top five important plants were Su-lan-tSao (*Dendrobium nobile*), Sa-ga (*Zingiber officinale*), Soe-lu (*Capsicum annuum*), Snying-pa (*Citrus medica*) and Kham-pu (*Prunus persica*), according to the quantitative analysis. The most cited use-category was “vegetable”, followed by “Ornamental plant”, “medicine” and “fruit”. The altitude might be the most important impact factor of the plant diversity and composition of home gardens, and the traffic conditions, local terrain, was also impact the plant diversity and composition of home gardens.

Conclusion: In remote areas such as the Yarlung Tsangpo Grand Canyon, the plants in home gardens are important sources of plant products such as foods, herbal medicines and fibers to support daily lives. The local home gardens in Tsang-la communities had high diversity of plants, and these plants provided many functions and services to support daily lives of local people. Local plant knowledge, including the features, life forms, habits, habitats and use values of plants, were the summary of the understanding of local people to their surrounding plant worlds. Local people selected appropriate plants to cultivate and manage in their home gardens under the guidance of the local plant knowledge. That is the answer to the question “why local people selected these plants?”

Key Words:

Home garden; Local Knowledge; Yarlung Tsangpo Grand Canyon; Tibetan; Southwest China

1 Background

Home garden is identified as a kind of small-scale land-use system which is used to manage and cultivate useful plants by local people, and home gardens can provide various plant products and services [1-3]. Besides, the economic plants cultivated in home gardens can bring extra income to local people [3]. In some parts of the world, home gardens also have important social and cultural significances [4]. Thus, investigating home gardens was regarded as an effective way to understand the biodiversity related local knowledge and culture of native people in Ethnobiology and Ethnoecology. In remote areas, the plants in home gardens are important sources of plant products such as foods, herbal medicines and fibers to support daily lives [1-10]. Some previous studies in less developed areas indicated that most of the nutrients of local people, such as Vitamins and minerals, were from the vegetables and fruits in home gardens [3]. In home gardens, the plant products which were rich in starch can help local people to fight against famine in remote areas [7, 9-10]. Therefore, home garden is important in less developed and remote areas.

Locate in the core region of the Himalaya biodiversity hotspot, the grand canyon of Yarlung Tsangpo is designed as one of the national natural reserves of China [11-12]. As one of the deepest and longest canyons in the world, Yarlung Tsangpo have rich diversity of habitats for its huge elevation drop. Thousands of plants and animals grow here. Thus, the grand canyon of Yarlung

Tsangpo were known as “the museum of vegetation” and “the paradise of lives” [13-14]. Two counties including Milin (Smin-gling Rdzong, 米林县) and Motuo (Me-dog Rdzong, 墨脱县) which belong to Linzhi Prefecture (Nying-khri, 林芝市) of Xizang Autonomous Prefecture locate in the grand canyon, and several Chinese ethnic groups settle in the region. Gongbo Tibetan, Tsang-la and Lho-pa people is the main population of the region. They have accumulated rich local knowledge of their surrounding natural world [15]. However, people who live in the region have to face the difficulties of transport caused by the rough terrain, which limit the exchanges of products and knowledge between local communities and outside. For example, Motuo County which were deep in the Grand Canyon had no complete motor way to outside until 2013, and some townships of the county even had no motor way until early 2018 [14]. Motuo County had been one of the most backward region of China, because of the terrible transport condition. Therefore, the region became one of the key regions of the poverty alleviation program of China [15].

Most of previous studies in the region were about biodiversity, and just few studies involved the relationship between local people and biodiversity such as the local knowledge of plants [13]. In our preliminary Ethnobotanical studies of local people in the region, we found rich local knowledge of plants [16-18]. The functions of the plants in home gardens played important roles in the daily lives of local people. And, some previous studies in the of Himalaya area outside China showed rich diversity and local knowledge of plants in traditional home gardens [19-21]. Thus, it is necessary to investigate the home gardens of local communities in the grand canyon of Yarlung Tsangpo.

Tsang-la people settle in the Yarlung Tsangpo basin of Xizang Autonomous Prefecture of China [22]. The Tsang-la language belongs to the Tibetan Language Branch of Chinese-Tibetan Language Family. Because the ancestors of Tsang-la people originated from the “Menyu (门隅)” district of Tibetan Plateau, they were also identified as “Motuo Menba (墨脱门巴)”, a branch of the Chinese Menba nationality (门巴族). “Motuo Menba” means “Menba people who live in Motuo County”. The history of Tsang-la people settling in Motuo County could be traced to over 300 hundreds years ago (early Qing Dynasty of China). Therefore, Tsang-la people were also called as “Pad-ma-rkos-pa” in Tibetan language, which meant “people of Pad-ma-rsko”. “Pad-ma-rkos (白玛岗)” was the ancient name of Motuo County in Qing Dynasty of China [22-23].

As mentioned above, investigating home gardens was regarded as an effective way to understand the biodiversity related local knowledge and culture of native people in Ethnobiology and Ethnoecology. Because the diversity and functions of the plants in local home garden was related closely to the local plants knowledge, which contained the understanding of local people to the plant world of their surrounding environments. Thus, the core aim of the present study was to answer the question: Why local people selected these plants? And, we wanted emphasize that the answer should be concluded from the local knowledge.

Specifically, the research questions of the present study are 1) How many and what plants are in the home gardens of local communities, and what are local knowledge and functions of these plants? 2) What are the most popular plants and why they are important for local communities? 3) Why people select these plants, and what are the impact factors of the plant composition in local home gardens? The main works of the present study include 1) collecting, recording and documenting the plants in the local home gardens, 2) visiting the home gardens and interviewing the owners of them,

then recording the local knowledge of plants, and 3) documenting the local knowledge of the functions of home gardens and attempting to answer that why local people selected these plants?

2 Material and Methods

2.1 study area and communities

The study area was in Beibeng Township (Hbras-spung Shang, 背崩乡) of Motuo County. Beibeng Township were deep in the grand canyon of Yarlung Tsangpo. Almost all of the permanent population in Beibeng Township is Tsang-la people [24]. Beibeng Township was the remotest township of Motuo County, and the motor ways linked among the communities were not fully available until early 2018, when the field works of the present study were being performed in the township.

From 400m to 3260m, the huge variations of altitude and variety of complex topography resulted diversification of habitat types in the township. From the bottom of the canyon to the top of mountains, the vegetation was changing from tropical rain forest and subtropical evergreen board-leaved forest to alpine coniferous forest or Rhododendron shrubs. These factors gave the township rich biodiversity [12-13].

The basic unit of Tsang-la community was “Gyong-tsho”. A “Gyong-tsho” usually consisted several families related to each other. These families usually originated from the same ancestor. Therefore, in the present study, we took “Gyong-tsho” as the unit of local communities. Beibeng Township contained nine “Gyong-tshos” which located at the both sides of Yarlung Tsangpo River. Each Gyong-tsho was identified as an administrative village by Chinese authority. These communities distributed in different places of the township. These communities had different elevations and surrounding vegetation.

2.2 Field Study and Data Collection

The field study was carried out in the nine communities of Beibeng Township from 2017 to 2018, and we visited the communities in different seasons. First, we visited the local community committee for getting field study permission. We explained our purpose to the community leaders and requested assistance from them. The assistance included providing local guides, introducing us to the community members and other necessary helps. All of our field studies were carried out with informed consent.

In the present study, we set “a home garden” as a plot of land which was used to cultivate useful plants except large-scale planted crops and separated by artificial barrier with other lands in the communities (Fig 2). Observation and semi-structure interviews were used to collect data in field study, and the semi-structure interviews were carried out with the cooperation of local guide. First, we went to visit a community member who was familiar with our local guide, and we asked him whether we could visit his/her home garden, after the guide introduced us and explained our purpose. Then, if he/she allowed us to visit the home garden, the observation and semi-structure interviews

were carried out. The observation was used to record the species, sizes and frequency of the plants in the home gardens, and the semi-structure interviews were used to get the information of the local knowledge of the plants. The local knowledge here were the knowledge provided by local people, including vernacular names, uses, features, life forms, habits, habitats, economic values and culture significances of plants. The informants of the interviews were the owners of the home gardens. The semi-structure interviews were performed based on the following questions:

Do you mind to introduce the plants you cultivated in your garden? And how many kinds of plants you had cultivated in your garden?

What the purpose you cultivate it? And how to use/cook/prepare it? (For one plant)

Do you mind to tell us some stories of the plant? If it is important to you/your family/your community/your religion, could you please told us more details about the importance. (For ritual plants or the plants with special or symbolic significance)

After we finished the study of one informant's home gardens, we would ask him/her to introduce us to his/her neighbors or relatives in the same community. Then we went to visit next informant. The observation and interviews would be attempt to perform in consistent strategy in each home garden.

2.3 Information documentation and plant specimen identification and preservation

All information was collected and organized, then documented based on "ethno-species" as a fundamental unit. All of the "ethno-species" were finally identified into the scientific taxa of plants. Ethno-species was defined as a "species entity" which matched a plant product with local name, and an ethno-species was usually identified according to particular usages or features such as colors, shapes and tastes of it by local people [25-26]. Ethno-species was used as the primary unit of analysis in Ethnobotany, and it could reflect the local knowledge about the plants of local people in the study area.

The interviews were performed in Mandarin and Tsang-la Language, then Tsang-la language were translated to Mandarin by the local guidance. The vernacular names of plants were recorded in Tsang-la Language which was usually written in Tibetan script [27]. In this paper, we recorded the vernacular names in the Latin form of Tibetan script, due to that most of the computer systems could not display Tibetan script correctly without installing special language packs. The spelling rules of Latin form of Tibetan script could be found in the related website of omniglot.com (<https://www.omniglot.com/writing/tibetan.htm>) [28].

In the field study, we collected the voucher specimen and took photos to the plants. For the native and local featured plants, we collected the voucher specimen, and for the common cultivated plants such as *Brassica rapa*, *Raphanus sativus* and *Capsicum annuum*, we took photos as the vouchers. The specimen and photos were collected with the informed consent and permission from local administrative department, and these plant specimens were identified at Herbarium of Kunming

Institute of Botany (KUN). The plant specimens were identified follow Flora of China and Flora of Xizang [29-30]. The species names were checked with The Plant List [31].

2.4 Quantitative Analysis

All of the information of local use and knowledge were organized as the list of “use-report” for quantitative analysis [32]. A use-report (UR) is counted each time an informant mentions the use of an ethno species in one of the categorized uses.

The local uses of plants were merged into 14 use categories containing fruit, ornamental plants, ritual plants, grain, wood, wine brewing, knitting, dye plants, vegetable, forage plants, spice, medicine, oil plants and others.

2.4.1 Frequency of citation (FC) and relative frequency of citation (RFC)

Frequency of citation (FC) was used to calculation the percentage of each kind of uses and other indices. The FC is calculated as is the sum of informants that cite a use for a particular species. Relative frequency of citation (RFC) was used to show the importance of each ethno species in the study area [32]. The values of RFC were calculated according to the formula:

$$RFC = \frac{FC_s}{N}$$

Where FCs is the Frequency of Citation (the number of informants who mention the use of a species) and N is the total number of informants in the survey. A high RFC value for a species indicates that the species is used both frequently and by a high proportion of informants in the study area.

2.4.2 Cultural importance index

The cultural importance index (CI) was used to reflect the prevalence of use and diversity of use of each ethno species [32]. The value of CI was the sum of proportion of informants that mention each of the use categories for a given ethno species. The CI was defined by the following formula:

$$CI = \sum_{u=1}^{NC} \sum_{i=1}^N \frac{UR_{ui}}{N}$$

Where UR_{ui} was the total number of use-report for each use category of a given ethno species, N was the total number of informants and NC was the total number of the use categories.

2.4.3 Cultural value index

Cultural value index (CV) was developed by Reyes-Garcia et al [33], was the product of three

factors which were calculated follow the formula:

$$CV_s = \left[\frac{NU_s}{NC} \right] \times \left[\frac{FC_s}{N} \right] \times \left[\sum_{u=1}^{NC} \sum_{i=1}^N \frac{UR_{ui}}{N} \right]$$

Factor 1: the number of use categories for the given ethno species (NUs) divided the total number of all use categories (NC). Factor 2: the relative frequency of citation of the given ethno species (RFC). Factor 3: The cultural importance index of the given ethno species (CI).

The values of CV varied from 0 to FC. A high value of CV meant that most of the informants thought the ethno species was preferred to use.

2.4.4 Jaccard Index and multidimensional scaling

The Jaccard Index (J) is also known as the Jaccard similarity coefficient, is a statistic index for comparing the similarity and diversity between two samples [34]. In the present study, the Jaccard Index was used to compare the similarity in plant species selection among different communities in the study area. The Index was calculated following the formula:

$$J(A, B) = \frac{A \cap B}{A \cup B - A \cap B}$$

Where A and B was the set of species in the home gardens of community A and B, $A \cap B$ was the number of the species found in both community A and B (the intersection of A and B), and $A \cup B$ was the total number of the species found in community A and B (the union of A and B). A high value of Jaccard Index meant that the species of the home gardens were similar between two communities.

The values of Jaccard Index were converted to the Jaccard Distance (JD) which was the complementary set of Jaccard Index.

$$JD(A, B) = 1 - J(A, B)$$

The purpose of the use of the values of the Jaccard Distance was to visualize the values of the Jaccard Index with multidimensional scaling. First, we took the values into the dissimilarity matrix. Then, we use the multidimensional scaling with ALSCAL in SPSS 19 to draw scatter plot graph, with the model of Euclidean Distance.

3 Results

3.1 Characteristics of the communities

A total of 78 home gardens in the nine communities of Beibeng Township were visited (Fig 1). The numbers and plant diversity of the home gardens in each community varied with the community

population, terrain, elevation, and surrounding vegetation (Table 1). Hbras-spung was the administrative center of the township.

Table 1 the study communities

Fig 1 the location of study communities

Fig 2 the home gardens in the Tsang-la communities in Beibeng Township

Notes: 1: A typical small “private” home garden in Spa-gdeng, the home garden belonged to just one household. 2: A typical home garden. 3: The biggest “public” home garden in Spa-gdeng, it was shared by over ten households.

3.2 The diversity of the plants and their uses

In the present study, 196 ethno-species were collected, and they were identified into 188 Botanical taxa, which were identified two to subspecies level and 181 to species level and five to the genera level, and they belonged to 160 genus and 78 families (Appendix 1). The most cited family was Asteraceae (12 species), followed by Orchidaceae (10 species), Cucurbitaceae (nine species) and Poaceae (nine species), and Forty-four families just only included one species. On genera level, the most frequent genus were *Cymbidium*, *Dioscorea* and *Solanum*. Forty species were estimated and catalogued by the International Union for Conservation of Nature (IUCN), and four of them were categorized as endangered species. They were *Coptis teeta* (EN), *Cephalotaxus hainanensis* (EN), *Schlumbergera truncate* (VU) and *Juniperus pingii* (NT) [41]. Among them *S. truncate* were exotic cultivated flowers while others were native plants introduced from local forests. All of the plants of Orchidaceae were recorded as endangered plants in the Information System of Chinese Rare and Endangered Plants (ISCREP) [42].

One hundred and twelve ethno-species were herbaceous plants, and 40 were trees and 26 were climbers and 18 were shrubs. Over half (122) of these ethno-species were introduced from native place or cultivated for a long time. Thirty-three ethno-species were introduced from Lhasa, 32 from other provinces of china, and seven from foreign countries (one from Bhutan and six from India).

A total of 87 home garden owners as informants were interviewed in the present study, and they provided 625 use-reports to us. These use-reports were categorized into 14 use-categories. The most cited use-category was “vegetable” (218 use-reports), followed by “Ornamental plant” (108 use-reports), “medicine” (92 use-reports) and “fruit” (90 use-reports). The use-reports in these four use-categories accounted for most (81.3%) of the total numbers of use-reports. Twenty-nine ethno-species were used in over two use-categories ($NU/NC > 0.14$), and among them, six ethno-species were used in three use-categories ($NU/NC = 0.21$). No ethno-species was used in over three use-categories.

3.3 The most popular plants

In total 196 ethno-species, fifteen of them had over 10 use-reports, while 85 of them had just cited one times (one use-report) respectively by the informants. The ethno-species had over 10 use-reports

were also had the highest values of CI and RFC. The Cultural value index (CV) combined the values of CI, RFC and NU/NC. Thus, from the values of CV of the ethno-species, the top five most popular ethno-species were Su-lan-tsao (*Dendrobium nobile*, CV=0.0064), Sa-ga (*Zingiber officinale*, CV=0.0044), Soe-lu (*Capsicum annuum*, CV=0.0028), Snying-pa (*Citrus medica*, CV=0.0020) and Kham-pu (*Prunus persica*, CV=0.0019) (Table 2). These plants were all cultivated for a long time by local people.

Table 2 the Top five popular ethno-species

3.4 Comparison of the species among the communities

The results of Jaccard Index were in Table 3 and Fig 2.

Table 3 the Jaccard Index of comparison the similarity in plant species selection among different communities in the study area

Fig 3 the multidimensional scaling (ALSCAL) based on Jaccard Distance

4 Discussion

4.1 Diversity and uses of the plants in home gardens

4.1.1 The important plants in home gardens

Based on the quantitative analysis, we found that some plants were important. These plants were cultivated commonly in the home gardens or cited frequently by the informants (Fig 3).

Fig 4 the Top 5 important plants in home gardens

Notes: 1: *Dendrobium nobile*, 2: *Zingiber officinale*, 3: *Capsicum annuum*, 4: *Citrus medica*, 5: *Prunus persica*, 6: *Prunus mira*

Su-lan-tsao (*Dendrobium nobile*) was the most frequently cited and cultivated plants in the home gardens. The plant was cultivated as featured product in the tropical and subtropical regions of Motuo County. *D. nobile* was the original plant of the Traditional Chinese Medicine “Shi-hu (石斛)” and the Traditional Tibetan Medicine “Bu-shes-tse” [35-36], and it could be sold for a high price in traditional medicine markets. Thus, *D. nobile* was popular to cultivated by local people for selling. Besides, another purpose of Tsang-la people to cultivate the plant was for decorating home gardens, Because of its beautiful flowers.

Sa-ga (*Zingiber officinale*) was multipurpose in local communities. The tender rhizomes of *Z. officinale* were seasonal vegetable in springs, and the mature rhizomes which were rich in aromatic oil were used as spice in cooking. Local Tsang-la people also used the rhizome as medicine to treat

cold. *Z. officinale* was the original plant of Traditional Tibetan Medicine “Ga-kya” which was used to treat digestion problems and the complications of “rlung (wind)” and “bad-gan (fluid)” diseases [36].

Capsicum annuum was the most important spice flavor in typical Tsang-la daily cuisines, and it was used in almost every local dish. Two ethno-species of *Capsicum annuum* were cultivated by Tsang-la people in the study area. The two ethno-species were identified based on the pungency level and origin places. One was named as “Soe-lu” which had temperate pungency, and the other one was named as “Bon-ga-soe-lu” which had strong pungency. “Soe-lu” was native while “Bon-ga-soe-lu” was introduced from India. We found that the morphological feature of “Bon-ga-soe-lu” was similar with “Bhut Jolokia (India's ghost pepper)” of India.

Nying-pa (*Citrus medica*) was common in local home gardens. Tsang-la people cultivated it for collecting fruits as medicine, traditionally. Tsang-la people collected the fruits of *C. medica*, then sliced up and dried the fruits in shade. The dried fruits were used as herbal tea to treat cold in local Tsang-la medicine. Local people named *C. medica* as “Motuo Da Ningmeng (墨脱大柠檬, Big Lemon of Motuo County)” in Chinese language, because the smell of its fruits were very similar with lemon. Besides, *C. medica* was the origin plant of Traditional Chinese Medicine “Xiang-yuan (香橼)” [35], so local people also collected its fruits and sold to traditional druggists.

Kham-pu (*Prunus persica*) was another highly frequent ethno-species in local home gardens. *P. persica* was cultivated as fruiter. Besides, Kham-pu was an important ritual plant in traditional culture which symbolized auspiciousness in Xizang district [37]. Another *Prunus* species, *P. mira* was also called as Kham-pu by local people, but local people thought it was not the same ethno-species with *P. persica*. Because the species was used for other purpose and not harvesting fruits. *P. mira* was common wild plant in Yarlung Tsangpo Canyon, and local people introduced it into home gardens as rootstocks for grafting *P. persica*.

4.2.2 Local knowledge of the identification of ethno-species

The diversity of ethno-species could indicate the local knowledge of understanding plants. In the study area, Tsang-la people identified the ethno-species from the use purpose and origin of plants. The ethno-species could mostly match to the botanical species, but sometimes under-differentiated or over-differentiated to botanical species [25-26]. For example, *Brassica rapa* (total UR=31) was the most frequent botanical species in the present study. *Brassica rapa* was identified as four ethno-species by local people, according to the differences of their cooking methods and morphological characteristics among them. For another example, Tsang-la people divided the plants of genus *Musa* into two ethno-species as “Tsun-ma-lai-su”, of which the tender shoots and flowers were edible as vegetable, and “A-ni-lai-su”, of which the fruits were edible as banana, although we had found that at least four botanical species of the genus were used by local people. Further, Tsang-la people divided the local cultivated populations of “A-ni-lai-su” into “Krun-go-a-ni-lai-su”, which was native or introduced from other regions of China, and “Ni-ra-a-ni-lai-su”, which was introduced from India, based on their origin places.

4.2 The important functions of the local home gardens

4.2.1 The self-service “food supermarkets” and “herb shops” of local people

Home gardens not only had high plant diversity, but also many functional plants including edible and medicinal plants [1-2]. From the previous studies, food plants were the most frequently cited in most of the home gardens through the world [1-3]. The results of our study was no exception. The plants used as vegetable (218 use-reports) were cited most frequently in the home gardens in the present study. The edible plants could provide carbohydrates, proteins, vegetable oil, vitamins and minerals [3]. Our previous study in Yunnan province of China showed that the edible plants in home gardens and wild could help people to fight against famine [38]. Nepali experts also found that home gardens were important to food supply in remote mountain area of Nepal [7].

Asteraceae was the most frequently cited families containing 12 botanical species in the present study, and seven of them were used vegetables. The plants of genus *Dioscorea* and *Solanum* were cited frequently, these species were used as food and medicine.

Thirty-three ethno-species were cultivated for medicinal uses. These medicinal plants were used to treat common ailments by local people, and some of them were sold to the druggists. The previous studies in Yunnan Province of China and Chiang Mai of Thailand also showed similar results [5-6].

Home gardens were also used as “crops seedling bases” by local people. Rice was the staple crop in the low altitude regions of Beibeng Township (the Tsang-la language “Hbras-spung” meant “rice field”). Local people raised the rice seedlings in the home gardens then planted the seedlings into the field in every spring. Beside rice seedlings, the seedlings of vegetables, fruit trees and some medicinal plants were also raised in home gardens. Local people usually built small greenhouses for crop seedling in their home gardens.

Besides, the home gardens were also the bases of local people for introducing alien plants from other places, and most of these plants were edible or medicinal plants. These plants enriched the food tables and herb chests of local people.

4.2.2 The base of local native endangered plants conservation and sustainable use

Four species in the home gardens were designed as endangered plants by IUCN, and three of them including *Coptis teeta* (EN), *Cephalotaxus hainanensis* (EN) and *Juniperus pingii* (NT) were native species [41]. All of the plants of Orchidaceae were recorded as endangered plants in the Information System of Chinese Rare and Endangered Plants (ISCREP) [42]. These plants were important for local medicine, ornament and religion (table 5). Local people introduced them from forests and cultivated and bred them carefully in their home gardens. Some previous studies indicated that this phenomenon could help conserve the local endangered useful plants, and it was an efficient way to rare plant resources sustainable uses [1-4]. It was worth mentioning that local people told us that the reason why they cultivated these endangered plants was for increasing the available resources of these plants. “These plants are extremely rare”, a home garden owner told us,

“if we always collect these plants in the forest rather than attempt to cultivate them, they will disappear”.

Table 4 the endangered plant species in the home gardens

4.2.3 The important part of beautiful homes

In the present study, we recorded 49 ethno-species which were cultivated as ornamental plants, and 32 of them were called as “Mong-nang” by local people. “Mong-nang” meant “beautiful flower” in Tsang-la language. In Tsang-la communities, “Mong-nangs” were important decorations in home gardens and balconies. Local people collected ornament plants from native forests and other places, then introduced to their home gardens. The informants told us that a big part of the “Mong-nangs” (we had recorded 29 ethno-species) were introduced from Lhasa, the center city of Xizang Autonomous Region, and they got the seedlings in Lhasa, and secured them carefully on the long way home. The informants claimed that beautiful flowers were essential in their lives. Interestingly, local people did not seem to know the Tsang-la name of each “Mong-nang”, but called them collectively as “Mong-nang”. It might be because that most of these plants were introduced from other places, they had no Tsang-la names. According to the study on traditional culture of the ethnic groups in Tibetan plateau, the ideal homes should be surrounded by beautiful gardens with many blooms [39].

Interestingly, the ornamental plants of Orchidaceae especially Genus *Cymbidium* were cited frequently and cultivated carefully in the home gardens by local people. Local people named them as “Lan-tshao”. We inferred that the name was from Mandarin “Lan-cao (兰草)”. The ornamental Orchidaceae plants were important in traditional Chinese culture, and they could fetch a good price in flower markets [40].

4.3 The impact factors of the plant diversity and arrangement in home gardens

From the results of multidimensional scaling (ALSCAL) (Fig 3), we could divide the communities into four groups, with their positions in the four quadrants of Fig 3. We found that all of the communities in quadrant 1 and 2 located on the west side of Yarlung Tsangpo River while most of the communities except Spa-gdeng in quadrant 3 and 4 located on the east side of the river. And, the altitudes of all of the communities in quadrant 1 and 4 were over 1000m while the altitudes of most of the communities except Spa-gdeng in quadrant 2 and 3 were below 1000m.

Spa-gdeng shared many home garden plants with the low altitude communities on the east side of Yarlung Tsangpo River especially Hbras-spung ($J=0.19$, the second-highest of the values of Jaccard Index), although it was a high altitude community located on the west side of the river. Why Spa-gdeng was so different with other communities? We inferred that it might be because Spa-gdeng was near the main road, which provided many chances for the people of Spa-gdeng to exchange plants and information with people from other places.

Therefore, we inferred that the altitude might be the most important impact factor of the plant

diversity and composition of home gardens, and the traffic conditions also impact the plant diversity and composition.

Besides, other factors, such as local terrain, was also impact the plant diversity of home gardens. Hbras-spung and Gder-kong had not only the most number of home gardens but also the highest diversity of plants in the home gardens. We inferred that the possible reason was that the two communities located at mesas or broad valleys, and they had more available flat fields for cultivating than other communities which usually located at rough terrain places. One home garden usually belonged to one household, in general. But in the local communities, we found that some home gardens were not private but shared by several households. Because of the limited available flat fields, the households usually shared home gardens, especially in the communities located at rough terrain places, such as Spo-gdong, Spa-gdeng and Jang-shing. Thus, we categorized the local home gardens into “private” and “public” home gardens (Fig 2). A typical example was that, in Spa-gdeng, most of the households shared two big public home gardens, while just few households could have their own private small home gardens.

4.4 Issues of the impact of the old houses reconstruction to the local home garden

Dig-gdong was a large community included 130 households, and the community had enough flat fields. But the big project of old houses reconstruction were in progress in the community, and most of the old facilities including houses and home gardens were dismantled. Thus, just only four survive home gardens were recorded. Interestingly, we recorded the largest number of home gardens and richest diversity of plants in Hbras-spung, of which the old houses reconstruction projects had been finished several years ago. Because the information of the plants in the home gardens in Hbras-spung before the old houses reconstruction projects was not available, we could not do further analysis. Therefore, it was difficult to know how the projects impact the diversity and composition of plants in the home gardens of local communities.

With the rapid economic development, the old houses reconstruction had implemented in more and more rural communities of developing regions [23]. However, few studies focused on that how the old houses reconstruction change the landscape patterns including the plant composition of home gardens of rural communities. Future researches should pay more attention to the issues.

5 Conclusion

In remote areas such as the Yarlung Tsangpo Grand Canyon, the plants in home gardens are important sources of plant products such as foods, herbal medicines and fibers to support daily lives. The local home gardens in Tsang-la communities had high diversity of plants, and these plants provided many functions and services to support daily lives of local people. A total of 78 home gardens in the nine communities of Beibeng Township were visited, and 196 ethno-species were collected. These ethno-species were identified into 188 Botanical taxa, which were identified two to subspecies level and 181 to species level and five to the genera level, and they belonged to 160

genus and 78 families. The top five important plants were Su-lan-tsao (*Dendrobium nobile*), Sa-ga (*Zingiber officinale*), Soe-lu (*Capsicum annuum*), Snying-pa (*Citrus medica*) and Kham-pu (*Prunus persica*), according to the CV values. The most cited family was Asteraceae, followed by Orchidaceae, Cucurbitaceae and Poaceae. A total of 87 home garden owners as informants were interviewed in the present study, and they provided 625 use-reports to us. These use-reports were categorized into 14 use-categories. The most cited use-category was “vegetable”, followed by “Ornamental plant”, “medicine” and “fruit”. We inferred that the altitude might be the most important impact factor of the plant diversity and composition of home gardens, and the traffic conditions also impact the plant diversity and composition. Other factors such as local terrain, was also impact the plant diversity and composition of home gardens.

Local plant knowledge, including the features, life forms, habits, habitats and use values of plants, were the summary of the understanding of local people to their surrounding plant worlds. Local people got the information of plants by observation, intuition, experience and trial, then summarized them as local knowledge by their own language. Finally, local people selected appropriate plants to cultivate and manage in their home gardens under the guidance of the local plant knowledge. These plants which had various functions provided necessary products and services to support the daily lives of local people. That is the answer to the question “why local people selected these plants?”

Reference

1. Kumar B M, Nair P K R. Tropical Homegardens. Springer Netherlands; 2006.
2. Huai HY, Hamilton A. Characteristics and functions of traditional homegardens: a review. *Frontiers of Biology in China*. 2009; 4(2): 151-157
3. Mohri H. et al. Ecosystem Services of Traditional Homegardens in South and Southeast Asia. In: Takeuchi K., Saito O., Matsuda H., Mohan G. (eds) *Resilient Asia*. Tokyo, Science for Sustainable Societies. Springer; 2018
4. Konoutan Médard Kafoutchoni, Idohou R , Egeru A , et al. Species richness, cultural importance, and prioritization of wild spices for conservation in the Sudano-Guinean zone of Benin (West Africa). *Journal of Ethnobiology and Ethnomedicine*. 2018, 14(1):1-30.
5. Prateep, Panyadee, Henrik, et al. Medicinal plants in homegardens of four ethnic groups in Thailand. *Journal of Ethnopharmacology*. 2019, 15,239,111927.
6. Yang L , Ahmed S , Stepp J , et al. Comparative homegarden medical ethnobotany of Naxi healers and farmers in Northwestern Yunnan, China. *Journal of Ethnobiology and Ethnomedicine*. 2014; 10, 6.
7. Gotor E. Home gardens in Nepal. *Biodiversity International*; 2013, 4p.
8. Abebe T , Wiersum KF , Bongers FJJM , et al. Diversity and dynamics in homegardens of Southern Ethiopia. *Tropical Homegardens*. 2006, 3(2):123-142.

9. Gautam R , Sthapit B , Subedi A , et al. Home gardens management of key species in Nepal: a way to maximize the use of useful diversity for the well-being of poor farmers. *Plant Genetic Resources*. 2009; 7, 02,142.
10. Enu Kwesi L, Gyasi E A, Vordzogbe V V. Managing the home garden for food security and as a germplasm bank.// *Lessons from West Africa in Sustainable Use of Biodiversity and Related Natural Resources Managing Agrodiversity the Traditional Way*. 2004.
11. Myers N , Mittermeier R A , Mittermeier C G , et al. Biodiversity hotspot for conservation priorities. *Nature*. 2000; 403, 6772, 853-858.
12. Deng LB, Chen DL, Deng LQ, Ecological Evaluation of Yarlung Zangbo Grand Canyon National Nature Reserve in Tibet . *SCIENTIA SILVAE SINICAE*. 2011, 47, 5, 1-6.
13. Sun H and Zhou ZK. The Phytogeographical Affinities and Nature of the Big Bend Gorge Of the Yalu Tsangpo River, SE Tibet, E. Himalayas. *Chinese Journal of Applied and Environmental Biology*. 1997; 3(2):184-190.
14. The People's Government of Linzhi. Introduction to Linzhi Prefecture-level City. <http://www.linzhi.gov.cn/linzhi/zmlz/zmlz.shtml> Access 01 JAN 2020.
15. Liu YH. Study on industrial development in motuo county, Tibet (in Chinese). *China Tibetology*. 2013; 03, 32-36.
16. Li S., Zhang Y., Guo Y. et al. Monpa, memory, and change: an ethnobotanical study of plant use in Mêdog County, South-east Tibet, China. *J Ethnobiology Ethnomedicine*. 2020; 5, 16.
17. Yang J, Chen WY, Fu Y, et al., Medicinal and edible plants used by the Lhoba people in Medog County, Tibet, China . *Journal of Ethnopharmacology*. 2020, 249,112430.
18. Chen WY, Wang YH, Wang YH. *Xizang Motuo Xian Luoba Zu Chuantong Liyong Zhiwu (Traditional Useful Plants of Luoba Ethnic Nationality in Motuo County, Xizang, China) (in Chinese)* . Yunnan Science and Technology Press, Kunming, China, 2019.
19. Barbhuiya A R, Sahoo U K, Upadhyaya K. Plant Diversity in the Indigenous Home Gardens in the Eastern Himalayan Region of Mizoram, Northeast India. *Economic Botany*. 2016; 70, 2,115-131.
20. Zaman, Sourovi, Siddiquee, Salah Uddin, Katoh, Masato. Structure and Diversity of Homegarden Agroforestry in Thakurgaon District, Bangladesh. *Open Forest Science Journal*, 2010; 3, 38-44.
21. MELKANIA, N. P. Biodiversity in Forest and Rangeland Ecosystems in Indian North-eastern Himalayan Region. *Indian Forester*. 2007; 12, 1609-1635.
22. Wu CZ. The Historical Evolution of the Menba People in Motuo County, Xizang (in Chinese). *Journal of Minzu University of China*. 1987; 1, 37-41.

23. Research on Traditional Settlements Space of the Menba Nationality in Medog County . Master Dgree, Beijing University of Civil Engineering and Architecture. 2017
24. Fan Xiaopeng, Deng Xiaocong. The Fieldwork of Border Settlement of the Motuo Area in Tibet. *Research on Heritages and Preservation*. 2016; 1, 4 31-36.
25. Berlin B: *Ethnobiological Classification: Principles of Categorization of Plants and Animals in Traditional Societies*. NJ: Princeton University Press Princeton; 1992.
26. Otieno, J., Abihudi, S., Veldman, S. et al. Vernacular dominance in folk taxonomy: a case study of ethnospecies in medicinal plant trade in Tanzania. *Journal of Ethnobiology and Ethnomedicine*. 2015; 11, 10.
27. Pudo. Menba Nationality in Linzhi (in Chinese). *Tibetan Folklore*, 2002; 2:22-23.
28. Simon Ager. Tibetan writing system in Omniglot: an encyclopedia of writing systems and languages. <https://www.omniglot.com/writing/tibetan.htm>, Access in 2019
29. Editorial Committee of Flora of China. *Flora of China*. Beijing, Science Press; 2013.
30. Editorial Committee of Flora of Xizang. *Flora of Xizang*. Beijing, Science Press; 1983.
31. The Plant List. Version 1.1. Published on the Internet; <http://www.theplantlist.org/>, accessed 1st January, 2013.
32. Tardío J, Pardo-de-Santayana M. Cultural importance indices, a comparative analysis based on the useful wild plants of Southern Cantabria, Northern Spain. *Economic Botany*. 2008; 62:24–39.
33. Victoria Reyes-García, Tomás Huanca, Vadez V , et al. Cultural, practical, and economic value of wild plants: A quantitative study in the Bolivian Amazon. *Economic Botany*. 2006, 60(1):62-74.
34. Jaccard P, THE DISTRIBUTION OF THE FLORA IN THE ALPINE ZONE.1. *New Phytologist*. 1912; 11: 37-50.
35. The Pharmacopoeia Committee Editorial Department (Eds.): *Pharmacopoeia of the People's Republic of China*. Beijing, Chemical Industry Press; 2010.
36. State Administration of Traditional Chinese Medicine: *Chinese materia medica*. Shanghai, Shanghai Scientific and Technical Publishers; 2002.
37. Robert Beer. *The hand book of Tibetan Buddhist symbols*. Boston, Shambhala Publications, 2003.
38. Zhang, L., Chai, Z., Zhang, Y. et al. Ethnobotanical study of traditional edible plants used by the Naxi people during droughts. *Journal of Ethnobiology and Ethnomedicine*. 2016; 12, 39.

39. Zhu HH. The state of Zen and Buddhism Nature: A comparison of Aesthetic of Han and Tibetan Classic gardens (in Chinese). *Journal of Central South Minzu University*. 2008; 28, 1, 68-78.
40. Zhou JZ. *Lan Wenhua (Orchid Culture)*. Beijing, China Agriculture Press; 2001.
41. IUCN. The IUCN Red List of Threatened Species. Version 2019-2. <https://www.iucnredlist.org> Accessed in 2019.
42. Institute of Botany, Chinese Academy of Sciences. The Information System of Chinese Rare and Endangered Plants. <http://www.iplant.cn/rep/> Accessed in 2019.

Acknowledgement

We are very grateful to the informants for sharing their knowledge with us. We also are very glad to say “Bha-dra” (Tsang-la language: thank you) to all of the local people who had helped us in the field works. We thank for technical guidance of prof. Pei Shengji. Besides, we thank to Mr. Xu and Mr. Shen as auto drivers in the wild works, Mr. Quansheng as Tsang-la translator, and the self-study course of Tibetan spelling from Beijing GaKaWa Tibetan Language learning organization.

Funding

The study was funded by “the second Tibetan Plateau Scientific Expedition and research (No.2019QZKK0502)” Project of China.

Author information

Affiliations

1 Yunnan Key Laboratory for Wild Plant Resources, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming, China;

2 Key Laboratory of Economic Plants and Biotechnology, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming, China;

3 University of Chinese Academy of Sciences, Beijing, China;

4 Germplasm Bank of Wild Species of China, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming, 650201, Yunnan, China;

5 Key Laboratory for Microbial Resources of the Ministry of Education, Yunnan Institute of Microbiology, School of Life Sciences, Yunnan University, Kunming, 650091, China;

6 State Key Laboratory of Photochemistry and Plant Resources in West China, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming, China

Authors' contribution

WYH organized the study team, and provided the technical supports and guidance. ZY and YLX designed and executed the research plan. ZY wrote the manuscript. LMX and LS recorded and organized the data. GYJ and ZY identified the specimen and checked the information. All authors took part in the field works. All authors were involved in the drafting and revising of the manuscript and approved the final revision.

Corresponding author

Correspondence to Yu-Hua Wang.

Ethics declarations

Ethics approval and consent to participate

The authors asked for permission from the local authorities and the people interviewed to carry out the study.

Consent for publication

The people interviewed were informed about the study's objectives and the eventual publication of the information gathered, and they were assured that the informants' identities would remain undisclosed.

Competing interests

The authors declare that they have no competing interests.

Availability of data and materials

Please contact author for data requests.