

Needs Assessment of Community Health Workers to enhance efficient delivery of their services for Community Case Management of Malaria in Kenya

Michelle D. S. Boakye

Pennsylvania State University <https://orcid.org/0000-0002-5641-0645>

Collins J. Owek

Centre for the Study of Adolescence, Nairobi

Elizabeth Oluoch

Mild May International, Kisumu

Sefa Bonsu Atakora

University of Ghana

Juddy Wachira

Moi University, Eldoret

Yaw A. Afrane (✉ yaw_afrane@yahoo.com)

University of Ghana <https://orcid.org/0000-0001-6576-523X>

Research

Keywords: Community Case Management of Malaria, Community Health Workers, Needs assessment

Posted Date: January 21st, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-20695/v2>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published on February 18th, 2021. See the published version at <https://doi.org/10.1186/s12936-021-03640-2>.

Abstract

Background

Malaria continues to be the leading cause of morbidity and mortality in Africa. Community Case Management of malaria (CCMm) through engaging Community Health Workers (CHWs) to effectively address management of malaria cases in some endemic communities was explored in this study. We assessed the needs of CHWs that would help sustain and retain their services to enhance the efficient delivery of CCMm.

Methods

Data on the needs of CHWs was gathered through a qualitative study consisting of in-depth interviews and focus group discussions (FGDs) conducted among study participants in five districts in western Kenya using a semi-structured questionnaire. The study participants comprised of 100 CHWs, 100 mothers of children under five years and 25 key informants made up of public health officers and clinicians involved in the CCMm. The interviews were conducted in English and Swahili or Dholuo, the local language. The recorded audio interviews were transcribed later. The analysis was done using NVivo version 7 software and transcripts were coded after which themes related to the objectives of the study were identified.

Results

All the study participants recognized the need to train and update CHWs on their work as well as remunerating them for their services to enhance efficient delivery of services. The CHWs on their part perceived the provision of gloves, RDTs, lancets, cotton wool and ethanol, bins (to dispose of RDTs and lancets), together with drugs for treating clients as their essential needs to undertake CCMm in the communities. Other logistical needs and incentives mentioned by CHWs and key informants for the successful delivery of CCMm included: gumboots, raincoats, torch lights, mobile phones, means of transportation (bicycles and motorbikes), uniforms and ID cards for identification.

Conclusion

CHWs would perform tasks better and their services retained for a sustainable CCMm if properly incentivized, offered refresher trainings (and updates) on malaria and equipped with the requisite tools identified in this study.

Background

There are a number of ongoing malaria intervention programs in Africa, yet, malaria is reported to be the leading cause of morbidity and mortality in this region with 93% of the 228 million malaria cases reported worldwide occurring in this region in 2018. Out of the 405,000 estimated malaria deaths, 94% occurred in Africa in the same year (1). Of all the age groups, children under the age of five are considered the most

vulnerable to malaria as 67% of malaria mortalities occurred in that group (1). Severe malaria among children under five is a common occurrence in western Kenya (2).

Prompt and accurate diagnosis of malaria together with appropriate treatment is key in malaria case management (3). In Kenya, Community Case Management of malaria (CCMm) is undertaken by community health workers (CHWs) in malaria endemic regions in an effort to combat the burden of the disease (3). CHWs are health volunteers who are selected from their communities and trained to work within their villages (4). In CCMm, CHWs diagnose malaria with rapid diagnostic tests (RDTs) and treat positive malaria cases with artemisinin-based combination therapy (ACT) (5) thus, enhancing access to health care in these communities (6).

Studies have reported the effectiveness of CHWs in reducing child mortalities associated with malaria in several countries (3,7,8). However, previous studies have identified challenges that could undermine the sustainability of CCMm (9–11) These include the lack of basic supplies and equipment to undertake their work, support and non-recognition of their services from clinicians, among others (11). The services of CHWs are crucial to the success of the CCMm and it is important to identify any gaps in relation to their needs in order to create an environment that would commit them to maintain their dedication to the programme. It is against this background that we conducted a qualitative study on the needs of CHWs that would help sustain and retain them to enhance efficient delivery of their services for CCMm.

Methodology

Methods

Study sites

The study was conducted in five districts in western Kenya where CHWs are undertaking CCMm. These include Kisumu West, Nyando and Muhooni sub-counties that are found in the malaria endemic lowland regions and Kisii and Kenyenyua sub-counties found in the epidemic-prone highlands of western Kenya (12).

Study participants

The study participants were made up of mothers/guardians of under-five children who have received services from the CHWs in the previous one year. CHWs were selected from Community Units (CU) in the study sites where CCMm was being undertaken to participate in focus group discussions. Key informants for the study were Community Health Extension Workers (CHEWs) who supervise the work of CHWs, heads of health facilities which the CHWs are affiliated with, Clinicians within these health facilities, Public Health Officers and District and County Medical Officers of Health.

Study design

This study employed a qualitative design using a semi-structured questionnaire for in-depth interviews and focus group discussions (FGDs). CHWs were conveniently sampled from two CUs in the selected study sites where CCMm was being practiced. A pilot study was conducted in a community unit in the Kisumu West district. The questionnaire captured questions on the perception of mothers/guardians of under-five children and other key informants on CCMm as well as the appropriateness and attitude of the CHWs who deliver CCMm, among others. The interview with the CHWs focused on their perception of the work and attitude of others towards them. The study was conducted from May to September 2014.

Sample size

Ten (10) CHWs who had been trained on CCMm from each community unit was selected for the FGDs. Ten (10) mothers/guardians of children under the age of five years who had enjoyed the services of the CHWs in the previous year, were selected for the study from each community units. In all, 100 community members were selected for the study from the ten community units within the five districts.

Key informant interviews were conducted with CHEWs who supervise the CHWs within each community unit. In each community unit, the Clinician-in-charge of the health facility affiliated with the CCMm program were chosen for key informant interviews. This came to a total of ten (10) CHEWs and ten (10) clinicians as key informants. For each of the five districts, the Public health officer (PHO) was also interviewed as a key informant.

Data collection

The interview guide for the study was pre-tested in a community unit in Kisumu West sub-county. The FGDs and key informant interviews were to assess the perception of the different categories of people interviewed. However, the questions for the FGDs and interviews were different depending on who is being interviewed in order to capture different themes and to answer different questions. The questionnaires for the FGDs were first written in English, and then translated into the local language (Dholuo or Swahili) for the FGDs. It was later back-translated into English by another person to ensure accuracy.

FGDs were conducted at health facilities for mothers/caregivers of under-five children who were conveniently sampled from households within the same CU that have been served by a CHW. FGDs for the CHWs were held at a central point accessible from affiliated health facilities. The CHEWs were interviewed individually as key informants at the health facility.

The remaining key informants such as heads of health facilities, clinicians, Public Health Officers and District and County Medical Officers of Health were also interviewed using semi-structured questionnaires. The questionnaire was on assessment of the needs of CHWs to undertake CCMm. All study participants signed a consent form after an explanation of the study purpose.

Data management and analysis

All the interviews were audio-recorded, transcribed, translated and de-identified. The transcripts were read as individual wholes to achieve an overarching understanding of the data. The data was coded and finally organized into themes based on the answers of the participants using Nvivo version 7 software. The assessment of the needs of the CHWs was explored based on a common induction approach to coding based on phenomenology rather than pre-existing theory (13). The results of the coding were compared to ensure consistency of text segmentation and the application of codes. Periodic checks were carried out to ensure intercoder agreement.

Results

Demographic characteristics of respondents

The mothers that were interviewed in the FGDs were between 19 to 36 years with a median age of 24 years. The ages of CHWs that were interviewed ranged from 25 to 59 years of age with a median age of 32. CHEWs were aged between 32 and 47 years with a median age of 37, while the ages of the clinicians and district/county director of health ranged from 28 to 52 years. Whilst most (56%) of the CHWs were females, none of the CHEWs was female. Among the clinicians and PHOs that were interviewed, 40% (8/20) of them were female. In terms of the educational level, majority of the CHWs (64%) had finished secondary school, the rest only having finished primary education. With the mothers, only 22% had finished secondary school, with 54% completing primary education while 24% had no education. None of the participants declined to be part of the study.

Perception of CHWs on their needs

Working materials

In assessing needs that facilitate effective services for malaria diagnoses and treatment in the community, the CHWs alluded to provision of the basic essential supplies being the most important. These include drugs for treating clients who are sick of malaria and more pain killers, gloves, Rapid diagnostic test kits (RDTs), cotton wool and ethanol, bins to dispose of RDTs and lancets for pricking their clients. They assert that sometimes their RDTs could run out for weeks and when they go to the health facility, they are told there is none at the facility itself to be given to them due to stock-outs. Whilst some CHWs from particular CUs assert that they have never been given drugs since their training, others indicated they were given drugs immediately after their training but these were not replenished when they ran out. They therefore have no option but to keep referring all clients whom they test and are positive to the health facility. None of the CHWs reported having been given items like ethanol to clean the finger before pricking the clients or dustbins to dispose of their rubbish when they finish.

On logistical support to enhance their work, the CHWs mentioned gumboots to be used when it is rainy and muddy. All of the CHWs assert that sometimes accessing a client's house can be difficult when it rains since it becomes muddy and difficult to walk through the mud without a gumboot. Other logistical supplies mentioned included bicycles or motorbikes for movement to distant areas to visit clients and

uniforms and ID cards for identification. In reality, each CHW is assigned one hundred (100) households to work with. This consist of approximately 500 people. These households are could be located in 1-3 villages. Also, the households are far apart and not clustered together. They all also think that they need a mobile phone for their clients to reach them and vice versa. They reported that sometimes clients would like to talk to them on the phone and if they do not have a mobile phone then it is difficult for clients to reach them at night. They assert that clients sometimes have to send someone to walk to their house to call them if there are issues.

"We need the drugs and the RDTs. We also need raincoats, gumboot and uniforms, at the same time we can say that we need bicycles to help us in moving around." CHW from FGD in Chemelil, Kisumu County.

"We asked for even gumboots since we walk in muddy areas, but they did not bring." CHW from FGD in Ojolla B, Kisumu County

Trainings

The CHWs perceived other needs to be training/updates on malaria to enable them provide better support to their clients. They referred to a three-day training on malaria diagnosis using RDTs and how to dispense anti-malarial drugs to clients as being the only training intervention received. The CHWs say that a local NGO facilitated this training in collaboration with the Ministry of Health. This was three years back at the time of the interview. The training was followed up with a week's attachment at the laboratory of a health facility. The criterion for qualification was the person being able to diagnose a case in the lab correctly. They assert that since that time they have not had any refresher courses/training on malaria diagnosis, treatment and prevention practices. The CHWs think that more NGOs should come in to help in their training and that the government can also run some refresher trainings for them.

Before their training on malaria, they had previously been trained on Community Health Strategy (CHS) by another NGO when they were recruited. This was a week-long training. The CHS concept has been described above in this report.

"...we need refresher training on malaria and maybe something on nutrition. Because if we go out there, we find that the children are malnourished and they are also sick of malaria so when you have knowledge about nutrition you will advise them on how to go about this." CHW from FGD in Kenya, Kisii

Remunerations

The CHWs also perceived remuneration as one of their key needs. They reported that they receive a monthly stipend of Kenya shillings (Ksh) 2000 (USD 22, at the time of interview) which comes from an NGO who partner with the Ministry of Health. Per their assertion, this is not only for the services they provide on malaria but for the general services they offer their clients. They are of the opinion that this amount is not commensurate with their efforts and time spent going around the village to care for sick people. Coupled with this inadequacy is an inconsistency in the disbursement of this fund. There were complaints that sometimes this could be delayed for several months. The stipend is needed to enable

them to meet basic needs and as compensation for the time used to support their clients. Compensation for their time affords them the ability to engage others to assist them in their farms while they are out to offer services to clients. They think that since they are part of the health care system, they should be recognized as such and be paid. Pressed further on how much they think they are worth, the CHWs said the minimum amount they should be paid should be Kenyan shillings 10,000 (USD 113) per month. They also think that they could be mobilized and given some piece of land to farm on as a cooperative so that they can get income from this venture.

"...we are sometimes called at night, and you have to leave everything and go. You stop working on your farm so that you can see clients every morning, we also need to be given something reasonable." CHW from FGD in Kisii.

"..... we need to be given the materials that can enable us to work, another thing is that money they give us, it should be something that can sustain us and should be on time too. The minimum should be about Ksh. 10, 000"

Recognition from staff of health facilities

The CHWs complained that they do not receive the necessary recognition from some healthcare workers which tend to undermine their efforts and the CCMm program. They indicated that sometimes when they take a client already tested and found to be positive for malaria by their system to a health facility, the clinician will still insist on the client being tested again from the laboratory of the health facility. As a result, some clients lose faith and interest in their services. Again, some of them assert that some technicians at some of the health facilities specifically tell clients not to trust the results of CHWs because they do not have the requisite training.

Perception of key informants (CHEWs, Clinician Health facility in-charges and PHOs) on CHWs' needs

Training

The health workers perceived the needs of the CHWs to be additional/ refresher trainings on the work that they are engaged in. The Clinicians assert that the three-day malaria training given to CHWs is not adequate to empower them to perform at an excellent level and that, each year, they should have the chance to update themselves with new regimen for testing, general diagnosis, and treatment of malaria cases, as is done for clinicians and technicians in the health facilities. One Clinician asserted that sometimes when the malaria parasitaemia is very low, the RDT might not pick up the parasites to show positivity in the reading but the person might be able to be diagnosed with malaria from the signs and symptoms but the person might be able to be diagnosed with malaria from the signs and symptoms. She asserts that a CHW might not know this if they are not given refresher courses and might just give antibiotics when the person actually has clinical malaria. The key informants think that since the CHWs are part of the health care system, the government should take up their training and not leave it alone to

NGOs. This, they say, is more sustainable than if it is left with NGOs whose funding might run out at some point.

".....they need more training and the government must step in to offer this or they should be told to stop and tell the people to come to the hospital. But in my opinion, they should be trained since their work is very important in the communities." Health facility in Charge in a KII at Chemelil.

Incentives

The key informants unanimously alluded to the need to have the CHWs adequately remunerated to retain their services in order to sustain the CCMm. One clinician particularly suggested that the government should go beyond branding CHWs as volunteers (and not remunerating them accordingly), and consider putting them on its payroll to assure CHWs of regular income, no matter how small the funds may be. The CHEWs also think the best way to remunerate the CHW is to support them to establish income generating activities (IGAs) such as small-scale farming where the proceeds will go to the CHWs. They stressed that this IGA idea is being tested in a few community units to see its viability. Also, the CHEWs think that the government could incentivize the CHWs by offering their families and themselves free medical care.

"In my opinion, according to the work load they do, the two thousand they are given is small, they should be given four thousand". CHEW in a KII

The best way to motivate the CHWs is to bring them together and initiate an income generating activity for them." CHEW in a KII

"You know they are volunteers, they are working without pay and at least they work. The time has reached for them to be motivated better, because as you know work without motivation is not effective." PHO in a KII

Working materials

The key informants also recognized the need for CHWs to be given some working materials such as means of transport to reach their clients, torchlight to be used when it is dark and raincoat or umbrella when it is raining. The clinicians think that CHWs should be given bicycles to facilitate them in moving around. CHEWs reported that sometimes clients of CHWs call them at night and since it is dark and most villages do not have electricity (and therefore street lights), CHWs need to have a torchlight to aid them to reach the client's place. They also have to walk through the rain sometimes to reach the health facility with a client or to see a client, and a raincoat or an umbrella will do them good. Some of the CHEWs also think that the CHWs could be given a simple mobile phone that their clients could reach out to them on when they need their services, especially at night. A CHEW thought that since these people are walking from house to house, they need to be given an identity card or wear particular uniform or T-shirt to identify them otherwise people will take them to be thieves walking around in the community.

Perception of caregivers of children under 5 years of age and pregnant women

The community members that included pregnant women advocated that every effort should be made to empower the CHWs to carry out their work. This would include training and/or other support given to clinicians to do their work.

"They are doing a good work and for me personally they are helping me and so when someone can be taken for training or given some money or items to help them in their work so that they can even take care of the children and ourselves, just as nurses do then that should be done to help them." Expectant mother from an FGD in Kenyanya, Kisii district.

Discussion

CHWs represent an important health resource with great potential for providing and extending basic healthcare to underserved populations (4). In the present study, the needs of CHWs undertaking CCMm in malaria endemic districts in Kenya were assessed in a qualitative study. This is particularly important as their needs will likely affect their performance and job satisfaction. Their needs from the perception of other stakeholders such as healthcare workers and mothers with children under five would provide a holistic view to help shape and promote efficient delivery of CCMm.

In this study it was found that the CHWs perceived their working materials as their primary needs for them to undertake services for malaria diagnoses and treatment in the community. Without RDTs and ACTs, the CHWs will not be able to work, yet most of the workers asserted they usually run out of RDTs and drugs for weeks. This seems a common phenomenon in Sub-Saharan African countries (14–16). All the CHWs interviewed reported that they had never received ethanol rub or even dustbins to dispose of rubbish. It is imperative this need is addressed; the ethanol will make their work safe and could prevent cross-infection. The dust bins on the other hand would ensure a more efficient disposal and prevent needle pricks from used RDTs.

Another area of concern to the CHWs was the lack of recognition, respect and negative attitude of some staff of health facilities. The fact that some of the healthcare workers do not appreciate their services in the communities can affect the zeal and willingness to operate as voluntary health workers in the communities. The CCMm system should involve the clinicians so that the CHWs are well incorporated into the system for effective management of malaria in communities. The respect and support from qualified health workers and the health system is essential to the success and retention of health volunteers (4,16–18).

Again, other logistical needs mentioned by CHWs and CHEWS, key informants as well as clinicians included gumboots, bicycles or motorbikes, mobile phones, torchlight, raincoats uniforms and ID cards for identification. These non-monetary incentives mentioned by the participants were also suggested in similar studies as ways of sustaining CCMm (6,19) The gumboots, raincoats, bicycles or motorbikes will enable them to reach community members even when it is raining. Also, the mobile phones will improve

effective communication between the community members and the CHWs. The uniforms and ID cards will enable the community members to readily recognize them as health workers in the communities (20) and not mistake them as intruders. This will give them some recognition and prestige in the communities and subsequently serve as a source of motivation.

The CHWs, pregnant women and key informants expressed the need of training and updating the CHWs on malaria as changes (to treatment etc) might have cropped up since the inception of the CCMm or supposed training of the CHWs. The key informants on the other hand thought the trainings should not only be left to the NGOs alone, but that the government should also take it up since the CHWs are also part of the healthcare system in Kenya.

The CHWs are part of the health system but are not on government pay-roll like the other formally-trained personnel. They therefore receive an allowance but not a salary with benefits such as pension and health insurance. Although the CHWs are considered volunteers, the CHWs revealed that the Ksh. 2000 (USD 22) they received from an NGO was grossly inadequate for the services they provide the community members including CCMm. The result of this study confirms previous findings of variations in incentives in low- and middle-income countries. These incentives varied from monetary incentives such as reasonable payment to nonmonetary incentives like community recognition, and acquisition of value skills as reported by Ormel et al (21). A study in Tanzania also found that CHWs are motivated to volunteer to work and apply knowledge gained to their own problems and those of their families and communities (22). The CHWs mentioned a minimum amount of Ksh. 10,000 (USD 113) as a financial incentive that will compensate them for their time and also help them engage others to assist them on their farm as they attend to community members. To sustain CCMm, remuneration will be needed to compensate CHWs as volunteerism cannot be relied on entirely to sustain the program (21).

In order for the volunteers to remain focused on their duties in promptly managing malaria in communities, efforts should be made to support them on their farms as earlier studies have revealed volunteers who do not get support from community members pertaining to their farming activities are more likely to drop out of their volunteer work and concentrate on the farming (23) as that is their source of livelihood.

Conclusion

CHWs undertaking CCMm have a valuable role to play in controlling malaria in developing countries such as Kenya. However, their needs such as logistical supplies, means of transportation, incentives, recognition from health care professionals must be met. These needs must be addressed by the government through the ministry of health, national malaria control programs, civil society organizations that are working on malaria prevention and control to motivate and retain the CHWs in efficient delivery of their services in communities. This study adds to the evidence that the CHWs together with other stakeholders perceive monetary as well as non-monetary incentives as crucial to their services. The gaps

in requisite needs identified including refresher training programs must be taken into consideration for the efficient delivery of CCMm in communities.

List Of Abbreviations

ACT	Artemisinin-based combination therapy
CCMm	Community Case Management of malaria
CHEWs	Community Health Extension Workers
CHS	Community Health Strategy
CHWs	Community Health Workers
CU	Community Units
FGDs	Focus group discussions
IGAs	Income generating activities
KEMRI	Kenya Medical Research Institute
KII	Key informant interview
Ksh	Kenyan shillings
NGO	Non-governmental organization
PHO	Public health officer
PHO	Public Health Officer
RDTs	Rapid diagnostic tests

Declarations

Ethics approval and consent to participate

Ethical approval for the study was obtained from the Scientific Steering Committee and the Ethical Review Committee of the Kenya Medical Research Institute (KEMRI) before starting the study. Approval was also sought from the county commissioner of Kisumu and Kisii Counties of western Kenya and all district commissioners in the selected districts. Written informed consent was obtained from all the study participants after explaining the objectives of the study in the local language to them before they were interviewed.

Consent for publication

Not applicable.

Availability of data and materials

Data for the manuscript are available in the Kenya Medical Research data repository to anyone who might need it.

Competing interests

The authors declare that they have no competing interests.

All authors declare that we all had full access to all of the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis.

Funding

This research is supported by grants from the World Health Organization: Joint Afro-TDR (AFHSS1207000—SMALL GRANTS PROGRAMME).

Authors' contributions

MDSB, CJO and EO participated in data collection, management, and analysis, drafting and producing the final manuscript; SBA, JW and YAA designed the study, supervised data collection, and helped in drafting and producing the final manuscript. All authors read and approved the final manuscript.

Acknowledgements

The authors wish to thank all field assistants who helped in data collection for the study.

References

1. World Health Organization. World Malaria Report 2019. Geneva. [Internet]. 2019. Available from: <https://apps.who.int/iris/handle/10665/330011>
2. Akech S, Chepkirui M, Ogero M, Agweyu A, Irimu G, English M, et al. The Clinical Profile of Severe Pediatric Malaria in an Area Targeted for Routine RTS,S/AS01 Malaria Vaccination in Western Kenya. *Clin Infect Dis* [Internet]. 2020 Jul 11 [cited 2021 Jan 12];71(2):372–80. Available from: <https://academic.oup.com/cid/article/71/2/372/5554744>
3. Siekmans K, Sohani S, Kisia J, Kiilu K, Wamalwa E, Nelima F, et al. Community case management of malaria: A pro-poor intervention in rural Kenya. *Int Health*. 2013 Sep;5(3):196–204.
4. Lehmann U, Sanders D. Community Health Workers: What Do We Know About Them? The State of the Evidence on Programmes, Activities, Costs and Impact on Health Outcomes of Using Community

- Health Workers. [Internet]. Geneva, Switzerland: World Health Organization; 2007. Available from: www.usaid.gov
5. World Health Organization. Global report on diabetes [Internet]. Geneva PP - Geneva: World Health Organization; 2016. Available from: <https://apps.who.int/iris/handle/10665/204871>
 6. Glenton C, Colvin CJ, Carlsen B, Swartz A, Lewin S, Noyes J, et al. Barriers and facilitators to the implementation of lay health worker programmes to improve access to maternal and child health: Qualitative evidence synthesis. Vol. 2013, Cochrane Database of Systematic Reviews. John Wiley and Sons Ltd; 2013.
 7. Johnson AD, Thomson DR, Atwood S, Alley I, Beckerman JL, Koné I, et al. Assessing Early Access to Care and Child Survival during a Health System Strengthening Intervention in Mali: A Repeated Cross Sectional Survey. Eisele T, editor. PLoS One [Internet]. 2013 Dec 11 [cited 2021 Jan 12];8(12):e81304. Available from: <https://dx.plos.org/10.1371/journal.pone.0081304>
 8. Kisia J, Nelima F, Otieno DO, Kiilu K, Emmanuel W, Sohani S, et al. Factors associated with utilization of community health workers in improving access to malaria treatment among children in Kenya. *Malar J*. 2012 Jul 30;11:248–248.
 9. Elmardi KA, Malik EM, Abdelgadir T, Ali SH, Elsyed AH, Mudather MA, et al. Feasibility and acceptability of home-based management of malaria strategy adapted to Sudan's conditions using artemisinin-based combination therapy and rapid diagnostic test. *Malar J* [Internet]. 2009 Dec 9 [cited 2021 Jan 12];8(1):39. Available from: <https://malariajournal.biomedcentral.com/articles/10.1186/1475-2875-8-39>
 10. Akweongo P, Agyei-Baffour P, Sudhakar M, Simwaka BN, Konaté AT, Adongo PB, et al. Feasibility and acceptability of ACT for the community case management of malaria in urban settings in five African sites. *Malar J*. 2011;10.
 11. Boakye MDS, Owek CJ, Oluoch E, Wachira J, Afrane YA. Challenges of achieving sustainable community health services for community case management of malaria 11 Medical and Health Sciences 1117 Public Health and Health Services. *BMC Public Health* [Internet]. 2018 Oct 1 [cited 2021 Jan 12];18(1):1150. Available from: <https://bmcpublichealth.biomedcentral.com/articles/10.1186/s12889-018-6040-2>
 12. Zhou G, Afrane YA, Vardo-Zalik AM, Atieli H, Zhong D, Wamae P, et al. Changing patterns of malaria epidemiology between 2002 and 2010 in western kenya: The fall and rise of malaria. *PLoS One*. 2011;6(5).
 13. Corbin J, Strauss A. Basics of qualitative research: Techniques and procedures for developing grounded theory. : 3rd ed. Thousand Oaks, CA: SAGE Publications, Inc.; 2008.
 14. Counihan H, Harvey SA, Sekeseke-Chinyama M, Hamainza B, Banda R, Malambo T, et al. Community health workers use malaria rapid diagnostic tests (RDTs) safely and accurately: Results of a longitudinal study in Zambia. *Am J Trop Med Hyg*. 2012 Jul;87(1):57–63.
 15. Kangwana BB, Njogu J, Wasunna B, Kedenge S V., Memusi DN, Goodman CA, et al. Malaria Drug Shortages in Kenya: A Major Failure to Provide Access to Effective Treatment. *Am J Trop Med Hyg*.

2009;80(5):737–8.

16. Druetz T, Ridde V, Kouanda S, Ly A, Diabaté S, Haddad S. Utilization of community health workers for malaria treatment: Results from a three-year panel study in the districts of Kaya and Zorgho, Burkina Faso. *Malar J*. 2015 Feb 13;14(1).
17. Chatio S, Akweongo P. Retention and sustainability of community-based health volunteers' activities: A qualitative study in rural Northern Ghana. Mistry N, editor. *PLoS One* [Internet]. 2017 Mar 15 [cited 2021 Jan 13];12(3):e0174002. Available from: <https://dx.plos.org/10.1371/journal.pone.0174002>
18. Rahman SM, Ali NA, Jennings L, Seraji MHR, Mannan I, Shah R, et al. Factors affecting recruitment and retention of community health workers in a newborn care intervention in Bangladesh. *Hum Resour Health*. 2010 May 3;8.
19. Strachan DL, Källander K, Ten Asbroek AHA, Kirkwood B, Meek SR, Benton L, et al. Interventions to improve motivation and retention of community health workers delivering integrated community case management (iCCM): Stakeholder perceptions and priorities. *Am J Trop Med Hyg*. 2012;87(SUPPL.5):111–9.
20. Ozano K, Simkhada P, Thann K, Khatri R. Improving local health through community health workers in Cambodia: challenges and solutions. *Hum Resour Health* [Internet]. 2018 Jan 6 [cited 2021 Jan 13];16(1):2. Available from: <https://human-resources-health.biomedcentral.com/articles/10.1186/s12960-017-0262-8>
21. Ormel H, Kok M, Kane S, Ahmed R, Chikaphupha K, Rashid SF, et al. Salaried and voluntary community health workers: Exploring how incentives and expectation gaps influence motivation. *Hum Resour Health* [Internet]. 2019 Jul 19 [cited 2021 Jan 13];17(1):59. Available from: <https://human-resources-health.biomedcentral.com/articles/10.1186/s12960-019-0387-z>
22. Greenspan JA, McMahon SA, Chebet JJ, Mpunga M, Urassa DP, Winch PJ. Sources of community health worker motivation: A qualitative study in Morogoro Region, Tanzania. *Hum Resour Health*. 2013 Oct 10;11(1).
23. Bhattacharyya K, Winch P, LeBan K, Tien M. *Community Health Worker Incentives and Disincentives: How They Affect Motivation, Retention, and Sustainability*. Arlington, Virginia: the Basic Support for Institutionalizing Child Survival Project (BASICS II) for the United States Agency for International Development; 2001.