

# Ruscogenin alleviates intestinal bleeding and blood flow induced by Dasatinib through ROCK/MLC pathway

**Yuankai Liu**

China Pharmaceutical University

**Yujie Dai**

China Pharmaceutical University

**Han Xu**

China Pharmaceutical University

**Qianliu Zhou**

China Pharmaceutical University

**Fang Li**

China Pharmaceutical University

**Boyang Yu**

China Pharmaceutical University

**Yuanyuan Zhang**

China Pharmaceutical University

**junping kou** (✉ [junpingkou@cpu.edu.cn](mailto:junpingkou@cpu.edu.cn))

China Pharmaceutical University <https://orcid.org/0000-0001-8377-915X>

---

## Research Article

**Keywords:** Dasatinib, Intestinal bleeding, Endothelial barrier disruption, ROCK-MLC signaling pathway, Ruscogenin

**Posted Date:** August 12th, 2021

**DOI:** <https://doi.org/10.21203/rs.3.rs-207273/v1>

**License:** © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

# Abstract

As a second-generation broad-spectrum tyrosine kinase inhibitor, Dasatinib has the antitumor effect of inhibiting tyrosine kinases such as BCR-ABL and SRC. It is mainly used to treat chronic and acute patients with chronic myelogenous leukemia. However, Dasatinib has many side effects, including gastrointestinal bleeding, respiratory infections, and renal failure, and effective means to improve the side effects of drugs are lacking. The occurrence of gastrointestinal bleeding is mainly due to the destruction of the vascular endothelial barrier by Dasatinib. For the first time, we administered a large dose of oral administration and monitored the intestinal bleeding with a laser Doppler flowmeter. We found that Ruscogenin (RUS) can improve dasatin Side effects caused by tinib. Through Doppler flowmeter detection, it was found that after Dasatinib was administered, blood flow in mice decreased, and intestinal Evans blue leakage increased. Western blot results showed that connexin (ZO-1, VE-cadherin, Occludin) is destroyed, the activation of ROCK increases, and the phosphorylation of MLC increases. Ruscogenin can reverse the above phenomenon and improve the side effects of gastrointestinal bleeding caused by Dasatinib. *In vitro*, giving Ruscogenin in advance can protect the endothelial barrier damage caused by Dasatinib, reduce the remodeling of F-actin, and restore the expression of connexin (ZO-1, VE-cadherin), inhibit the activation of ROCK, and reduce Phosphorylation of MLC. This study provides a new direction for improving the side effects caused by clinical drugs.

## Highlights

- Intestinal bleeding and barrier destruction caused by gavage of Dasatinib in mice
- Ruscogenin protects the endothelial barrier function and reduces vascular leakage.
- Ruscogenin reverses the side effects of Dasatinib by inhibiting the ROCK/MLC signaling pathway

## Introduction

In clinical treatment, drug-induced side effects not only aggravate the patient's condition, but also bring a great financial burden to the patient<sup>[1]</sup>. A survey data shows that 71.3% of patients who require hospitalization due to acute bleeding caused by drugs<sup>[2]</sup>. Therefore, prevention and treatment of side effects caused by drugs have a positive effect on the treatment of patients.

As a second-generation broad-spectrum tyrosine kinase inhibitor, Dasatinib exerts an anti-tumor effect by inhibiting tyrosine kinases such as BCR-ABL and SRC<sup>[3-4]</sup>. Clinically use mainly in the treatment of chronic myeloid leukemia and acute lymphoblastic leukemia<sup>[5]</sup>, as well as leukemia patients resistant to Imatinib, Dasatinib inhibits wild-type BCR-ABL much stronger than the first-line treatment drug Imatinib At 325 times<sup>[6-7]</sup>. Dasatinib's global sales in 2016 were \$1.8 billion. Although Dasatinib is very effective, it often shows serious side effects. The most common side effects in clinical reports include gastrointestinal symptoms<sup>[8]</sup>, hemorrhagic events, hepatotoxicity<sup>[9]</sup>, pleural effusion<sup>[10-12]</sup>, pulmonary hypertension<sup>[13-14]</sup>,

and respiratory infections with neutral granules. Adverse events such as cell reduction and renal failure<sup>[15]</sup>.

Clinical investigations shown that 23% of patients taking Dasatinib have developed bleeding, and gastrointestinal bleeding is a relatively high incidence of adverse reactions. Among patients with gastrointestinal bleeding, most patients received high doses of Dasatinib (> 100 mg) or received twice daily doses<sup>[16-18]</sup>. It can be judged that oral Dasatinib can cause gastrointestinal bleeding. The possible mechanism of Dasatinib's destruction of the endothelial barrier is due to the activation of ROCK / MLC signals, which causes actin polymerization and the breakdown of connexins in vascular endothelial cells. Intervention in the ROCK / MLC signaling pathway has been recognized for protecting the blood-brain barrier<sup>[19-20]</sup>. Therefore, intervention of the endothelial ROCK-MLC signaling pathway and improvement of barrier disorders is to alleviate Dasatinib-induced intestinal bleeding.

At present, there are not many effective monitoring methods to examine the patient's bleeding. Some Articles have shown that an increase in organ bleeding can cause a decrease in blood flow<sup>[21-22]</sup>. Most of the models used to study the side effects caused by Dasatinib are intraperitoneal injection or oral low-dose long-term administration<sup>[15,23-24]</sup>. Animal models of large-dose oral administration are lacking. In clinical use, large-dose oral administration is also common. Therefore, large-dose oral administration was selected.

Ruscogenin is one of the effective active ingredients of *Ophiopogon japonicus*, which has effect on anti-inflammatory, anti-thrombosis and reduces capillary permeability<sup>[25-26]</sup>. Ruscogenin can alleviate endothelial cell injury, protect blood-brain barrier function, and relieve pulmonary hypertension caused by monocrotaline<sup>[27-28]</sup>. In the early stage of the laboratory, It was proved that Ruscogenin can up-regulate the expression of adhesion proteins VE-cadherin and p120-catenin, inhibit the nuclear translocation of VE-cadherin, thereby improving the adhesion of endothelial cell adhesion junction and reducing the damage of the pulmonary vascular endothelium induce by Lipopolysaccharide<sup>[29]</sup>.

Based on the above data, we hypothesized that RUS can be used as a clinical intervention to improve the side effects of Dasatinib causing endothelial dysfunction.

## Materials And Methods

### 2.1 Materials

Ruscogenin, 98%, from J&K® (Beijing China). Flux of Evans blue dye-labeled albumin (EB-albumin, Sigma, USA).

### 2.2 Animals

All animal care and experimental procedures were carried out according to the current European Communities Council-ECC guidelines for the care of laboratory animals and ethical guidelines for

investigations of experimental pain in conscious animals. Animal studies were reported in compliance with the ARRIVE guidelines. Male C57BL/6J mice (20–25 g; 8 weeks; specified pathogen free) were provided by the Model Animal Research Centre of Yangzhou University (Yangzhou, Jiangsu, China) and kept in cages containing standard bedding, with at least five mice per cage. Mice were housed in a specific pathogen-free facility with 12 h light/dark cycle [07 to 19 h, temperature ( $22 \pm 2^\circ\text{C}$ ), humidity (40–70%), controlled ventilation] and with sterile water and irradiated food available ad libitum. Animals were allowed to acclimatize to their housing environment for at least 7 days prior to experimentation and to the experimental room for 1 h before experiments.

### 2.3 Drug treatments

Ruscogenin was weighed and dissolved in absolute ethanol to prepare a mother liquor. When used, the corresponding concentration was adjusted with physiological saline. Dasatinib was dissolved in physiological saline at a concentration of 140 mg/kg and administered intragastrically. 30 minutes after the administration of Ruscogenin, Dasatinib was administered by gavage. After 30 minutes, the mice were killed.

### 2.4 Blood flow monitoring

Before administering Dasatinib, mice were given Ruscogenin in 30 min, and 2 hours later anesthetized by Pellto-barbitalum Natricum. The small intestine was removed and placed on a Doppler flowmeter for scanning to obtain images.

### 2.5 Western blot analysis

Western blotting analysis was performed as previously described. The cells were lysed and centrifuged. Equal amounts of proteins (40  $\mu\text{g}$ ) were examined using 10% SDS-PAGE and subsequently transferred to PVDF membranes (Millipore, USA) using electrophoresis. The membrane was blocked with 5% BSA for 1.5 h and incubated overnight at  $4^\circ\text{C}$  with primary antibodies against ZO-1 (1:200, Abcam, USA), VE-cadherin (1:1000, Santa Cruz, USA), Occludin (1:200, Abcam, USA), ROCK1 (1:500, Santa Cruz, USA), MLC (1:1000, CST, USA), phospho-MLC (1:1000, CST, USA),  $\beta$ -actin (1:10000, Bioworld, USA), followed by incubation with peroxidase-conjugated secondary antibodies (1:8000, Bioworld, Louis Park, USA) and visualization using enhanced chemiluminescence (ECL, Vazyme Biotech, Nanjing, China). The results were quantified using the Image Lab™ software (version 4.1, Bio-Rad, California, USA).

### 2.6 Histology and immunohistochemistry

The excised vessel was dehydrated with 40% sucrose, embedded in optimal cutting temperature (OCT) and frozen at  $-80^\circ\text{C}$ . The jejunum was sectioned into slices of  $4\mu\text{m}$  of thickness with a cryotome (Leica, Mannheim, Germany). Specimens were washed in PBS and stained with Haematoxylin and Eosin (H&E). For immunohistochemical staining, slides were incubated with primary antibodies at  $4^\circ\text{C}$  overnight. Alexa Fluor 488-/594-labelled antibodies were used as secondary antibodies. The nuclei were stained with

DAPI(Beyotime Biotechnology, Shanghai, China). Pathological changes in jejunum were observed under a fluorescence microscope (Leica, Mannheim, Germany).

## 2.7 Cell culture

HUVECs were purchased from the ZhongQiaoXinZhou Co. Ltd. (Shanghai, China) and cultured in RPMI 1640 (Invitrogen, Carlsbad, CA, USA) supplemented with 10% fetal bovine serum (FBS, Sigma, St. Louis, MO, USA), 100 U/ml penicillin and 100U/ml streptomycin (Amersco, OH, USA) at 37 °C in a humidified atmosphere of 5% CO<sub>2</sub> and 95% air.

## 2.8 Dasatinib-induced disruption of HUVECs barrier

In the control group, serum-free RPMI-1640 medium was added to the chamber for 2 h. Dasatinib (final concentration 150 nM) was added to the model group for 2 h. The administration group was added with 0.01, 0.1, 1 μM RUS for 1 h, and then 150 nM Dasatinib was added for 2 h.

## 2.9 Statistical analysis.

All results are expressed as the means±SD. Statistical analysis was performed using Student's two-tailed t test for comparison between two groups and one-way analysis of variance (ANOVA) followed by Dunnett's test when the data involved three or more groups.  $P < 0.05$  was considered statistically significant. All analyses were performed using GraphPad Prism Version 5.01 (GraphPad Software Inc.USA).

# Results

## 3.1 Oral Dasatinib reduced intestinal blood flow in mice

Clinically, Dasatinib is used for oral treatment of chronic myeloid leukemia. In this section, the time points of blood flow reduction in the small intestine caused by Dasatinib were observed by different doses of Dasatinib (70, 140, 210 mg/kg). As shown in Fig. 1, administration of 70 mg/kg Dasatinib did not caused evident decrease blood flow in intestinal. Compared with the control group, the blood flow in small intestine was significantly decreased ( $P < 0.01$ ) when the gavage of 140 mg/kg Dasatinib was administered for 90 minutes, and the blood flow of the small intestine continued to decrease at 135 min ( $P < 0.01$ ). When the mice were given by gavage 210 mg/kg Dasatinib for 45 min, the blood flow in the small intestine was significantly decreased ( $P < 0.01$ ), and the blood flow in the small intestine was continuously decreased at 90 and 135 min ( $P < 0.01$ ).

## 3.2 RUS reverted the Intestinal barrier leakage in C57BL/6 subjected to Dasatinib

We found that oral Dasatinib caused a decrease in intestinal blood flow. Therefore, we used the Evans blue method to detect the effects of Dasatinib-induced small intestinal vascular leakage in mice. The results are shown in Fig. 2A. Compared with the control group, the leakage rate of Evans blue in the small

intestine of the model group was significantly increased ( $P < 0.001$ ), and the inhibitory effect of RUS at the high dose was the best ( $P < 0.001$ ). This indicates that RUS has an obvious effect on intestinal vascular leakage caused by Dasatinib. As shown in Fig. 2C, 2D, RUS at 30  $\mu\text{g}/\text{kg}$  reversed the blood flow reduction caused by Dasatinib at 90 min ( $P < 0.05$ ). HE staining showed that RUS alleviated Dasatinib-induced intestinal barrier destruction and hemorrhage.

### **3.3 RUS ameliorated the expression of junction and adhesion proteins in C57BL/6 subjected to Dasatinib**

We investigated the effect of RUS on connexin expression by Western blot. The results of protein imprinting showed that compared with the control group, the expression of junction proteins ZO-1, Occludin and VEcadherin in the model group were significantly decreased ( $P < 0.001$  &  $P < 0.05$ ), and RUS at high dose significantly inhibited Dasatinib-induced ZO-1 and Occludin. VEcadherin expression decreased ( $P < 0.001$  &  $P < 0.01$ ). Immunofluorescence results showed that RUS increased the expression of ZO-1.

### **3.4 RUS ameliorated the side effects of Dasatinib by inhibiting the ROCK1 / MLC signaling pathway**

Hemorrhagic complications of Dasatinib mainly phosphorylate MLC through the ROCK pathway, resulting in cytoskeletal remodeling; therefore, we investigated whether RUS can reduce the phosphorylation of MLC by inhibiting the activation of ROCK-1, thereby protecting the intestinal barrier. Western blot results showed that high dose RUS significantly inhibited Dasatinib-induced ROCK-1 activation and phosphorylation of MLC ( $P < 0.001$  &  $P < 0.05$ ).

### **3.5 RUS reverted the barrier leakage in HUVECs subjected to Dasatinib.**

As the Fig. 5 shows that the HUVECs were treated with RUS (0.01-1  $\mu\text{M}$ ), subsequently exposed to 2 h of Dasatinib (150 nM), and the barrier protective effect of RUS was evaluated using EB albumin assays in HUVECs. The results demonstrated that treatment with RUS can ameliorate endothelial barrier function after 2 h of Dasatinib ( $P < 0.01$ ). We also observed the effect of RUS on the changes in F-actin using an immunofluorescence assay. The results demonstrated that RUS stabilized the cytoskeleton and reduced the incidence of cell contraction leading to cell shrinkage in HUVECs subjected to Dasatinib.

### **3.6 RUS improved the expression of junction and adhesion proteins in HUVECs cells subjected to Dasatinib**

First, we investigated the effects of RUS on the expression of junction and adhesion proteins after 2 h of Dasatinib. The results of the western blot analysis indicated that treatment with RUS (0.1  $\mu\text{M}$  & 1  $\mu\text{M}$ ) up-regulated the decreased expression of ZO-1 and VEcadherin after 2 h of Dasatinib in HUVECs. These findings suggested that RUS improved the Dasatinib-induced abnormal expression of junction and adhesion proteins in HUVECs ( $P < 0.001$  &  $P < 0.01$ ).

Considering the role of the ROCK/MLC pathway in microvascular endothelial barrier function, we also used western blotting to evaluate the effect of RUS on the expression of proteins associated with the

ROCK/MLC pathway. The results showed that treatment with RUS (0.1 $\mu$ M and 1 $\mu$ M) inhibited ROCK1 inhibited, and reduced MLC phosphorylation ( $P < 0.01$  &  $P < 0.05$ ). These results indicated that the ROCK/MLC pathway was also involved in the RUS-induced improvement in microvascular endothelial barrier function.

## Discussion

The side effects of clinical drugs have always been a major problem for patients' rehabilitation. Drugs that cause serious adverse reactions, even included some commonly used clinical drugs, such as oral antiplatelet drugs, warfarin, oral hypoglycemic drugs, etc[2]. Common adverse reactions include bleeding events, hypoglycemia, arrhythmia, and so on, lack effective preventive intervention and monitor methods[30]. Therefore, it is of great significance to find and develop bioactive molecules that can prevent adverse drug reactions.

In order to find a way to ameliorate the hemorrhagic side effects induced by Dasatinib in clinical practice, we selected Ruscogenin, an active ingredient derived from the natural drug, *Ophiopogon japonicus*. Experiments have shown that RUS can significantly inhibit Dasatinib-induced blood flow decreased at intestinal, through by Evans blue shown that RUS can improved intestinal vascular leakage, and up-regulated the expression of connexin ZO-1, Occludin, VEcadherin, and inhibited ROCK activation and MLC phosphorylation. RUS Inhibit the remodeling of the endothelium of blood vessels, exerting the protective effect of the endothelium and inhibiting the adverse reactions caused by Dasatinib.

In order to simulate the clinical use of Dasatinib better[31], we observed it by high-dose oral administration Dasatinib and found that RUS can display a certain protective effect. In patients with intracerebral hemorrhage, monitoring found that the patient's cerebral blood flow decreased[22], so we suspected that monitoring intestinal blood flow can reflect intestinal bleeding. The experimental results showed that after administration of Dasatinib by gavage, the intestinal blood flow of mice significantly decreased. Through this way, we provide a monitoring method for patients who need long-term to take Dasatinib or similar drugs that can cause adverse events like bleeding. After oral administration of Dasatinib, the small intestine structure of the mice was destroyed, bleeding increased, and vascular leakage increased. Pre-administration of RUS reversed the decrease in blood flow and reduced leakage of Evans Blue.

Only recently, the intestinal vascular endothelium is recognized as an important barrier in the intestine[32–33]. Vascular endothelial cell is one of the main components of intestinal mucosa. Protecting the intestinal endothelial barrier and maintain vascular stability is become a new way to treat intestinal diseases[34–36]. The destruction of connexin will affect the integrity of intestinal blood vessels, increase the permeability of blood vessels, and cause leakage of blood vessels. Therefore, connexin plays a significant role in stabilizing intestinal permeability. Tight junctions are essential for regulating intestinal permeability[37–38]. The tight junction proteins between endothelial cells also play important functions. Common tight junction proteins include ZO-1, claudin-5, occludin and so on. Existing studies have

shown that related proteins such as ZO-1, occludin, claudin-5 can be used as criteria for evaluating tight junctions. When these proteins are significantly reduced, it indicates that the tight junction function is destroyed[39]. Among them, ZO-1 is a junctional adaptor protein that interacts with multiple other junctional components. It can regulate the formation of blood vessels in vitro and in vivo, maintain the connection tension, and at the same time, ZO-1 binds to F-actin and is involved in the regulation of cytoskeleton by actomyosin[40]. The endothelial cytoskeleton plays an important role in maintaining the structure of endothelial cells. There is a specifically expressed protein in endothelial cells: VE-cadherin, which plays an important role in maintaining the adhesion connection and is also an important indicator for evaluating the integrity of the adhesion connection. The steady-state dynamics of VE-cadherin at AJs is a critical determinant of AJ integrity[41]. There is a close relationship between VE-cadherin and ZO-1[40]. A recent study shows that protecting the endothelial barrier has become a new strategy for treating inflammatory bowel disease[42–43]. All in all, adherens junctions and tight junctions play an important role in maintaining the endothelial barrier. Early administration of RUS can protect the expression of connexin. Compared with the model group, the expressions of ZO-1, VE-cadherin, and Occludin are all increased, and the results are the same in HUVECs in vitro.

The cytoskeletal protein F-actin polymerization produces a mechanical force to change the cell morphology. Therefore, the integrity of the endothelial cytoskeletal protein F-actin structure is the basic structural for maintaining the barrier function of endothelial cells. Dasatinib phosphorylates MLC via the ROCK pathway[24,44], induces the polymerization of F-actin structure, remodels the cytoskeleton, and destroys the barrier function of endothelial cells. In view of the function of RUS to protect endothelial cells, we speculate that RUS can inhibit ROCK activation, reduce MLC phosphorylation, reduce cytoskeleton remodeling, restore connexin expression, and play a protective role by interfering with the ROCK-MLC signaling pathway.

Dasatinib has excellent efficacy in the clinical treatment of leukemia, but as a protein kinase inhibitor, it is bound to affect the function of the endothelial barrier. It is well known that many different phosphatases and kinases can act as regulators of endothelial barrier function[45–47]. After taking Dasatinib, it destroys the original regulatory function, causing damage to the endothelium, which causes numerous adverse reactions. Therefore, before we use these drugs with a high risk of side effects, we can match to some drugs with hemostasis, barrier protection and regulation related signaling pathways to bring more benefits to patients. Some natural products have good development potential[48].

Ruscogenin is a kind of saponins, which is more common in the traditional Chinese medicine *Ophiopogon japonicus*. Existing studies have shown that Ruscogenin can inhibit venous thrombosis and arterial thrombosis, reduce inflammation[49], and protect the barrier function., anti-thrombosis and protective barrier function[19]. Due to the lack of effective interventions for the bleeding side effects caused by Dasatinib, we considered that Ruscogenin has the function of protecting the endothelial barrier, and there are no clear side effects reported, so we assumed that Ruscogenin can improve the side effects caused by Dasatinib through protects barrier. The use of traditional Chinese medicine to regulate the side effects of other drugs has been initially recognized as a new treatment strategy[50].

# Declarations

## Conflicts of interest

The authors declare that there is no conflict of interest regarding the publication of this paper.

## Authors and Contributors

Yuanyuan Zhang, Junping Kou and Boyang Yu designed experiments. Yuankai Liu, Yujie Dai and Han Xu performed experiments. Yuankai Liu wrote the manuscript. Yuanyuan Zhang and Junping Kou polished the language. All authors contributed toward editing the manuscript.

## Funding

This study was supported by the "Double First-Class" University project (CPU2018GF07), the Fundamental Research Funds for the Central Universities (2632019ZD17).

## Data availability statement

Data are available on request due to privacy or other restrictions

# Abbreviations

BSA (Bovine serum albumin); ECL (Enhanced chemiluminescence); FBS (Fetal bovine serum); HRP (Horse radish peroxidase); HUVECs (Human umbilical vein endothelial cells); MLC (Myosin Light Chain); ROCK (Rho-associated kinase); PVDF (Polyvinylidene fluoride); SDS (Sodium dodecylsulfate); AJs (Adherens junction); TJ (Tight junctions).

# References

1. Ubel PA, Abernethy AP, Zafar SY (2013) Full Disclosure - Out-of-Pocket Costs as Side Effects [J]. *N Engl J Med* 369(16):1484–1486
2. Budnitz DS, Lovegrove MC, Shehab N, et al (2011) Emergency Hospitalizations for Adverse Drug Events in Older Americans [J]. *N Engl J Med* 365(21):2002–2012
3. Ren M, Qin H, Ren R et al (2011) Src activation plays an important key role in lymphomagenesis induced by FGFR1 fusion kinases [J]. *Can Res* 71(23):7312–7322
4. Eide CA, Adrian LT, Tyner JW et al (2011) The ABL switch control inhibitor DCC-2036 is active against the chronic myeloid leukemia mutant BCR-ABL T315I and exhibits a narrow resistance profile [J]. *Can Res* 71(9):3189–3195
5. Duque AJ, Lin CH, Han K et al (2018) CBP modulates sensitivity to Dasatinib in Pre-BCR acute lymphoblastic leukemia [J]. *Can Res* 78(22):6497–6508


6. Oliver O, Giuseppe S, Apperley JF et al (2018) Long-term efficacy and safety of Dasatinib in patients with chronic myeloid leukemia in accelerated phase who are resistant to or intolerant of Imatinib[J]. *Blood Cancer* 8(9):88
7. Huang XJ, Jiang Q, Hu JD et al (2019) Four-year follow-up of patients with Imatinib-resistant or intolerant chronic myeloid leukemia receiving Dasatinib: efficacy and safety[J]. *Frontiers of Medicine* 13(3):344–353
8. Steegmann JL, Cervantes F, Le Coutre P et al (2012) Off-target effects of BCR–ABL1 inhibitors and their potential long-term implications in patients with chronic myeloid leukemia. [J] *Leukemia Lymphoma* 53(12):2351–2361
9. Justice JN, Nambiar AM, Tchkonja T et al (2019) Senolytics in idiopathic pulmonary fibrosis: Results from a first-in-human, open-label, pilot study[J]. *EBIO Medicine* 40:554–563
10. Cortes JE, Saglio G, Kantarjian HM et al (2016) Final 5-Year study results of DASISION: The Dasatinib versus Imatinib study in treatment-Naïve chronic myeloid leukemia patients trial[J]. *J Clin Oncol* 34(20):2333–2340
11. Hughes TP, Laneuville P, Rousselot P et al (2019) Incidence, outcomes, and risk factors of pleural effusion in patients receiving Dasatinib therapy for Philadelphia chromosome-positive leukemia[J]. *Haematologica* 104(1):93–101
12. Andrew MB, Daniel BC, Rebecca SH (2013) Treatment-Related Toxicities in a Phase II Trial of Dasatinib in Patients with Squamous Cell Carcinoma of the Lung[J]. *Journal of Thoracic Oncology* 8(11):1434–1437 t,et al.
13. Daccord C, Letovanec I, Yerly P et al. First histopathological evidence of irreversible pulmonary vascular disease in Dasatinib-induced pulmonary arterial hypertension[J]. *European Respiratory Journal*.2018;51(3)
14. Weatherald J, Chaumais MC, Savale L,et al. Long-term outcomes of Dasatinib-induced pulmonary arterial hypertension: a population-based study[J]. *The European Respiratory Journal*.2017;50(1)
15. Calizo RC, Bhattacharya S, van Hasselt JGC et al (2019) Disruption of podocyte cytoskeletal biomechanics by Dasatinib leads to nephrotoxicity[J]. *Nat Commun* 10(1):2061
16. Quintás-Cardama A, Kantarjian H, Ravandi F et al (2009) Bleeding diathesis in patients with chronic myelogenous leukemia receiving Dasatinib therapy[J]. *Cancer* 115(11):2482–2490
17. Brave M, Goodman V, Kaminskas E et al (2008) Sprycel for Chronic Myeloid Leukemia and Philadelphia Chromosome-Positive Acute Lymphoblastic Leukemia Resistant to or Intolerant of Imatinib Mesylate[J]. *Clin Cancer Res* 14(2):352–359
18. Outmanoeuvre A, ColomFernández B, Jiménez AM et al (2017) Dasatinib Reversibly Disrupts Endothelial Vascular Integrity by Increasing Non-Muscle Myosin II Contractility in a ROCK-Dependent Manner[J]. *Clin Cancer Res* 23(21):6697–6707
19. Cao GS, Jiang N, Hu Y (2016) **et al. Ruscogenin attenuates cerebral ischemia-induced blood-brain barrier dysfunction by suppressing TXNIP/NLRP3 inflammasome activation and the MAPK pathway**[J]. *Int J Mol Sci* 17(9):1418–1435

20. Shi Y, Zhang L, Pu H, *et al.* Rapid endothelial cytoskeletal reorganization enables early blood–brain barrier disruption and long-term ischaemic reperfusion brain injury[J]. *Nat Commun*, 2016, 7:10523–10541
21. Liu Liu, Ping Zhang, Zhaosi, Zhang, **et al.** *LXA4 ameliorates cerebrovascular endothelial dysfunction by reducing acute inflammation after subarachnoid hemorrhage in rats*[J]. *Neuroscience*, 2019, 408:105–114.
22. Vaibhav K, Braun M, Khan MB (2018) **et al. Remote ischemic post-conditioning promotes hematoma resolution via AMPK-dependent immune regulation**[J]. *J Exp Med* 215(10):2636–2654
23. Phan Carole, Jutant Etienne-Marie, Tu Ly, **et al.** *Dasatinib increases endothelial permeability leading to pleural effusion*[J]. *The European respiratory journal*, 2018, 51(1):218–222
24. Dasgupta SK, Le A, Vijayan KV *et al* (2017) Dasatinib inhibits actin fiber reorganization and promotes endothelial cell permeability through RhoA-ROCK pathway[J]. *Cancer Med* 6(4):809–818
25. Yuanyuan Z, Yuwei H, Yazheng Z, *et al.* DT-13 Ameliorates TNF- $\alpha$ -Induced Vascular Endothelial Hyperpermeability via Non-Muscle Myosin IIA and the Src/PI3K/Akt Signaling Pathway[J]. *Front Immunol*, 2017, 8:925–936
26. Li Y, Liu Y, Yan X, *et al.* Pharmacological Effects and Mechanisms of Chinese Medicines Modulating NLRP3 Inflammasomes in Ischemic Cardio/Cerebral Vascular Disease[J]. *The American Journal of Chinese Medicine*, 2018, 46(08):1727–1741
27. Guan T, Liu Q, Qian Y, *et al.* Ruscogenin reduces cerebral ischemic injury via NF- $\kappa$ B-mediated inflammatory pathway in the mouse model of experimental stroke[J]. *Eur J Pharmacol*, 2013, 714(1–3):303–311
28. Ghulam Jilany Khan, Mohsin Rizwan, Muhammad Abbas, *et al.* *Pharmacological effects and potential therapeutic targets of DT-13*[J]. *Biomedicine & Pharmacotherapy*, 2018, 97:255–263.
29. Liu H, Yu X, Yu S *et al* (2015) Molecular mechanisms in lipopolysaccharide-induced pulmonary endothelial barrier dysfunction[J]. *Int Immunopharmacol* 29(2):937–946
30. Nishiwaki S, Maeda M, Yamada M *et al* (2017) Clinical efficacy of fecal occult blood test and colonoscopy for Dasatinib-induced hemorrhagic colitis in CML patients[J]. *Blood* 129(1):126–128
31. Jain P, Kantarjian H, Alattar ML (2015) **et al. Long-term molecular and cytogenetic response and survival outcomes with Imatinib 400 mg, Imatinib 800 mg, Dasatinib, and nilotinib in patients with chronic-phase chronic myeloid leukaemia: retrospective analysis of patient data from five clinical trials**[J]. *The Lancet Haematolog* 2(3):e118–e128
32. Spadoni I, Zagato E, Bertocchi A *et al* (2015) A gut-vascular barrier controls the systemic dissemination of bacteria[J]. *Science* 350(6262):830–834
33. Spadoni I (2017) Fornasa G, and Rescigno M. Organ-specific protection mediated by cooperation between vascular and epithelial barriers[J]. *Nat Rev Immunol* 17(12):761–773
34. Danese S, Sans M, de la Motte C, *et al.* Angiogenesis as a Novel Component of Inflammatory Bowel Disease Pathogenesis[J]. *Gastroenterology*. 2006;130(7):2060–2073

35. Danese S (2014) Panés J. Development of Drugs to Target Interactions Between Leukocytes and Endothelial Cells and Treatment Algorithms for Inflammatory Bowel Diseases[J]. *Gastroenterology* 147(5):981–989
36. He Y, Yuan XM, Zuo H, et al. Berberine Exerts a Protective Effect on Gut-Vascular Barrier via the Modulation of the Wnt/Beta-Catenin Signaling Pathway During Sepsis[J]. *Cellular Physiology and Biochemistry*. 2018; 49(4):1342–1351
37. Kim Y, West GA, Ray G (2017) et al. **Layilin is critical for mediating hyaluronan 35 kDa-induced intestinal epithelial tight junction protein ZO-1 in vitro and in vivo**[J]. *Matrix Biol* 66:93–109
38. Guo J, Cai H, Ma J (2017) et al. **Long non-coding RNA NEAT1 regulates permeability of the blood-tumor barrier via miR-181d-5p-mediated expression changes in ZO-1, occludin, and claudin-5**[J]. - *Molecular Basis of Disease* 1863(9):2240–2254 *Biochimica et Biophysica Acta (BBA)*
39. Tornavaca O, Chia M, Dufton N (2015) et al. **ZO-1 controls endothelial adherens junctions, cell-cell tension, angiogenesis, and barrier formation**[J]. *The Journal of Cell Biology* 208(6):821–838
40. Komarova YA, Kruse K, Mehta D et al (2017) Protein Interactions at Endothelial Junctions and Signaling Mechanisms Regulating Endothelial Permeability[J]. *Circ Res* 120(1):179–206
41. Bazzoni G, Dejana E (2004) Endothelial Cell-to-Cell Junctions: Molecular Organization and Role in Vascular Homeostasis[J]. *Physiol Rev* 84(3):869–901
42. Langer V, Vivi E, Regensburger D (2019) et al. **IFN- $\gamma$  drives inflammatory bowel disease pathogenesis through VE-cadherin-directed vascular barrier disruption**[J]. *Journal of Clinical Investigation* 129(11):4691–4707
43. Dejana E, Orsenigo F, Lampugnani MG (2008) The role of adherens junctions and VE-cadherin in the control of vascular permeability[J]. *J Cell Sci* 121(13):2115–2122
44. Gover PA, Granot G, Pasmanik CM et al (2019) Bosutinib, Dasatinib, Imatinib, nilotinib, and ponatinib differentially affect the vascular molecular pathways and functionality of human endothelial cells[J]. *Leuk Lymphoma* 60(1):189–199
45. Polacheck WJ, Kutys ML, Yang J (2017) et al. **A non-canonical Notch complex regulates adherens junctions and vascular barrier function**[J]. *Nature* 552(7684):258–262
46. Han Ji, Zhang G, Welch EJ, et al. **A critical role for Lyn kinase in strengthening endothelial integrity and barrier function**[J]. *Blood*. 2013;122(25):4140–4149
47. Gorovoy M, Han J, Pan H (2009) et al. **LIM Kinase 1 Promotes Endothelial Barrier Disruption and Neutrophil Infiltration in Mouse Lungs**[J]. *Circ Res* 105(6):549–556
48. Teng JF, Qin DL, Mei QB (2019) et al. **Polyphyllin VI, a saponin from *Trillium tschonoskii* Maxim. induces apoptotic and autophagic cell death via the ROS triggered mTOR signaling pathway in non-small cell lung cancer**[J]. *Pharmacol Res* 147:104396
49. Huang YL., Kou JP, Ma L (2008) et al. **Possible mechanism of the anti-inflammatory activity of ruscogenin: role of intercellular adhesion molecule-1 and nuclear factor-kappaB**[J]. *Journal of Pharmacological Sciences* 108(2):198–205


50. Parim B, Sathibabu Uddandrao VV, Saravanan G (2019) **Diabetic cardiomyopathy: molecular mechanisms, detrimental effects of conventional treatment, and beneficial effects of natural therapy**[J]. Heart Fail Rev 24(2):279–299

## Figures





**Figure 1**

The change of intestinal blood flow in mice with Dasatinib .Mice were orally administered with Dasatinib (70×140×210 mg/kg). The representative images of intestinal blood flow in different groups. The magnitude of IBF is represented by different colors, with blue to red denoting low to high,Quantitative analysis of IBF in different groups. Data are expressed as the mean±SD, n=6. #P<0.05 vs.Control,##P<0.01 vs.Control,###P<0.001 vs.Control; \*P<0.05 vs. Model, \*\*P<0.01 vs.Model,\*\*\*P<0.001 vs.Model.




**Figure 2**

Protective effect of RUS on side effects caused by Dasatinib. Mice were orally administered with RUS (3–10–30  $\mu\text{g}/\text{kg}$ ) prior to Dasatinib (140  $\text{mg}/\text{kg}$ , 90 min, po). A-B) Representative gross appearance of EB-stained intestinal from mouse. Quantitative analysis of EB extravasation by spectrophotometry. Data are expressed as the mean  $\pm$  SD,  $n=6$ .  $###P<0.01$  vs. Control mice;  $**P<0.01$  vs. Model mice. C) Hematoxylin-eosin stained slides of mouse intestinal sections in different groups were examined under a light microscope, and representative stained sections showed RUS improves Dasatinib-induced bleeding. Scale bar = 100  $\mu\text{m}$ .  $n=3$ . D) The representative images of intestinal blood flow in different groups. The magnitude of IBF is represented by different colors, with blue to red denoting low to high. Quantitative analysis of IBF in different groups. Data are expressed as the mean  $\pm$  SD,  $n=6$ .  $\#P<0.05$  vs. Control,  $##P<0.01$  vs. Control,  $###P<0.001$  vs. Control;  $*P<0.05$  vs. Model,  $**P<0.01$  vs. Model,  $***P<0.001$  vs. Model.




**Figure 3**

Effect of RUS on the expression of junction proteins in mice with Dasatinib. Mice were orally administered with RUS (3,10,30μg/kg) prior to Dasatinib (140 mg/kg, po). A-C) Representative Western blots and the quantitative analysis of the ratio of ZO-1,Occludin,VE-cadherin. Data are expressed as the mean±SD, n=4. #P<0.05 vs.Control,##P<0.01 vs.Control,###P<0.001 vs.Control; \*P<0.05 vs. Model, \*\*P<0.01 vs.Model,\*\*\*P<0.001 vs.Model. D) Representative confocal microscopy images showing the immunostaining of ZO-1 in mice.Scale bar =100 μm. n=3.




**Figure 4**

Effect of RUS on the ROCK1/MLC signaling pathway in mice with Dasatinib. Mice were orally administered with RUS (3,10,30μg/kg) prior to Dasatinib (140 mg/kg, po). A) Western blots and the quantitative analysis of the ratio of ROCK-1, B) Western blots and the quantitative analysis of the ratio of P-MLC. Data are expressed as the mean±SD, n=4. #P<0.05 vs.Control,##P<0.01 vs.Control,###P<0.001 vs.Control; \*P<0.05 vs. Model, \*\*P<0.01 vs.Model,\*\*\*P<0.001 vs.Model.



**Figure 5**

Effects of RUS on barrier function in HUVECs subjected to Dasatinib. The HUVECs were treated with RUS (0.1-1  $\mu$ M) 1h, exposed to 2 h of Dasatinib. (A, B) The barrier-protective effect of RUS was evaluated using EB albumin assays and Representative confocal microscopy images showing the immunostaining of F-actin in HUVECs. The Data are expressed as the mean $\pm$ SD, n=3. #P<0.05 vs.Control,##P<0.01 vs.Control,###P<0.001 vs.Control; \*P<0.05 vs. Model, \*\*P<0.01 vs.Model,\*\*\*P<0.001 vs.Model.



**Figure 6**

Effects of RUS on the expression of junction proteins and ROCK1/MLC signaling pathway in HUVECs subjected to Dasatinib. (A,B) The HUVECs were treated with RUS(0.01-1 $\mu$ M)1h, followed by 2 h of Dasatinib. The expression levels of ZO-1, VE-cadherin, Occludin levels were measured using Western blotting. The band intensities were assessed using scanning densitometry, and the data were normalized against the  $\beta$ -actin signal. The Data are expressed as the mean $\pm$ SD, n=4. #P<0.05 vs.Control,##P<0.01 vs.Control,###P<0.001 vs.Control; \*P<0.05 vs. Model, \*\*P<0.01 vs.Model,\*\*\*P<0.001 vs.Model. (C,D) The HUVECs were treated with RUS(0.01-1 $\mu$ M)1h, followed by 2 h of Dasatinib. The expression levels of ROCK-1,P-MLC levels were measured using Western blotting. The expression levels of ROCK1, MLC and p-MLC were determined using Western blotting analysis. The band intensities were assessed through scanning densitometry. The Data are expressed as the mean $\pm$ SD, n=4. #P<0.05 vs.Control,##P<0.01 vs.Control,###P<0.001 vs.Control; \*P<0.05 vs. Model, \*\*P<0.01 vs.Model,\*\*\*P<0.001 vs.Model.

## Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [GraphicalAbstract.png](#)