Air Pollution and Racial Disparities in Pregnancy Outcomes in the United States: A Systematic Review

Bonaventure Suiru Dzekem

University of Chicago

Briseis Aschebrook-Kilfoy

University of Chicago

Christopher O. Olopade (✉ solopade@bsd.uchicago.edu)

University of Chicago

Research Article

Keywords: Air pollution, Racial disparities, Pregnancy outcomes

Posted Date: February 23rd, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-208924/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Abstract

Background

Exposure to air pollutants and other environmental factors increases the risk of adverse pregnancy outcomes. There is growing evidence that adverse outcomes related to air pollution disproportionately affect racial and ethnic minorities. This paper explores the importance of race as a risk factor for air pollution-related poor pregnancy outcomes.

Methods

We systematically reviewed epidemiologic studies investigating the effects of exposure to air pollution on pregnancy outcomes in the United States.

Results

Findings across all reviewed studies show more adverse pregnancy outcomes (preterm birth, small for gestational age, low birth weight and stillbirths) due to air pollution among Blacks and Hispanics than among non-Hispanic Whites.

Conclusion

This paper enhances our general understanding of the impact of air pollution on birth outcomes and, specifically, of disparities in birth outcomes for infants born to Black and Hispanic mothers.

Introduction

According to the World Health Organization (WHO), 91% of the world's population lives in places where air pollution exceeds recommended limits (1). Globally, about 4.2 million deaths are attributed to exposure to ambient air pollution and an additional 3.8 million deaths to exposure to household air pollution, with a disproportionate burden among women and children (2, 3).

There is evidence that exposure to air pollution adversely affects human health (4–8). Additionally, in the last two decades, evidence has emerged that exposure to a variety of pollutants in the air and other environmental factors increase the risk of adverse pregnancy outcomes (9–12). Further, evidence is accumulating that adverse outcomes such as low birth weight (LBW), preterm birth (PTB) and small for gestational age (SGA) are common in racial minority groups (13–15). These pregnancy outcomes have been shown to have serious health consequences during the neonatal period and infancy (16–18), childhood (19–21), and adulthood (22–24). Public health studies in the United States often incorporate race and ethnicity as factors to adjust for. Alternatively, results are reported stratified by race/ethnicity. The adverse impact of air pollution in racial and ethnic minorities is attributed to factors including age, geographical location, socioeconomic status and education (25–27). Although this accumulating research offers solid evidence that exposure to air pollution increases the risk of adverse pregnancy outcomes in general, there is still a need to assess its specific impact on pregnancy outcomes in minority populations, in light of growing racial disparities.

Here, we reviewed the epidemiologic studies investigating the effects of exposure to air pollution on pregnancy outcomes in the United States with results stratified by race and ethnicity to provide a comprehensive view of existing research, including types of measurement and study designs, to inform future research directions and to deepen insight into this important topic.

Methods

We searched the PubMed database for eligible studies that explored the effects of pregnancy outcomes on women across all races by combining Medical Subject Headings (MeSH). Key search terms used for the search included a combination of “air pollution” and one or more of the following; “pregnancy outcomes,” “birth outcomes,” “complications of pregnancy,” “race,” “ethnicity,” “black,” “white,” “African American” and “continental population groups.” Searches with these terms yielded 14,661 articles. After filtering for studies in the English language that examined effects of air pollution on pregnancy outcomes, we retained 124 articles. Titles and abstracts of all retained studies were reviewed for eligibility criteria and the full articles for eligible studies were accessed via journal sites. Eligible studies were those that explored air pollution and racial disparities in pregnancy outcomes. We excluded articles that did not have racial comparisons and studies that did not specifically explore the effects of air pollution. Of the 124 studies retained, 16 compared pregnancy outcomes among two or more racial groups. For each, we report the study design, sample size, type of pollutant and method used to capture exposure. We also report the type of adverse pregnancy outcome; specifically, preterm births (PTB), small for gestational age (SGA), low birth weight (LBW), and stillbirths. We compared findings across non-Hispanic white, Hispanic and Black race/ethnicity, Table 1.

Table 1
Air Pollution and Racial Disparities in Pregnancy Outcomes in the United States

Reference	Study design	Study years	Country	Study population /sample size	Total black	Total white	Total Hispanic	Method used to capture exposure	Outcome measure	Adverse Pregnancy outcome Total	
Benmarhnia T et al	Retrospective cohort	2005–2010	California USA	1,066,783	175,297	891,486	Not specified	Measurement of PM2.5 and NO2 by EPA	PTB, SGA	Prevalence PTB = 10.9%, SGA = 2.2%	F
Gray SC et al	Retrospective cohort	2001–2006	North Carolina USA	457,642	22.8%	62.4%	14.8%	Bayesian Measurement of PM2.5 and O3	LBW, SGA, PTB	LBW = 7.3%; SGA = 10.2%; PTB = 9.8%	C 2 F 1
Wallace ME et al	Retrospective cohort study	2000–2008	Multi national	223,375	50,255	110,541	38,811	CMAQ model	PROM	PROM = 7%	F
Hao H, et al	Retrospective cohort	2002–2006	Georgia USA	511,658	161,583	329,152	Not Specified	CMAQ model CO, SO2, O3, NO2, PM2.5	PTB	PTB = 9.3%	F
Woodruff TJ et al	Retrospective cohort	1998–1999	California USA	4,098,740	15.0%	61.2%	18.8%	U.S. EPA Ambient air quality measurement PM10, O3, CO, NO2, SO2	SGA, PTB	SGA = 9.0%, PTB = 9.9%	A 2 C
Kingsley SL et al	Retrospective cohort	2002–2012	Rhode Island USA	61,640	4,706	36,510	Not Specified	land-use regression and satellite remote sensing. PM2.5, BC	PTB		F
DeFranco E et al	Retrospective cohort	2006–2010	Ohio USA	351,036	46.6%	44.8%	5.8%	U.S. EPA Local measurement of PM2.5	Stillbirths	Stillbirths = 0.5%	S 1
Nobles CJ et al	Retrospective cohort	2002–2010	Maryland USA	109,126	216	43,055	5,328	CMAQ model, SO2, NO2, PM10	FGR	FGR = 1.5% (825)	F
Le HQ et al	Retrospective cohort	1990–2001	Michigan USA	164,905	93,078	68,164	Not specified	Ambient air monitors, SO2, O3CO, NO2,, PM10	PTB, SGA	nSGA = 13,754 nPTB = 24,954	S F
Miranda ML et al	Retrospective cohort	2004–2008	North Carolina USA	468,517	23.3%	60.3%	16.4%	Proximity to major road	LBW, PTB, SGA	LBW = 6.8%, PTB = 10.5%, 10.2%	L F S
Laurent O et al	Retrospective cohort	1997–2006	California USA	70,000	6,261	30,548	23,678	Mixed method NOx, CO, O3, PM	LBW		L
Bell ML et al	Retrospective cohort	1999–2002	California USA	358,504	10.7%	83.4%	Not specified	US EPA measurements on NO2, SO2, PM, CO	LBW	LBW = 4.01%	L t
Rich DQ et al	Retrospective cohort	1999–2003	New Jersey USA	350,107	38,978	84,747	54,473	US EPA measurements PM2.5, NO2, SO2, CO	SGA	SGA = 8%, VSGA = 2%	S
Geer LA et al	Retrospective cohort	1998–2004	Texas USA	1,548,904	10.6%	34.7%	50.7%	US EPA measurements PM, SO2 NO2, CO	LBW	LBW = 2.8%	L t

Reference	Study design	Study years	Country	Study population /sample size	Total black	Total white	Total Hispanic	Method used to capture exposure	Outcome measure	Adverse Pregnancy outcome Total	
Ghosh JK et al	Case control	2003	CaliforniaUSA	1,761	7%	10.9%	73.6%	Air quality monitoring data	LBW		L
Kahr MK et al	Retrospective cohort	2011–2014	Texas USA	9004	10.7%	30.5%	69.5%	People commuting to work	PTB	PTB = 10%	F

Results

All 16 studies that met the review criteria were conducted between 1990 and 2014. Of these, 15 were retrospective cohort studies using birth registries and one was a case control study (Table 1). There was representation of states across regions of the United States. Sample sizes ranged from 1,761 births to 1,548,904 births. Several different methods were used to estimate levels of air pollution exposure, including Bayesian measurement, Community Multi-scale Air Quality (CMAQ) model, data from United States Environmental Protection Agency (EPA), installation of ambient air monitors, proximity to major roads, and mixed methods. While most studies looked at a single pregnancy outcome, others considered multiple outcomes. Specific details of racial disparities in pregnancy outcomes across race are described below.

Air pollution and Preterm birth

Preterm birth was assessed in eight out of 16 of the studies we reviewed. Benmarhnia et al. (28) used decomposition analysis to understand the racial disparities in PTB in California. Two pollutants were included: fine-grained particulate matter (PM_{2.5}) and nitrogen dioxide (NO₂). A higher prevalence of PTB was observed for non-Hispanic Black (NHB) mothers when compared with non-Hispanic White mothers. The predicted difference in probability of PTB between Black and White infants was 0.056 (95% CI: 0.054, 0.058). All included predictors explained 37.8% of the Black–White disparity. Overall, individual variables (17.5% for PTB) such as age and level of education, and neighborhood-level variables (16.1% for PTB) such as socioeconomic environment explained a greater proportion of the Black–White difference in birth outcomes than air pollution (5.7% for PTB). Gray and colleagues reported that infants born to NHB mothers were at increased odds of PTB (adjusted OR 1.46, 95% CI: 1.42 to 1.50). (27)

Using the CMAQ model and measurements from stationary monitors, Hao et al. (29) investigated the association between 11 ambient air pollutants and the risk of PTB in the state of Georgia. They observed that all traffic-related pollutants (carbon monoxide (CO), NO₂, PM_{2.5}, elemental carbon) were associated with PTB [e.g., odds ratios for interquartile range increases in CO during the first, second, and third trimesters and total pregnancy were 1.005 (95% CI: 1.001, 1.009), 1.007 (95% CI: 1.002, 1.011), 1.010 (95% CI: 1.006, 1.014), and 1.011 (95% CI: 1.006, 1.017), respectively]. Associations were higher for mothers with low educational attainment and African American mothers. There were no clear explanations for the observed associations. Other studies had similar findings (29–32).

Woodruff et al. (31) using data from the EPA Ambient Air Monitoring observed that Hispanic, African-American, and Asian/Pacific Islander mothers experienced higher mean levels of air pollution and were more than twice as likely to live in the most polluted counties compared with White mothers. In addition, there was an increase in the odds of PTB (adjusted odds ratio (AOR) = 1.05; 95% CI, 0.99–1.12) in counties with higher air pollution.

Le et al. (32) reported a stronger association between air pollutants and PTB for Blacks than Whites. PTB was associated with SO₂ (OR 1.07, 1.01 – 1.14) exposure in the last month of pregnancy, while O₃ exposures exceeding 92 parts per billion (OR 1.08, 1.02 – 1.14) were associated with PTB in the first months of pregnancy.

Air pollution and Small for Gestational Age

Of all 16 studies reviewed, six assessed SGA. Benmarhnia et al. (28) explored racial disparities in SGA in California. The predicted difference in probability of SGA between Black and White infants was 0.084 (95% CI: 0.081, 0.087). Together, individual demographics, neighborhood socioeconomic environment (such as unemployment and poverty rates), and neighborhood air pollution explained 37.8% of the Black-White disparity. There was a higher prevalence of SGA among non-Hispanic Blacks (18.3%) as compared to non-Hispanic Whites (9.9%).

In North Carolina (27), infants born to NHB mothers and Hispanic mothers were at an increased odds of being small for gestational age with (AOR = 2.18, 95% CI: 2.12 to 2.24 and AOR = 1.21, CI: 1.17 to 1.26, respectively), compared to NHW mothers. After controlling for race and individual and area-level socio-economic status, this difference persisted, suggesting that air pollution is an additional contributor to the observed outcomes.

In a retrospective cohort study of 4,098,750 births in California, Woodruff et al. (31) observed that Hispanic, African-American, and Asian/Pacific Islander mothers experienced higher mean levels of air pollution, determined from strategically-placed EPA monitors. They were also more than twice as likely to live in the most polluted counties compared with White mothers, after controlling for maternal risk factors, region, and educational status [Hispanic mothers: AOR = 4.66; 95% CI: 1.92–11.32; African-American mothers: AOR = 2.58; 95% CI: 1.00–6.62; Asian/Pacific Islander mothers: AOR = 2.82; 95% CI, 1.07–7.39]. However, there was no significant increase in the odds of SGA (AOR = 0.96; 95% CI, 0.86–1.07) in counties with higher air pollution.

In a cohort of 164,905 births in Michigan, Le et al. (32) showed that there was an association between term SGA with exposure to CO and NO₂ during the first trimester of pregnancy. They also reported an association between term SGA and exposure to O₃ and PM₁₀ during the later stages of pregnancy. There was

evidence of stronger associations between CO and term-SGA, NO₂ and term-SGA and SO₂ and term-SGA for infants of Black mothers as compared to White mothers.

Rich et al. (33) found significantly increased risk of SGA associated with first and third trimester exposures to PM_{2.5} and increased risk of very small for gestational age (VSGA) associated with first, second and third trimester exposures to high NO₂ concentrations. According to this study, mothers of SGA and VSGA infants were more likely to be less than 25 years old and less likely to have completed high school, compared to mothers of appropriate-size births. They were also more likely to be single, African American, and to have smoked during pregnancy.

Air Pollution and Low Birth Weight

Low birth weight was assessed in five of the 14 studies reviewed. Gray et al. (27) used multivariate analysis of factors, including PM_{2.5} and O₃ to assess their association with weight differences in grams and 95% CI for all births. PM_{2.5} exposures were associated with LBW among infants born to NHB and Hispanic mothers more than those born to NHW mothers (-187.5g, 95% CI: -183.6 to -191.4 and -46.8g, 95% CI: -41.8 to -51.7, respectively).

Miranda et al. (34) characterized maternal exposure to traffic-related air pollution during pregnancy by using residential proximity to major road ways as a proxy. Women residing within 250 meters of a major roadway were at a 3-5% increased odds of having a LBW baby than women residing more than 250 meters away (p<0.05). The mean birth weight was 3376g for NHW, 2114g for NHB, and 3330g for Hispanics.

In a retrospective cohort study in California (35), African Americans, despite being a significantly lower percentage of the study population, had 3.19% of the LBW infants, compared to 1.23% for whites and 1.59% for Hispanics. Bell et al. (36) also reported that the association between air pollutants (especially PM_{2.5}) and LBW for infants of Black mothers was stronger than for White mothers.

In a large cohort study of 1,548,904 births in Texas, Geer et al. (37) reported that interquartile increases in ambient air pollutant concentrations of SO₂ and O₃ were associated with a 4.99g (95% CI, 1.87–8.11) and 2.72g (95% CI, 1.11–4.33) decreases in birth weight, respectively. Lower birth weight was associated with exposure to O₃ in the first and second trimester, whereas results for other pollutants did not differ significantly by trimester.

In a case-control study, Ghosh et al. (38), observed that women who were exposed to secondhand smoke at home had increased odds of term LBW (AOR = 1.36; 95% CI: 0.85, 2.18) compared to unexposed women. Blacks and Hispanic had higher odds of having LBW babies compared to White mothers.

Air pollution and stillbirths

We found only one cohort study that assessed stillbirths in relation to air pollution. According to DeFranco et al. (11) high average PM_{2.5} exposure through pregnancy was not associated with a significant increase in stillbirth risk (AOR 1.21; 95% CI: 0.96,1.53). There was also no higher risk of stillbirth associated with exposure in either the first or second trimester. However, exposure to high levels of PM_{2.5} in the third trimester of pregnancy was associated with 42% increased risk of stillbirth, (AOR = 1.42; 1.06, 1.91). Stillbirth rates were higher among mothers older than 40 years (11.6 per 1000), NHB mothers (8.6 per 1000), and mothers with lower education level and tobacco use.

Discussion

This review suggests that air pollution is associated with adverse pregnancy outcomes in the United States and higher rates of adverse outcomes among people of color, especially Black women, than among Whites. These findings are consistent with studies conducted in other countries (13, 14, 39). A multitude of factors other than exposure to air pollution that have also been shown to be associated with poor pregnancy outcomes, including socioeconomic status, level of education, maternal smoking (40–42). Almost all the studies we reviewed were large retrospective cohort studies, therefore, such other factors likely at least partially explain the disparities observed in the studies reviewed. We point, however, to evidence from molecular epidemiology studies that suggest biological mechanisms to explain the effects of air pollution on pregnancy outcomes (43–45). There is little data to suggest that Black mothers have any greater genetic predisposition to adverse effects of air pollution than Whites (26, 28, 45). This could be the case, however, and is a potential area for future research. Finally, while studies suggest that PTB is associated with SO₂, and O₃ exposure, the effects of such pollutants may in part be due to their roles in exacerbating other disease conditions that are known to cause PTB; e.g., hypertension, diabetes, asthma and preeclampsia (46, 47). This is a further area for future research.

Even though none of the studies reviewed explored household air pollution, there is evidence that it also contributes to poor pregnancy outcomes, especially in developing countries where cooking is done by burning of biomass in poorly ventilated homes (12, 48–50). A number of measures can be taken to improve air quality and reduce the negative impacts of air pollution on overall health and on gestational health in particular. At individual and household levels, these measures include: avoidance of cooking with biomass fuel indoors, ensuring adequate household ventilation, quitting smoking, use of chimneys and the use of air filters (51). At state and national levels, policy measures could include: carbon tax and subsidies for alternative energy sources (52).

A potential shortcoming in our review is the limited number of available published studies. We cannot rule out that many studies on this topic with null results may have remained unpublished. We also could not assess how differences in the methods by which air pollution was captured in different studies may have contributed to the reported results. Potential exposure misclassification is certainly an important possibility to consider when evaluating research in this area of investigation, if we are to build a corpus of studies reliable enough to guide changes in practice and policy.

In summary, we have reviewed existing evidence that suggests a need for further investigation of the impact of air pollution on birth outcomes and how these vary by race and ethnicity. Despite the limitations noted, we believe that this review of 16, U.S.-based studies increases our understanding of the impact of air

pollution on maternal-child health and of disparities in maternal-child health across races and ethnicities. In conclusion, with the goal of enhancing the literature and filling gaps in our knowledge, on the basis of this review we suggest future studies that:

- Explore the biologic factors underlying air pollution's adverse effects.
- incorporate measurement of household air pollution and personalized exposure measurement.
- Establish the causal mechanisms of air pollution's impact on specific outcomes, including prematurity, birthweight, IUGR and fetal deaths.
- Determine the period during pregnancy in which women are most likely to be affected by air pollutants and to establish the specific effects of different pollutants and pollutant mixtures.
- Explore differential effects of specific air pollutants on Blacks versus Whites.

Declarations

- Ethics approval and consent to participate: N/A
- Consent for publication: All authors have read and approved the manuscript for publication
- Availability of data and materials: N/A
- Competing interests: None
- Funding: None
- Authors' contributions: CO and BD and BA conceived the idea for this review, BD drafted the manuscript. All authors critically reviewed and approved the final version.

Acknowledgements: This research paper is made possible through the support of the University of Chicago Center for Global Health and the Graduate Program in Health Administration and Policy (GPHAP). Special thanks to Laura Botwinick for mentorship and guidance during this project and to Susan Duncan for editorial assistance.

References

1. Global ambient air pollution [Internet]. [cited 2019 Apr 19]. Available from: <http://maps.who.int/airpollution/>
2. WHO | Air pollution [Internet]. WHO. [cited 2019 Apr 19]. Available from: <http://www.who.int/airpollution/en/>
3. Oluwale O, Arinola GO, Ana GR, Wiskel T, Huo D, Olopade OI, et al. Relationship between household air pollution from biomass smoke exposure, and pulmonary dysfunction, oxidant-antioxidant imbalance and systemic inflammation in rural women and children in Nigeria. *Glob J Health Sci*. 2013 Mar 18;5(4):28–38.
4. Shima M. Health Effects of Air Pollution: A Historical Review and Present Status. *Nihon Eiseigaku Zasshi Jpn J Hyg*. 2017;72(3):159–65.
5. Mannucci PM, Franchini M. Health Effects of Ambient Air Pollution in Developing Countries. *Int J Environ Res Public Health*. 2017 12;14(9).
6. Coker E, Kizito S. A Narrative Review on the Human Health Effects of Ambient Air Pollution in Sub-Saharan Africa: An Urgent Need for Health Effects Studies. *Int J Environ Res Public Health*. 2018 01;15(3).
7. Fowler D, Dise N, Sheppard L. Committee on air pollution effects research: 40 years of UK air pollution. *Environ Pollut Barking Essex 1987*. 2016 Jan;208(Pt B):876–8.
8. Khilnani GC, Tiwari P. Air pollution in India and related adverse respiratory health effects: past, present, and future directions. *Curr Opin Pulm Med*. 2018;24(2):108–16.
9. Glinianaia SV, Rankin J, Bell R, Pless-Mulloli T, Howel D. Particulate air pollution and fetal health: a systematic review of the epidemiologic evidence. *Epidemiol Camb Mass*. 2004 Jan;15(1):36–45.
10. Wu J, Laurent O, Li L, Hu J, Kleeman M. Adverse Reproductive Health Outcomes and Exposure to Gaseous and Particulate-Matter Air Pollution in Pregnant Women. *Res Rep Health Eff Inst*. 2016 Jul;(188):1–58.
11. DeFranco E, Hall E, Hossain M, Chen A, Haynes EN, Jones D, et al. Air Pollution and Stillbirth Risk: Exposure to Airborne Particulate Matter during Pregnancy Is Associated with Fetal Death. *PLoS ONE* [Internet]. 2015 Mar 20 [cited 2019 Feb 2];10(3). Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4368103/>
12. Arinola GO, Dutta A, Oluwale O, Olopade CO. Household Air Pollution, Levels of Micronutrients and Heavy Metals in Cord and Maternal Blood, and Pregnancy Outcomes. *Int J Environ Res Public Health*. 2018 17;15(12).
13. Grobman WA, Parker CB, Willinger M, Wing DA, Silver RM, Wapner RJ, et al. Racial Disparities in Adverse Pregnancy Outcomes and Psychosocial Stress. *Obstet Gynecol*. 2018 Feb;131(2):328–35.
14. Zhang S, Cardarelli K, Shim R, Ye J, Booker KL, Rust G. Racial Disparities in Economic and Clinical Outcomes of Pregnancy Among Medicaid Recipients. *Matern Child Health J*. 2013 Oct;17(8):1518–25.
15. Jones MR, Diez-Roux AV, O'Neill MS, Guallar E, Sharrett AR, Post W, et al. Ambient air pollution and racial/ethnic differences in carotid intima-media thickness in the Multi-Ethnic Study of Atherosclerosis (MESA). *J Epidemiol Community Health*. 2015 Dec;69(12):1191–8.
16. Paneth NS. The problem of low birth weight. *Future Child Cent Future Child David Lucile Packard Found*. 1995;5(1):19–34.
17. Kavut S, Celik K. Incidence and risk factors of postnatal growth restriction in preterm infants. *J Matern-Fetal Neonatal Med Off J Eur Assoc Perinat Med Fed Asia Ocean Perinat Soc Int Soc Perinat Obstet*. 2018 Apr;31(8):1105–7.

18. Catov JM, Scifres CM, Caritis SN, Bertolet M, Larkin J, Parks WT. Neonatal outcomes following preterm birth classified according to placental features. *Am J Obstet Gynecol*. 2017;216(4):411.e1-411.e14.
19. Jefferis BJMH, Power C, Hertzman C. Birth weight, childhood socioeconomic environment, and cognitive development in the 1958 British birth cohort study. *Br Med J*. 2002;325(7359):305–8.
20. Korten I, Ramsey K, Latzin P. Air pollution during pregnancy and lung development in the child. *Paediatr Respir Rev*. 2017 Jan;21:38–46.
21. Moreira RS, Magalhães LC, Alves CRL. Effect of preterm birth on motor development, behavior, and school performance of school-age children: a systematic review. *J Pediatr (Rio J)*. 2014 Apr;90(2):119–34.
22. Kaijser M, Bonamy A-KE, Akre O, Cnattingius S, Granath F, Norman M, et al. Perinatal risk factors for ischemic heart disease: Disentangling the roles of birth weight and preterm birth. *Circulation*. 2008;117(3):405–10.
23. Crump C, Sundquist K, Sundquist J. Adult outcomes of preterm birth. *Prev Med*. 2016;91:400–1.
24. Raju TNK, Buist AS, Blaisdell CJ, Moxey-Mims M, Saigal S. Adults born preterm: a review of general health and system-specific outcomes. *Acta Paediatr Oslo Nor 1992*. 2017 Sep;106(9):1409–37.
25. Blumenshine P, Egerter S, Barclay CJ, Cubbin C, Braveman PA. Socioeconomic disparities in adverse birth outcomes: a systematic review. *Am J Prev Med*. 2010 Sep;39(3):263–72.
26. McKinnon B, Yang S, Kramer MS, Bushnik T, Sheppard AJ, Kaufman JS. Comparison of black–white disparities in preterm birth between Canada and the United States. *CMAJ Can Med Assoc J*. 2016 Jan 5;188(1):E19–26.
27. Gray SC, Edwards SE, Schultz BD, Miranda ML. Assessing the impact of race, social factors and air pollution on birth outcomes: a population-based study. *Environ Health*. 2014 Jan 29;13:4.
28. Benmarhnia T, Huang J, Basu R, Wu J, Bruckner TA. Decomposition Analysis of Black–White Disparities in Birth Outcomes: The Relative Contribution of Air Pollution and Social Factors in California. *Environ Health Perspect [Internet]*. 2017 Oct 4 [cited 2019 Jan 25];125(10). Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5933346/>
29. Hao H, Chang HH, Holmes HA, Mulholland JA, Klein M, Darrow LA, et al. Air Pollution and Preterm Birth in the U.S. State of Georgia (2002–2006): Associations with Concentrations of 11 Ambient Air Pollutants Estimated by Combining Community Multiscale Air Quality Model (CMAQ) Simulations with Stationary Monitor Measurements. *Environ Health Perspect*. 2016 Jun;124(6):875–80.
30. Kahr MK, Suter MA, Ballas J, Ramphul R, Lubertino G, Hamilton WJ, et al. Preterm birth and its associations with residence and ambient vehicular traffic exposure. *Am J Obstet Gynecol*. 2016 Jul;215(1):111.e1-111.e10.
31. Woodruff TJ, Parker JD, Kyle AD, Schoendorf KC. Disparities in exposure to air pollution during pregnancy. *Environ Health Perspect*. 2003 Jun;111(7):942–6.
32. Le HQ, Batterman SA, Wirth JJ, Wahl RL, Hoggatt KJ, Sadeghnejad A, et al. Air pollutant exposure and preterm and term small-for-gestational-age births in Detroit, Michigan: Long-term trends and associations. *Environ Int*. 2012 Sep;44:7–17.
33. Rich DQ, Demissie K, Lu S-E, Kamat L, Wartenberg D, Rhoads GG. Ambient air pollutant concentrations during pregnancy and the risk of fetal growth restriction. *J Epidemiol Community Health*. 2009 Jun;63(6):488–96.
34. Miranda ML, Edwards SE, Chang HH, Auten RL. Proximity to roadways and pregnancy outcomes. *J Expo Sci Environ Epidemiol*. 2013 Jan;23(1):32–8.
35. Laurent O, Wu J, Li L, Chung J, Bartell S. Investigating the association between birth weight and complementary air pollution metrics: a cohort study. *Environ Health*. 2013 Feb 17;12:18.
36. Bell ML, Ebisu K, Belanger K. Ambient Air Pollution and Low Birth Weight in Connecticut and Massachusetts. *Environ Health Perspect*. 2007 Jul;115(7):1118–24.
37. Geer LA, Weedon J, Bell ML. Ambient air pollution and term birth weight in Texas from 1998 to 2004. *J Air Waste Manag Assoc* 1995. 2012 Nov;62(11):1285–95.
38. Ghosh JKC, Wilhelm M, Su J, Goldberg D, Cockburn M, Jerrett M, et al. Assessing the Influence of Traffic-related Air Pollution on Risk of Term Low Birth Weight on the Basis of Land-Use-based Regression Models and Measures of Air Toxics. *Am J Epidemiol*. 2012 Jun 15;175(12):1262–74.
39. Morello-Frosch R, Shenassa ED. The environmental “riscscape” and social inequality: implications for explaining maternal and child health disparities. *Environ Health Perspect*. 2006 Aug;114(8):1150–3.
40. Šrám RJ, Binková B, Dejmeek J, Bobak M. Ambient Air Pollution and Pregnancy Outcomes: A Review of the Literature. *Environ Health Perspect*. 2005 Apr;113(4):375–82.
41. Victora CG, Barros F, Huttly SR, Teixeira AMB, Vaughan P. Early childhood mortality in a brazilian cohort: The roles of birthweight and socioeconomic status. *Int J Epidemiol*. 1992;21(5):911–5.
42. How Racism May Cause Black Mothers To Suffer The Death Of Their Infants [Internet]. NPR.org. [cited 2019 May 18]. Available from: <https://www.npr.org/sections/health-shots/2017/12/20/570777510/how-racism-may-cause-black-mothers-to-suffer-the-death-of-their-infants>
43. Song J, Chen Y, Wei L, Ma Y, Tian N, Huang SY, et al. Early-life exposure to air pollutants and adverse pregnancy outcomes: protocol for a prospective cohort study in Beijing. *BMJ Open [Internet]*. 2017 Sep 3 [cited 2019 May 18];7(9). Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5588991/>
44. Dejmeek J, Solanský I, Benes I, Leníček J, Šrám RJ. The impact of polycyclic aromatic hydrocarbons and fine particles on pregnancy outcome. *Environ Health Perspect*. 2000 Dec;108(12):1159–64.
45. Perera F, Hemminki K, Jedrychowski W, Whyatt R, Campbell U, Hsu Y, et al. In utero DNA damage from environmental pollution is associated with somatic gene mutation in newborns. *Cancer Epidemiol Biomark Prev Publ Am Assoc Cancer Res Cosponsored Am Soc Prev Oncol*. 2002 Oct;11(10 Pt 1):1134–7.

46. Padula AM, Huang H, Baer RJ, August LM, Jankowska MM, Jellife-Pawlowski LL, et al. Environmental pollution and social factors as contributors to preterm birth in Fresno County. *Environ Health* [Internet]. 2018 Aug 29 [cited 2019 May 18];17. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6114053/>
47. Lavigne E, Yasseen AS, Stieb DM, Hystad P, van Donkelaar A, Martin RV, et al. Ambient air pollution and adverse birth outcomes: Differences by maternal comorbidities. *Environ Res*. 2016;148:457–66.
48. Alexander D, Northcross A, Wilson N, Dutta A, Pandya R, Ibigbami T, et al. Randomized Controlled Ethanol Cookstove Intervention and Blood Pressure in Pregnant Nigerian Women. *Am J Respir Crit Care Med*. 2017 15;195(12):1629–39.
49. Ghosh JKC, Wilhelm M, Ritz B. Effects of Residential Indoor Air Quality and Household Ventilation on Preterm Birth and Term Low Birth Weight in Los Angeles County, California. *Am J Public Health*. 2013 Apr;103(4):686–94.
50. Khan MN, B Nurs CZ, Mofizul Islam M, Islam MR, Rahman MM. Household air pollution from cooking and risk of adverse health and birth outcomes in Bangladesh: a nationwide population-based study. *Environ Health Glob Access Sci Source*. 2017 13;16(1):57.
51. New Guidance for Residential Air Cleaners - ProQuest [Internet]. [cited 2020 Jun 30]. Available from: <https://search.proquest.com/openview/411efdabd7ff51ce0fbe480b039313e0/1?pq-origsite=gscholar&cbl=41118>
52. US EPA O. Economic Incentives [Internet]. US EPA. 2014 [cited 2020 Jun 30]. Available from: <https://www.epa.gov/environmental-economics/economic-incentives>

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Table1review.docx](#)