

Proportion And Factors Of Death Among Preterm Neonates Admitted In Aksum University Compressive Specialized Hospital Neonatal Intensive Care Unit Northern Ethiopia 2019.

Teferi Gebru Gebremeskel (✉ teferigebru12@gmail.com)

Aksum University <https://orcid.org/0000-0002-8276-5685>

SINTAYEHU MAMO

Aksum University

Research

Keywords:

Posted Date: April 7th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-21042/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background Birth-related complications are the single most direct cause of national neonatal death, 35% of the world's 3.1 million deaths a year and the second most common cause of under-five deaths associated with the cost of living worldwide. To achieve the goals of designing an effective strategy to manage the problem assessing proportion and factor of death among preterm neonates are important. Methods An Institution-based cognitive research study was conducted at Aksum Referral and General specialty hospital, Northern Ethiopia. All pre-natal isolates that were allowed to enter into a high-growth union during childbirth. The data collected will be cleared and entered into the Epi-info version 7 and exported to STATA version 12 for further analyses. bivariable and multivariable analysis performed to determine the relationship between baseline and explanatory variables. Variables with a p-value of less than 0.05 will be the cut-off point. Result overall mortality rate was 22.2%, with 95% CI (17.9-26.9), of which 18 (23.4%) died within the 1 st 48 hrs and 59(76.6%) died the 1 st 7 days of life. Place of delivery(AOR:5.13(3.46-10.04)), Birth weight (AOR: 4.69 (2.29-9.63)), kangaroo mother care (AOR: 2.71 (1.20 - 6.11)), Hypoglycemia(AOR: 2.47 (1.30- 4.72)), Breastfeeding initiated (AOR: 6.35(4.62-11.98)), APGAR score(AOR: 8.63 (3.52 - 21.17)), Age (AOR: 4.19(2.86-12.46)) were predictor for the occurrence of preterm neonate death. Conclusion the proportion of preterm birth mortality was high. Provide quality care in maternity care, access the intensive care unit at maternity and professional health care institutions and take precautionary measures to prevent complications. And collaboration with the woreda health office improves breastfeeding and kangaroo mother care should be better.

Background

Globally, 35% of all prenatal problems are a major cause of national prenatal problems associated with the cost of living worldwide [1]. In developing regions, the likelihood of dying in childbirth is six times higher than in developed countries, eight times higher than in developed countries [2]. Premature birth can have serious health consequences for the child, and many of the surviving children, often at the expense of family and community, are respiratory problems, mental illness, and mental disorders neurological problems[3, 20]. Globally in 2016, 2.6 million newborns were killed. Of these, 99% of maternal deaths occur in low-income and middle-income countries. 39% were reported in South Asia, 39% in South Africa. Ethiopia is one of the five countries that have died in infants, children, Pakistan, Nigeria, and the Democratic Republic of the Congo [5]. May according to the 2016 Ethiopian demographic health survey, the death to all from childbirth is 29 out of 1000 births [6]. Studies reported that preterm neonatal mortality in Ethiopia there is regional variation 28.28% University of Gondar [7], Tigray region Northern Ethiopia 34% [8], and Jimma University Specialized Hospital, Ethiopia 34.9% [9], 47% rural Ethiopia[10], 63% Northern Ethiopia[11]. According to different kinds of literature, factors that are associated with preterm neonatal mortality include; Maternal Age, Residence, Place of delivery, Birth weight, Gestational age at birth, APGAR score, breastfeeding upon birth, kangaroo mother care (KMC), Antenatal care, Pregnancy complications, Labor and delivery complications and so on. [12–15, 5, 16–21, 7–11]. Proper management and the use of timely guidelines improve effectively after childbirth. The Sustainable

Development Goal of Child Deaths by 2030, it is possible to protect newborns, and children under 5 years of age, all of which are intended to reduce newborns mortality less than 12 death per 1000 live birth and death of under-five less than 25 death per 1000 live birth [5]. Therefore, this study aims to assess the proportion and factors of death among preterm neonates admitted to Aksum University specialized hospital NICU.

Methods

An Institution-based cognitive research study was conducted at Aksum Referral and General specialty hospital, Northern Ethiopia from June 2008 to May 2011 E.C. Aksum General specialty hospital offers higher education to more than seven million people in Ethiopia. The northwest country lies 1,024 km from Addis Ababa capital city Ethiopia and 246 km from Mekele capital city of Tigray Region. Neonatology is part of the department of pediatrics and provides services for the treatment of pediatrics patients. All pre-natal intensive care units will provide intensive care, especially for newborns, due to a large number of studies. All pre-natal isolates that were allowed to enter into a high-growth union during childbirth were excluded from the study as they were not readily available in the table room. The sample size was determined by comparing into the classification because the estimated risk of premature mortality was 28.8% [7] using a study by Gondar university using an accurate approach for population, Z-value of 1.96 at 95% confidence level and margin of error of 5% will be considered, but during the study, we made it easier to record 346 standardized checklists and used previously checked checklists. Death: Death from any cause of a preterm neonate. Preterm: neonates who delivered less than 37 completed weeks of gestation.

All necessary adult client information to collect from hospital records is recorded from a table with qualified Midwives. Data quality will be insured by closing data closures and reviewing available data and checking for inconsistencies, code errors, and range completion, accuracy, clarity, missing value and appropriate corrections. The data collected will be cleared and entered into the Epi-info version 7 and exported to STATA version 12 for further analyses. Descriptive and summary statistics will be carried out and presented using texts, graphs, and tables. bivariable and multivariable analysis performed to determine the relationship between baseline and explanatory variables. Variables with a p-value of less than 0.05 in the bivariable analysis will be the cut-off point for selection in the final analysis and the 95% CI is presented and interpreted.

Results

Socio-demographic and obstetric characteristics of mothers

There were 346 participants in this study and a 100% response rate. 346 pre-existing neurological data were included in the analysis. 58.67% of the mothers in the city. Two-thirds (63%) of the mothers were between the age of 29 and 34 years. More than 90% of neonates delivered from the health facility. More than 252 (72%) mothers have more than 4 times the regular maternity care (Table 1).

Table 1 Sociodemographic and obstetric characteristics of mothers of preterm neonates admitted in NICU at Aksum university comprehensive specialized Hospital from June 2008 to May 2011 E.C (n = 346)

Serial no	Characteristics		Frequency	Percentage (%)	Remark
1	Maternal age	<20 year	72	20.8	
		20-34 year	218	63	
		>=35	56	16.2	
2	Residence	Urban	203	58.67	
		Rural	143	41.33	
3	Place of delivery	Home	34	9.83	
		Health center	150	43.35	
		Hospital	162	46.82	
4	Type of pregnancy	Singleton	329	95.09	
		Multiple	17	4.91	
5	Frequency of ANC visit	<4 times	94	27.17	
		>=4 times	252	72.83	
6	Parity (number of births)	I	52	15.03	
		II-IV	272	78.61	
		≥ V	22	6.36	
7	Previous bad obstetrics history	Yes	34	9.83	
		No	312	90.17	
8	Onset of labor	Spontaneous	325	93.93	
		Induced	21	6.07	
9	Mode of delivery	SVD	332	95.95	
		CS	7	2.02	
		INSTRUMENTAL	7	2.02	
10	Cause of labor	Spontaneous	325	93.94	
		Induced	21	6.06	
11	History of PROM	Yes	54	15.61	
		No	292	84.39	
12	History of PIH	Yes	21	6.07	
		No	325	93.93	

Characteristics of the preterm neonates

Of the 346 preterm infants, 180(52%) were female and 6(1.73%) were less than 1 kg. Approximately 21(6.06%) were employed and 110(31.8%) seniors were cared for by the kangaroo mother. Of the 276 (79.8%) who started breastfeeding immediately. 56(16.18%) had an APGAR score of less than or equal to 6 (Table 2).

Table 2: Characteristics of preterm neonates admitted in NICU at Aksum University comprehensive specialized Hospital from June 2008 to May 2011 E.C (n = 346).

Serial no	Characteristics	Frequency	Percentage (%)	Remark
1	Sex of the neonate	Male	166	47.98
		Female	180	52.02
2	Weight for gestational age at birth	1.5-2.5	270	78.03
		1-1.5	70	20.23
		<1	6	1.73
3	Gestational age (weeks)	< 32	42	12.14
		32-35	116	33.53
		35-37	188	54.34
4	Was the newborn develop low oxygen saturation	Yes	74	21.39
		No	272	78.61
5	First minute APGAR score	<=6	56	16.18
		>=7	290	83.22
6	Fifth minute APGAR score	<=6	7	2.02
		>=7	339	97.98
7	Was the baby start breastfeed	Yes	276	79.77
		No	70	20.23
8	Treatment for hypoglycemia	Yes	28	8.09
		No	318	91.91
9	Maternal fever during labor	<36.5	104	30.06
		36.5-37.5	235	67.92
		>37.5	7	2.02
10	Newborn received kangaroo mother care	Yes	110	31.79
		No	236	68.21

The proportion of preterm neonatal death

In my study, the overall mortality rate was 22.2%, with 95% confidence interval (17.9-26.9), of which 18 (23.4%) died within the first 48 hours and 59(76.6%) died the first 7 days of life(newborn death). The median survival time was 20 days. The causes of death are numerous, but the majority causes are complications prematurity 54%, PNA 22.1%, sepsis 14.3% and other 9.1% (figure 1).

Associated factors of proportions of death for preterm neonates

In binary logistic regression: preterm neonates who were delivered from mothers above 35 years, neonates who were delivered at home, neonates who were delivered from mothers had a history of PROM and PIH, neonates delivered from mothers having less than 4 ANC follow up, neonates whose birth weight was between 1 to 1.5kg, neonates who were not received KMC, neonates who were treated for hypoglycemia, neonates who were not initiated breastfeed following delivery, neonates who were delivered from mothers who undergo induction and neonates whose APGAR score less than or equal to six were significantly associated with the death of preterm neonates

However, in the multi-variable analysis neonates whose birth weight was between 1 to 1.5kg, neonates who were not received KMC, neonates who were treated for hypoglycemia, neonates whose APGAR score

was less than or equal to six, preterm neonates who have not started breastfeeding, neonates who were delivered at home and neonates who were delivered from mothers above 35 years have remained statistically significant factors.

The odds of death among preterm neonates whose birth weight was between 1 to 1.5kg was Five times [AOR = 4.69; 95% CI (2.29-9.63)] more likely as compared to neonates whose birth weight was between 1.5-2.5k.g.

The risk of death among neonates who were delivered from mothers above the of 35 years 4 times [AOR = 4.19, 95% CI (2.86, 12.46)] more vulnerable than neonates who were delivered from mothers in the age group of between 20-34 years.

The odds of death for preterm neonates who had < 7 APGAR score at birth was nine times [AOR = 8.63; 95% CI (3.52 - 21.17))] risk than those who had ≥ 7 APGAR score at birth

Providing KMC for all preterm neonates reduces the odds of death by 96.5% as compared to neonates who do not provide KMC [AOR = 2.71; 95% CI (1.20 - 6.11)].

The odds of death for preterm neonates who were delivered at home were five times [AOR = 5.13, 95% CI (3.46, 10.04)] risk than neonates who were delivered at the hospital.

The risk of death for preterm neonates who was not started breastfeeding was six times [AOR = 6.35, 95% CI (4.62, 11.98)] risk than their counterparts.

The odds of death for preterm neonates who were treated for hypoglycemia was 2 times [AOR = 2.47; 95% CI (1.30- 4.715)] more vulnerable than those who were not treated (**Table 3**).

Table 3 Bivariate and Multivariate analysis of neonatal and obstetric factors predicting preterm birth outcomes.

*P-value <0.05, ** P-value <0.001, COR- crude odds ratio, AOR- Adjusted odds ratio

Discussion

Ongoing problems in developing country the recurrent problem are the lack of general information and general reports on the burden of preterm infants [22]. In contrast to developed communities where reliable statistics are available and provide specific information, it directs every decision regarding early childhood care. The mortality rate among preterm infants was 22.2% at maternity university General specialty hospital. The cause of death is not a single problem leading to death, with the major causes being the first 54.55%, PH 22.1%, and sepsis 14.3%. In several birth analyses, Trinidad and Tobago were 12.4% [15], Cameroon 15.7% [18] in different parts of Ethiopia [9–11].

The birth rate and preterm infant were significantly higher (22.25%). This may be due to the number of studies that have been exposed only to the study stations and to the most vulnerable conditions. In

Variables		Status		COR (CI)	AOR(CI)
		Survived	Died		
Age	<20	55(76.4%)	17(23.6%)	2.56(1.35-4.86)	1.31(0.43-3.94)
	20-34	197(90.4%)	21(9.6%)	1	1
	>35	17(30.4%)	39(69.6%)	11.04(6.47-18.85)	4.19(2.86-12.46)*
Place of delivery	home	7(16.3%)	36(83.7%)	7.05(4.17-11.92)*	5.13(3.46-10.04)*
	HC	128(90.8%)	13(9.2%)	0.7(0.41-1.29)	0.84(0.75-1.64)
	Hospital	134(82.7%)	28(17.3%)	1	1
PROM	Yes	19(35.2%)	35(64.8%)	6.496(4.13-10.21)	1.17 (0.67-2.01)
	No	250(85.6%)	42(14.4%)	1	1
PIH	Yes	10(47.6%)	11(52.4%)	4.25(2.112-7.60)	1.38 (0.77, 2.47)
	No	259(79.7%)	66(20.3%)	1	1
Frequency of ANC	<4	59(62.8%)	35(37.2%)	2.40(1.53-3.76)	1.13(0.65-1.99)
	>=4	210(83.3%)	42(16.7%)	1	1
Birth weight	1.5-2.5	240(88.6%)	31(11.4%)	1	1
	1-1.5	29(38.7%)	46(61.3%)	7.1(4.48-11.19)	4.69 (2.29-9.63)**
KMC	Yes	75(68.2%)	35(31.8%)	1	1
	No	194(82.2%)	42(17.8%)	3.52(1.33-7.81)	2.71 (1.20 - 6.11)*
Hypoglycemia	Yes	7(25%)	21(75%)	5.58(3.37-9.25)	2.47 (1.30- 4.72)*
	No	262(82.4%)	56(17.6%)	1	1
Breastfeeding initiated	Yes	248(89.9%)	28(10.1%)	1	1
	No	21(30%)	49(70%)	9.22(5.78-14.71)	6.35(4.62-11.98)*
Cause of labor	Spontaneous	262(80.6%)	63(19.4%)	1	1
	Induced	7(33.3%)	14(66.7%)	4.69(2.62-8.41)	3.08(0.83-11.52)
1 st minute APGAR score	<7	28(50%)	28(50%)	3.26(2.05-5.21)	8.63 (3.52 - 21.17)**
	>=7	241(83.1%)	49(16.9%)	1	1
5 th minute APGAR score	<7	0(0%)	7(100%)	4.8 (2.2 - 10.48)	1.28(0.55-3.06)
	>=7	269(79.4%)	70(20.6%)	1	1

contrast, the finding was below the level of research conducted in a different part of the world

To save the life and growth of children worldwide reported that preterm neonatal mortality was 36.15 and 29.3 percent respectively [12, 23], Jimma University special Hospital, Ethiopia 34.9% [9] Gondar University Hospital, Ethiopia 28.28%[7]. This may be due to the decrease in maternal mortality, increased health care services, pre-referral, community health search and use behavior, and access to trained healthcare providers.

Neonates whose birth weight was between 1-1.5 kg, neonates who were not received KMC, neonates who were

treated for hypoglycemia and neonates whose APGAR score was less than or equal to six have remained statistically significant factors.

The odd of death among preterm neonates whose birth weight was between 1-1.5 kg was Five times more likely as compared to neonates whose birth weight was between 1.5–2.5 kg. This result compares with studies conducted in Iran (36) and Nigeria (37).

Hypoglycemia is significantly associated with premature mortality in premature infants [AOR = 2.47; 95% CI 2.76 (1.30–4.71)]. This result is consistent with studies conducted at the university in Gondar. This is due to the possibility of adipose fat tissue, which serves as a source of glucose until they can feed on additional embryos.

The odds of death were increased by nine times for a neonate with the first minute APGAR score < 7 at birth as compared to the counterpart with. The results were obtained from the Iran[24], Nigeria[25], Tikur Anbesa specialized hospital, Ethiopia[26], University of Gondar hospital. Ethiopia[7].

This finding showed that the difference in mortality among older siblings decreased by 96.5% in kangaroo mother care [AOR = 2.71; 95% CI 2.41 (1.20–6.11)]. This is similar to study conduct at the University of Gondar. This may be KMC that prevents hypothermia by reducing the body surface area to the outside surface and helps easily accessing breast milk.

According to several studies in different areas, the risk of death among maternal-born 35 and above year peers is 4 times higher than that of 20 to 34 years. The world has reported that maternal mortality over 35 years is a prerequisite for premature deaths [12, 13, 15–17, 10, 11, 27, 28].

The mortality rate of premature infants born at home is five times as high as those at the same hospital supported by studies conducted in Johannesburg, South Africa [18, 28] and Jimma, Ethiopia [8].

The rate of death for premature infants who do not start breastfeeding is six times higher than their units. Therefore, the early onset of childbirth has been reported to be a preventive measure for premature death.

Conclusions

In this study the proportion of preterm birth mortality was high. The Median survival time was 20 days. In neonates whose birth weight was between 1-1.5kg, neonates who were not received KMC, neonates who were treated for hypoglycemia, neonates whose APGAR score was less than or equal to six, preterm neonates who have not started breastfeeding, neonates who were delivered at home and neonates who were delivered from mothers above the of 35 years remained statistically significant factors. Therefore, Aksum university specialized hospitals must provide quality care in maternity care, access the intensive care unit at maternity and professional health care institutions and take precautionary measures to prevent complications. The Aksum specialty Hospital in collaboration with the woreda health office should improve breastfeeding and kangaroo mother care. The main limitation of this study was a secondary source of information may not have some important variables that affect the output variables. It was a series-section study and may not reveal the cause and effect relationship.

Abbreviations

AOR- Adjusted Odd Ratio, APGAR- Appearance Pulse Grimace Activity Respiration, CI- Confidence Interval, COR- Crude Odd Ratio, EDHS- Ethiopian Demographic and Health Survey, GA- Gestational Age, KMC- Kangaroo Mother Care, NICU- Neonatal Intensive Care Unit, NGOs- Non-Governmental Organizations, NMR-Neonatal Mortality Rate, PIH- Pregnancy Induced Hypertension, PNA- Prenatal Asphyxia, PROM- Premature Rupture of Membrane, SPSS- Statically Package Software for Social science, WHO-World Health Organization

Declarations

Ethics approval and consent to participant

Ethical clearance was obtained from the Institutional Review Committee (IRC), College of Medicine and Health Sciences, University of Aksum. A permission letter was received from those administrative bodies of Aksum comprehensive specialized hospital, then proceeded with data collection.

Consent for publication

Not applicable

Availability of data and materials

All relevant data are within the manuscript and its Supporting Information files.

Computing interest

The authors declare that they have no competing interests.

Funding

"The authors received no specific funding for this work".

Authors' contributions

TG and SM designed the study, performed the statistical analysis, drafted the paper, data analysis and read and approved the final paper.

Acknowledgments

We are highly indebted to all participants of the study, supervisors of data collection and data collectors for their worthy efforts and participation in this study.

Authors' details

Sintayew Mamo: School of medicine, College of Health Sciences, Aksum University, Aksum, Ethiopia.
Email: sintiebdr@gmail.com

References

1. ASIA JABIRI HLW, AVELINA SEMIONO ETL: Prevalence and factors associated with neonatal sepsis among neonates in Temeke and Mwananyamala Hospitals in Dar es Salaam, Tanzania. *Tanzania Journal of Health Research* 2016, 18(4).
2. Haftom Gebrehiwot misgna hBG, mulugeta Molla Birhanu: Knowledge, Practice and associated factors of the essential newborn care at home among mothers in Gulomekada District, Eastern Tigray, Ethiopia, 2014. *BMC Pregnancy and Child Health*
3. ELF, *Preterm birth and the lungs*.
4. James N. Martin, M., FACOG Presiden, *Prenatal Care Is Important to Healthy Pregnancies*. American Congress of Obstetricians and Gynecologists, 2012.
5. TU: Child Mortality. *Child Mortal.* 2017, 3.
6. Central Statistical Agency: the DHS Program. Ethiopia Demographic and Health survey. 2016.
7. Yismaw, A.E. and Tarekegn, A.A., 2018. Proportion and factors of death among preterm neonates admitted to the University of Gondar comprehensive specialized hospital neonatal intensive care unit, Northwest Ethiopia. *BMC research notes*, 11(1), p.867.
8. Mengesha HG, Wuneh AD, Lerebo WT, Tekle TH. Survival of neonates and predictors of their mortality in the Tigray region, Northern Ethiopia: a prospective cohort study. *BMC Pregnancy Childbirth*. 2016;16(1):202.
9. Debelew GT, Afework MF, Yalew AW. Determinants and causes of neonatal mortality in Jimma zone, southwest Ethiopia: a multilevel analysis of prospective follow up study. *PLoS One*. 2014;9(9):e107184.
10. Weldearegawi B, Melaku YA, Abera SF, Ashebir Y, Haile F, Mulugeta A, et al. Infant mortality and causes of infant deaths in rural Ethiopia: a population-based cohort of 3684 births. *BMC public health*. 2015;15(1):770.
11. Mengesha HG, Lerebo WT, Kidanemariam A, Gebrezgiabher G, Berhane Y. Pre-term and post-term births: predictors and implications on neonatal mortality in Northern Ethiopia. *BMC Nurs*. 2016;15(1):48.
12. Lawn JE, Cousens S, Zupan J. 4 million neonatal deaths: When? Where? Why? *The Lancet*. 2005;365(9462):891-900.
13. Jehan I, Harris H, Salat S, Zeb A, Mobeen N, Pasha O, et al. Neonatal mortality, risk factors and causes: a prospective population-based cohort study in urban Pakistan. *Bulletin of the World Health*

- Organization. 2009;87(2):130-8.
14. Gizaw M, Molla M, Mekonnen W. Trends and risk factors for neonatal mortality in Butajira District, South Central Ethiopia,(1987-2008): a prospective cohort study. *BMC pregnancy and childbirth*. 2014;14(1):64.
 15. Risso SdP, Nascimento LFC. Risk factors for neonatal death in neonatal intensive care unit according to survival analysis. *Revista Brasileira de terapia intensiva*. 2010;22(1):19-26.
 16. Dhaded SM, Somannavar MS, Vernekar SS, Goudar SS, Mwenche M, Derman R, et al. Neonatal mortality and coverage of essential newborn interventions 2010-2013: a prospective, population-based study from low-middle income countries. *Reproductive health*. 2015;12(2): S6.
 17. Razeq NMA, Khader YS, Batiha AM. The incidence, risk factors, and mortality of preterm neonates: A prospective study from Jordan (2012-2013). *Turkish journal of obstetrics and gynecology*. 2017;14(1):28.
 18. Ballot DE, Chirwa TF, Cooper PA. Determinants of survival in very low birth weight neonates in a public sector hospital in Johannesburg. *BMC pediatrics*. 2010;10(1):30.
 19. Cupen K, Barran A, Singh V, Dialsingh I. Risk Factors Associated with Preterm Neonatal Mortality: A Case Study Using Data from Mt. Hope Women's Hospital in Trinidad and Tobago. *Children*. 2017;4(12):108.
 20. Ndombo PK, Ekei QM, Tochie JN, Temgoua MN, Angong FTE, Ntock FN, et al. A cohort analysis of neonatal hospital mortality rate and predictors of neonatal mortality in a sub-urban hospital of Cameroon. *Italian journal of pediatrics*. 2017;43(1):52.
 21. Mwaniki MK, Gatakaa HW, Mturi FN, Chesaro CR, Chuma JM, Peshu NM, et al. An increase in the burden of neonatal admissions to a rural district hospital in Kenya over 19 years. *BMC Public Health*. 2010;10(1):591.
 22. Howson C, Kinney M, Lawn J. Born too soon: the global action report on preterm birth. Geneva: World Health Organization. 2012:1-126.
 23. WHO U. Countdown to 2015: maternal, newborn and child survival. Decade Report (2000-2010): taking stock of maternal, newborn and child survival Geneva: WHO, UNICEF. 2010.
 24. Ladan H, Marzieh N, Behnaz M, Zahra N [Survival predictors of preterm neonates: Hospital-based study in Iran \(2010-2011\)](#)
 25. [Mustapha Bello, Simon Pius, Bello Abdullahi Ibrahim](#) Characteristics and predictors of outcome of care of preterm newborns in resource constraints setting, Maiduguri, Northeastern Nigeria 2019
 26. Tekleab AM, Amaru GM, Tefera YA. Reasons for admission and neonatal outcome in the neonatal care unit of a tertiary care hospital in Addis Ababa: a prospective study. *RESEARCH AND REPORTS IN NEONATOLOGY*. 2016;6:17-23.
 27. Selemani M, Mwanyangala MA, Mrema S, Shamte A, Kajungu D, Mkopi A, et al. The effect of mother's age and other related factors on neonatal survival associated with first and second birth in rural, Tanzania: evidence from Ifakara health and demographic surveillance system in rural Tanzania. *BMC pregnancy and childbirth*. 2014;14(1):240.

Figures

Pie chart shows proportions for the causes of death




Figure 1

cause of preterm neonatal death