Development of 105 SNP markers in endangered turtle species *Pelodiscus sinensis* using RAD-seq

Qiming Xie (✉ 2188149989@qq.com)

School of Animal Science, Anhui Agricultural University, Hefei, China

Fan Liu

Anhui Agricultural University

Jilong Zhang

Anhui Agricultural University

Xilei Li

Anhui Agricultural University

Tiantian Chen

Anhui Agricultural University

Guoxia Fang

Feixi County Animal Disease Prevention and Control Center

Rensheng Ma

Feixi County Animal Disease Prevention and Control Center

Shi-Ping Su

Anhui Agricultural University

Research Article

Keywords: *Pelodiscus sinensis*, RAD-seq, SNP, Conservation genetics

Posted Date: February 11th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-213566/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

In this study, Restriction site-associated DNA sequencing (RAD-seq) was used to develop single-nucleotide polymorphism (SNP) markers for *Pelodiscus sinensis*, which is a commercially important species in China. We collected 15 individuals from Anhui Province and RAD-seq data of 20 individuals from NCBI: PRJNA530350 to screen high quality SNP maker through genome-wide. A total of 105 high-quality SNP markers were identified. The population genetic statistics showed that the minor allele frequency varied from 0.0571 to 0.3382, and the observed heterozygosity and expected heterozygosity ranged from 0.1143 to 0.6765 and 0.1078 to 0.4477, respectively. These novel SNP markers will serve as a valuable genetic tool for future conservation and population genetics studies in *P. sinensis*.

Main Text

The Chinese soft-shelled turtle (*Pelodiscus sinensis*) is widely distributed in China, Taiwan, Korea, and North Vietnam and has been used as food and medicine for thousands of years in China (Haitao et al. 2008). *Pelodiscus sinensis* is regarded as one of the commercially most important aquatic reptile species worldwide and is also an important model organism for evolutionary and developmental biology owing to its soft shell (Cordeiro et al. 2020; Wang et al. 2013). However, in the past decades, the number of *P. sinensis* has markedly reduced owing to human overexploitation and habitat degeneration, and the species is now listed on the IUCN Red List. At present, in China, there are massive *P. sinensis* farms that meet the market demand; however, the animals exhibit several genetic issues such as inbreeding, fixation of deleterious alleles, and loss of genetic diversity. Development of reliable molecular markers is essential for understanding *P. sinensis* population diversity and genetic structure, which are the basis of farm management and genetic resource conservation.

Previous genetic studies of *P. sinensis* have focused on using a few mitochondrial DNA (mtDNA) (Xiong et al. 2019; Xiong et al. 2010) and microsatellite markers (Bu et al. 2014; Zhu et al. 2012). Compared with mtDNA and microsatellite markers, single nucleotide polymorphisms (SNPs) are more repeatable across laboratories, widespread in the genome, flexible for analyzing both neutral and adaptive genetic variation, and easy to acquire and apply using high-throughput sequencing technologies (Kruck et al. 2013). Restriction site-associated DNA sequencing (RAD-seq) can identify large polymorphic genetic markers across the genome in a single, straightforward, and cost-effective experiment (Davey et al. 2011). Presently, the knowledge about genome-wide germplasm resources of soft-shelled turtle remains limited, and genome-wide SNP markers are lacking.

In this study, we used 15 *P. sinensis* individuals collected from Anhui Province and used the RAD-seq data of 20 individuals from NCBI (PRJNA530350) (Liang et al. 2019). Muscle tissue samples from the 15 individuals were obtained and stored at -80°C . RAD library preparation and sequencing were performed by BGI Tech Solutions CO., Ltd. (1010 Room, Building No.7, BGI Parl, No.21, Hongan 3rd Street, Yantian District, Shenzhen, Guangdong Province, China, 518000). The raw read filtration criteria were as follows: remove (a) reads without barcodes, (b) reads without the enzyme cut site, (c) reads with the adapter, (d)

reads with $\geq 40\%$ low-quality bases (quality value ≤ 15), and (e) reads with $\geq 10\%$ unidentified bases. *SAMtools* of the Burrows-Wheeler Aligner (BWA) tool (Li and Durbin 2009) was used to map the filtered reads to the Chinese soft-shelled turtle reference genome sequence (GCF_000230535.1), and *bcftools* was used to call SNP variants. To improve the quality of SNPs, *vcftools* (Danecek et al. 2011) was used to filter the raw SNP set with the following criteria: (a) depth of loci ≤ 15 and ≥ 3 , (b) GQ of loci ≥ 30 , (c) shared by 95% sample, and (d) minor allele frequency ≥ 0.05 . Other complex variant events such as InDels were ignored. *PLINK* was used to exclude those SNP sites that failed the Hardy-Weinberg equilibrium (*HWE*) test with a *P* value of 0.000001 and calculate observed heterozygosity (*Ho*), expected heterozygosity (*He*), and minor allele frequency (*MAF*). *bedtools* (Quinlan and Hall 2010) was used to extract a 300 bp flanking sequence from the reference genome according to the SNP sites, and *batchprimer3* (You et al. 2008) was used to design primers. Full details of the methods in this study can be found in supplementary material 1.

After quality control, a total of 105 high-quality SNP markers were identified from 10952494 raw SNP sites. The population genetic statistics for each SNP were analyzed (Table 1), and primers were designed for future utilization. The observed heterozygosity and expected heterozygosity varied from 0.1143 to 0.6765 and 0.1078 to 0.4477, respectively. The minor allele frequency ranged from 0.0571 to 0.3382. The detailed information of the 105 SNP markers can be found in supplementary material 2. These novel SNP markers could be a valuable tool for investigating the genetic diversity and population structure of *P. sinensis*, which could facilitate the improvement of farm management and germplasm resource conservation.

Declarations

Funding

This work was supported by the Hefei independent innovation policy “loan to supplement” project (No. J2019G37).

Conflict of Interest

The authors declare that they have no conflict of interest.

Ethics Approval

This study was followed by the ethical standard of China (GB/T 35892-2018).

Consent to participate

Not applicable.

Consent for publication

Not applicable.

Data Availability

The RAD-seq data are openly available in NCBI at <https://www.ncbi.nlm.nih.gov/sra>, PRJNA680995.

Code Availability

Not applicable.

Supplementary Information

The detailed process of data analysis and genetic information of 105 SNP markers can be found in Supplementary material 1 and 2.

ESM Competing Interests

The authors declare that they have no conflict of interest.

Author Contributions

Qiming Xie and Shiping Su conceived the presented idea. Qiming Xie and Fan Liu carried out the experiment. Qiming Xie developed the theory and performed the computations. Jilong Zhang and Xilei Li verified the analytical methods. Qiming Xie wrote the manuscript with support from Fan Liu, Jilong Zhang, and Xilei Li. Guoxia Fang and Rensheng Ma provided the experimental samples. Shiping Su supervised the findings of this work. All authors discussed the results and contributed to the final manuscript.

Acknowledgments

We thank Yang Li for providing computer equipment.

References

- Bu X, Liu L, Nie L (2014) Genetic diversity and population differentiation of the Chinese soft-shelled turtle (*Pelodiscus sinensis*) in three geographical populations. *Biochem Syst Ecol* 54:279-284
doi:10.1016/j.bse.2014.02.022
- Cordeiro IR, Yu R, Tanaka M (2020) Regulation of the limb shape during the development of the Chinese softshell turtles. *Evol Dev* 22:451-462 doi:10.1111/ede.12352
- Danecek P et al. (2011) The variant call format and VCFtools. *Bioinformatics* 27:2156-2158
doi:10.1093/bioinformatics/btr330
- Davey JW, Hohenlohe PA, Etter PD, Boone JQ, Catchen JM, Blaxter ML (2011) Genome-wide genetic marker discovery and genotyping using next-generation sequencing. *Nature Reviews Genetics* 12:499-510
doi:10.1038/nrg3012

- Haitao S, Parham JF, Zhiyong F, Meiling H, Feng Y (2008) Evidence for the massive scale of turtle farming in China. *Oryx* 42:147-150 doi:10.1017/S0030605308000562
- Kruck NC, Innes DI, Ovenden JR (2013) New SNPs for population genetic analysis reveal possible cryptic speciation of eastern Australian sea mullet (*Mugil cephalus*) *Molecular Ecology Resources* 13:715-725 doi:10.1111/1755-0998.12112
- Li H, Durbin R (2009) Fast and accurate short read alignment with Burrows-Wheeler transform *Bioinformatics* 25:1754-1760 doi:10.1093/bioinformatics/btp324
- Liang HW, Wang LH, Sha H, Zou GW (2019) Development and Validation of Sex-Specific Markers in *Pelodiscus Sinensis* Using Restriction Site-Associated DNA Sequencing *Genes* 10 doi:10.3390/genes10040302
- Quinlan AR, Hall IM (2010) BEDTools: a flexible suite of utilities for comparing genomic features *Bioinformatics* 26:841-842 doi:10.1093/bioinformatics/btq033
- Wang Z et al. (2013) The draft genomes of soft-shell turtle and green sea turtle yield insights into the development and evolution of the turtle-specific body plan *Nature Genetics* 45:701-+ doi:10.1038/ng.2615
- Xiong G, Wang X-Q, Zhou X-W, Zeng D, Chen Z-N, Wang P, Kang L (2019) Genetic variation in the Chinese soft-shell turtles (*Pelodiscus spp.*) revealed by sequences of mitochondrial Cytb gene *Mitochondrial DNA Part A* 30:874-879 doi:10.1080/24701394.2019.1693551
- Xiong L, Nie L, Li X, Liu X (2010) Comparison research and phylogenetic implications of mitochondrial control regions in four soft-shelled turtles of *Trionychia* (*Reptilia*, *Testudinata*) *Genes & Genomics* 32:291-298 doi:10.1007/s13258-010-0015-8
- You FM et al. (2008) BatchPrimer3: a high throughput web application for PCR and sequencing primer design *BMC Bioinformatics* 9:253 doi:10.1186/1471-2105-9-253
- Zhu L, Li Z, Li J, Feng X, Xie N (2012) Genetic Diversity of Farmed Chinese Soft-Shelled Turtle (*Pelodiscus sinensis*) Evaluated from Microsatellite Analysis *Journal of Animal and Veterinary Advances* 11:1217-1222

Tables

Table 1 Summary statistics of 105 single nucleotide polymorphism (SNP) markers of *Pelodiscus sinensis*

Position	REF	ALT	<i>MAF</i>	<i>Ho</i>	<i>He</i>
NW_005852246.1:55	A	G	0.2794	0.5588	0.4027
NW_005866747.1:181	T	C	0.1176	0.2353	0.2076
NW_005857380.1:3906	G	A	0.3382	0.6765	0.4477
NW_005854040.1:9429	A	T	0.0857	0.1714	0.1567
NW_005858158.1:32026	T	C	0.1029	0.2059	0.1847
NW_005853731.1:38813	C	T	0.0735	0.1471	0.1362
NW_005851636.1:42529	T	C	0.0882	0.1765	0.1609
NW_005854351.1:55917	A	C	0.0882	0.1765	0.1609
NW_005854898.1:73656	C	T	0.0571	0.1143	0.1078
NW_005855443.1:82027	A	C	0.0571	0.1143	0.1078
NW_005853897.1:98550	C	T	0.0882	0.1765	0.1609
NW_005851906.1:111998	G	A	0.0588	0.1176	0.1107
NW_005856908.1:129462	C	T	0.0735	0.1471	0.1362
NW_005856362.1:129953	A	G	0.1176	0.2353	0.2076
NW_005856316.1:164154	G	A	0.1471	0.2941	0.2509
NW_005858876.1:203033	G	A	0.0588	0.1176	0.1107
NW_005856075.1:210584	C	T	0.1176	0.2353	0.2076
NW_005858997.1:239774	A	G	0.0735	0.1471	0.1362
NW_005857910.1:242076	A	C	0.0588	0.1176	0.1107
NW_005854296.1:252072	C	T	0.1029	0.2059	0.1847
NW_005857744.1:297199	G	C	0.0735	0.1471	0.1362
NW_005856504.1:319845	G	A	0.1618	0.3235	0.2712
NW_005857680.1:323433	T	G	0.1176	0.2353	0.2076
NW_005855650.1:384770	G	A	0.0857	0.1714	0.1567
NW_005857694.1:407992	C	A	0.1029	0.2059	0.1847
NW_005854923.1:416335	T	G	0.0588	0.1176	0.1107
NW_005854923.1:416341	C	G	0.0588	0.1176	0.1107
NW_005857006.1:421367	A	G	0.1765	0.3529	0.2907

NW_005858887.1:475260	A	G	0.0588	0.1176	0.1107
NW_005854747.1:476067	C	T	0.0735	0.1471	0.1362
NW_005871043.1:523453	G	T	0.1029	0.2059	0.1847
NW_005871043.1:523454	A	C	0.1029	0.2059	0.1847
NW_005856824.1:546807	C	T	0.0588	0.1176	0.1107
NW_005854417.1:562053	A	G	0.0571	0.1143	0.1078
NW_005856333.1:562648	T	A	0.0735	0.1471	0.1362
NW_005853643.1:608336	G	T	0.0588	0.1176	0.1107
NW_005855688.1:615542	G	T	0.1618	0.3235	0.2712
NW_005854399.1:618147	T	C	0.2714	0.5429	0.3955
NW_005854399.1:618222	A	G	0.2794	0.5588	0.4027
NW_005854399.1:618253	T	G	0.2794	0.5588	0.4027
NW_005855929.1:624215	T	G	0.0571	0.1143	0.1078
NW_005855694.1:659040	T	C	0.0882	0.1765	0.1609
NW_005858300.1:719759	C	T	0.0882	0.1765	0.1609
NW_005855698.1:795091	A	T	0.0588	0.1176	0.1107
NW_005857290.1:823432	A	T	0.2059	0.4118	0.3270
NW_005853196.1:884143	G	T	0.0882	0.1765	0.1609
NW_005854405.1:896447	T	C	0.0882	0.1765	0.1609
NW_005853644.1:917097	G	A	0.0735	0.1471	0.1362
NW_005855218.1:1025924	T	C	0.0588	0.1176	0.1107
NW_005851747.1:1115210	G	T	0.0588	0.1176	0.1107
NW_005854428.1:1223918	T	C	0.2647	0.5294	0.3893
NW_005853167.1:1227139	A	T	0.1618	0.3235	0.2712
NW_005855792.1:1262551	G	A	0.0588	0.1176	0.1107
NW_005856802.1:1268217	A	G	0.0588	0.1176	0.1107
NW_005871004.1:1270305	T	C	0.1912	0.3824	0.3093
NW_005852159.1:1315537	A	G	0.1618	0.3235	0.2712
NW_005856823.1:1330905	C	A	0.0588	0.1176	0.1107

NW_005854781.1:1427409	T	A	0.1471	0.2941	0.2509
NW_005855476.1:1508482	G	A	0.1176	0.2353	0.2076
NW_005858753.1:1620758	G	A	0.0882	0.1765	0.1609
NW_005859202.1:1628000	G	A	0.1029	0.2059	0.1847
NW_005871004.1:1630732	T	G	0.0588	0.1176	0.1107
NW_005855236.1:1678320	C	T	0.0735	0.1471	0.1362
NW_005853122.1:1680360	C	G	0.0735	0.1471	0.1362
NW_005852052.1:1813181	G	A	0.2059	0.4118	0.3270
NW_005852052.1:1813185	C	T	0.2059	0.4118	0.3270
NW_005857440.1:1876366	G	A	0.1912	0.3824	0.3093
NW_005853738.1:1881846	C	T	0.1286	0.2571	0.2241
NW_005855397.1:1897760	T	C	0.0588	0.1176	0.1107
NW_005855397.1:1897783	T	C	0.0735	0.1471	0.1362
NW_005851563.1:2179842	T	C	0.1029	0.2059	0.1847
NW_005856064.1:2201529	G	A	0.0735	0.1471	0.1362
NW_005856064.1:2201558	C	T	0.0735	0.1471	0.1362
NW_005856064.1:2201567	A	G	0.0735	0.1471	0.1362
NW_005858084.1:2204830	C	T	0.0571	0.1143	0.1078
NW_005857732.1:2240113	A	C	0.0571	0.1143	0.1078
NW_005857401.1:2240213	G	A	0.0735	0.1471	0.1362
NW_005852467.1:2246898	G	A	0.0882	0.1765	0.1609
NW_005852467.1:2246902	T	C	0.0882	0.1765	0.1609
NW_005853176.1:2506285	G	A	0.1324	0.2647	0.2297
NW_005852769.1:2548930	T	A	0.1324	0.2647	0.2297
NW_005859159.1:2600139	A	T	0.0588	0.1176	0.1107
NW_005854634.1:2883497	A	G	0.1618	0.3235	0.2712
NW_005853523.1:2999985	C	T	0.0588	0.1176	0.1107
NW_005851929.1:3140999	C	T	0.3382	0.6765	0.4477
NW_005855792.1:3166874	C	G	0.0588	0.1176	0.1107

NW_005854896.1:3706567	C	T	0.0882	0.1765	0.1609
NW_005851925.1:3737622	C	A	0.0588	0.1176	0.1107
NW_005853657.1:3836504	A	T	0.0588	0.1176	0.1107
NW_005855489.1:4002735	A	G	0.0882	0.1765	0.1609
NW_005855637.1:4174003	T	C	0.1176	0.2353	0.2076
NW_005871043.1:4270476	G	C	0.0882	0.1765	0.1609
NW_005871043.1:4270505	C	T	0.0882	0.1765	0.1609
NW_005858987.1:4749868	G	T	0.0571	0.1143	0.1078
NW_005858573.1:5726004	T	C	0.0588	0.1176	0.1107
NW_005858573.1:6235628	G	A	0.1176	0.2353	0.2076
NW_005858573.1:6235633	A	G	0.1176	0.2353	0.2076
NW_005855477.1:7581107	A	G	0.1324	0.2647	0.2297
NW_005853697.1:8051021	C	T	0.0588	0.1176	0.1107
NW_005871009.1:8713326	C	T	0.0588	0.1176	0.1107
NW_005853106.1:8919331	C	T	0.0588	0.1176	0.1107
NW_005854622.1:9141592	G	A	0.0714	0.1429	0.1327
NW_005854622.1:9141593	C	T	0.0571	0.1143	0.1078
NW_005871025.1:9733061	T	A	0.0588	0.1176	0.1107
NW_005855477.1:11005396	G	A	0.0882	0.1765	0.1609

Note: REF, major allele; ALT, minor allele; *MAF*, minor allele frequency; *Ho*, observed heterozygosity; *He*, expected heterozygosity.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Supplementarymaterials1.docx](#)
- [Supplementarymaterials2.xlsx](#)