Relationship Between Oxygen Radical Absorbance Capacity (ORAC) Index, Body Composition and Blood Biochemical Markers in Overweight /Obese Compared to Normal Weight Subjects

Zahra Madani

Islamic Azad University Science and Research Branch

Maryam Moussavi Javardi

Islamic Azad University Science and Research Branch

Abolghassem Djazayeri

Tehran University of Medical Sciences, School of Nutrition and Dietetics

Majid Karandish

Nutrition and Metabolic Disease Research Center, Ahvaz Jundishapour University of Medical Sciences, Ahvaz

Ariyo Movahedi (✉ amm35@mail.aub.edu)

Islamic Azad University Science and Research Branch <https://orcid.org/0000-0003-3017-8945>

Research

Keywords: Oxygen Radical Absorption Capacity (ORAC), Lipid Profile, Fasting blood Glucose, Obesity, Overweight

Posted Date: April 23rd, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-21395/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background and aim: Obesity is an important preventable disease, which promotes the development of multiple chronic disorders by altering several factors including oxidative stress. Dietary antioxidants protect the body against oxidative stress. The purpose of this study was to evaluate the potential association of oxygen radical absorbance capacity (ORAC) index with obesity and its associated biochemical factors in adults.

Methods: The present study was a cross-sectional study of 157 adult females and males in two groups of normal weight and overweight /obesity. General characteristics were registered and a 147-item food frequency questionnaire was completed for every participant. Thereafter, anthropometric measurements were conducted. Biochemical indices were obtained from the Electronic Health Clinic Database of science and Research branch of Islamic Azad University (SRBIAU) of Tehran, Iran. The amount of the dietary ORAC index was estimated using the data provided by the USDA Foods Table.

Results: The results showed that there was a significant difference between the normal and overweight/obese groups in terms of weight, body mass index (BMI), waist to hip ratio (WHR), visceral fat level, body fat mass and body fat percentage ($P=0.0001$). Participants with normal weight consumed more fruits and vegetables than individuals in another group. Also, the dietary ORAC index in normal individuals was higher than the case group, but there was not significant difference ($P=0.222$). There was also a significant inverse correlation between dietary ORAC and BMI and WHR, whereas a significant positive correlation between dietary ORAC and plasma levels of HDL was observed ($P<0.05$).

Conclusion: Our findings suggest that consumption of antioxidants from dietary sources is inversely associated with BMI and other measurements of adiposity, fasting blood glucose (FBG), triglyceride (TG), total Cholesterol (TC) and LDL in both groups. It seems that following a diet rich in antioxidants can counteract obesity and its associated comorbidities.

Introduction

Obesity is a complex and multifactorial disease with a high prevalence around the world [1,2]. The worldwide prevalence of overweight and obesity has doubled since 1980 and about a third of the world's population is classified as overweight or obese [2]. At least 2.8 million deaths worldwide are related to obesity and overweight and the prevalence of the disease is increasing annually. The disease results from complex phenotypes associated with excess fat in the body that affect other metabolic factors in addition to body size [1]. Obesity is associated with numerous disorders, including diabetes, lipid profile abnormalities, liver disease, immune dysfunction and oxidative stress [3–5].

Oxidative stress is defined by excessive endogenous oxidative species; especially reactive oxygen species (ROS), such as superoxide, hydrogen peroxide and hydroxyl radical ions which damage cells and manipulate signal pathways [6,7]. Fat accumulation stimulates lipid peroxidation and damage caused by free fatty acids and cytokines, such as TNF- α , leading to the exacerbation of the oxidative damage [8].

Subjects with obesity and metabolic diseases have also shown lower levels of antioxidant activity [9]; additionally, studies have shown that obesity increases the risk of metabolic syndrome in obese and overweight people due to increased oxidative stress [5,10,11]. On the other hand, oxidative stress disrupts insulin secretion from pancreatic beta cells [12] and induces insulin resistance [13], as well as impaired glucose transport to the muscles and the adipose tissue [12]. Hyperglycemia produces alterations in various metabolic and cellular functions; including dyslipidemia, hypertension, endothelial dysfunction, and oxidative stress [14,15].

In recent years, antioxidants have been used to overcome the harmful effects of ROSs. Natural antioxidants found in healthy foods, include vitamins E and C, coenzyme Q10, alpha-lipoic acid, lycopene, and polyphenols [16]. The results of the studies indicate that there is an inverse relationship between body fat, central fat and antioxidant capacity [17].

Various methods are available for measuring the dietary total antioxidant capacity (TAC) [18,19]. According to the United States Department of Agriculture (USDA) database, the oxygen radical absorbance capacity (ORAC) index is an indicator that measures the antioxidant power of foods and other chemicals in the laboratory, and it is a preferable method because of its biological relevance to the *in vivo* antioxidant efficacy [20]. Recent studies have shown that fruits and vegetables with high ORAC, such as spinach and blueberries, prevent chronic diseases [20–22].

Since oxidative stress, directly and indirectly, causes many diseases, including obesity and metabolic syndrome, the purpose of the present study was to investigate the relationship between the dietary ORAC index and fasting blood glucose and lipid profile in overweight or obese subjects compared to normal weight subjects.

Methods

Study Type and Participants

This study was a cross-sectional descriptive-analytic case-control study. The subjects were normal weight and overweight/obese adults who were randomly selected among the students and staff of Science and Research branch of Islamic Azad University (SRBIAU) of Tehran. All the basic required information, including BMI and the latest blood test, was available in the University Electronic Health Clinic Database. After choosing possible eligible samples from the main data bank, some of the staff members were invited to volunteer in this study. 70 individuals in each group were considered to participate in this study who has BMI > 18.5 and age ranged between 18–65 years. All volunteers should not be under lipid and glucose control drug or be pregnant. All as well as unwillingness to cooperate during the study indicated by leaving more than 50% of the questions in the food frequency questionnaire (FFQ) unanswered.

Study implementation

This study was approved by the ethical Iran National Committee for Ethics in Biomedical Research under code IR.IAU.SRB.REC.1396.66. All the eligible volunteers were informed about the details of the study and written consent were taken. On the day of referral to the SRBIAU Clinic, information about demographic characteristics was obtained through face-to-face interviews using validated questionnaires.

Anthropometric measurements, including weight and body composition, were conducted using the bioelectric impedance analyzer (INBODY, Model 270, South Korea); height was measured using a digital freestanding stadiometer (BSM-170, InBody Co. Ltd, South Korea). Dietary assessment was also carried on using a validated semi-quantitative FFQ containing 148 food items [23].

Foods' ORAC measurement

Dietary antioxidant consumption was estimated based on the ORAC index of selected foods reported by the Nutrient Data Laboratory of USDA, and expressed as μmol of Trolox Equivalents per 100 grams of foods ($\mu\text{molTE}/100\text{ g}$) [20].

Statistical analysis method

Independent T-test or Mann-Whitney test (in the case of nonparametric data) were used to compare the mean of quantitative outcomes between the two groups. Correlation models were used to find out the relationship of the variables of the study. SPSS software version 25 was used for data analysis and P-value < 0.05 was considered as statistically significant.

Results

In the present study, the relationship between dietary ORAC index with lipid profile and fasting blood glucose in obese/overweight population and normal weight population (mean age of 38.73 ± 9.65 years in both groups) was examined. After calculating the sample size and simple random sampling, 180 adults, 90 overweight/obese and 90 with normal weight, were recruited based on the inclusion and exclusion criteria. Amongst these, 23 volunteers were excluded because of incomplete questionnaires (more than half of the items were not completed). Finally, the study was conducted with 71 normal weight and 86 overweight or obese subjects.

Basic information of two groups is presented in Table 1. As shown in Table 1, participants with normal weight had a lower visceral fat mass (five kilograms; $P < 0.01$) and a lower body fat percentage (1%; $P < 0.01$). Waist to hip ratio (WHR) was significantly lower in the normal weight group as compared to overweight/obese group ($P < 0.01$). As expected, fasting blood glucose (FBG), total cholesterol (TC), triglyceride (TG), and low-density lipoprotein (LDL) were higher in overweight/obese volunteers than normal volunteers were, whereas HDL was greater in the normal weight group; these differences were not significant.

Table 1
Basic information of two groups

Variable	Normal weight	Overweight/obese	P-Value
N	71	86	
Sex (M/F)	60/11	67/19	0.731
Marital status (S/M)	55/16	50/36	0.648
Age (year)	38.90 ± 10.976	38.60 ± 9.394	0.854
Anthropometric indices, mean ± SD			
Weight (Kg)	73.45 ± 10.66	90.05 ± 13.22	0.0001*
Height (Cm)	172.54 ± 7.58	172.20 ± 8.33	0.789
BMI (Kg/m ²)	24.57 ± 2.32	30.28 ± 3.16	0.0001*
WHR	0.090 ± 0.04	0.95 ± 0.06	0.0001*
Visceral fat	7.55 ± 2.10	12.30 ± 3.59	0.0001*
Body fat mass (Kg)	19.76 ± 10.76	28.77 ± 7.48	0.0001*
Body fat percentag (%)	2.14 ± 0.78	3.14 ± 2.77	0.004*
Biochemical parameters, mean ± SD			
FBG(mg/dl)	89.01 ± 10.73	92.09 ± 16.33	0.175
TG(mg/dl)	152.52 ± 85.38	164.99 ± 84.67	0.362
TC (mg/dl)	172.17 ± 35.30	175.06 ± 29.90	0.580
LDL (mg/dl)	98.44 ± 29.71	100.22 ± 25.75	0.687
HDL (mg/dl)	44.51 ± 8.41	42.40 ± 9.67	0.151
Abbreviations: BMI, Body mass index; WHR, Waist to hip ratio; F, Female; M, Male; FBG, Fasting blood glucose ; TG, Triglyceride; TC, Total Cholesterol; LDL, low-density lipoproteins; HDL, high-density lipoproteins.			
* $P < 0.05$			

Dietary data analyses are presented in Table 2. The consumption of vegetable, fruits, vitamin C, and vitamin E was higher in the normal weight group than the other group; the difference was only significant in fruit consumption ($P = 0.035$). Moreover, the dietary ORAC index was higher in normal weight volunteers than the overweight/obese group; the difference, though, was not significant ($P = 0.222$).

Table 2
Comparison of dietary intake and ORAC in two groups of normal weight and overweight/obese

Variable	Normal weight (n = 71)	Overweight/obese (n = 86)	P-Value
Vegetable	14.116 ± 1.713	13.529 ± 1.872	0.817
Fruits	18.943 ± 2.515	15.781 ± 2.302	0.035*
Vitamin.C	140.60 ± 87.96	123.07 ± 74.85	0.081
Vitamin.E (alpha.tocopherol)	11.98 ± 10.42	10.45 ± 9.44	0.064
ORAC	46002.65 ± 14651.20	17747.27 ± 8049.51	0.222
Abbreviations: ORAC, Oxygen radical absorbance capacity.			
*P < 0.05			

The correlation between the dietary ORAC index with anthropometric indices and biochemical parameters in the two groups are presented in Table 3. According to Table 3, ORAC index was inversely correlated with all anthropometric indices and biochemical parameters except for HDL. Significant correlation between ORAC with BMI, WHR and HDL was observed (P < 0.05).

Table 3

Correlation coefficient between anthropometric indices and blood biochemical markers with dietary ORAC index

Variable	Normal weight (n = 71)		Overweight/obese (n = 86)		Total (n = 157)	
	CC	P-Value	CC	P-Value	CC	P-value
BMI (Kg/m ²)	-0.768	0.047*	-0.876	0.010*	-0.831	0.036*
WHR	-0.796	0.042*	-0.869	0.023*	-0.680	0.043*
Body fat mass (Kg)	-0.188	0.117	-0.196	0.068	-0.035	0.661
Body fat percentage (%)	-0.083	0.489	-0.201	0.062	-0.012	0.879
Visceral fat	-0.094	0.435	-0.172	0.111	-0.071	0.377
FBG (mg/dl)	-0.041	0.734	-0.008	0.938	-0.033	0.677
TG (mg/dl)	-0.031	0.796	-0.023	0.829	-0.020	0.802
TC (mg/dl)	-0.204	0.087	-0.066	0.544	-0.076	0.343
LDL (mg/dl)	-0.194	0.105	-0.008	0.944	-0.089	0.266
HDL (mg/dl)	0.773	0.049*	0.948	0.012*	0.890	0.034*
Abbreviations: BMI, Body mass index; WHR, Waist to hip ratio; FBG, Fasting blood glucose;						
TG, Triglyceride; TC, Total Cholesterol; LDL, low-density lipoproteins; HDL, high-density lipoproteins; CC, Correlation coefficient						
* $P < 0.05$						

Discussion

In the present study, we were able to show that the dietary ORAC index was higher in normal weight volunteers as compared to the overweight/obese group. In addition, there was an inverse relationship between the dietary ORAC index and anthropometric indices, FBG, TG, TC and LDL. To the best of our knowledge, this is the first study investigating the association between dietary ORAC index, body composition and plasma biochemical markers in both overweight/obese and normal weight groups. Previous studies have shown an inverse relationship between the consumption of dietary antioxidants and several outcomes such as stroke [24], hypertension [25], myocardial infarction [26], cancer [27] and mortality [28].

In the present study, the dietary ORAC index was inversely correlated with BMI. Excessive fat accumulation can lead to obesity [29]. With increased obesity, the extra stored fat increases the production of oxygen radicals and inflammatory factors, as well as cholesterol and blood lipids [30].

Excess calories inhibit the production of adiponectin [31]. In the short time, the body tries to modify these damaging anomalies, but in the long run the immune system's ability to function is attenuated and oxidative stress can damage cells. Additionally, increased secretion of leptin usually accompanied by obesity leads to activation of T-lymphocytes leading to the production of proinflammatory factors, such as interleukin-2 and TNF- α which in turn can cause inflammation and oxidative stress in the body [32]. The body responds to these changes by using natural antioxidants, such as vitamins E and C. If the body is unable to eliminate these factors, excessive accumulation of oxidants can lead to impaired protein and fat function, cellular dysfunction and, ultimately, to inflammation [5,33,34]. Obesity-induced inflammation and oxidative stress increase insulin resistance[6] by reducing the expression of insulin receptor substrate-1 (IRS-1) [35]. As a result, promoting the body's antioxidant capacity can greatly prevent the complications of diabetes [36].

Consuming antioxidant-rich foods with a high capacity to absorb oxygen radicals will give the immune system a chance to counteract inflammation[37]. Despite the type of the present study as a short study, similar to the findings in the current study, the higher dietary ORAC index has been an indication of higher quality of the diet and was related to higher consumption of fruits and vegetables. The favorable effects of these antioxidant-rich foods on improvement of fasting blood glucose, lipid profile and obesity have been investigated in some pre-clinical and clinical studies[38–42]. In recent years, chemical antioxidants, such as “phytochemicals”, are suggested as plausible candidates in the prevention and/or treatment of specific diseases, including obesity and obesity-related metabolic disorders [43].

Parallel with our findings, many studies have shown that the dietary antioxidant capacity of volunteers with normal weight is higher as compared to overweight/obese ones. In a study, Hermsdorff et al demonstrated that dietary antioxidant capacity was negatively and significantly correlated with waist circumference (WC) [17]. Also another study showed that high WHR is linked with increased oxidative stress [44]. Similarly, an increase in dietary antioxidant was shown to reduce the incidence of abdominal obesity [45]. In line with our findings, previous studies showed that obesity is related with unhealthy diet, which is full of carbohydrate, fat, simple sugars and sweet beverages, whereas a protective effect against obesity is observed in the consumption of a diet rich in fruits and vegetables [46,47] and a diet high in vegetables and fruits has been positively related to the consumption of antioxidant vitamins [47].

The present study showed that there was an inverse association between the dietary ORAC index with FBG, TG, LDL and TC, whereas a positive correlation between ORAC and HDL was observed. As well as, Many studies have supported these findings [48–50]. Some studies have also reported that antioxidant capacity in obese and overweight volunteers is promoted which was postulated to be due to immune system's trying to override the present inflammation [51,52].

Conclusion

Our findings suggest that consumption of dietary antioxidant was inversely associated with BMI and other anthropometric measurements, FBG, TG, LDL and TC in both groups. It seems that following a diet

rich in ORAC can counteract obesity and its associated comorbidities.

Declaration

Ethical Approval & Consent to participate

This study was approved by the ethical Iran National Committee for Ethics in Biomedical Research under code IR.IAU.SRB.REC.1396.66

Consent for publication

All the eligible volunteers were informed about the details of the study & written consent were taken.

Availability of supporting data

Not applicable

Competing interests

Authors declare that there is no competing interests what so ever.

Funding

Not applicable

Authors' contributions

We certify that we have participated sufficiently in the intellectual content, conception & design of this work or the analysis & interpretation of the data, as well as the writing of the manuscript, to take public responsibility for it & have agreed to have our name listed as a contributor.

Acknowledgements

Not applicable. We have not received substantial contributions from non-contributors & no contributor has been omitted.

Authors' information

1. Zahra Madani, Master in Public Health Nutrition, Department of Nutrition, Science and Research Branch, Islamic Azad University, Tehran, Iran 2. Maryam Moussavi Javardi, Master in Public Health Nutrition, Department of Nutrition, Science and Research Branch, Islamic Azad University, Tehran, Iran 3. Abolghassem Djazayeri, PhD. in Nutritional Science, Department of Community Nutrition, School of Nutritional Sciences and Dietetics, Tehran University of Medical Sciences, Tehran, Iran 4. Prof. Majid Karandish, PhD. in Nutritional Science, Nutrition and Metabolic Diseases Research Center, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran 5. Dr. Ariyo Movahedi*, PhD. in Clinical Nutrition, Department of Nutrition, Science and Research Branch, Islamic Azad University, Tehran, Iran.

Competing interests:

Authors declare that there is no competing interests what so ever.

References

1. 10 facts on obesity. World Heal. Organ. [Internet]. World Health Organization; 2017; Available from: <http://www.who.int/features/factfiles/obesity/en/>
2. Chooi YC, Ding C, Magkos F. The epidemiology of obesity. *Metabolism*. W.B. Saunders; 2019;92:6–10.
3. Jung UJ, Choi MS. Obesity and its metabolic complications: The role of adipokines and the relationship between obesity, inflammation, insulin resistance, dyslipidemia and nonalcoholic fatty liver disease. *Int. J. Mol. Sci.* MDPI AG; 2014. p. 6184–223.
4. Uranga RM, Keller JN. The Complex Interactions Between Obesity, Metabolism and the Brain. *Front. Neurosci.* [Internet]. Frontiers Media S.A.; 2019;13:1–21. Available from: <https://www.frontiersin.org/article/10.3389/fnins.2019.00513/full>
5. Manna P, Jain SK. Obesity, Oxidative Stress, Adipose Tissue Dysfunction, and the Associated Health Risks: Causes and Therapeutic Strategies. *Metab. Syndr. Relat. Disord.* [Internet]. Mary Ann Liebert, Inc.; 2015 [cited 2019 Jan 7];13:423–44. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26569333>
6. Sies H. The Concept of Oxidative Stress After 30 Years. *Biochem. Oxidative Stress* [Internet]. Cham: Springer International Publishing; 2016. p. 3–11. Available from: http://link.springer.com/10.1007/978-3-319-45865-6_1
7. Schieber M, Chandel NS. ROS function in redox signaling and oxidative stress. *Curr. Biol.* Cell Press; 2014.
8. Marseglia L, Manti S, D'Angelo G, Nicotera A, Parisi E, Di Rosa G, et al. Oxidative stress in obesity: a critical component in human diseases. *Int. J. Mol. Sci.* Multidisciplinary Digital Publishing Institute (MDPI); 2014;16:378–400.

9. An H, Du X, Huang X, Qi L, Jia Q, Yin G, et al. Obesity, altered oxidative stress, and clinical correlates in chronic schizophrenia patients. *Transl. Psychiatry*. Nature Publishing Group; 2018;8.
10. Roberts CK, Sindhu KK. Oxidative stress and metabolic syndrome. *Life Sci*. 2009;84:705–12.
11. Rani V, Deep G, Singh RK, Palle K, Yadav UCS. Oxidative stress and metabolic disorders: Pathogenesis and therapeutic strategies. *Life Sci*. Elsevier Inc.; 2016;148:183–93.
12. Tangvarasittichai S. Oxidative stress, insulin resistance, dyslipidemia and type 2 diabetes mellitus. *World J. Diabetes* [Internet]. 2015;6:456. Available from: <http://www.wjgnet.com/1948-9358/full/v6/i3/456.htm>
13. Hurrell S, Hsu WH. The etiology of oxidative stress in insulin resistance. *Biomed. J. Elsevier B.V.*; 2017. p. 257–62.
14. Laakso M, Kuusisto J. Insulin resistance and hyperglycaemia in cardiovascular disease development. *Nat. Rev. Endocrinol*. Nature Publishing Group; 2014;10:293–302.
15. Bornfeldt KE, Tabas I. Insulin resistance, hyperglycemia, and atherosclerosis. *Cell Metab*. 2011;14:575–85.
16. Pizzino G, Irrera N, Cucinotta M, Pallio G, Mannino F, Arcoraci V, et al. Oxidative Stress: Harms and Benefits for Human Health. *Oxid. Med. Cell. Longev*. Hindawi Limited; 2017;2017.
17. Hermsdorff HHM, Puchau B, Volp ACP, Barbosa KB, Bressan J, Zulet MÁ, et al. Dietary total antioxidant capacity is inversely related to central adiposity as well as to metabolic and oxidative stress markers in healthy young adults. *Nutr. Metab. (Lond)*. BioMed Central; 2011;8:59.
18. Martínez-Tomé M, Murcia MA, Mariscal M, Lorenzo ML, Gómez-Murcia V, Bibiloni M, et al. Evaluation of antioxidant activity and nutritional composition of flavoured dehydrated soups packaged in different formats. Reducing the sodium content. *J. Food Sci. Technol*. Springer India; 2015;52:7850–60.
19. Soccio M, Laus MN, Flagella Z, Pastore D. Assessment of antioxidant capacity and putative healthy effects of natural plant products using soybean lipoxygenase-based methods. An overview. *Molecules*. MDPI AG; 2018;23.
20. Haytowitz D, Bhagwat S. USDA Database for the Oxygen Radical Absorbance Capacity (ORAC) of Selected Foods, Release 2. US Dep. Agric. [Internet]. 2010;10–48. Available from: http://www.pureacai berry.com/testimonials/ORAC_R2.pdf
21. Galasko DR, Peskind E, Clark CM, Quinn JF, Ringman JM, Jicha GA, et al. Antioxidants for Alzheimer disease: A randomized clinical trial with cerebrospinal fluid biomarker measures. *Arch. Neurol*. 2012;69:836–41.
22. Carocho M, Ferreira ICFR, Morales P, Soković M. Antioxidants and Prooxidants: Effects on Health and Aging. *Oxid. Med. Cell. Longev*. [Internet]. Hindawi Limited; 2018;1–2. Available from: <https://www.hindawi.com/journals/omcl/2018/1472708/>
23. Mirmiran P, Hosseini Esfahani F, Mehrabi Y, Hedayati M, Azizi F. Reliability and relative validity of an FFQ for nutrients in the Tehran Lipid and Glucose Study. *Public Health Nutr*. 2010;13:654–62.

24. Colarusso L, Serafini M, Lagerros YT, Nyren O, La Vecchia C, Rossi M, et al. Dietary antioxidant capacity and risk for stroke in a prospective cohort study of Swedish men and women. *Nutrition*. Elsevier Inc.; 2017;33:234–9.
25. Villaverde P, Lajous M, MacDonald C-J, Fagherazzi G, Bonnet F, Boutron-Ruault M-C. High dietary total antioxidant capacity is associated with a reduced risk of hypertension in French women. *Nutr. J.* 2019;18:31.
26. Rautiainen S, Levitan EB, Orsini N, Åkesson A, Morgenstern R, Mittleman MA, et al. Total antioxidant capacity from diet and risk of myocardial infarction: A prospective cohort of women. *Am. J. Med.* 2012;125:974–80.
27. Vece MM, Agnoli C, Grioni S, Sieri S, Pala V, Pellegrini N, et al. Dietary total antioxidant capacity and colorectal cancer in the Italian epic cohort. *PLoS One*. Public Library of Science; 2015;10.
28. Bastide N, Dartois L, Dyèvre V, Dossus L, Fagherazzi G, Serafini M, et al. Dietary antioxidant capacity and all-cause and cause-specific mortality in the E3N/EPIC cohort study. *Eur. J. Nutr. Dr. Dietrich Steinkopff Verlag GmbH and Co. KG*; 2017;56:1233–43.
29. Ellulu MS, Patimah I, Khaza'ai H, Rahmat A, Abed Y. Obesity and inflammation: the linking mechanism and the complications. *Arch. Med. Sci. [Internet]*. 2017;13:851–63. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28721154>
30. Fernández-Sánchez A, Madrigal-Santillán E, Bautista M, Esquivel-Soto J, Morales-González A, Esquivel-Chirino C, et al. Inflammation, oxidative stress, and obesity. *Int. J. Mol. Sci. Multidisciplinary Digital Publishing Institute (MDPI)*; 2011;12:3117–32.
31. Nigro E, Scudiero O, Monaco ML, Palmieri A, Mazzarella G, Costagliola C, et al. New insight into adiponectin role in obesity and obesity-related diseases [Internet]. *Biomed Res. Int. Hindawi Publishing Corporation*; 2014 [cited 2017 May 2]. p. 658913. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25110685>
32. Procaccini C, De Rosa V, Galgani M, Carbone F, La Rocca C, Formisano L, et al. Role of adipokines signaling in the modulation of T cells function. *Front. Immunol. [Internet]*. 2013;4:332. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24151494>
33. Aggarwal BB. Targeting inflammation-induced obesity and metabolic diseases by curcumin and other nutraceuticals. *Annu. Rev. Nutr. [Internet]*. NIH Public Access; 2010 [cited 2019 Jan 7];30:173–99. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/20420526>
34. Abdali D, Samson SE, Grover AK. How effective are antioxidant supplements in obesity and diabetes? *Med. Princ. Pract. [Internet]*. Karger Publishers; 2015 [cited 2019 Jan 7];24:201–15. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25791371>
35. Jager J, Grémeaux T, Cormont M, Le Marchand-Brustel Y, Tanti JF. Interleukin-1 β -induced insulin resistance in adipocytes through down-regulation of insulin receptor substrate-1 expression. *Endocrinology*. 2007;148:241–51.
36. Dal S, Sigrist S. The Protective Effect of Antioxidants Consumption on Diabetes and Vascular Complications. *Diseases*. MDPI AG; 2016;4:24.

37. Hussain T, Tan B, Yin Y, Blachier F, Tossou MCB, Rahu N. Oxidative Stress and Inflammation: What Polyphenols Can Do for Us? *Oxid. Med. Cell. Longev.* [Internet]. 2016;1–9. Available from: <https://www.hindawi.com/journals/omcl/2016/7432797/>
38. Solverson PM, Henderson TR, Debelo H, Ferruzzi MG, Baer DJ, Novotny JA. An Anthocyanin-Rich Mixed-Berry Intervention May Improve Insulin Sensitivity in a Randomized Trial of Overweight and Obese Adults. *Nutrients* [Internet]. 2019;11. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/31775396>
39. Fraga CG, Croft KD, Kennedy DO, Tomás-Barberán FA. The effects of polyphenols and other bioactives on human health. *Food Funct. Royal Society of Chemistry*; 2019;10:514–28.
40. Papada E, Kaliora AC. Antioxidant and anti-inflammatory properties of mastiha: A review of preclinical and clinical studies. *Antioxidants*. 2019.
41. Bahadoran Z, Golzarand M, Mirmiran P, Shiva N, Azizi F. Dietary total antioxidant capacity and the occurrence of metabolic syndrome and its components after a 3-year follow-up in adults: Tehran Lipid and Glucose Study. *Nutr. Metab.* 2012;9:1–9.
42. Azzini E, Giacometti J, Russo GL. Antiobesity Effects of Anthocyanins in Preclinical and Clinical Studies. *Oxid. Med. Cell. Longev. Hindawi*; 2017.
43. Mopuri R, Islam MS. Medicinal plants and phytochemicals with anti-obesogenic potentials: A review. *Biomed. Pharmacother. Elsevier Masson SAS*; 2017;89:1442–52.
44. Siddiqui AH, Gulati R, Tauheed N, Pervez A. Correlation of waist-to-hip ratio and oxidative stress in patients of acute myocardial infarction. *J. Clin. Diagnostic Res.* 2014;8:4–7.
45. Raja Kumar S, Mohd Ramli ES, Abdul Nasir NA, Ismail NHM, Mohd Fahami NA. Preventive Effect of Naringin on Metabolic Syndrome and Its Mechanism of Action: A Systematic Review. *Evidence-based Complement. Altern. Med. Hindawi Limited*; 2019.
46. Yannakoulia M, Ntalla I, Papoutsakis C, Farmaki AE, Dedoussis G V. Consumption of vegetables, cooked meals, and eating dinner is negatively associated with overweight status in children. *J. Pediatr.* 2010;157:815–20.
47. Karfopoulou E, Brikou D, Mamalaki E, Bersimis F, Anastasiou CA, Hill JO, et al. Dietary patterns in weight loss maintenance: results from the MedWeight study. *Eur. J. Nutr. Dr. Dietrich Steinkopff Verlag GmbH and Co. KG*; 2017;56:991–1002.
48. van der Schaft N, Schoufour JD, Nano J, Kieft-de Jong JC, Muka T, Sijbrands EJG, et al. Dietary antioxidant capacity and risk of type 2 diabetes mellitus, prediabetes and insulin resistance: the Rotterdam Study. *Eur. J. Epidemiol. Springer Netherlands*; 2019;34:853–61.
49. Kim K, Vance TM, Chun OK. Greater Total Antioxidant Capacity from Diet and Supplements Is Associated with a Less Atherogenic Blood Profile in U.S. Adults. *Nutrients. Multidisciplinary Digital Publishing Institute (MDPI)*; 2016;8.
50. Casas-Grajales S. Antioxidants in liver health. *World J. Gastrointest. Pharmacol. Ther. Baishideng Publishing Group Inc.*; 2015;6:59.

51. Savu O, Ionescu-Tirgoviste C, Atanasiu V, Gaman L, Papacoea R, Stoian I. Increase in Total Antioxidant Capacity of Plasma despite High Levels of Oxidative Stress in Uncomplicated Type 2 Diabetes Mellitus. *J. Int. Med. Res.* SAGE Publications; 2012;40:709–16.
52. Petelin A, Tedeschi P, Maietti A, Jurdana M, Brandolini V, Pražnikar ZJ. Total Serum Antioxidant Capacity in Healthy Normal Weight and Asymptomatic Overweight Adults. *Exp. Clin. Endocrinol. Diabetes.* 2017;125:470–7.