Hotter and Drier Climate Made the Mediterranean Europe and Northern Africa Region a Shrubber Landscape

Wei Fang

Pace University

Chuixiang Yi (✉ cyi@qc.cuny.edu)

CUNY Queens College: Queens College <https://orcid.org/0000-0001-8546-6157>

Deliang Chen

University of Gothenburg: Goteborgs Universitet

Peipei Xu

Anhui Normal University

George Hendrey

CUNY Queens College: Queens College

Nir Krakauer

CUNY City College: The City College of New York

Katherine Jensen

CUNY City College: The City College of New York

Shan Gao

Chinese Academy of Sciences

Zihan Lin

Michigan State University

Gabriella Lam

Pace University

Qin Zhang

Dalian University of Technology

Tao Zhou

Beijing Normal University

Research Article

Keywords: LAI deficit, GPP deficit, climate drivers, legacy effect, landcover transition

Posted Date: February 19th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-214050/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Oecologia on October 13th, 2021. See the published version at <https://doi.org/10.1007/s00442-021-05041-3>.

Hotter and Drier Climate Made the Mediterranean Europe and Northern Africa Region a Shrubber Landscape¹

Running Title: Mediterranean Area has Become Shrubber

Wei Fang^{1,2}, Chuixiang Yi^{*2,3}, Deliang Chen⁴, Pei Xu^{2,5}, George Hendrey^{2,3}, Nir Krakauer^{3,6}, Katherine Jensen^{3,6}, Shan Gao^{2,7}, Zihan Lin⁸, Gabriella Lam¹, Qin Zhang^{2,9}, Tao Zhou¹⁰

¹Department of Biology, Pace University, New York, NY 10038, USA

²School of Earth and Environmental Sciences, Queens College, City University of New York, Flushing, NY 11367, USA

³The Graduate Center, City University of New York, New York, NY 10016, USA

⁴Department of Earth Sciences, University of Gothenburg, 40530 Gothenburg, Sweden

⁵School of Geography and Tourism, Anhui Normal University, Wuhu 241002, China

⁶The City College of New York, City University of New York, New York, NY 10031, USA

⁷Institute of Tibetan Plateau Research, Chinese Academy of Sciences, Beijing 100101, China

⁸Center for Global Change and Earth Observations, Michigan State University, East Lansing, Michigan 48823, USA

⁹Dalian University of Technology, Dalian 116024, China.

¹⁰Faculty of Geographical Science, Beijing Normal University, Beijing 100875, China

Corresponding author: chuixiang.yi@qc.cuny.edu

¹ Author Contributions: CY and WF conceived and formulated the idea. WF, KJ, ZL processed the satellite data, WF, SG performed statistical analysis, WF, DL, PX wrote the manuscript; other authors provided editorial advice.

Abstract: A shift to warmer temperatures has left the Mediterranean Europe and Northern Africa (MENA) region more vulnerable to drought and land degradation. We used LAI and GPP deficit, the difference between actual and historical-maximum values, to describe vegetation structural and functional dynamics and consequential landcover change in response to climate variability during 2001-2019 in the area (20°W-45°E, 20°N-45°N). We found that (1) the vegetation responses varied significantly among eight landcover types with the following ranked importance: forests, savannas, a mosaic of cropland and natural vegetation (CNV), croplands, permanent wetlands, urban land, grasslands, and shrublands, each with distinctive yet overlapping signatures over the spectrums of the climate conditions considered. (2) Forests, occupying the coolest and wettest niche of the MENA region, showed the strongest and most dominating response to severe drought with a lag of 1-3 years and a legacy of 10 years. (3) The total areas of savannas and CNV mosaics in MENA increased by 394,994 km² and 404,592 km² respectively while that of forests decreased by 33,091 km² despite of the fertilizer effect of elevated ambient CO₂. Shrublands, occupying the hottest and driest niche of MENA, extended by 287,134 km² while grasslands and croplands retreated by 490,644 km² and 225,263 km². The area of permanent wetlands increased by 49,192 km², and that of urban land increased by 39,570 km². A net total of 57,649 km² of barren land became vegetated over the years. With warmer temperature and more extended period of drought, MENA has evolved towards a shrubbier landscape.

Keywords: LAI deficit, GPP deficit, climate drivers, legacy effect, landcover transition

1. Introduction

Landcover, especially vegetation dynamics over land, plays a critical role in atmospheric processes and the water and carbon cycles globally (Foley, et al, 1996; Sitch et al, 2003; Assal et al, 2016). An analysis of FLUXNET data revealed that the exchanges of carbon, water, and energy between terrestrial ecosystems and the atmosphere are limited primarily by water availability when the mean annual temperature is above a threshold of 16°C (Yi *et al.*, 2010). Climate records from the past six decades show that the annual mean temperature of a significant part of the Mediterranean, Europe, and Northern Africa (MENA) region have shifted from below to above this threshold of 16 °C (Yi *et al.*, 2014; Allen et al., 2015; Huang et al., 2015). This shift to higher temperatures has left the largely arid and semi-arid MENA region more vulnerable to drought and land degradation (Somot et al., 2008; Segui et al., 2010; Friend, 2010; IPCC, 2013; Zhang et al., 2020). Studies have reported drought-induced forest impacts and diebacks in the Mediterranean region (Allen et al., 2010; Carnicer et al., 2011; Peñuelas et al., 2001; Martínez-Vilalta and Piñol, 2002; Matusick et al., 2013), as well as shifts in vegetation composition (Jump and Peñuelas, 2005; Anderegg et al., 2013).

Climate-induced variability of vegetation in the MENA region is a matter of both ecological and economic concern, as strong sensitivity to climate can result in rapid land use change (Vanacker *et al.* 2005; Serra et al, 2008). Over longer time scales, relatively small shifts in background climate can impact the distribution of ecosystems and potentially the viability of agricultural and pastoral systems (Ciais *et al.*, 2005; Vicente-Serrano *et al.*, 2012). Interest in the climate sensitivities of vegetation in semiarid regions is evident in the substantial body of research devoted to characterizing the relationships between precipitation, soil type, land management, and vegetation growth in these water-stressed regions (Zaitchik et al, 2007, Vicente-Serrano et al, 2012). Previous studies have shown that, since 1948, warming climate has moved the 16 °C isotherm poleward, leading to a predicted northward shift of vegetation

distribution of the Northern hemisphere (Yi et al, 2014). Vicente-Serrano et al (2012) also showed a decline of the average vegetation cover in the semiarid Mediterranean environments during 1984-2008.

At regional to continental scales, satellite observation offers a feasible and effective approach for monitoring vegetation dynamics. One of the most common observations of vegetation dynamics is based on the normalized difference vegetation index (NDVI). The NDVI, computed as a normalized ratio of reflectance in the near-infrared and red portions of the electromagnetic spectrum, provides a measure of chlorophyll abundance at the Earth's surface and offers an indirect measure of energy absorption and vegetation density (Myneni *et al.*, 1995; Kerr et al., 2003).

A number of vegetation indices with direct ecological meanings are derived from NDVI using additional algorithms and information. One of them is the leaf area index (LAI). LAI is a fundamental parameter that reflects vegetation structure in the processes of fixing atmospheric CO₂ into organic matter. LAI is defined as the one-sided green leaf area per unit ground area in broadleaf canopies and as half the total needle surface area per unit ground area in coniferous canopies. Another widely used vegetation index is gross primary production (GPP), which is a measure of the growth of the terrestrial vegetation that reflects vegetation function in the processes of fixing atmospheric CO₂ into organic matter. GPP correlates closely with LAI where LAI is ~4 or less, suggesting that leaf area is a critical determinant of GPP in most of the MENA region (Chapin III et al, 2011a).

This study takes advantage of interannual climatic variability during 2001-2019 (Sprintson et al., 2009; Fischer and Knutti, 2015) to characterize the climatic sensitivities of vegetation in the MENA region. Time series data from the Moderate Imaging Spectrometer (MODIS) are used to describe interannual patterns in vegetation. Data sources, processing steps,

and methods are detailed below. We applied the “perfect deficit” approach (Yi *et al.*, 2012) to identify the primary climatic drivers of LAI variability over vegetation across the entire MENA domain aggregated by landcover class.

We used both LAI deficit and GPP deficit as direct measures of climate and non-climate stress on ecosystem structure and function experienced by the vegetation of the semiarid MENA area. A focus is on climate stress expressed by a number of climate indices describing thermal and water conditions and their combined effects. We attempt to establish empirical connections of LAI deficit and GPP deficit with a few climate variables/indices (i.e. annual average temperature (T), annual average precipitation (mm), Dryness Index, Temperature-Precipitation (TP) Index, and Standardized Precipitation-Evapotranspiration Index (SPEI)). The aim is to understand how landcover changes, especially vegetation structural and functional changes and variabilities, respond to these climate variables: Specifically, we will find out which variable(s) are the dominant driver(s) at the time scale of a decade. When vegetation types are considered individually, we hypothesize that LAI deficit and GPP deficit of different vegetation types would respond to climate variables differently, and the distribution of vegetation types may show a shift over the years or may extend or shrink in response to climate change. The direction and intensity of landcover shift over a short-term (i.e. after the 2002 severe drought) and a long-term (2001-2018) will be further explored with landcover transition matrices. Such ecological assessment at the temporal scale of decades is much needed for the strategic planning of resource management, for optimizing vegetation productivity and ecosystem services.

2. Materials and Methods

2.1 Climate and remote sensing data

Remote sensing data such as MODIS LAI, GPP and landcover, and various climate measures were collected for the MENA region, defined as the bounding box of 20°N - 45°N & 20°W - 45°E. This is a box-shaped area around the Mediterranean Forests, Woodlands & Scrub biome defined by the World Wide Fund for Nature (WWF) in the MENA region (Fig.1).

2.1.1 Remote sensing data

Nineteen years (01-Jan-2001 – 31-Dec-2019) of global, 8-day composite MODIS LAI (MODIS/006/MOD15A2H), GPP (MODIS/006/MOD17A2H), and yearly landcover data (MODIS/006/MCD12Q1) at 500m resolution were obtained from the NASA Land Processes Distributed Active Archive Center (LP DAAC), USGS/Earth Resources Observation and Science (EROS) Center within Google Earth Engine. Annual landcover classification is defined with the International Geosphere Biosphere Programme (IGBP) global vegetation classification scheme. It describes the landcover properties derived from observations spanning a year of Terra and Aqua MODIS data. The primary landcover scheme identifies 17 landcover classes defined by IGBP, which includes 11 natural vegetation classes, 3 developed and mosaicked land classes, and 3 non-vegetated land classes. The landcover types were further grouped into eight categories of vegetated cover: forests (including class 1-5, evergreen needleleaf forest, evergreen broadleaf forest, deciduous needleleaf forest, deciduous broadleaf forest and mixed forest), shrublands (including class 6-7, closed shrublands and open shrublands), savannas (classes 8-9 woody savannas and savannas), grasslands (class 10), permanent wetlands (class 11), croplands (class 12), urban and built-up (class 13) and crop+natural vegetation (CNV) mosaic (class 14). Pixels with landcover changed over the years were coded white in the map (Fig. 1) and were excluded for the no-landcover-change-based analyses along with the non-vegetated areas (including snow and ice-class 15, barren land-class 16 and water-class 17).

2.1.2. Climate indices

Two primary instrumental measures (temperature and precipitation) and three compound climate indices (one with a linear function, one with an exponential function, and one being standardized) were chosen for this study.

Monthly global surface air temperature (average air temperature at 2m height) and precipitation (monthly sum) data (01-Jan-2001 – 31-Dec-2019) in 0.25 degrees resolution were provided by ECMWF/Copernicus Climate Change Service within Google Earth Engine. ERA5 is the fifth generation ECMWF atmospheric reanalysis of the global climate that combines model data with observations from across the world into a globally complete and physically consistent dataset.

To calculate dryness index, monthly average net radiation (W m^{-2}) data (01-Jan-2001 – 31-Dec-2019) in 0.1 degrees resolution were collected from the FLDAS (Famine Early Warning System Network (FEWS NET) Land Data Assimilation System) dataset (McNally et al, 2017) provided by the NASA GES DISC at NASA Goddard Space Flight Center within Google Earth Engine. This FLDAS dataset uses Noah version 3.6.1 surface model with CHIRPS-6 hourly precipitation that has been downscaled using the NASA Land Surface Data Toolkit.

The radiation data were first resampled with the same projection system and resolution of the temperature and precipitation data. Net radiation (R_n) is defined as the difference between net shortwave radiation and net longwave radiation.

The dryness index of Budyko was calculated by

$$\text{Dryness} = \frac{R_n}{LP}, \quad (1)$$

where R_n ($\text{MJ m}^{-2} \text{ yr}^{-1}$) and P (mm yr^{-1}) are respectively monthly mean net radiation and precipitation for each grid cell and L is a constant 2.5 MJ kg^{-1} , the enthalpy of vaporization (Budyko, 1961). Since temperature is often positively correlated with R_n , Dryness increases in response to increasing T or decreasing P . When there is no precipitation in a month, the dryness index value could be infinitely large. Since on the global scale, the average net radiation was between 98 and 112.6 W m^{-2} (Liang, 2017), the dryness index values (annual average of the monthly values) were capped under 100 for later analysis.

Temperature-Precipitation (TP) Index was derived initially from a model of soil nitrogen variations (Jenny, 1980) and further adapted to model vegetation responses to a changing climate (Yi et al., 1996). It is calculated by

$$TP = e^{0.08T} (1 - e^{-0.005P}) \quad (2)$$

where T ($^{\circ}\text{C}$) and P (mm yr^{-1}) are annual mean temperature and precipitation for each grid cell respectively (Yi et al., 1996). TP Index increases in response to decreasing T or increasing P .

SPEI (Standardized Precipitation Evapotranspiration Index) was a measure of water balance, calculated by the standardization of water deficit D , while

$$D = P - PET, \quad (3)$$

where P (mm yr^{-1}) and PET (mm yr^{-1}) are annual precipitation and potential evapotranspiration for each grid cell respectively (Vicente-Serrano et al, 2010). The SPEI can measure drought severity according to its intensity and duration. It can identify the onset and end of drought episodes as well. The SPEI is a standardized variable, and it can therefore be compared with other SPEI values over time and space. An SPEI of 0 indicates a value corresponding to 50% of the cumulative probability of D , according to a log-logistic distribution.

A globally-gridded (0.05° resolution) SPEI index calculated from the CRU TS3.23 precipitation and reference evapotranspiration, was provided by the SPEI website

(<https://digital.csic.es/handle/10261/202305>). The SPEI used in the analyses was at the 12-month temporal scale ending December of the previous year.

2.2 Perfects, deficits and relative deficits

Many large-scale vegetation and landcover studies are formulated as statistical analysis of climatological anomalies. This study takes a vegetation-centric approach, focusing on the differences of vegetation performance under optimal and suboptimal conditions.

Yi *et al* (2012) introduced a “perfect-deficit” approach for quantifying links between climate extremes and variations in carbon storage. We apply this approach in developing a climate stress indicator based on our MODIS LAI and GPP data in the MENA area for every 8-day period throughout a year, then converted to monthly values. For each pixel, the “perfect” LAI value of the month is defined as the maximum LAI value for this month over the 19 years, and the “deficit” LAI value of the month for each year is defined as the difference between the observed value from the “perfect” value of that month. Therefore, for the entire studied time span (year 2001-2019) for the i th month,

$$\text{LAI}_{\text{Perfect}}(i) = \text{Maximum}\{ \text{LAI}_{\text{Observed}}(i)_{2001}, \text{LAI}_{\text{Observed}}(i)_{2002}, \dots, \text{LAI}_{\text{Observed}}(i)_{2019} \}, \quad (4a)$$

$$\text{GPP}_{\text{Perfect}}(i) = \text{Maximum}\{ \text{GPP}_{\text{Observed}}(i)_{2001}, \text{GPP}_{\text{Observed}}(i)_{2002}, \dots, \text{GPP}_{\text{Observed}}(i)_{2019} \}, \quad (4b)$$

and for the i th month of the j th year,

$$\text{LAI}_{\text{Deficit}}(i,j) = \text{LAI}_{\text{Perfect}}(i) - \text{LAI}_{\text{Observed}}(i,j), \quad (5a)$$

$$\text{GPP}_{\text{Deficit}}(i,j) = \text{GPP}_{\text{Perfect}}(i) - \text{GPP}_{\text{Observed}}(i,j). \quad (5b)$$

Essentially, we are looking at how the observed LAI for a given month under suboptimal climate and ecological conditions departed from the highest value observed for that time of the year over the 19-year period. The perfect value is presumed to correspond to the optimal growing

conditions of climate, edaphic features, and other impinging ecological factors for that particular area.

The monthly LAI or GPP deficit values were first averaged within each year to give us an annual mean deficit for each pixel, then were averaged spatially (pixel by pixel) across each of the landcover classes for each year.

2.3 Data Analyses

There were four sequential steps of data analyses as shown in Fig. 2. Step 1, eight vegetated landcover types with no change over 18 years were extracted to conduct multiple regression analyses; Step 2, regressions of vegetation structural (LAI deficit) and functional (GPP deficit) responses over climate drivers (including temperature, precipitation, TP index, dryness index, and SPEI) were conducted over all vegetated pixels and within each of 8 vegetation types; Step 3, results of 10 regression analyses and probability density function (PDF) curves of 7 variables all pointed to the strong constraints within each vegetation type, which shed new light on landcover changes over the years. Step 4, using the characteristic order along the spectrums of climate variables and vegetation response variables, direction and intensity of landcover change were quantified using time series and transition matrices.

All LAI, GPP and landcover data were retrieved within Google Earth Engine from MODIS products (MOD15A2 and MOD17A2, MCD12Q1). All climate datasets (global surface air temperature, precipitation, and net shortwave radiation data with temporal resolution of one month) were resolved to a common projection at 0.25° resolution. All annual LAI deficits, annual GPP deficits, and all climate indices were calculated within Google Earth Engine using JavaScript. Since we are looking for long-term large-scale patterns, all data were aggregated to annual averages of eight landcover types. All processed spatial data were then exported to

Geotiff files with WGS84 and 500m resolution for further statistical tests and graphing in R-Studio (version 1.2.1335). All maps were made in ArcGIS Desktop 10.7.1.11595.

3. Results

3.1 Climate drivers

The first purpose of the study was to find the climate indices that the vegetation structure indicator LAI and function indicator GPP responded to the most, and how those relationships vary across different vegetation types. We found that overall LAI deficit was negatively correlated with annual average temperature (Fig.3a1, Adjusted $R^2=0.529$, $p<0.001$) and Dryness (Fig.3a4, Adjusted $R^2=0.588$ for linear and 0.770 for curvilinear relationship, $p<0.001$), positively correlated with precipitation (Fig. 3a2, Adjusted $R^2=0.613$, $p<0.001$) and TP Index (Fig.3a4, Adjusted $R^2=0.491$, $p<0.001$), and not correlated with SPEI (Fig.3a5, Adjusted $R^2=0.007$, $p=0.155$) (Table 1). When the vegetation was divided into eight categories, only forest consistently showed a similar pattern to the overall vegetation structural responses while shrubland showed an opposite trend or no trend.

Similarly, we found GPP deficit of overall vegetation was negatively correlated with annual average temperature (Fig. 3b1, Adjusted $R^2=0.553$, $p<0.001$) and dryness (Fig. 3b4, Adjusted $R^2=0.624$ for linear and 0.850 for curvilinear relationship, $p<0.001$), positively correlated with accumulative precipitation (Fig. 3b2, Adjusted $R^2=0.548$, $p<0.001$), TP Index (Fig. 3b3, Adjusted $R^2=0.509$, $p<0.001$), and weakly with SPEI (Fig. 3b5, Adjusted $R^2=0.041$, $p=0.007$) (Table 1).

When divided into eight landcover categories: only forest showed an almost consistently similar pattern to the overall vegetation structural responses while shrubland showed an opposite trend or no trend. The only exception was in precipitation. Within each vegetation type, GPP

deficit decreased when precipitation increased. In another word, more precipitation, less reduction of GPP within each vegetation type as we would normally expect. The overall correlation between GPP deficit and precipitation, however, is counter-intuitively positive. When the timing of landcover change and PDF curves were closely examined in later part of this study, it became clear that the overall pattern was more driven by intrinsic productivities of each vegetation type rather than local responses to climate variables alone.

3. 2 Landcover-specific signatures over the spectrums of climate conditions

Based on the probability density function (PDF) curves for both dependent variables (i.e. vertical PDF graphs for LAI and GPP deficits) and independent variables (i.e. horizontal PDF graphs for climate indices) (Fig 3), each vegetation cover demonstrated unique and sometimes overlapping distribution signatures over the spectrums of structural and functional responses, and climate conditions.

For example, forests had by far the highest LAI deficits ($11.28 \pm 1.12 \text{ m}^2 \text{ m}^{-2}$), followed by savannas ($7.09 \pm 0.74 \text{ m}^2 \text{ m}^{-2}$), CNV mosaic ($6.81 \pm 0.59 \text{ m}^2 \text{ m}^{-2}$), croplands ($6.75 \pm 0.50 \text{ m}^2 \text{ m}^{-2}$), wetlands ($6.05 \pm 0.52 \text{ m}^2 \text{ m}^{-2}$), urban land ($3.94 \pm 0.31 \text{ m}^2 \text{ m}^{-2}$), grasslands ($3.30 \pm 0.38 \text{ m}^2 \text{ m}^{-2}$) and shrublands ($1.48 \pm 0.21 \text{ m}^2 \text{ m}^{-2}$) in a descending order (Table 2). The LAI deficit of shrublands was less than half of that of grasslands, which was the second lowest among all. Likewise, forests had the highest GPP deficits, followed by the same descending order (Table 2). This order reflected the differential responses, both structural and functional, to these climate variables among these vegetation types, and their differential responses to climate stress.

When looking into the spectrums of climate conditions, stratification of landcover classes by climate conditions is clearly evident in the data record. Forests occupied the coolest ($10.00 \pm 0.47 \text{ }^\circ\text{C}$) and wettest ($1102.31 \pm 87.31 \text{ mm}$) niche while shrublands occurred in the hottest

(17.46±0.32 °C) and driest (342.24±49.24 mm) area, polar opposite to forests. Each vegetation landcover type is constrained within a limited range of climate conditions but each has a certain level of overlapping with others. Within a constrained range of climate conditions, alternative steady states (vegetation types) may exist.

3.3 Landcover change over time

3.3.1 Remotely sensed areal change of landcover

According to the MODIS annual landcover product during 2001-2018, the total areas of savannas and CNV mosaics in MENA increased by 394,994 km² (5.26% of savanna area in 2001) and 404,592 km² (29.70% of CNV area in 2001), with a significantly steady increasing trend ($R^2 = 0.9295$ and 0.9129 , respectively), while the area of forests decreased by 33,091 km² (1.56%) (Fig. 4a, 4b & 4c) despite the fertilizer effect of elevated ambient CO₂. Forest response to elevated CO₂ is conserved across a broad range of productivity (Norby, et al, 2005). What was initially observed in the temperate FACE experiments may not be representative of other regions (Hickler et al., 2008), and belowground response could be more continuous than aboveground response to CO₂ enrichment (Jackson et al, 2009) which cannot be detected by satellite data.

More specifically, the area identified as forest in this analysis decreased by 41,401 km² (a 2.92% drop) over a period of 3 years following the drought/heat wave of 2002, and didn't recover to the predrought level until 2014, 11 years after the drought. A second major drop happened between 2016 and 2017, leaving an overall areal decline between 2001 and 2018.

Over the 18 years, shrublands, occupying the hottest and driest niche of MENA, extended by 287,134 km² (11.53%) while grasslands and croplands retreated by 490,644 km² (4.62%) and 225,263 km² (1.22%) (Fig. 4d, 4e & 4f). Shrubs showed a 5-

year increase after the 2002 drought and an overall increasing trend in shrublands (Fig. 4f), while grasses showed an initial 3-year increase after the 2002 drought then continuous decline afterwards for 13 years (Fig. 4e). Croplands occupied the largest area in 2003 (18,582,372 km²) both temporally across 18 years and spatially across 8 vegetation types, right after the 2002 drought, then declined afterwards with fluctuations.

The area of permanent wetlands increased by 49,192 km² (24.88%), and that of urban land increased by 39,570 km² (2.90%) (Fig. 4g & 4h).

3.3.2 Matrices of landcover change

To quantify the direction and intensity of change, we used the area of each vegetation type that had changed to make transition matrices. A transition matrix could illustrate the detailed vegetation transition during either a short-term period (i.e. during 2002-2003 right after the 2002 heatwave/drought, Table 3a) or a long-term period (i.e. during 2001-2018, Table 3b, Fig. 5).

In Table 3a and 3b, the diagonal cell areas remained in the same vegetation types. The upper right triangle portion indicates transitions to a less productive vegetation type, while the lower left triangle indicates transitions to a more productive vegetation type, based on what we have learned from PDF curves of LAI and GPP deficits.

During 2002-2003, 17,775 km² of forested area became less productive and remote sensing identified them as savannas. While a total of 22,680 km² of barren land became vegetated (turned into grassland and shrubland). As a result, a total of 52,986 km² became less productive based on the four categories with over 5,000 km² (i.e. from forest to savanna, from savanna to grassland, from cropland to grassland, and from grassland to shrubland) while a total

of 56,814 km² became more productive based the five categories listed in the lower triangle (i.e. from grassland to savanna, from grassland to cropland, from shrubland to grassland, and from barren land to either grassland or shrubland).

During 2001-2018, 55,948 km² of forested area became less productive savannas (Fig.5, visible in southern France and western Portugal) while 54,475 km² of savannas had the reversed transition, leaving a net loss of forests by a mere 1,473 km². A total of 85,257 km² of barren land became vegetated (mostly turned into grassland and shrubland) and 27,609 km² of vegetated pixels became barren, leaving a net gain of 57,649 km² of vegetated area from barren land (0.913%). Such green-up is visible in parts of northern Algeria and Tunisia, eastern Iran, northern Saudi Arabia and northern Egypt (Fig. 5). There was a total area of 413,922 km² that became more productive and 260,560 km² became less productive (based only on categories of transition with 20,000 km² or more).

4. Discussion

The overall aim of the study was to use LAI and GPP deficits to identify and better understand impacts of climate on landcover in the MENA region during 2001-2018. Specifically, we would like to 1) determine critical climate drivers for internal variability of the landcover as a whole and for several ecosystem types individually, 2) explore Landcover-specific signatures over the spectrums of climate conditions, and 3) identify landcover changes over time and relate these changes to climate changes (and other non-climatic drivers).

4.1 Vegetation-centric approach of LAI and GPP deficits

LAI and GPP deficits, the vegetation-centric approach, provide a different perspective into the climate-vegetation interaction, contrast to the traditional climate anomaly approach.

Through examining the responses of LAI deficit and GPP deficit of all vegetation types combined and individually to various climate indices (including T, P, TP Index, Dryness and SPEI), using univariate regression analyses, both LAI and GPP deficits proved to be effective indicators of how ecosystem photosynthetic structures and functions respond to climate stress. This study took this as a starting point to further explore the responses of the two deficits to climatic conditions for the MENA region.

In general, regressions of LAI deficits and GPP deficits behaved very similarly, except to precipitation (Fig. 3 a2 & b2). Both Forest and overall LAI deficits ($\text{m}^2 \text{m}^{-2}$) had a positive correlation with annual precipitation (mm), which was quite counter-intuitive. To understand this, let's start with Fig. 3b2.

Within each vegetation type, GPP deficits (kg C m^{-2}) decreased when annual precipitation (mm) increased, which meant less GPP loss within the same year with more precipitation. This is consistent with existing literature (Wu et al, 2011; Nielsen et al., 2015). The overall regression of GPP deficits over precipitation, however, had a positive slope, which was overwhelmingly dominated by intrinsic variation among different vegetation types (i.e. forests generally have the highest GPP base, therefore its absolute increase in response to increase of precipitation should be the highest).

Likely, as in Fig. 3a2, the overall regression of LAI deficits over precipitation had a positive slope, with the same reason as for GPP deficits—they are tightly constrained within each vegetation type. However, within forests, the regression slope was also positive. To understand this pattern, we should also look at what has happened to the area (Fig. 4a and Table 3a). In 2002, the MENA region experienced a widespread severe drought. The total area of forests decreased drastically from 2,102,011 km^2 (in 2002) to 2,040,610 km^2 (in 2003), and it continued to decline for two more years (with 2005 having the smallest area of forests). It took

another 10 years (2005-2014) for the area of forests to restore to the pre-drought level. In other words, there were 3 years of lag and 10 years of legacy effects in terms of the distribution area of forests. Similarly, forest area was at its highest in 2017, then declined drastically in 2018, possibly due to another extreme event.

If we look at the time series of LAI deficits, the similar lag and legacy effects exist. The decline of LAI happened in the following few years, therefore LAI was at its highest when precipitation is at its lowest, or LAI deficit was the lowest when climate conditions (such as precipitation) suddenly became less optimal. Therefore, we see this counter-intuitive positive slope for regression of LAI deficits over annual precipitation.

4.2 Five climate indices

There are many climate indices available. For this study, we used two primary instrumental measures (temperature and precipitation) and three compound climate indices, each presenting a different group of indices. Dryness index is expressed in a linear relationship between net radiation and precipitation; TP index is expressed with an exponential formula, containing temperature and precipitation; SPEI is a standardized index allowing flexibility of temporal scales.

Since Dryness, TP index and SPEI all contain the information of precipitation and are not independent from each other, multiple univariate regressions might be easier to interpret than a single multivariate regression.

When response variables LAI deficit and GPP deficit were plotted against each of the five climate indices, all except SPEI were tightly clustered within each vegetation type (Fig.3 and Table 2). Because SPEI is a standardized index, site specificity has probably been removed during the calculation. On the other hand, these tightly distributed clusters may indicate strong

resistance to transitions between vegetation types and alternative stable states caused by the changing climate (Hirota et al, 2011; Chapin et al, 2011b). Although they currently seem to be within the resilience ranges, further studies with finer spatial and temporal scales, and with finer categorization of vegetation types and functional groups could tell a different story. Patterns operating at finer scales will be different from what has been looked at (bigger scales), but this may not necessarily lead to better understanding of the processes operating at coarser resolution we have studied.

4.3 Eight vegetation covers and their PDFs

Despite their structural and functional responses to climate and ecological conditions, most vegetation types are tightly constrained, representing steady states for specific combinations of climate conditions and their immediate alternative steady states if strong enough forces change them one way or another (Fig. 3a-h).

Opposite to patterns in the forests, shrubland LAI deficit was positively correlated with Dryness. However, that does not necessarily mean if the area experienced prolonged directional change related to temperature, water availability and solar radiation, a transition of vegetation coverage between forests and shrubland could happen. When we looked deeply into the ranges of annual average temperature (10.00 ± 0.47 vs 17.46 ± 0.32 °C), precipitation (918.59 ± 72.76 vs 285.20 ± 41.03 mm), TP index (0.51 ± 0.03 vs 0.19 ± 0.01), Dryness (2.37 ± 0.54 vs 20.77 ± 3.95) and SPEI (-0.34 ± 0.61 vs -0.64 ± 0.62) of forest and shrubland (Table 1), forest and shrubland are experiencing very different local climate changes, thus led to polar opposite responses. The results are consistent with the “16°C threshold” prediction (Yi *et al.*, 2010), which stated that the exchanges of carbon, water, and energy between terrestrial ecosystems and the atmosphere are limited primarily by water availability when mean annual temperature is above a threshold of

16°C and by temperature when below 16°C. Other studies also have looked into the changes of landcover under the influence of climate change. For example, Macua et al. (2017) studied factors affecting forest dynamics in the Iberian Peninsula from 1987 to 2012 based on Landsat scenes, and found that the geographical transition from shrubland to forests is closely related to higher soil moisture (Topographic Wetness Index-TWI) and lower winter solar radiation. Meanwhile, strategies aiming at decreasing risk of decline and promoting resistance to abrupt stress in the short-term may not enhance long-term resilience (Vilà-Cabrera, et al., 2018).

4.4 Landcover changes and their concerning implications

Combining with transition maps, transition matrix is a powerful tool to specify the direction and intensity of landscape changes over the region. Comparing short-term and long-term transition matrix can also indicate how vegetation responds to extreme events (such as 2002 drought) immediately and in a long run.

Even though the percentage of vegetated area for forest (4.66%) is far less than cropland (41.19%), grassland (23.88%), savanna (17.38%) and shrubland (5.95%), the patterns of LAI deficit of forest were always consistent with the LAI and GPP deficit of all vegetation, which indicates the strong forest LAI response to climate and its dominating influence over the response of the integrated vegetated landcover in our analysis. The MENA area was still benefiting from the increase of ambient temperature and CO₂ during 2001-2018 indicated in the green-up of barren land and the gradual recovery of forested areas after the 2002 drought. Overall, forests, cropland and grassland were more vulnerable to climate stress and had declined over the past 18 years), while savanna and shrubland are more resilient and their distribution expanded.

One of most striking patterns was the increase of shrubs, both from upstream of cropland and grasses and downstream of barren land. Based the areal changes of vegetation covers during 2001-2018 (partly in Table3b), a total area of 38,938 km² converted from grassland to shrubland and 22,897 km² from cropland to shrubland, while a total area of 37,853 km² of shrubland converted back to grassland and 14,163km² back to cropland. A total area of 52,376 km² converted from barren land to shrubland, while 15,232 km² of shrubland converted back to barren land. There have been various reports on local shrubland die-back in the region (e.g. Lloret et al., 2016 in south Spain), but the overall trend based on satellite data for entire MENA region showing an increase. With higher temperature and more extended period of drought, MENA is becoming less productive (with less forests and croplands) with more shrubby vegetation covers, although a fraction of barren land (less than 1%) became vegetated.

The two dominant vegetated landcovers in this area are cropland (41.19%) and grassland (23.88%) (Table 2). Areas of grassland and cropland fluctuated in opposite patterns over the years (Fig. 4). This is likely due to the fact that the expansion of cropland was usually an encroachment into grassland, while fallow croplands generally returned to grassland.

The average location and altitude of forest vegetation has not yet changed despite the changes in temperature and dryness. On the other hand, the mean and median latitude of cropland decreased over the last decade, likely due to human effort improving irrigation systems for agriculture purposes. The mean latitude and median elevation of grassland also decreased over the last decade, probably benefiting from the fertilization effect of elevated ambient CO₂ concentration. This might have a tight association with the escalating incidence of wildfire near arid and semiarid areas throughout the world, some of which have caused devastating losses.

Over the decade of 2001-2018 in MENA, the mean latitude ($R^2=0.864$, $p<0.001$) and median latitude ($R^2=0.513$, $p<0.05$) of cropland both decreased significantly. Such a counter-

intuitive change is probably due to huge effort of human intervention: establishing irrigation systems and building dams for agriculture (United Nations, 1999; FAO, 2008).

The mean latitude ($R^2=0.779$, $p<0.001$) and median elevation ($R^2=0.473$, $p<0.05$) of grassland also decreased significantly. Prior field manipulation experiments have shown that elevated ambient CO₂ concentration can stimulate the growth and accumulation of standing biomass of grassland in semiarid areas, acting like a carbon fertilizer and a booster of water use efficiency (Dijkstra et al, 2010). Grasses die back during the dry seasons, which naturally turn into standing fuel, causing the increasing rate of wildfire and devastating loss. This has been happening throughout the world, especially in Mediterranean ecoregions such as California, US (Abatzoglou and Williams, 2016, Parks et al., 2017, Goss et al., 2020).

The area of permanent wetlands increased by 49,192 km² (24.88%) most likely due to sea level rise, and that of urban land increased by 39,570 km² (2.90%) (Fig. 4g & 4h) due to human land use.

4.5 Conclusions

- LAI and GPP deficits, the vegetation-centric approach, provided a different perspective into the climate-vegetation interaction, contrast to the traditional climate anomaly approach.
- Vegetation responses varied among eight landcover types with distinctive yet overlapping signatures over the spectrums of climate conditions considered, which shed a new light to the direction of landcover change (alternatives of the steady state).
- Transition matrices (short-term and long-term) of landcover types could quantify the direction and intensity of landcover changes after extreme events and over a period of time.
- The climate-vegetation interaction in the MENA region was leading to a shrubbier landscape.

496 **Acknowledgment:** Not applicable

497 **Declarations**

498 **Funding:** This research was supported by NSF 81620108010 to WF, PSC-CUNY ENHC-48-33
499 and CUNY CIRG- 80209-08 22 to CY, and Federal Work-Study Undergraduate Research
500 Assistantship to GL.

501 **Conflict of interest:** The authors declare that they have no conflict of interest.

502 **Ethical approval:** This article does not contain any studies with human participants or animals
503 performed by any of the authors.

504 **Consent to participate:** All patients included in this study gave written informed consent to
505 participate in this research.

506 **Consent for publication:** All patients included in this research gave written informed consent to
507 publish the data and images contained within this case report.

508 **Availability of data and code:** The datasets and code used and/or analyzed during the current
509 study are available from the corresponding author on reasonable request.

510

References

- Allen, C. D., Breshears, D. D., & McDowell, N. G. (2015) On underestimation of global vulnerability to tree mortality and forest die-off from hotter drought in the Anthropocene. *Ecosphere*, 6(8), art129. doi:<https://doi.org/10.1890/ES15-00203.1>
- Allen, C. D., Macalady, A. K., Chenchouni, H., Bachelet, D., McDowell, N., Vennetier, M., . . . Cobb, N. (2010) A global overview of drought and heat-induced tree mortality reveals emerging climate change risks for forests. *Forest Ecology and Management*, 259(4), 660-684. doi:10.1016/j.foreco.2009.09.001
- Anderegg, W. R., Kane, J. M., & Anderegg, L. D. (2013) Consequences of widespread tree mortality triggered by drought and temperature stress. *Nature climate change*, 3(1), 30-36.
- Assal, T. J., Anderson, P. J., & Sibold, J. (2016) Spatial and temporal trends of drought effects in a heterogeneous semi-arid forest ecosystem. *Forest Ecology and Management*, 365, 137-151. doi:<https://doi.org/10.1016/j.foreco.2016.01.017>
- Berry, S. L., & Roderick, M. L. (2002) Estimating mixtures of leaf functional types using continental-scale satellite and climatic data. *Global Ecology and Biogeography*, 11(1), 23-39. doi:<https://doi.org/10.1046/j.1466-822X.2002.00183.x>
- Budyko, M. I. (1961) The Heat Balance of the Earth's Surface. *Soviet Geography*, 2(4), 3-13. doi:10.1080/00385417.1961.10770761
- Carnicer, J., Coll, M., Ninyerola, M., Pons, X., Sánchez, G., & Peñuelas, J. (2011) Widespread crown condition decline, food web disruption, and amplified tree mortality with increased climate change-type drought. *Proceedings of the National Academy of Sciences*, 108(4), 1474-1478. doi:10.1073/pnas.1010070108
- Chapin, F., Matson, P., & Mooney, H. (2002) *Principles of Terrestrial Ecosystem Ecology*. Paper presented at the Springer New York.
- Chapin, F. S., Matson, P. A., & Vitousek, P. M. (2011) Temporal Dynamics. In *Principles of Terrestrial Ecosystem Ecology* (pp. 339-367). New York, NY: Springer New York.
- Chapin III, F. S., Matson, P. A., Vitousek, P. (2011) *Principles of Terrestrial Ecosystem Ecology* (2 ed.): Springer-Verlag New York.
- Ciais, P., Reichstein, M., Viovy, N., Granier, A., Ogée, J., Allard, V., . . . Valentini, R. (2005) Europe-wide reduction in primary productivity caused by the heat and drought in 2003. *Nature*, 437(7058), 529-533. doi:10.1038/nature03972
- Dijkstra, F., Blumenthal, D., Morgan, J., LeCain, D., & Follett, R. (2010) Elevated CO₂ effects on semi-arid grassland plants in relation to water availability and competition. *Functional Ecology*, 24(5), 1152-1161. doi:10.1111/j.1365-2435.2010.01717.x
- Foley, J., Prentice, I., Ramankutty, N., Levis, S., Pollard, D., Sitch, S., & Haxeltine, A. (1996) Foley JA, Prentice IC, Ramankutty N, Levis S, Pollard D, Sitch S, Haxeltine A. 1996. An integrated biosphere model of land surface processes, terrestrial carbon balance, and vegetation dynamics. Global Biogeochem Cycles. *Global Biogeochem. Cycles*, 10, 603-628. doi:10.1029/96GB02692
- Food and Agriculture Organization of the United Nations. 2008. The near East and North Africa: growing demand, limited resources. FAO Newsroom.

553 Friend, A. D. (2010) Terrestrial plant production and climate change. *Journal of Experimental*
554 *Botany*, 61(5), 1293-1309. doi:10.1093/jxb/erq019

555 Goss, M., Swain, D. L., Abatzoglou, J. T., Sarhadi, A., Kolden, C. A., Williams, A. P., &
556 Diffenbaugh, N. S. (2020) Climate change is increasing the likelihood of extreme autumn
557 wildfire conditions across California. *Environmental Research Letters*, 15(9), 094016.
558 doi:10.1088/1748-9326/ab83a7

559 HICKLER, T., SMITH, B., PRENTICE, I. C., MJÖFORS, K., MILLER, P., ARNETH, A., &
560 SYKES, M. T. (2008) CO₂ fertilization in temperate FACE experiments not
561 representative of boreal and tropical forests. *Global change biology*, 14(7), 1531-1542.
562 doi:https://doi.org/10.1111/j.1365-2486.2008.01598.x

563 Hirota, M., Holmgren, M., Van Nes, E. H., & Scheffer, M. (2011) Global resilience of tropical
564 forest and savanna to critical transitions. *Science*, 334(6053), 232-235.
565 doi:10.1126/science.1210657

566 Huang, K., Yi, C., Wu, D., Zhou, T., Zhao, X., Blanford, W. J., . . . Li, Z. (2015) Tipping point
567 of a conifer forest ecosystem under severe drought. *Environmental Research Letters*,
568 10(2), 024011. doi:10.1088/1748-9326/10/2/024011

569 Jackson, R. B., Cook, C. W., Pippen, J. S., & Palmer, S. M. (2009) Increased belowground
570 biomass and soil CO₂ fluxes after a decade of carbon dioxide enrichment in a warm-
571 temperate forest. *Ecology*, 90(12), 3352-3366. doi:10.1890/08-1609.1

572 Jenny, H. (1980) The soil resource. Origin and behavior. *Vegetatio*, 57(2), 102-102.
573 doi:10.1007/BF00047304

574 Jump, A. S., & Peñuelas, J. (2005) Running to stand still: adaptation and the response of plants
575 to rapid climate change. *Ecology letters*, 8(9), 1010-1020.

576 Kerr, J. T., & Ostrovsky, M. (2003) From space to species: ecological applications for remote
577 sensing. *Trends in ecology & evolution*, 18(6), 299-305.

578 Lloret, F., Enrique, G., Pérez-Ramos, I. M., Marañón, T., Saura-Mas, S., Díaz-Delgado, R., &
579 Villar, R. (2016) Climatic events inducing die-off in Mediterranean shrublands: are
580 species' responses related to their functional traits? *Oecologia*, 180(4), 961-973.

581 Liang, S. (2017) Remote Sensing of Earth's Energy Budget: An Overview of Recent Progress.
582 *Comprehensive Remote Sensing Vol. 5: Earth's Energy Budget*, 1-31.

583 Martínez-Vilalta, J., & Piñol, J. (2002) Drought-induced mortality and hydraulic architecture in
584 pine populations of the NE Iberian Peninsula. *Forest Ecology and Management*, 161(1-3),
585 247-256.

586 Matusick, G., Ruthrof, K. X., Brouwers, N. C., Dell, B., & Hardy, G. S. J. (2013) Sudden forest
587 canopy collapse corresponding with extreme drought and heat in a mediterranean-type
588 eucalypt forest in southwestern Australia. *European Journal of Forest Research*, 132(3),
589 497-510.

590 McNally, A., Arsenault, K., Kumar, S., Shukla, S., Peterson, P., Wang, S., . . . Verdin, J. P.
591 (2017) A land data assimilation system for sub-Saharan Africa food and water security
592 applications. *Scientific data*, 4(1), 1-19.

593 Myneni, R., Maggion, S., Jaquinta, J., Privette, J., Gobron, N., Pinty, B., . . . Williams, D. (1995)
594 Optical remote sensing of vegetation: modeling, caveats, and algorithms. *Remote sensing*
595 *of environment*, 51(1), 169-188.

596 Myneni, R. B., Hall, F. G., Sellers, P. J., & Marshak, A. L. (1995) The interpretation of spectral
597 vegetation indexes. *IEEE Transactions on Geoscience and Remote Sensing*, 33(2), 481-
598 486.

599 Nielsen, U. N., & Ball, B. A. (2015) Impacts of altered precipitation regimes on soil
600 communities and biogeochemistry in arid and semi-arid ecosystems. *Glob Chang Biol*,
601 21(4), 1407-1421. doi:10.1111/gcb.12789

602 Norby, R. J., Delucia, E. H., Gielen, B., Calfapietra, C., Giardina, C. P., King, J. S., . . . Oren, R.
603 (2005) Forest response to elevated CO₂ is conserved across a broad range of productivity.
604 *Proc Natl Acad Sci U S A*, 102(50), 18052-18056. doi:10.1073/pnas.0509478102

605 Parks, S. A., Holsinger, L. M., Miller, C., & Parisien, M. A. (2018) Analog-based fire regime
606 and vegetation shifts in mountainous regions of the western US. *Ecography*, 41(6), 910-
607 921.

608 Peñuelas, J., Lloret, F., & Montoya, R. (2001) Severe drought effects on Mediterranean woody
609 flora in Spain. *Forest Science*, 47(2), 214-218.

610 Seguí, P. Q., Ribes, A., Martin, E., Habets, F., & Boé, J. (2010) Comparison of three
611 downscaling methods in simulating the impact of climate change on the hydrology of
612 Mediterranean basins. *Journal of hydrology*, 383(1-2), 111-124.

613 Serra, P., Pons, X., & Saurí, D. (2008) Land-cover and land-use change in a Mediterranean
614 landscape: a spatial analysis of driving forces integrating biophysical and human factors.
615 *Applied Geography*, 28(3), 189-209.

616 Sitch, S., Smith, B., Prentice, I. C., Arneth, A., Bondeau, A., Cramer, W., . . . Sykes, M. T.
617 (2003) Evaluation of ecosystem dynamics, plant geography and terrestrial carbon cycling
618 in the LPJ dynamic global vegetation model. *Global change biology*, 9(2), 161-185.

619 Somot, S., Sevault, F., Déqué, M., & Crépon, M. (2008) 21st century climate change scenario for
620 the Mediterranean using a coupled atmosphere–ocean regional climate model. *Global and*
621 *Planetary Change*, 63(2-3), 112-126.

622 Sprintsin, M., Karnieli, A., Berliner, P., Rotenberg, E., Yakir, D., & Cohen, S. (2009) Evaluating
623 the performance of the MODIS Leaf Area Index (LAI) product over a Mediterranean
624 dryland planted forest. *International journal of Remote sensing*, 30(19), 5061-5069.

625 Stocker, T. F., Qin, D., Plattner, G. K., Tignor, M. M., Allen, S. K., Boschung, J., ... & Midgley,
626 P. M. (2014). Climate Change 2013: The physical science basis. Contribution of working
627 group I to the fifth assessment report of IPCC the intergovernmental panel on climate
628 change.

629 United Nations. Economic Commission for Africa. Subregional Office for North Africa, (1999)
630 Relevant water management and irrigation issues in North Africa. [Addis Ababa] © UN.
631 ECA.

632 Vanacker, V., Linderman, M., Lupo, F., Flasse, S., & Lambin, E. (2005) Impact of short-term
633 rainfall fluctuation on interannual land cover change in sub-Saharan Africa. *Global*
634 *Ecology and Biogeography*, 14(2), 123-135.

- Vicente-Serrano, S. M., Beguería, S., & López-Moreno, J. I. (2010) A multiscalar drought index sensitive to global warming: the standardized precipitation evapotranspiration index. *Journal of climate*, 23(7), 1696-1718.
- Vicente-Serrano, S. M., Beguería, S., Lorenzo-Lacruz, J., Camarero, J. J., López-Moreno, J. I., Azorin-Molina, C., . . . Sanchez-Lorenzo, A. (2012) Performance of drought indices for ecological, agricultural, and hydrological applications. *Earth Interactions*, 16(10), 1-27.
- Vicente-Serrano, S. M., Zouber, A., Lasanta, T., & Pueyo, Y. (2012) Dryness is accelerating degradation of vulnerable shrublands in semiarid Mediterranean environments. *Ecological Monographs*, 82(4), 407-428.
- Vidal-Macua, J. J., Ninyerola, M., Zabala, A., Domingo-Marimon, C., & Pons, X. (2017) Factors affecting forest dynamics in the Iberian Peninsula from 1987 to 2012. The role of topography and drought. *Forest Ecology and Management*, 406, 290-306.
- Vilà-Cabrera, A., Coll, L., Martínez-Vilalta, J., & Retana, J. (2018) Forest management for adaptation to climate change in the Mediterranean basin: A synthesis of evidence. *Forest Ecology and Management*, 407, 16-22.
- Wu, Z., Dijkstra, P., Koch, G. W., Peñuelas, J., & Hungate, B. A. (2011) Responses of terrestrial ecosystems to temperature and precipitation change: a meta-analysis of experimental manipulation. *Global change biology*, 17(2), 927-942. doi:<https://doi.org/10.1111/j.1365-2486.2010.02302.x>
- Yi, C., Kaiyu Lui, and Tiangjie Li,. (1996) Research on relations of soil zonal distributions with climate in the monsoon region of the eastern part of China. *ACTA PEDOLOGICA SINICA*, 32, 385-390.
- Yi, C., Ricciuto, D., Li, R., Wolbeck, J., Xu, X., Nilsson, M., . . . Arain, M. A. (2010) Climate control of terrestrial carbon exchange across biomes and continents. *Environmental Research Letters*, 5(3), 034007.
- Yi, C., Rustic, G., Xu, X., Wang, J., Dookie, A., Wei, S., . . . Nagy, Z. (2012) Climate extremes and grassland potential productivity. *Environmental Research Letters*, 7(3), 035703.
- Yi, C., Wei, S., & Hendrey, G. (2014) Warming climate extends dryness-controlled areas of terrestrial carbon sequestration. *Scientific Reports*, 4, 5472.
- Zaitchik, B. F., Evans, J. P., Geerken, R. A., & Smith, R. B. (2007) Climate and vegetation in the Middle East: Interannual variability and drought feedbacks. *Journal of climate*, 20(15), 3924-3941.
- Zhang, P., Jeong, J. H., Yoon, J. H., Kim, H., Wang, S. S., Linderholm, H. W., Fang, K., Wu, X., Chen, D. (2020) Abrupt shift to hotter and drier climate over inner East Asia beyond the tipping point. *Science*, 370(6520), 1095-1099. doi:10.1126/science.abb3368

Table 1 Regressions of annual LAI deficit and GPP deficit over climate drivers

(*** indicates $p < 0.001$; ** indicates $p < 0.01$; * indicates $p < 0.05$; ns indicates $p > 0.05$)

Dependent Variable (Y)	Independent Variable (X)	Adjusted Coefficient of Determination (R^2)	Degree of freedom (df1, df2)	Significance (p)	Intercept	Slope	
LAI Deficit	Temperature	0.53	1,150	<0.0001	18.39	-0.95	***
	Precipitation	0.62	1,150	<0.0001	-1.73	0.01	***
	TP Index	0.50	1,150	<0.0001	-2.18	21.74	***
	Dryness Index	0.59	1,150	<0.0001	8.98	-0.37	***
	SPEI	0.01	1,150	0.15	6.12	0.64	ns
GPP Deficit	Temperature	0.55	1,151	<0.0001	2925.10	-129.56	***
	Precipitation	0.55	1,151	<0.0001	255.63	1.23	***
	TP Index	0.51	1,151	<0.0001	122.75	2952.20	***
	Dryness Index	0.62	1,151	<0.0001	1644.70	-50.74	***
	SPEI	0.04	1,151	0.01	1283.74	159.26	**

TP Index: Temperature-Precipitation Index;

SPEI: Standardized Precipitation-Evapotranspiration Index;

LAI: Leaf Area Index;

GPP: Gross Primary Productivity ($\text{gCm}^{-2}\text{yr}^{-1}$)

681 **Table 2** Vegetation responses, climate conditions and area for each landcover type

	Year (n)	Forests		Savannas		CNVmosaic		Grasslands		Croplands		Shrublands		Wetlands		Urban	
		Mean	Stdev	Mean	Stdev	Mean	Stdev	Mean	Stdev	Mean	Stdev	Mean	Stdev	Mean	Stdev	Mean	Stdev
Average LAI Deficit	19	11.28	1.12	7.09	0.74	6.81	0.59	3.30	0.38	6.75	0.50	1.48	0.21	6.05	0.52	3.94	0.31
GPP Deficit ($\text{gCm}^{-2}\text{yr}^{-1}$)	19	1688.77	215.26	1489.04	199.60	1414.60	184.56	955.82	126.61	1413.75	167.39	509.14	66.47	1140.73	141.38	1082.17	130.98
Average Temperature ($^{\circ}\text{C}$)	19	10.00	0.47	12.04	0.37	12.26	0.38	11.46	0.48	13.60	0.36	17.46	0.32	14.74	0.36	14.23	0.37
Precipitation (mm)	19	1102.31	87.31	1008.09	88.36	896.59	82.67	748.08	49.67	643.01	50.51	342.24	49.24	736.53	66.18	736.76	61.23
TP Index	19	0.51	0.03	0.41	0.02	0.39	0.02	0.45	0.03	0.35	0.02	0.19	0.01	0.31	0.01	0.34	0.02
Dryness Index	19	2.37	0.54	4.68	0.88	2.11	0.34	11.87	2.05	9.54	1.11	20.77	3.95	9.17	1.69	7.75	1.06
SPEI	19	-0.34	0.61	-0.45	0.52	-0.42	0.60	-0.31	0.47	-0.47	0.42	-0.64	0.62	-0.49	0.49	-0.51	0.41
Area ($100,000 \text{ km}^2$)	18	20.72	0.37	77.21	1.21	14.83	1.55	106.04	2.03	182.91	1.51	26.42	0.78	2.19	0.18	13.80	0.12

682

683 LAI: Leaf Area Index;

684 GPP: Gross Primary Productivity ($\text{gCm}^{-2}\text{yr}^{-1}$)

685 TP Index: Temperature-Precipitation Index;

686 SPEI: Standardized Precipitation-Evapotranspiration Index;

687

Table 3a Area (km²) of vegetation type changed from 2002 (in rows) to 2003 (in columns)

(Note: only categories with more than 5,000 km² are listed)

2002\2003	Forest	Savanna	CNVmosaic	Cropland	Wetland	Urban	Grassland	Shrubland	Barren	Water
Forest	539687	17775								
Savanna	(3854)	845142					17186			
CNVmosaic			89324							
Cropland				1518885			10727			
Wetland					16949					
Urban						104781				
Grassland		9834		15729			1025717	7298		
Shrubland							8570	343054		
Barren							9879	12801	6367401	
Water										11527805

Table 3b Area (km²) of vegetation type changed from 2001 (in rows) to 2018 (in columns)

(Note: only categories with more 20,000 km² are listed)

2001\2018	Forest	Savanna	CNVmosaic	Cropland	Wetland	Urban	Grassland	Shrubland	Barren	Water
Forest	500508	55948								
Savanna	54475	716599	21938				57586			
CNVmosaic			71082							
Cropland		22846	29875	1390808			63253	22897		
Wetland					14708					
Urban						104735				
Grassland		106902		77943			814238	38938		
Shrubland							37853	280748		
Barren							31652	52376	6309837	
Water										11525705

Figure Legends:

Figure 1 A map of vegetated areas with no change of landcover types during 2001-2018 (500m resolution) in Mediterranean Europe and North Africa (MENA, 20°W-45°E & 20°N-45°N). Pixels that experienced landcover change were in white color. Annual landcover classification is defined by the MODIS Landcover CMG (MCD12Q1) product's International Geosphere Biosphere Programme (IGBP) global vegetation classification scheme (https://lpdaac.usgs.gov/products/modis_products_table/mcd12c1). The primary landcover scheme identifies 17 classes, and were grouped into 8 categories: forests (class 1-5), shrubland (class 6-7), savannas (class 8-9), grasslands (class 10), permanent wetlands (class 11), croplands (class 12), urban and build-up (class 13) and crop+natural vegetation mosaic (class 14). The non-vegetated areas (water and barren land) were excluded from relevant analyses.

Figure 2 Conceptual workflow of this study. Step 1, eight vegetated landcover types with no change over 18 years were extracted to conduct multiple regression analyses; Step 2, regressions of vegetation structural (LAI deficit) and functional (GPP deficit) responses over climate drivers (including temperature (°C, T), precipitation (mm, P), TP index, Dryness index, SPEI (Standardized Precipitation Evapotranspiration Index)) were conducted over all vegetated pixels and within each vegetation type; Step 3, 10 regression analyses and probability density function (PDF) curves of 7 variables shedding new light on future landcover changes. Step 4, direction and intensity of landcover change were quantified using time series and transition matrices.

Figure 3 Regressions of vegetation structural (LAI deficit) and functional (GPP deficit) responses over climate drivers (i.e. annual average temperature (°C, T), accumulative precipitation (mm, P), Temperature-Precipitation (TP) index, Dryness index, SPEI (Standardized Precipitation Evapotranspiration Index)) for all vegetated pixels (black lines, solid if $p < 0.01$, dotted if $p > 0.05$) and within each of 8 vegetation types (color coded with 95% confidence intervals in gray); For each panel, probability density function (PDF) curves were plotted with the same scheme of color code, horizontal on the top for the climate variables (T, P, TP index, Dryness or SPEI) and vertical on the right for the response variables (LAI or GPP deficit).

Figure 4 Total area (km²) of eight vegetation types over the period of Jan. 1, 2001-Dec. 31, 2018 in Mediterranean Europe and North Africa (MENA, 20°W-45°E & 20°N-45°N).

Figure 5 A map of vegetated areas with changes of landcover types during 2001-2018 (500m resolution) in Mediterranean Europe and North Africa (MENA, 20°W-45°E & 20°N-45°N).

Figure 1

730

731

732 **Figure 2**

733

Fig.3 a1

Fig.3 b1

Fig.3 a2

Fig.3 b2

Fig.3 a3

Fig.3 b3

Fig.3 a4

Fig.3 b4

Fig.3 a5

Fig.3 b5

Fig. 4

Figure 5

Figures

Figure 1

A map of vegetated areas with no change of landcover types during 2001-2018 (500m resolution) in Mediterranean Europe and North Africa (MENA, 20°W-45°E & 20°N-45°N). Pixels that experienced landcover change were in white color. Annual landcover classification is defined by the MODIS Landcover CMG (MCD12Q1) product's International Geosphere Biosphere Programme (IGBP) global vegetation classification scheme (https://lpdaac.usgs.gov/products/modis_products_table/mcd12c1). The primary landcover scheme identifies 17 classes, and were grouped into 8 categories: forests (class 1-5), shrubland (class 6-7), savannas (class 8-9), grasslands (class 10), permanent wetlands (class 11), croplands (class 12), urban and build-up (class 13) and crop+natural vegetation mosaic (class 14). The non-vegetated areas (water and barren land) were excluded from relevant analyses.

Figure 2

Conceptual workflow of this study. Step 1, eight vegetated landcover types with no change over 18 years were extracted to conduct multiple regression analyses; Step 2, regressions of vegetation structural (LAI deficit) and functional (GPP deficit) responses over climate drivers (including temperature ($^{\circ}\text{C}$, T), precipitation (mm, P), TP index, Dryness index, SPEI (Standardized Precipitation Evapotranspiration Index)) were conducted over all vegetated pixels and within each vegetation type; Step 3, 10 regression analyses and probability density function (PDF) curves of 7 variables shedding new light on future landcover changes. Step 4, direction and intensity of landcover change were quantified using time series and transition matrices.

Fig.3 a1

Fig.3 b1

Fig.3 a2

Fig.3 b2

Fig.3 a3

Fig.3 b3

Fig.3 a4

Fig.3 b4

Fig.3 a5

Fig.3 b5

Figure 3

Regressions of vegetation structural (LAI deficit) and functional (GPP deficit) responses over climate drivers (i.e. annual average temperature ($^{\circ}\text{C}$, T), accumulative precipitation (mm, P), Temperature-Precipitation (TP) index, Dryness index, SPEI (Standardized Precipitation Evapotranspiration Index)) for all vegetated pixels (black lines, solid if $p < 0.01$, dotted if $p > 0.05$) and within each of 8 vegetation types (color coded with 95% confidence intervals in gray); For each panel, probability density function (PDF)

curves were plotted with the same scheme of color code, horizontal on the top for the climate variables (T, P, TP index, Dryness or SPEI) and vertical on the right for the response variables (LAI or GPP deficit).

Figure 4

Total area (km²) of eight vegetation types over the period of Jan. 1, 2001-Dec. 31, 2018 in Mediterranean Europe and North Africa (MENA, 20°W-45°E & 20°N-45°N).

Figure 5

A. map of vegetated areas with changes of landcover types during 2001-2018 (500m resolution) in Mediterranean Europe and North Africa (MENA, 20°W-45°E & 20°N-45°N).