

The Impact of Ossification Spread on Cervical Spine Function in Patients With Ossification of The Posterior Longitudinal Ligament

Keiichi Katsumi (✉ kkatsu_os@yahoo.co.jp)

Spine Center, Department of Orthopedic Surgery, Niigata Central Hospital

Takashi Hirai

Tokyo Medical and Dental University

Toshitaka Yoshii

Tokyo Medical and Dental University

Satoshi Maki

Chiba University

Kanji Mori

Shiga University of Medical Science

Narihito Nagoshi

Keio University

Soraya Nishimura

Keio University

Kazuhiro Takeuchi

Okayama Medical Center

Shuta Ushio

Tokyo Medical and Dental University

Takeo Furuya

Chiba University

Kei Watanabe

Niigata University Medical and Dental Hospital

Norihiro Nishida

Yamaguchi University

Kota Watanabe

Keio University

Takashi Kaito

Osaka University

Satoshi Kato

Kanazawa University

Katsuya Nagashima

University of Tsukuba

Masao Koda

University of Tsukuba

Kenyu Ito

Nagoya University

Shiro Imagama

Nagoya University

Yuji Matsuoka

Tokyo Medical University

Kanichiro Wada

Hirosaki University

Atsushi Kimura

Jichi Medical University

Tetsuro Ohba

University of Yamanashi

Hiroyuki Katoh

Tokai University

Yukihiro Matsuyama

Hamamatsu University School of Medicine

Hiroshi Ozawa

Tohoku Medical and Pharmaceutical University

Hiroataka Haro

University of Yamanashi

Katsushi Takeshita

Jichi Medical University

Masahiko Watanabe

Tokai University

Morio Matsumoto

Keio University

Masaya Nakamura

Keio University

Masashi Yamazaki

University of Tsukuba

Atsushi Okawa

Tokyo Medical and Dental University

Yoshiharu Kawaguchi

University of Toyama

Research Article

Keywords: ossification of posterior longitudinal ligament, progression, JOACMEQ, cervical spine function, whole spine, bridge formation, neck visual analog scale, vertebral body

Posted Date: February 22nd, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-216266/v1>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Scientific Reports on July 12th, 2021. See the published version at <https://doi.org/10.1038/s41598-021-93602-0>.

Abstract

Ossification of the posterior longitudinal ligament (OPLL) is a progressive disease. The bridging of ossified lesions to the vertebral body gradually increases, thereby decreasing the mobility of the cervical spine; thus, cervical spine function may decrease over time. However, cervical spine function in patients with cervical OPLL has not been evaluated in large prospective studies. Therefore, we conducted a prospective multicenter study to clarify whether ossification spread can influence cervical spine function and quality of life (QOL) in patients with cervical OPLL. In total, 238 patients (162 men, 76 women; mean age, 63.9 years) were enrolled from 16 institutions. Each patient underwent whole spine computed tomography and was evaluated cervical spine function and QOL using the Japanese Orthopaedic Association Cervical Myelopathy Evaluation Questionnaire (JOACMEQ). In the multivariate regression analysis, a higher neck VAS score and a larger number of bridge formations of OPLL in the whole spine were significant predictors of adverse outcomes related to cervical spine function. This is the first prospective multicenter study to reveal the impact of ossification spread on cervical spine function. These findings are important to understand the natural course of OPLL and can serve as controls when evaluating postoperative cervical spine function.

Introduction

Ossification of the posterior longitudinal ligament (OPLL) was first described by Key in 1838,¹ and the concept of OPLL has been widely used since Tsukimoto published an autopsy case report on this subject in 1960.² OPLL is characterized by the replacement of ligamentous tissue by ectopic bone formation and has been recognized as one of the main causes of cervical myelopathy.²⁻⁴ Although several reports have demonstrated the pathophysiology of OPLL, its clinical features, and outcome data of surgical treatments,³⁻⁵ there are few reports on cervical spine function in patients with OPLL. Because OPLL is a progressive disease, the bridging of ossified lesions to the vertebral body gradually increases, thereby decreasing the mobility of the cervical spine; thus, cervical spine function and quality of life (QOL) may decrease over time. Understanding cervical spine function in patients with OPLL who had been treated conservatively is important in evaluating the natural history of OPLL and the side effects of surgical treatment, in particular, of posterior surgery.

Although the Japanese Orthopaedic Association (JOA) score⁶ is used as a functional assessment of cervical myelopathy worldwide, the scoring system does not include cervical spine function and neck pain, both of which seriously impact patients' QOL (Supplemental Table 1). In 2007, the JOA established a self-administered questionnaire, the Japanese Orthopaedic Association Cervical Myelopathy Evaluation Questionnaire (JOACMEQ), as a new outcome measurement tool for patients with cervical myelopathy, which included measurements of cervical spine function such as neck pain, stiff neck, disability, and QOL (Supplemental Table 2).⁷

In this study, we conducted a prospective multicenter study of patients with cervical OPLL who had been treated conservatively to evaluate the existence and distribution of OPLL in the whole spine using

computed tomography (CT). Furthermore, we investigated cervical spine function and QOL using the JOACMEQ. The purpose of this study was to clarify whether ossification spread can influence cervical spine function and QOL in patients with cervical OPLL.

Materials And Methods

This prospective multicenter study was conducted at the 16 institutions of the Japanese Multicenter Research Organization for Ossification of the Spinal Ligament established by the Japan Ministry of Health, Labour and Welfare. This study was approved by the institutional review board of each institution and performed in accordance with approved guidelines and in compliance with the principles of the Declaration of Helsinki. Informed consent was obtained from all patients before enrollment. This study included patients with cervical OPLL diagnosed by neck radiographs who had symptoms such as neck pain, numbness in the upper or lower extremities, clumsiness, or gait disturbance. Patients who had a history of previous cervical surgeries and were younger than 20 years were excluded. Each patient underwent whole spine CT and was evaluated based on the JOACMEQ. A total of 238 patients (162 men, 76 women; mean age, 63.9 years; range, 36–92 years) were enrolled prospectively from September 2015 to December 2017.

Cervical spine function and radiological examinations including the extent of ossified lesions obtained from CT images of the whole spine in conservatively treated OPLL patients were conducted. In addition, we examined the effects of the spread of ossification on cervical spine function.

Clinical examinations.

Age, sex, height, weight, body mass index (BMI), and presence of diabetes mellitus (DM) were recorded as basic clinical data. The clinical status was evaluated using the JOA score and JOACMEQ. The JOA score is a 17-point instrument where points are assigned based on the ratings of motor function, sensory function, and urinary bladder function that is used as a functional assessment of cervical myelopathy worldwide. The JOACMEQ, which is a self-administered questionnaire, includes 24 questions corresponding to five domains: Q1 (cervical spine function), Q2 (upper extremity function), Q3 (lower extremity function), Q4 (bladder function), and Q5 (quality of life). The score of each domain ranges from 0 to 100 and is proportional to the patients' clinical conditions; a normal score is 100 points. Cervical spine function was evaluated using four questions: Q1-1 and Q1-2 (neck extension), Q1-3 (neck rotation), and Q1-4 (neck flexion) from domain Q1 of the JOACMEQ (Table 1). The score for the cervical spine function domain was calculated as follows (points): $Q1-1 \times 20 + Q1-2 \times 10 + Q1-3 \times 15 + Q1-4 \times 5 - 50$. The 100-mm visual analog scale (VAS) was used to evaluate the degree of pain or stiffness in the neck or shoulders.

Radiological examinations.

The OPLL incidence in the whole spine from the clivus to S1 was evaluated using mid-sagittal CT images. Observed ossified lesions were recorded for each vertebral body and at the intervertebral disc

level. The ossification index for OPLL (OP-index) was defined as the sum of the levels of ossification at the vertebral bodies and the intervertebral discs according to the method described in previous reports.⁸⁻¹⁰ We recorded the OP-index not only for the cervical spine (0–14; C1, C1/C2...C7, C7/T1) but also for the whole spine (0–49; C1... S1). Using a previously reported method,⁸ we categorized patients into three groups according to the cervical OP-index: Grade 1 (cervical OP-index, ≤ 5), Grade 2 (cervical OP-index, 6–9), and Grade 3 (cervical OP-index, ≥ 10). Furthermore, when the ossified lesions formed a bridge with the posterior border of the adjacent vertebral body, it was judged as “bridge formation.”⁹ We examined the bridge formation of OPLL from the cervical spine (0–7; C1/C2...C7/T1) to the whole spine (0–24; C1/C2...L5/S1) (Figure 1). All CT data were evaluated by six experienced spine surgeons (KK, KM, SM, NN, SU, and KT). Before the analysis, all observers read the same CT images of 20 patients to confirm interobserver intra-class correlation coefficients (ICCs). The mean interobserver ICC was 0.83 (0.79–0.85), indicating a substantial agreement and consistency with previous results.⁸

Statistical analyses.

Data were expressed as mean and standard deviations. Differences between groups were evaluated using Mann-Whitney U-tests. Kruskal-Wallis analysis was applied to compare the four questions of the cervical spine function domain in the JOACMEQ, the three groups according to cervical OP-index grade, and the five different age groups. Spearman’s rank correlation coefficients between the clinical or radiological findings and cervical spine function were calculated. We categorized patients into two groups based on points for cervical spine function: good cervical function (≥ 50 points) and poor cervical function (< 50 points), and multiple logistic regression analysis was performed to identify variables independently associated with cervical spine function. First, the significance of the variables was evaluated using univariate analysis. Next, variables with $p < 0.1$ were subjected to multivariate analysis and $p < 0.05$ was considered significant. Statistical analyses were performed using SPSS software (version 25; IBM Corp., Armonk, NY, USA).

Results

Clinical examinations.

Height, weight, and BMI were 162.7 ± 9.7 cm (range, 138–187 cm), 69.1 ± 15.2 kg (range, 32–114 kg), and 25.9 ± 4.5 kg/m² (range, 16.4–45.2 kg/m²), respectively. Fifty-nine of the 238 patients (24.8%) had DM, and the cervical JOA score was 12.3 ± 3.4 points (range, -2–17 points). The prevalence of neck pain was 60.0% (142/238 patients), and the neck or shoulder VAS score was 38.8 ± 31.3 (range, 0–100). The scores for each question of Q1-1, Q1-2, Q1-3, and Q1-4 were 2.4 ± 0.7 , 2.5 ± 0.7 , 2.0 ± 0.8 , and 2.6 ± 0.7 , respectively (Figure 2). The score for Q1-3 was significantly lower than that of the other questions ($p < 0.05$). The score for the domain of cervical spine function was 65.9 ± 28.5 points. According to the cervical OP-index grading, the scores of cervical spine function were 67.8 ± 27.1 in Grade 1, 66.8 ± 29.4 in Grade 2, and 57.0 ± 30.4 in Grade 3. There were no significant differences between the cervical OP-index grading groups.

Radiological examinations.

The OP-index of the cervical spine and whole spine was 6.0 ± 3.2 (range, 1–14) and 8.6 ± 6.4 (range, 1–38), respectively. The cervical OP-index was Grade 1 in 116 patients (48.7%), Grade 2 in 87 patients (36.6%), and Grade 3 in 35 patients (14.7%). The number of bridge formations of OPLL in the cervical spine and the whole spine were 0.7 ± 1.1 (range, 0–6) and 1.4 ± 0.3 (range, 0–17), respectively.

Sex difference and age-related change.

The OP-index and bridge formations of OPLL for the whole spine of women were significantly higher than those of men (both $p < 0.05$) (Table 2). Patients were divided into five groups based on age categories: ≤ 49 years ($n=37$), 50–59 years ($n=48$), 60–69 years ($n=66$), 70–79 years ($n=71$), and ≥ 80 years ($n=16$). Because of the small number of patients, four men and one woman in their 30s and one man and one woman in their 90s were included in the groups of ≤ 49 years and ≥ 80 years, respectively. Cervical spine function was significantly lower in the 70–79-years and ≥ 80 -years age groups than in the ≤ 49 -years age group (both $p < 0.05$). In the questions of the cervical spine function domain, the score for each question tended to decline with age; however, only Q1-2 showed a significant difference ($p < 0.05$). Moreover, the score of Q1-3 was significantly lower than those of Q1-2 and Q1-4 in all age groups except in the ≥ 80 -years age group (all $p < 0.05$). Bridge formation of OPLL in the cervical spine was significantly higher in the ≥ 80 -years age group than in the ≤ 49 -years and 50–59-years age groups (both $p < 0.05$) (Table 3).

Correlation between cervical spine function and other clinical and radiological parameters.

Cervical spine function was negatively correlated with age and neck VAS ($r = -0.23$, $p < 0.001$; $r = -0.45$, $p < 0.001$; respectively) and positively correlated with the JOA score ($r = 0.31$, $p < 0.001$). In the radiological findings, there was a negative correlation between cervical spine function and bridge formation of OPLL in the whole spine ($r = -0.15$, $p < 0.05$).

Logistic regression analysis of factors adversely associated with cervical spine function.

The univariate analysis demonstrated that there were significant correlations with age, JOA score, neck VAS, cervical OP-index grade, OP-index, and bridge formation of OPLL in the cervical and whole spine (all $p < 0.05$) (Table 4). In the multivariate regression analysis, a higher neck VAS score (odds ratio [OR]=0.978, 95% confidence interval [CI]=0.967–0.988, $p = 0.000$) and a larger number of bridge formations of OPLL in the whole spine (OR=0.820, 95% CI=0.713–0.943, $p = 0.005$) were independent factors that adversely affected cervical spine function.

Discussion

The analysis of treatment outcomes using patient-based assessments is becoming standard practice. The JOACMEQ is a patient-based assessment for evaluating cervical compressive myelopathy, and several reports using this scoring system have been published to date.^{7,11-14} Regarding the QOL of

patients with OPLL, several reports on postoperative axial neck pain have been published^{12,15,16}; however, to the best of our knowledge, detailed reports of cervical spine function based on large prospective multicenter series are lacking. Comprehensive evaluations of cervical spine function in patients with OPLL who underwent conservative treatment are important to understand the natural course of OPLL and can serve as a control when evaluating cervical spine function after surgery.

In this study, the score of cervical spine function was 65.9 points (age, 63.9 years; JOA score, 12.3 points). Tanaka et al.¹⁷ examined the standard value of the JOACMEQ in 1629 healthy Japanese volunteers and reported that the score of cervical spine function was >90 points in individuals in their 20s to 60s, whereas the score decreased to 80 and 70 points, respectively, in those in their 70s and 80s. Moreover, Ohya et al.¹³ reported that cervical spine function in 44 patients with OPLL (age, 63.8 years; preoperative JOA score, 10.9 points) was 55.5 points preoperatively and 64.9 points postoperatively. Our results were also lower than those of the healthy volunteers reported by Tanaka et al.¹⁷ and similar to the findings of Ohya et al.¹³ Even in the patients aged ≤ 49 years, which was the group with the highest points in this study, their points were equivalent to those of healthy volunteers in their 80s as reported by Tanaka et al. In the present study, the score of Q1-3 was significantly lower than that of the other questions in the overall cohort and many age groups. In general, the range of motion (ROM) in the cervical spine decreases due to age-related and degenerative changes; however, flexion movement is considered to be less affected than extension movement.^{18,19} Yuan et al.²⁰ reported that patients with cervical OPLL had decreased cervical ROM including flexion, extension, and rotation compared to the normal controls. Our study revealed that rotation movement was the most limited, and flexion and extension movements were generally comparable. Whether this result is due to a decrease in ROM, neck pain, or neurologic compression remains to be solved.

This study also investigated age-related changes in parameters and found that cervical spine function tended to decline with age from middle age. Since the neck VAS scores did not differ significantly by age, we considered that a decline in cervical spine function was caused by the decreasing mobility of the cervical spine. As several studies have demonstrated the progression of ossification in many patients with OPLL,^{3-5,21} we considered that the OP-index and bridge formation of OPLL increased with age. However, although there appeared to be an increase in the tendency of bridge formations with age, a significant difference was found only in the bridge formations in the cervical spine. Thus, we speculated that factors other than age may be involved in the onset and progression of OPLL.

In the correlation analysis, age, neck VAS, and the JOA score were significantly correlated with cervical spine function. In addition, bridge formation of OPLL was significantly correlated with the whole spine but not with the cervical spine. Furthermore, following logistic regression analysis, only two independent factors were significantly associated with adverse cervical spine function namely, a higher neck VAS score and a larger number of bridge formations of OPLL in the whole spine.

From the results of this study, neck pain caused by degenerative changes and/or neurologic deficits had a greater effect on cervical spine function in the clinical findings. Furthermore, in the radiological findings,

it was also revealed that 1) the spread of bridge formations of the vertebrae rather than that of ossified lesions at the vertebra and intervertebral discs 2) in the whole spine rather than in the cervical spine had a greater effect on cervical spine function. Fujimori et al.²² evaluated the intervertebral segmental ROM in patients with OPLL at the intervertebral discs between bridging type (ossification bridging intervertebral segment with bony union) and non-bridging type using functional CT and reported that the intervertebral ROMs were 0.3° in the bridging type and 4.9° in the non-bridging type. We consider that the structural impairment caused by bridge formation of OPLL has an impact on the deterioration of cervical spine function. According to previous reports, cervical alignment is influenced by the global spine alignment, and concomitant thoracic spine mobility is necessary to produce the complete range of movements at the cervical spine.^{23,24} Reduced mobility in the thoracic spine has also been reported to cause neck pain and dysfunction.²⁵ Therefore, when assessing the impact of OPLL on cervical spine function, we believe it is important to pay attention to bridge formation of OPLL, not only in the cervical spine but also in the whole spine. In summary, cervical spine dysfunction can be divided into two categories namely, functional impairment due to pain radiating to the neck and shoulders and structural impairment due to decreased mobility of the whole spine caused by bridge formations of ossified lesions spreading to the vertebral body. It is clinically important to recognize these impairments when evaluating cervical spine function in patients with OPLL.

This study has several limitations. First, although this study was a prospective nationwide multicenter survey, participants in the study were not randomly selected from the general population which could have created a bias. However, it is problematic to perform whole spine CT in asymptomatic healthy individuals due to radiologic exposure. Second, this study used a cross-sectional design; therefore, longitudinal studies are necessary to establish causal relationships between cervical spine function and the observed associations. Finally, we did not evaluate cervical spine radiographs in the radiological examination. Therefore, we could not examine the relationship between cervical spine function and cervical ROM and alignment. Further studies are necessary to clarify these issues. Nevertheless, we believe that our findings provide important information that highlights the impact of the spread of ossification on cervical spine function in patients with OPLL.

Conclusions

The present study revealed the impact of the spread of ossification on cervical spine function, including the impact of sex differences and age-related changes, in a prospective nationwide multicenter study. A higher neck VAS score and a larger number of bridge formations of OPLL in the whole spine were significant predictors of adverse outcomes related to cervical spine function. These findings are important to understand the natural course of OPLL and can serve as a control when evaluating cervical spine function after surgery.

Declarations

1. Ethics approval and consent to participate

Written informed consent was obtained from each study participant before enrollment at each institution. The study protocol was approved by the ethics committee at each participating institution i.e., Niigata Central Hospital : et2020-03, Niigata University Medical and Dental General Hospital : 2015-2525, Tokyo Medical and Dental University: M2000-2068, Chiba University Graduate School of Medicine: 1654, Shiga University of Medical Science: 25-108, Keio University: 20180096, National Hospital Organization Okayama Medical Center: H28-71, Yamaguchi University Graduate School of Medicine; H27-117, Osaka University Graduate School of Medicine: 15153-2, Kanazawa University: 2015-087, University of Tsukuba: H26-108, Nagoya University Graduate School of Medicine; 2005-0354, Hirosaki University Graduate School of Medicine; 2015-209, Jichi Medical University; A17-106, University of Yamanashi; 1239, Tokai University School of Medicine:13R-297, Hamamatsu University School of Medicine; 15-279, Tohoku Medical and Pharmaceutical University; 2016-2-041, University of Toyama, Toyama; 25-42

2. Consent to publish

Not applicable.

3. Availability of data and materials

The study data and details of materials used may be made available upon reasonable request by sending an e-mail to the first author.

4. Competing interests

The authors declare that they have no competing interests.

5. Funding

This work was supported by Health and Labour Science Research grants (201610008B) and by a grant from the Japan Agency for Medical Research and Development (16ek0109136h0002).

6. Author contributions

KK, TH, TY, KM, SU, SM, NN, MM, MN, KW, and YK designed the study; KK, TH, TY, KM, SU, SM, NN, KT, TF, KW, NN, KW, TK, SK, KN, MK, KI, SI, YM, KW, AK, TO, HK, YM, HO, and YK collected the data; KK, KM, SU, SM, NN and KT analyzed and interpreted the data; KK, KW, and YK wrote the initial draft of the manuscript; KK and TH performed the statistical analyses. KK, TH, TY, SM, KM, NN, SN, KT, SU, TF, KW, NN, KW, TK, SK, KN, MK, KI, SI, YM, KW, AK, TO, HK, YM, HO, HH, KT, MW, MN, MY, AO, and YK participated in revising the manuscript. MM, MN, MY, AO, and YK supervised the study. MM, MY, and AO acquired the funding. All authors read and approved the final manuscript.

References

1. Key GA. On paraplegia depending on the ligament of the spine. *Guy's Hosp Rep.* 3, 17–34 (1838).
2. Tsukimoto H. An autopsied case of compression myelopathy with a callus formation in the cervical spinal canal. *Nihon Geka Hokan* 29, 1003–1007 (in Japanese) (1960).
3. Iwasaki M, et al. Surgical strategy for cervical myelopathy due to ossification of the posterior longitudinal ligament: Part 1: Clinical results and limitations of laminoplasty. *Spine* 32, 647–653, doi:10.1097/01.brs.0000257560.91147.86 (2007).
4. Matsunaga S, et al. Radiographic predictors for the development of myelopathy in patients with ossification of the posterior longitudinal ligament: A multicenter cohort study. *Spine* 33, 2648–2650, doi:10.1097/brs.0b013e31817f988c (2008).
5. Fragen KM, Cox JB, Hoh DJ. Does ossification of the posterior longitudinal ligament progress after laminoplasty? Radiographic and clinical evidence of ossification of the posterior longitudinal ligament lesion growth and the risk factors for late neurologic deterioration. *J Neurosurg Spine* 17, 512–524, doi:10.3171/2012.9.SPINE12548 (2012).
6. Japanese Orthopaedic Association. Scoring system for cervical myelopathy. *J Jpn Orthop Assoc.* 68, 490–503 (in Japanese) (1994).
7. Fukui M, et al. An outcome measure for patients with cervical myelopathy: Japanese Orthopaedic Association Cervical Myelopathy Evaluation Questionnaire (JOACMEQ): Part 1. *J Orthop Sci.* 12, 227–240, doi:10.1007/s00776-007-1118-1 (2007).
8. Hirai T, et al. Prevalence and distribution of ossified lesions in the whole spine of patients with cervical ossification of the posterior longitudinal ligament a multicenter study (JOSL CT study). *PLoS One* 11, e0160117, doi:10.1371/journal.pone.0160117 (2016).
9. Kawaguchi Y, et al. Ossification of the posterior longitudinal ligament in not only the cervical spine, but also other spinal regions: Analysis using multidetector computed tomography of the whole spine. *Spine* 38, E1477–E1482, doi:10.1097/BRS.0b013e3182a54f00 (2013).
10. Yoshii T, et al. Co-existence of ossification of the nuchal ligament is associated with severity of ossification in the whole spine in patients with cervical ossification of the posterior longitudinal ligament-A multi-center CT study. *J Orthop Sci.* 24, 35–41, doi:10.1016/j.jos.2018.08.009 (2019).
11. Yoon ST, et al. Predictive factors affecting outcome after cervical laminoplasty. *Spine* 38(22 Suppl 1), S232–S252, doi:10.1097/BRS.0b013e3182a7eb55 (2013).
12. Fujiwara H, et al. Impact of cervical sagittal alignment on axial neck pain and health-related quality of life after cervical laminoplasty in patients with cervical spondylotic myelopathy or ossification of the posterior longitudinal ligament: A prospective comparative study. *Clin Spine Surg.* 31, E245–E251, doi:10.1097/bsd.0000000000000619 (2018).
13. Ohya J, et al. Patient satisfaction with posterior decompression surgery for cervical ossification of the posterior longitudinal ligament: Prognostic radiographic factors and patient-reported outcomes for the effectiveness of surgical treatment. *World Neurosurg.* 96, 272–279, doi:10.1016/j.wneu.2016.09.011 (2016).

14. Chien A, et al. Translation, cross-cultural adaptation, and validation of a Chinese version of the Japanese Orthopaedic Association Cervical Myelopathy Evaluation Questionnaire. *Spine* 39, 963–970, doi:10.1097/BRS.0000000000000313 (2014).
15. Kawaguchi Y, et al. Relationship between postoperative axial symptoms and the rotation angle of the cervical spine after laminoplasty. *Eur J Orthop Surg Traumatol.* 23(Suppl 1), S53–S58, doi:10.1007/s00590-013-1219-9 (2013).
16. Wang SJ, Jiang SD, Jiang LS, Dai LY. Axial pain after posterior cervical spine surgery: A systematic review. *Eur Spine J.* 20, 185–194, doi:10.1007/s00586-010-1600-x (2011).
17. Tanaka N, et al. An outcome measure for patients with cervical myelopathy: The Japanese Orthopaedic Association Cervical Myelopathy Evaluation Questionnaire (JOACMEQ): An average score of healthy volunteers. *J Orthop Sci.* 19, 33–48, doi:10.1007/s00776-013-0494-y (2014).
18. Swinkels RA, Swinkels-Meewisse IE. Normal values for cervical range of motion. *Spine* 39, 362–367, doi:10.1097/BRS.0000000000000158 (2014).
19. Machino M, et al. Age-related and degenerative changes in the osseous anatomy, alignment, and range of motion of the cervical spine: A comparative study of radiographic data from 1016 patients with cervical spondylotic myelopathy and 1230 asymptomatic subjects. *Spine* 41, 476–482, doi:10.1097/BRS.0000000000001237 (2016).
20. Yuan W, et al. Postoperative three-dimensional cervical range of motion and neurological outcomes in patients with cervical ossification of the posterior longitudinal ligament: Cervical laminoplasty versus laminectomy with fusion. *Clin Neurol Neurosurg.* 134, 17–23, doi:10.1016/j.clineuro.2015.04.004 (2015).
21. Katsumi K, et al. Natural history of the ossification of cervical posterior longitudinal ligament: A three-dimensional analysis. *Int Orthop.* 42, 835–842, doi:10.1007/s00264-017-3667-z (2016).
22. Fujimori T, et al. Three-dimensional measurement of intervertebral range of motion in ossification of the posterior longitudinal ligament: are there mobile segments in the continuous type? *J Neurosurg Spine* 17, 74–81, doi: 10.3171/2012.3.SPINE111083 (2012).
23. Pepke W, et al. Cervical spine alignment following surgery for adolescent idiopathic scoliosis (AIS): a pre-to-post analysis of 81 patients. *BMC Surgery* 19, 7, doi:10.1186/s12893-019-0471-2 (2019).
24. Boissiere L, et al. Cervical spine balance: postoperative radiologic changes in adult scoliosis surgery. *Eur Spine J.* 24, 1356–1361, doi:10.1007/s00586-015-3854-9 (2015).
25. Joshi S, Balthillaya G, Neelapala YV. Thoracic posture and mobility in mechanical neck pain population: A review of the literature. *Asian Spine J.* 13, 849-860, doi:10.31616/asj.2018.0302 (2019).

Tables

Table 1. The domain of cervical spine function in the JOACMEQ

Questionnaire		Score
Q1-1	While in the sitting position, can you look up at the ceiling by tilting your head upward?	
	1) Impossible	1
	2) Possible to some degree (with some effort)	2
	3) Possible without difficulty	3
Q1-2	Can you drink a glass of water without stopping despite your neck symptoms?	
	1) Impossible	1
	2) Possible to some degree	2
	3) Possible without difficulty.	3
Q1-3	While in the sitting position, can you turn your head toward a person who is seated to the side but behind you and speak to that person while looking at his/her face?	
	1) Impossible	1
	2) Possible to some degree	2
	3) Possible without difficulty	3
Q1-4	Can you look at your feet when you go downstairs?	
	1) Impossible	1
	2) Possible to some degree	2
	3) Possible without difficulty	3

JOACMEQ, Japanese Orthopaedic Association Cervical Myelopathy Evaluation Questionnaire

The score for the cervical spine function domain was calculated as follows (points): $Q1-1 \times 20 + Q1-2 \times 10 + Q1-3 \times 15 + Q1-4 \times 5 = 50$.

Table 2. Sex differences in parameters

Characteristic	Male (n=162)	Female (n=76)	p-value
Age (years)	64.0 ± 12.7	63.6 ± 11.5	0.51
BMI (kg/m ²)	25.9 ± 4.3	26.1 ± 4.8	0.90
JOA score (points)	12.0 ± 3.5	12.8 ± 3.1	0.19
Neck VAS score	38.8 ± 32.3	38.8 ± 29.2	0.78
Cervical spine function (points)	65.1 ± 29.4	65.9 ± 27.5	0.96
Questions of Q1 domain			
Q1-1	2.3 ± 0.7	2.4 ± 0.7	0.25
Q1-2	2.5 ± 0.7	2.5 ± 0.7	0.61
Q1-3	2.1 ± 0.8	2.0 ± 0.8	0.83
Q1-4	2.6 ± 0.6	2.5 ± 0.8	0.09
C-OP-index	6.0 ± 3.1	6.0 ± 3.4	0.81
W-OP-index	7.5 ± 5.1	10.9 ± 8.1	0.006**
Cervical OP-index grade	1.7 ± 0.7	1.6 ± 0.8	0.44
Bridge formation of C-OPLL	0.7 ± 1.2	0.7 ± 1.1	0.76
Bridge formation of W-OPLL	1.1 ± 1.8	2.0 ± 3.0	0.014*

Data are presented as mean ± standard deviation.

BMI, body mass index; JOA, Japanese Orthopaedic Association; VAS, visual analog scale (100-mm)

Q1 domain, cervical spine function domain of the Japanese Orthopaedic Association Cervical Myelopathy Evaluation Questionnaire; C-, cervical; W-, whole spine; OP-index, Ossification index for OPLL; bridge formation, ossified lesions forming a bridge with the posterior border of the adjacent vertebral body

*p<0.05; **p<0.01

Table 3. Age-related changes in parameters

Characteristic	≤49 years (n=37)	50–59 years (n=48)	60–69 years (n=66)	70–79 years (n=71)	≥80 years (n=16)	p-value
Age (years)	43.8 ± 3.5	54.8 ± 2.8	65.4 ± 2.8	74.6 ± 2.7	84.3 ± 3.3	0.000 ^{***}
Male sex, no. (%)	26 (70.3)	32(66.7)	42(63.6)	50(70.4)	12(75.0)	0.87
BMI (kg/m ²)	28.9 ± 5.4	27.4 ± 4.3	25.7 ± 3.9	24.5 ± 3.6	22.2 ± 2.6	0.000 ^{***}
JOA score (points)	12.8 ± 2.7	12.5 ± 3.0	12.8 ± 3.4	12.0 ± 3.4	9.6 ± 5.0	0.07
Neck VAS score	40.5 ± 32.5	41.2 ± 31.5	32.0 ± 28.2	41.0 ± 31.9	46.1 ± 31.9	0.34
Cervical spine function (points)	78.2 ± 23.3	67.4 ± 28.8	67.1 ± 27.7	58.2 ± 30.8	54.4 ± 24.8	0.005 ^{**}
Questions of Q1 domain						
Q1-1	2.7 ± 0.5	2.2 ± 0.8	2.4 ± 0.7	2.3 ± 0.7 _a	2.2 ± 0.7	0.07
Q1-2	2.8 ± 0.6 ^a	2.7 ± 0.6 ^a	2.6 ± 0.6 ^a	2.3 ± 0.8 ^a	2.2 ± 0.8	0.001 ^{**}
Q1-3	2.3 ± 0.7 ^a	2.1 ± 0.8 ^a	2.1 ± 0.8 ^a	1.9 ± 0.8 ^a	1.8 ± 0.7	0.07
Q1-4	2.7 ± 0.6 ^a	2.7 ± 0.5 ^a	2.6 ± 0.6 ^a	2.4 ± 0.8 ^a	2.3 ± 0.8	0.11
C-OP-index	5.4 ± 3.2	5.8 ± 3.2	5.9 ± 3.0	6.1 ± 3.2	7.5 ± 3.4	0.33
W-OP-index	9.4 ± 7.7	9.1 ± 6.3	7.5 ± 5.0	8.7 ± 7.0	9.5 ± 5.4	0.70
Bridge formation of C-OPLL	0.5 ± 0.7	0.5 ± 1.0	0.8 ± 1.2	0.8 ± 1.2	1.5 ± 1.6	0.028 [*]
Bridge formation of W-OPLL	1.5 ± 2.3	1.1 ± 1.9	1.3 ± 1.8	1.6 ± 2.9	1.8 ± 1.9	0.36

Data are presented as mean ± standard deviation.

BMI, body mass index; JOA, Japanese Orthopaedic Association; VAS, visual analog scale (100-mm)

Q1 domain, cervical spine function domain of the Japanese Orthopaedic Association Cervical Myelopathy Evaluation Questionnaire; C-, cervical; W-, whole spine; OP-index, ossification index for OPLL;

bridge formation, ossified lesions forming a bridge with the posterior border of the adjacent vertebral body

^aSignificant differences were found among the questions in each age group (all $p < 0.05$)

* $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$

Table 4. Comparison between the groups with poor and good cervical function

Characteristic	Poor cervical function group (n=62)	Good cervical function group (n=176)	p-value
Age (years)	66.9 ± 11.2	62.8 ± 12.5	0.024*
Male sex, no. (%)	44 (71.0)	118 (67.1)	0.57
Body height	162.6 ± 9.4	162.7 ± 9.9	0.90
Body weight	69.5 ± 16.4	68.9 ± 14.8	0.79
BMI	26.1 ± 4.9	25.9 ± 4.3	0.70
Comorbid diabetes mellitus, no. (%)	17 (27.4)	42 (23.9)	0.58
JOA score	11.1 ± 3.7	12.7 ± 3.2	0.001**
Neck VAS score	55.1 ± 32.4	33.1 ± 28.8	0.000***
C-OP-index	6.8 ± 3.4	5.7 ± 3.1	0.018*
W-OP-index	10.0 ± 7.8	8.1 ± 5.7	0.046*
Cervical OP-index grade	1.8 ± 0.8	1.6 ± 0.7	0.024*
Bridge formation of C-OPLL	1.1 ± 1.4	0.6 ± 1.0	0.010*
Bridge formation of W-OPLL	2.2 ± 3.2	1.1 ± 1.8	0.003**

Data are presented as mean ± standard deviation.

BMI, body mass index; JOA, Japanese Orthopaedic Association; VAS, visual analog scale (100-mm); C-, cervical; W-, whole spine; OP-index, ossification index for OPLL; bridge formation, ossified lesions forming a bridge with the posterior border of the adjacent vertebral body

* $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$

Figures



Cervical OP-index	3	5	9
Bridge formation of cervical OPLL	0	1	3
The score of cervical spine function domain (points)	100	100	40

Figure 1

Representative reconstructive mid-sagittal computed tomography image of cervical OP-index and bridge formation of cervical OPLL. The score of the cervical spine function domains of the left, middle, and right images were 100 points, 100 points, and 40 points, respectively. OPLL, ossification of the posterior longitudinal ligament; OP-index, ossification index for OPLL




Figure 2

The scores for each question of Q1-1 and Q1-2 (neck extension), Q1-3 (neck rotation), and Q1-4 (neck flexion). The score of Q1-3 was significantly lower than that of the other questions. * $p < 0.05$

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupplementalTable1...docx](#)
- [SupplementalTable2...docx](#)