

Epidemiology and risk factors of rectal colonization of carbapenemase-producing *Enterobacteriaceae* among high-risk patients from ICU and HSCT wards in a university hospital

Li Yan

Chongqing Medical University First Affiliated Hospital

Jide Sun

Chongqing Medical University First Affiliated Hospital

Xiuyu Xu

Chongqing Medical University First Affiliated Hospital

Shifeng Huang (✉ sfhuang@hospital.cqmu.edu.cn)

Chongqing Medical University First Affiliated Hospital

Research

Keywords: fecal carriage, CRE, CPE, KPC, risk factors

Posted Date: April 13th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-21674/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Version of Record: A version of this preprint was published at Antimicrobial Resistance and Infection Control on September 23rd, 2020. See the published version at <https://doi.org/10.1186/s13756-020-00816-4>.

Abstract

Background: Nosocomial infections caused by carbapenemase-producing *Enterobacteriaceae* (CPE) constitute a major global health concern and are associated with increased morbidity and mortality. Rectal colonization with CPE is a risk factor for bacterial translocation leading to subsequent endogenous CPE infections. This prospective observational study was aimed to investigate the prevalence and epidemiology of rectal colonization of CPE, the carbapenemase genotypes, and to identify the independent risk factors for the acquisition of CPE colonization in high-risk patients from ICU and HSCT wards in a university hospital in China.

Methods: In a prospective cohort study, 150 fecal samples from rectal swabs were consecutively obtained for inpatients from the ICU and HSCT wards from November 2018 to May 2019, and screening test for CPE was conducted by using home-made trypsin soybean broth (TSB) selective media and MacConkey agar. Antimicrobial susceptibility was determined by the broth microdilution method and carbapenemase genes were characterized by both the GeneXpert Carba-R and PCR for *bla*_{KPC}, *bla*_{NDM}, *bla*_{IMP}, *bla*_{VIM} and *bla*_{OXA}. In order to further investigate the risk factors and clinical outcomes of CPE colonization, a prospective case-control study was also performed.

Results: 26 suspected CPE strains, including 17 *Klebsiella pneumoniae*, 6 *Escherichia coli*, 1 *Citrobacter freundii*, 1 *Enterobacter Kobe*, and 1 *Raoultella ornithinolytica*, were identified in 25 non-duplicated fecal samples from 25 patients, with a carriage rate of 16.67% (25/150). Through GeneXpert Carba-R and subsequent PCR and sequencing, all the suspected CPE isolates were identified to be positive for the carbapenemase genes, of which 17 were *bla*_{KPC}-carriers, and another 9 were *bla*_{NDM}-producers. Multivariate analysis indicated that urinary system diseases, operation of bronchoscopy, and combined use of antibiotics were independent risk factors for acquiring CPE colonization in high-risk patients from the ICU and HSCT wards.

Conclusions: This study revealed a high prevalence of rectal CPE colonization in high-risk patients from ICU and HSCT wards, and a predominant colonization of the KPC-producing *K. pneumoniae* strains. Stricter infection control measures are urgently needed to limit the dissemination of CPE strains, especially in patients who were afflicted by urinary system diseases, have underwent bronchoscopy, and were previously exposed to combined antibiotic use.

Background

Carbapenem-resistant *Enterobacteriaceae* (CRE) has become a serious public health threat worldwide [1, 2]. CRE infections have been increasing rapidly and posing serious challenges to the clinical management [3], thus, timely and efficient diagnosis, strict and evidence-based infection control measures, and prompt and effective therapy are of paramount importance. To help direct research and development efforts toward the production of novel drugs, CRE was recently listed as one of the three critical-priority pathogens by the World Health Organization (WHO) [4, 5].

Resistance to carbapenems in *Enterobacteriaceae* occurs via one or a combination of the following four mechanisms: carbapenemase production, production of ESBLs and/or AmpC in combination with porin loss/deficiency, carbapenem efflux, or mutations in penicillin-binding proteins (PBPs) [6, 7], among which carbapenemase gene acquisition is of greatest concern [8], as it most frequently occurs via transfer of plasmids, which usually co-harbor β -lactamases and other resistance determinants, rendering these carbapenemase-producing *Enterobacteriaceae* (CPE) strains multi-drug resistance (MDR) or extensively drug-resistant (XDR) [9].

CPE is the most problematic owing to higher-level antimicrobial resistance, clonal spread and horizontal gene transfer [10]. Carbapenemases include Ambler class A β -lactamases (e.g., KPC and GES), class B metallo- β -lactamases (MBLs; e.g., NDM, VIM, IMP, and SPM), and class D β -lactamases (e.g., OXA-48 and OXA-181) [6]. In the case of CPE, only some "second-line" drugs, such as polymyxins, tigecycline, and fosfomycin, and some "last resort" antibiotics, such as aztreonam/avibactam [11], ceftazidime/avibactam (CAZ/AVI) [12], meropenem/vaborbactam [13], imipenem/relebactam [14], and several newer investigational agents eravacycline [15], plazomicin [16], and cefiderocol [17], may be active, however, resistance to these new drugs also can be expected to emerge, as has been seen with aztreonam/avibactam [18] and ceftazidime-avibactam [19]. It is therefore crucial to implement efficient infection control measures to limit the spread of these pathogens.

The spread of CPE has been enabled by the lack of control measures directed at CPE carriers in healthcare settings. Both screening patients for asymptomatic CPE colonization and implementation of contact precautions will reduce patient-to-patient transmission. Although evidence-based guidelines on prevention and control of CPE are available, CPE still remains as a severe challenge in health care settings, and more studies on appropriate countermeasures are required especially for high-risk patients from ICU and HSCT wards. This prospective observational study was aimed to investigate the prevalence and epidemiology of fecal carriage of CPE, the carbapenemase genotypes, and to identify the independent risk factors for the acquisition of CPE colonization in high-risk patients in ICU and HSCT wards from a university hospital in China.

Methods

Setting and study design

The present study was a single-center prospective survey performed at the first Affiliated Hospital of Chongqing Medical University, China. Our hospital is a 3200-bed teaching hospital providing all types of acute care with around 200,000 admissions and 1000,000 hospital-days per year.

We implemented a prospective analysis of a cohort of CPE colonized patients detected from November 2018 to May 2019. All hospitalized patients in ICU wards (including central ICU, respiratory ICU, critical surgery ICU, neurology ICU, neurosurgery ICU, thoracic and cardiac surgery ICU) and hematopoietic stem cell transplantation (HSCT) wards during this period were included, due to the high prevalence of CRE carriage observed in these wards in the baseline survey, they are considered the highest-risk sites for CRE transmission.

Rectal swabs were obtained from all patients in a representative sample of wards in ICU and HSCT. Each cohort patient was systematically screened for CPE colonization on the day of admission. A total of 150 rectal swabs were collected from non-repetitive patients. Stool samples collected from rectal swab were cultured on selective TSB media (Pangtong, China), and the suspected CPE isolates were further identified at the species level by the VITEK MS (bioMérieux, MO, USA) system, and routine antimicrobial susceptibility testing was performed using the VITEK2 compact (bioMérieux, Inc., NC, USA) system.

Bacterial Identification and Antimicrobial Susceptibility Testing

A total of 26 clinical CRE strains were isolated and identified from November 2018 to May 2019 by using the VITEK MS (bioMérieux, Hazelwood, MO, United States) automated system at the department of laboratory medicine, all of which were resistant to at least one carbapenem on the basis of antimicrobial susceptibility testing results

determined by the broth microdilution method, with the criteria of MIC of ≥ 2 mg/L for ertapenem, ≥ 4 mg/L for imipenem, and ≥ 4 mg/L for meropenem.

Detection Of Antibiotic Resistance Genes

Both GeneXpert Carba-R (*bla*_{KPC}, *bla*_{NDM}, *bla*_{IMP}, *bla*_{VIM} and *bla*_{OXA}) and PCR were used to confirm the suspected resistance phenotypes. Moreover, the Carba NP test and sCIM were performed on all isolates to determine whether any bacteria produced carbapenemases by phenotypic methods but were negative by genotypic methods, or vice versa [20].

Risk Factors For Cpe Colonization

We conducted a prospective case-control study to explore the risk factors and clinical outcomes of patients colonized with CPE from November 2018 to June 2019 in Chongqing, China. Patients with CPE colonization were included as cases. Controls were identified as patients without CPE colonization with well-balanced demographic characteristics, pre-existing medical conditions, and immune-compromising comorbidities as compared with the cases. Clinical and epidemiological data, including the demographics, underlying diseases, the primary diagnosis at admission, invasive procedures prior to the isolation of CPE, previous exposures of antibiotic within 3 months, and the clinical outcomes, were extracted from the patients' electronic medical records system and clinical microbiology laboratory database.

Statistical Analysis

Data were collected using Excel software (Microsoft, Redmond, CA, USA), and were described using the mean with standard deviation (SD) for continuous variables and proportions (%) for qualitative variables.

All analyses were performed using SPSS v.22.0 software (SPSS Inc., Chicago, IL, United States). Univariate analyses were performed separately for each of the variables. Categorical variables were compared using a chi-square test or Fisher's exact test as appropriate. Continuous variables were compared using Student's t-test (normally distributed variables) and Wilcoxon rank-sum test (non-normally distributed variables) as appropriate. The odds ratio (OR) and 95% confidence interval (CI) were calculated to evaluate the strength of any association. Variables with $P \leq 0.05$ on univariate analysis were evaluated as potential covariates in a stepwise multivariate logistic regression model. All P-values were two sided, and $P < 0.05$ was considered significant.

Ethical Considerations

The data and samples analyzed in the present study were obtained in accordance with the standards and approved by the Chongqing Medical University Institutional Review Board and Biomedical Ethics Committee.

Results

Prevalence of positive CRE colonization among high-risk patients from ICU and HSCT wards

In the CRE prevalence survey, 150 patients from ICU and HSCT wards were prospectively recruited, and a total of 150 non-repetitive fecal swabs were collected, among which 26 CRE strains from 25 individual patients were detected as being CRE positive, with a carriage rate of 16.67% (25/150). Among the 26 CRE isolates, 22 CRE isolates (84.62%, 22/26) were isolated from 21 individual patients in the ICU wards, with a carriage rate of 18.58% (21/113); another 4 were from 4 patients in the HSCT wards, with a carriage rate of 10.81% (4/37). The 26 isolated CRE strains included 17 *Klebsiella pneumoniae* (65.38%), 6 *Escherichia coli* (23.08%), 1 *Citrobacter freundii* (3.85%), 1 *Enterobacter Kobe* (3.85%), and 1 *Raoultella ornithinolytica* (3.85%). The 26 CRE strains were isolated from 25 non-duplicated samples from 25 individual patients (14 male and 11 female; 21 from ICU and 4 from HSCT) whose mean age was 63.48 ± 22.81 years. Among these patients, 40.0% (10/25) were transferred from another hospital.

Confirmation of carbapenemase-producing phenotypes and molecular characterization of carbapenemase genes

Both Carba NP test and sCIM were performed on all isolates to determine whether the CRE isolates produced carbapenemases by phenotypic methods. Among the 26 CRE isolates, all were positive in both the two carbapenemase phenotype confirmatory tests, in concordance with our later molecular characterization results from both GeneXpert Carba-R and subsequent PCR and sequencing, which demonstrated 26 CPE isolates with carbapenemase genes. As was shown in Table 1, the most prevalent carbapenemase gene was *bla_{KPC}* (65.38%, 17/26), detected in all the 17 *K. pneumoniae* isolates; followed by *bla_{NDM}* (34.62%, 9/26), detected in all of the 6 *Escherichia coli* isolates, 1 *Citrobacter freundii*, 1 *Enterobacter Kobe*, and 1 *Raoultella ornithinolytica* isolate; *bla_{IMP}*, *bla_{VIM}*, and *bla_{OXA-48}* were not detected (Table 1). Notably, all the 17 *K. pneumoniae* isolates harbored *bla_{KPC}*, while all the 6 carbapenemase-producing *Escherichia coli* strains carried *bla_{NDM}*. It's also worth noting that one patient was demonstrated to co-harbor two CPE colonization isolates, with one *K. pneumoniae* strain producing KPC, and another *E. coli* isolate expressing NDM. Moreover, all the 26 CPE isolates showed resistance to all the three tested carbapenems (imipenem, meropenem, and ertapenem), showing a perfect correlation between CPE phenotype and genotype.

Table 1

Molecular characterization of carbapenemase genes in 26 CRE isolates as detected by GeneXpert Carba-R and PCR

Bacteria	Carbapenemases (isolate number)				
	<i>bla_{KPC}</i>	<i>bla_{NDM}</i>	<i>bla_{VIM}</i>	<i>bla_{IMP}</i>	<i>bla_{OXA-48}</i>
<i>Klebsiella pneumoniae</i>	17	-	-	-	-
<i>Escherichia coli</i>	-	6	-	-	-
<i>Citrobacter freundii</i>	-	1	-	-	-
<i>Enterobacter Kobe</i>	-	1	-	-	-
<i>Raoultella ornithinolytica</i>	-	1	-	-	-

Risk factors for the acquisition of rectal CPE colonization in high-risk patients from ICU and HSCT wards

The risk factors for the acquisition of rectal CPE colonization in high-risk patients from ICU and HSCT wards were shown in Table 2. The univariate analysis indicated that higher APACHE II Scores, diabetes mellitus, urinary system disease, coronary heart disease, endocrine system Diseases, multiple organ dysfunction syndrome, bronchoscopy, and combined use of antibiotics were significantly more frequent in patients with rectal CPE colonization ($P \leq 0.05$). Further multivariate logistic regression analysis demonstrated that urinary system disease (OR [Odd ratio]: 18.06, 95% CI [Confidence Interval]: 3.31–98.62, $p = 0.001$), operation of bronchoscopy (OR: 4.05, 95% CI: 1.30–12.60, $p = 0.01$), and previous exposure to combined antibiotics (OR: 3.60, 95% CI: 1.18–10.93, $p = 0.02$) were independent risk factors for the acquisition of rectal CPE colonization in high-risk patients from ICU and HSCT wards.

Table 2
Univariate and multivariate analyses of risk factors for patients with CPE colonization

Variable	CRE Colonization (n = 25)	Control (n = 75)	Univariate analysis		Multivariate analysis	
			OR (95% CI)	P-value	OR (95% CI)	P-value
Patient characteristics						
Man gender (Male)	14(56.00%)	48(64.00%)	0.65(0.21-2.00)	0.45		
Age	63.48 ± 22.81	60.76 ± 19.01	1.01(0.98-1.05)	0.36		
Transferring from another hospital	10 (40.00%)	37(49.33%)	0.92(0.29-2.92)	0.88		
Underlying diseases						
PITT Scores	2.28 ± 1.86	2.98 ± 2.09	0.79(0.54-1.15)	0.22		
APACHE II Scores	17.16 ± 7.18	15.74 ± 8.13	1.10(1.01-1.20)	0.05	0.40 (0.09-1.71)	0.22
Hypertension	13(52.00%)	38(50.66%)	0.62(0.18-2.15)	0.42		
diabetes	12(48.00%)	20(26.67%)	3.97(1.16-13.62)	0.03	4.30(0.67-27.306)	0.12
Tuberculosis	2(8.00%)	4(5.33%)	0.94(0.10-8.50)	0.95		
Hepatitis B	2(8.00%)	3(4.00%)	2.17(0.24-19.57)	0.09		
Tumour	12(48.00%)	21(28.00%)	2.88(0.85-9.72)	0.09		
Respiratory diseases	17(68.00%)	36(48.00%)	2.64(0.76-9.14)	0.13		
COPD	9(36.00%)	5(6.67%)	0.92(0.20-4.20)	0.91		
Hepatobiliary disease	8(32.00%)	26(34.67%)	0.63(0.18-2.10)	0.47		
Gastrointestinal disease	6(24.00%)	27(36.00%)	0.36(0.09-1.46)	0.15		
Chronic kidney disease	6(24.00%)	21(28.00%)	0.75(0.21-2.70)	0.66		
Urinary system disease	10(40.00%)	6(8.00%)	7.71(2.21-26.83)	0.001	18.06(3.31-98.62)	0.001
Cardiovascular disease	15(60.00%)	41(54.67%)	0.87(0.27-2.80)	0.82		

Variable	CRE Colonization (n = 25)	Control (n = 75)	Univariate analysis		Multivariate analysis	
			OR (95% CI)	P-value	OR (95% CI)	P-value
Coronary heart disease	7(28.00%)	10(13.33%)	4.26(1.12–16.28)	0.03	3.29(0.91–11.98)	0.07
Neurological disease	9(36.00%)	18(24.00%)	2.27(0.64–7.99)	0.20		
Disease of immune system	1(4.00%)	5(6.67%)	0.22(0.02–2.85)	0.25		
Hematological system diseases	4(16.00%)	16(21.33%)	1.47(0.32–6.72)	0.62		
Endocrine System Diseases	15(60.00%)	28(37.33%)	3.16(1.05–9.54)	0.04	1.62(0.33–7.99)	0.55
Surgery in the past 6 months	2(8.00%)	16(21.33%)	0.32(0.05–1.99)	0.22		
Multiple organ dysfunction syndrome	6(24.00%)	8(10.67%)	6.27(1.35–29.05)	0.02		
Transplant	4(16.00%)	11(14.67%)	1.82(0.43–7.64)	0.42		
Urinary tract infection	10(40.00%)	12(16.00%)	3.64(0.78–16.87)	0.09		
Respiratory infection	21(84.00%)	51(68.00%)	2.20(0.37–12.92)	0.38		
Abdominal infection	8(32.00%)	2(2.67%)	0.48(0.05–4.68)	0.53		
Septic shock	1(4.00%)	9(12.00%)	0.99(0.95–1.02)	0.53		
Severe anemia	5(20.00%)	17(26.15%)	1.06(0.27–4.13)	0.94		
Glucocorticoid	7(28.00%)	12(16.00%)	2.58(0.86–9.72)	0.21		
Cardiotonic	2(8.00%)	5(6.67%)	0.78(0.08–7.87)	0.84		
Receipt of total parenteral Nutrition (days)	39.00 ± 57.47	16.71 ± 25.33	0.99(0.95–1.02)	0.53		
Urinary catheter(days)	56.08 ± 60.35	21.80 ± 26.07	1.04(0.99–1.09)	0.08		
Nasal catheter(days)	35.38 ± 48.75	21.59 ± 36.98	0.97(0.92–1.04)	0.16		
Mechanical ventilation(days)	34.64 ± 48.61	15.15 ± 19.01	1.02(0.98–1.09)	0.16		

Variable	CRE Colonization (n = 25)	Control (n = 75)	Univariate analysis		Multivariate analysis	
			OR (95% CI)	P-value	OR (95% CI)	P-value
Trachea cannula	24.80 ± 48.02	12.65 ± 24.27	0.98(0.95–1.03)	0.55		
Tracheotomy	10(40.00%)	31(41.33%)	0.17(0.02–1.28)	0.08		
Bronchoscopy	12(48.00%)	13(17.33%)	8.15(1.61–41.22)	0.01	4.05(1.30–12.60)	0.01
Drainage tube	19(76.00%)	55(73.33%)	0.31(0.05–1.76)	0.18		
Penicillin	5(20.00%)	17(22.67%)	0.60(0.14–2.55)	0.48		
Cephalosporins	8(32.00%)	25(33.33%)	0.37(0.08–1.61)	0.19		
Carbapenem	10(40.00%)	24(32.00%)	1.12(0.32–3.84)	0.86		
Fluoroquinolone	5(20.00%)	11(14.67%)	0.87(0.18–4.30)	0.87		
Tetracycline	3(12.00%)	4(5.33%)	0.93(0.14–5.90)	0.92		
Glycopeptide	3(12.00%)	12(16.00%)	0.20(0.03–1.15)	0.07		
Tigecycline	4(16.00%)	3(4.00%)	1.94(0.27–13.98)	0.51		
Combined use of antibiotics	18(72.00%)	32(42.67%)	5.91(1.20–29.16)	0.03	3.60(1.18–10.93)	0.02

Discussion

CPE transmission has been enabled by the lack of control measures directed at CPE carriers in healthcare settings. Screening patients for asymptomatic colonization and implementation of contact precautions could reduce patient-to-patient transmission [21]. Investigation of the prevalence of the rectal CPE colonization among high-risk patients from ICU and HSCT wards can contribute to better infection control measures to limit CPE dissemination. Our study revealed that the rate of rectal CPE colonization in the high-risk patients from ICU and HSCT wards was 16.67%, with the sole carbapenemase gene being *bla*_{KPC} in *K. pneumoniae* isolates and *bla*_{NDM} in *E. coli*. Stricter infection control measures are urgently needed to limit the CPE dissemination, especially in patients who were afflicted by the urinary system diseases, have underwent bronchoscopy, and were previously exposed to combined antibiotic use.

While CRE infection rates were reported to be varied (from 7.6–44.4%) in individual studies for CRE-colonized patients [22, 23], a recent meta-analysis suggested an overall 16.5% risk of infection with CRE amongst patients

colonized with CRE [24]. From a clinical point of view, all patients in whom colonization could represent a risk factor for invasive disease should be screened [25]. Notably, chemotherapy for acute leukemia, solid organ transplantation, and ICU stay were identified as the most important patient-related conditions associated with a significant risk of CRE infections in colonized patients [26]. Accordingly, with this indication, ICUs, Transplant Units, Hematological Units, major surgical and Infectious Disease units represent the preferential setting for active targeting screening. We thus focused on the investigation of the prevalence of CPE colonization in patients from ICU and HSCT wards in our hospital. As expected, this study revealed a higher rate of fecal carriage of CPE (16.67%) in the high-risk patients from ICU and HSCT wards than that was reported for randomly recruited patients from several previous studies in Fujian (6.6%) [27], Hunan (8.5%) [28], and Thailand (8.7%) [29].

Rapid detection and characterization of carbapenemase types in colonized CPE strains according to the Ambler classification will lead to improved guidance on the implementation of infection control measures. In this study, we have demonstrated that carbapenem resistance in rectal *K. pneumoniae* and *E. coli* isolates was respectively solely associated with KPC and NDM production. On the other hand, as selective pressure from carbapenem use continues, multiple carbapenemases per pathogen, including pathogens carrying three different carbapenemase genes [30, 31], or multiple CPE pathogens per patient are increasingly common. In our screening, although non CPE isolates co-harboring double or triple carbapenemases were identified, we did witness one patient co-colonized by two CPE isolates, with one KPC-producing *K. pneumoniae* strain and another NDM-expressing *E. coli* strain. As these deleterious pathogens may spread throughout health care facilities, more close attention to increased infection control measures and stewardship of the carbapenem-containing drugs should be paid in order to control selection of even more detrimental pathogens.

High prevalence of CPE carriers may cause difficulties for patient flow, as well as high costs due to the need for contact isolation. In order to prevent colonization, the investigation of the clinical predictors and risk factors for rectal CPE colonization brooks no delay. A recently published case-control study demonstrated solid organ and stem cell transplantation, mechanical ventilation, fecal incontinence, and exposure in the prior 30 days to carbapenems, vancomycin, and metronidazole as independent factors associated with CRE colonization [32]. In this prospective case-control study, operation of bronchoscopy, urinary system diseases, and combined use of antibiotics were demonstrated to be independent associated with rectal CPE colonization in high-risk patients from the ICU and HSCT wards. One possible explanation for bronchoscopy operation as an independent risk factor may be due to the fact that invasive bronchoscopy operation might have destroyed patients' natural barrier functions, and possibly led to CPE colonization. Given that invasive procedures such as the use of urinary catheters were frequently performed in patients with urinary system diseases, it is also easy to understand urinary system diseases as another independent risk factor for rectal CPE colonization. As invasive procedures were potential risk factors, infection control measures preventing the microbial colonization of the insertion sites are necessary. In addition, our study identified combined use of antibiotics as being associated with rectal CPE colonization. One possibility is that inappropriate combined antibiotic use may disrupt the gastrointestinal microflora and eradicate susceptible competing strains, thus elevating the incidence of CPE colonization [33].

Our study had several limitations. Firstly, we did not perform pulsed-field gel electrophoresis (PFGE) to determine the clonal relationships among the isolated CPE strains. Secondly, this was a prospective single-center case-control study with relatively small sample size, and our results might not be applicable to other settings. Thirdly, we didn't investigate the subsequent CPE infection rates in these CPE-colonized patients.

Conclusion

In conclusion, this study provided a detailed report of the prevalence, molecular epidemiology, and risk factors for rectal CPE colonization in high-risk patients from ICUs and HSCT wards in one medical center. Our data showed a high prevalence of rectal CPE colonization in high-risk patients from ICU and HSCT wards, and a predominant colonization of the KPC-producing *K. pneumoniae* strains. A bundle of infection control and prevention measures with an anti-infective stewardship program is urgently needed to reduce the rectal CPE colonization, especially in patients who were afflicted by urinary system diseases, have underwent bronchoscopy, and were previously exposed to combined antibiotic use.

Abbreviations

CLSI: Clinical and Laboratory Standards Institute; CRE: Carbapenem-resistant *Enterobacteriaceae*; CPEO: Carbapenemase-producing *Enterobacteriaceae*; MIC: Minimum inhibitory concentration; WHO: World health organization

Declarations

Antimicrobials

ETP: Ertapenem; IMP: Imipenem; MEM: Meropenem

Acknowledgements

We would like to thank all the enrolled patients for their participation in our study.

Funding

This study was supported in part by the Science and Technology Research Program of Chongqing Municipal Education Commission (No. KJ1500235), and the Medical Research Program of Chongqing Health and Family Planning Commission (No. 2018MSXM009).

Availability of data and materials

All the data of this article is available from the corresponding author if reasonably requested.

Authors' contributions

All the authors contributed to data analysis, drafting and revising the article, gave final approval of the version to be published, and agree to be accountable for all aspects of the work. LY and SFH designed this study, participated in statistic calculations and interpretation of data, and drafted this manuscript. LY, JDS, and XYX collected the samples and performed the experiments.

Ethics approval and consent to participate

The data and samples analyzed in the present study were obtained in accordance with the standards and approved by the Chongqing Medical University Institutional Review Board and the Biomedical Ethics Committee. Informed consent was acquired from each patient.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

1. Durante-Mangoni E, Andini R, Zampino R. Management of carbapenem resistant Enterobacteriaceae infections. *Clin Microbiol Infect.* 2019; 25(8): 943-50.
2. Righi E, Peri AM, Harris PN, Wailan AM, Liborio M, Lane SW, et al. Global prevalence of carbapenem resistance in neutropenic patients and association with mortality and carbapenem use: systematic review and meta-analysis. *J Antimicrob Chemother.* 2017; 72(3): 668-77.
3. Potter RF, D'Souza AW, Dantas G. The rapid spread of carbapenem-resistant *Enterobacteriaceae*. *Drug Resist Updat.* 2016; 29: 30-46.
4. World Health Organization. Essential Medicines and Health Products. Prioritization of Pathogens to Guide Discovery, Research and Development of New Antibiotics for drug-Resistant Bacterial Infections, Including Tuberculosis; WHO: Geneva, Switzerland, 2017.
5. WHO publishes list of bacteria for which new antibiotics are urgently needed. WHO Media Centre. 2017; 1–4. doi:/entity/mediacentre/news/releases/2017/bacteria-antibiotics-needed/en/index.html.
6. Nordmann P, Dortet L, Poirel L. Carbapenem resistance in Enterobacteriaceae: here is the storm! *Trends Mol Med.* 2012; 18(5): 263-72.
7. Bush K, Bradford PA. Epidemiology of β -Lactamase-Producing Pathogens. *Clin Microbiol Rev.* 2020; 33(2): e00047-19.
8. Nordmann P, Poirel, L. The difficult-to-control spread of carbapenemase producers among *Enterobacteriaceae* worldwide. *Clin Microbiol Infect.* 2014; 20: 821–30.
9. Roberts LW, Harris PNA, Forde BM, Ben Zakour NL, Catchpoole E, Stanton-Cook M, et al. Integrating multiple genomic technologies to investigate an outbreak of carbapenemase-producing *Enterobacter hormaechei*. *Nat Commun.* 2020; 11(1): 466.
10. Lutgring JD, Limbago BM. The problem of carbapenemase-producing-carbapenem-resistant-*Enterobacteriaceae* detection. *J Clin Microbiol.* 2016; 54(3): 529-34.
11. Biedenbach DJ, Kazmierczak K, Bouchillon SK, Sahm DF, Bradford PA. In vitro activity of aztreonam-avibactam against a global collection of gram-negative pathogens from 2012 and 2013. *Antimicrob Agents Chemother.* 2015; 59(7):4239–4248.
12. Falcone M, Paterson D. Spotlight on ceftazidime/avibactam: a new option for MDR gram-negative infections. *J Antimicrob Chemother.* 2016; 71(10): 2713–22.
13. Lomovskaya O, Sun D, Rubio-Aparicio D, Nelson K, Tsivkovski R, Griffith DC, et al. Vaborbactam: spectrum of beta-lactamase inhibition and impact of resistance mechanisms on activity in *Enterobacteriaceae*. *Antimicrob Agents Chemother.* 2017; 61(11): e01443–e01517.
14. Haidar G, Clancy CJ, Chen L, Samanta P, Shields RK, Kreiswirth BN, et al. Identifying spectra of activity and therapeutic niches for ceftazidime-avibactam and imipenem-relebactam against carbapenem-resistant

- Enterobacteriaceae*. Antimicrob Agents Chemother. 2017; 61(9):e00642–e00717.
15. Monogue ML, Thabit AK, Hamada Y, Nicolau DP. Antibacterial efficacy of eravacycline in vivo against gram-positive and gram-negative organisms. Antimicrob Agents Chemother. 2016; 60 (8): 5001–5.
 16. Denervaud-Tendon V, Poirel L, Connolly LE, Krause KM, Nordmann P. Plazomicin activity against polymyxin-resistant *Enterobacteriaceae*, including MCR-1-producing isolates. J Antimicrob Chemother. 2017; 72(10): 2787–91.
 17. Hackel MA, Tsuji M, Yamano Y, Echols R, Karlowsky JA, Sahm DF. In vitro activity of the siderophore cephalosporin, cefiderocol, against carbapenem-nonsusceptible and multidrug-resistant isolates of gram-negative bacilli collected worldwide in 2014 to 2016. Antimicrob Agents Chemother. 2018; 62(2): e01968–17.
 18. Alm RA, Johnstone MR, Lahiri SD. Characterization of *Escherichia coli* NDM isolates with decreased susceptibility to aztreonam/avibactam: role of a novel insertion in PBP3. J Antimicrob Chemother. 2015; 70: 1420–8.
 19. Nelson K, Hemarajata P, Sun D, Rubio-Aparicio D, Tsivkovski R, Yang S, et al. Resistance to ceftazidime-avibactam is due to transposition of KPC in a porin-deficient strain of *Klebsiella pneumoniae* with increased efflux activity. Antimicrob Agents Chemother. 2017; 61: e00989-17.
 20. Wang Q, Wang X, Wang J, Ouyang P, Jin C, Wang R, et al. Phenotypic and genotypic characterization of carbapenem-resistant *Enterobacteriaceae*: data from a longitudinal large-scale CRE study in China (2012–2016). Clin Infect Dis. 2018; 67 (suppl_2): S196-S205.
 21. Del Bianco F, Morotti M, Zannoli S, Dirani G, Fantini M, Pedna MF, et al. Comparison of four commercial screening assays for the detection of blaKPC, blaNDM, blaIMP, blaVIM, and blaOXA48 in rectal secretion collected by swabs. Microorganisms 2019; 7(12): 704.
 22. Pisney LM, Barron MA, Kassner E, Havens D, Madinger NE. Carbapenem resistant *Enterobacteriaceae* rectal screening during an outbreak of New Delhi Metallo- β -lactamase-producing *Klebsiella pneumoniae* at an acute care hospital. Infect Control Hosp Epidemiol. 2014; 35: 434–6.
 23. Lubbert C, Becker-Rux D, Rodloff AC, Laudi S, Busch T, Kalsers UX BM. Colonization of liver transplant recipients with KPC-producing *Klebsiella pneumoniae* is associated with high infection rates and excess mortality: a case-control analysis. Infection. 2014; 42: 309–16.
 24. Tischendorf J, de Avila RA, Nasia SN. Risk of infection following colonization with carbapenem-resistant *Enterobacteriaceae*: a systematic review. Am J Infect Control. 2016; 44: 539–43.
 25. Ambretti S, Bassetti M, Clerici P, Petrosillo N, Tumietto F, Viale P, et al. Screening for carriage of carbapenem resistant *Enterobacteriaceae* in settings of high endemicity: a position paper from an Italian working group on CRE infections. Antimicrob Resist Infect Control. 2019; 8: 136
 26. Paño Pardo JR, Serrano Villar S, Ramos Ramos JC, Pintado V. Infections caused by carbapenemase-producing *Enterobacteriaceae*: risk factors, clinical features and prognosis. Enferm Infecc Microbiol Clin. 2014; 32 Suppl 4: 41–8.
 27. Zhao ZC, Xu XH, Liu MB, Wu J, Lin J, Li B. Fecal carriage of carbapenem-resistant *Enterobacteriaceae* in a Chinese university hospital. Am J Infect Control. 2014; 42(5): e61–e64.
 28. Liu Q, Liu L, Li Y, Chen X, Yan Q, Liu WE. Fecal carriage and epidemiology of carbapenem-resistant *Enterobacteriaceae* among hospitalized patients in a university hospital. Infect Drug Resist. 2019; 12: 3935–42.

29. Shanmugakani RK, Akeda Y, Sugawara Y, Laolerd W, Chaihongsa N, Sirichot S, et al. PCR-dipstick-oriented surveillance and characterization of *mcr-1*- and carbapenemase-carrying *Enterobacteriaceae* in a Thai hospital. *Front. Microbiol.* 2019; 10: 149.
30. Khan MA, Mohamed AM, Faiz A, Ahmad J. Enterobacterial infection in Saudi Arabia: First record of *Klebsiella pneumoniae* with triple carbapenemase genes resistance. *J Infect Dev Ctries.* 2019; 13(4): 334-41.
31. Vannice K, Benoliel E, Kauber K, Brostrom-Smith C, Montgomery P, Kay M, et al. Notes from the field: clinical *Klebsiella pneumoniae* isolate with three carbapenem resistance genes associated with urology procedures—King County, Washington, 2018. *MMWR Morb Mortal Wkly Rep* 2019; 68: 667–8.
32. Mills JP, Talati NJ, Alby K, Han JH. The epidemiology of carbapenem-resistant *Klebsiella pneumoniae* colonization and infection among long-term acute care hospital residents. *Infect Control Hosp Epidemiol.* 2016; 37: 55–60.
33. Li S, Jia X, Li C, Zou H, Liu H, Guo Y, et al. Carbapenem-resistant and cephalosporin-susceptible *Pseudomonas aeruginosa*: a notable phenotype in patients with bacteremia. *Infect Drug Resist.* 2018; 11: 1225-35.