

Determinants of physical, psychological, and social well-being in older adults residing in senior care facilities of Pakistan

Jawad Tariq

Forman Christian College (A Chartered University)

Rubeena Zakar

University of the Punjab

Mohammad Vaqas Ali

Forman Christian College (A Chartered University)

Muhammad Zakria Zakar

University of Poonch Rawalakot

Amal Sajjad

University of the Punjab

Florian Fischer (✉ florian.fischer1@charite.de)

Institute of Public Health, Charité – Universitätsmedizin Berlin

Research Article

Keywords: Relocation autonomy, well-being, older adult, loneliness, senior care facility, old age home

Posted Date: November 3rd, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-2196160/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Additional Declarations: No competing interests reported.

Version of Record: A version of this preprint was published at BMC Geriatrics on June 5th, 2023. See the published version at <https://doi.org/10.1186/s12877-023-04014-w>.

Abstract

Background

This study investigated the effects of relocation autonomy, loneliness, and satisfaction with services along with socio-demographic characteristics on physical, psychological, and social well-being of older residents living in senior care facilities of Punjab, Pakistan.

Methods

Data were collected from 270 older residents living in 18 senior care facilities across 11 districts of Punjab, Pakistan. The scales used in the study were assessed for their construct validity, convergence validity, and internal consistency. Three separate multiple regression analyses were carried out to predict physical, psychological, and social well-being of residents from socio-demographic variables, relocation autonomy, loneliness, and satisfaction with services.

Results

The results of multiple regression analyses showed that the models predicting physical ($R^2 = 0.579$), psychological ($R^2 = 0.654$), and social well-being ($R^2 = 0.615$) were statistically significant ($p < 0.001$). Number of visitors was a significant predictor of physical ($b = 0.82$, $p = 0.01$), psychological ($b = 0.80$, $p < 0.001$), and social ($b = 2.40$, $p < 0.001$) well-being. Loneliness significantly predicted physical ($b = -0.14$, $p = 0.005$), psychological ($b = -0.19$, $p < 0.001$), and social ($b = -0.36$, $p < 0.001$) well-being. Control over relocation process significantly predicted physical ($b = 0.56$, $p < 0.001$) and psychological ($b = 0.36$, $p < 0.001$) well-being. Satisfaction with services significantly predicted physical ($b = 0.07$, $p < 0.001$) and social ($b = 0.08$, $p < 0.001$) well-being.

Conclusion

Pragmatic, equitable and cost-effective interventions are needed to improve the wellbeing of older residents living in senior care facilities. Friendly behavior of mobilizing staff and adjusted residents to facilitate new residents, therapeutic interventions such as relocation support programs, reminiscence therapy and intergenerational support, and increasing their exposure and connection to the outside world, can raise their physical, psychological, and social well-being.

Background

Demographic changes such as the increase in people aged 60 years and above have created a structural lag in many developing countries of South-East Asia like Pakistan that lack organizational structures and institutional capacities to meet the demands of older persons [1, 2]. According to the latest available statistics, Pakistan was ranked the 5th worst country for older adults [3]. There are more than 15 million people aged 60 years and above in Pakistan, and the projections show that the number will become 40 million by 2050, making them 12% of the total population [4]. A similar trend can also be observed in other South-East Asian countries bringing challenges such as the provision of healthcare facilities, transportation, jobs, recreation, housing, and increasing the number of different types of senior care facilities such as old age homes, nursing homes, assisted living facilities, retirement homes, and residential care facilities for older adults [2, 5–7]. Some of the studies from this region conducted on senior care facilities have stressed to increase the number of such facilities due to the increasing population of older adults, demand for healthcare systems, housing, changing preferences of older persons to live in such settings to avoid loneliness, abandonment of parents by the children due to changing norms of filial piety and familial care-giving, and increased abuse and neglect of older adults in home

settings [1, 2, 6, 8–11]. Abuse and neglect of older adults is a significant public health problem. However, there is no nationwide screening system in Pakistan for such abuse available yet [12]. Some of the academic studies have reported high prevalence of abuse in older adults in Pakistan. For example, Hashmi et al. found that 53.3%, 39.3%, and 53.5% of the older adults in two hospitals in Lahore, Pakistan, reported abuse, neglect, and loneliness, respectively. Of those who reported abuse, 56.6% were females [12]. Likewise, a study by Zaidi et al. [13] reported appalling condition of older adults in relation to abuse. Mushtaq and Ali [14] found that 42% and 41% of the older adults in Quetta, Pakistan experienced neglect and abuse, respectively. A study in Nepal found domestic abuse to be one of the significant motivations for older adults to move to old age homes [8]. It is important to highlight that older adults may be at higher risk of experiencing abuse and lower well-being in these senior care facilities as described in many studies [15–19].

Old age homes are the common residential setting for older adults in South-East Asian countries providing accommodation, food, clothing, recreation, and basic healthcare. However, they are often stereotypically labeled as homes for older adults who have been neglected and abandoned by their families [2, 5, 20, 21]. Gerontologists and Geriatricians point out that low life satisfaction, lesser perceived control, and poor well-being may not be an outcome of aging as older adults in vulnerable settings are at a higher risk of experiencing these negative consequences as compared to other groups of older adults [1, 2, 5]. For example, studies have found that older adults who relocated to senior care facilities were at higher risk of having lower well-being and higher depression than community dwellers [22–24]. This can be theoretically explained by the continuity theory which posits that older adults seek stability and consistency in their environment instead of change. In this context, the physical environment such as one's residence represents a sense of self, personhood, identity, meaning, and a network of embedded relations [25, 26]. This argument gains strength from the person-environment model which explains that relocation to a new setting entails detaching oneself from the meanings that the previous residence held for the person, resulting in negative consequences such as loneliness, loss of social ties, maladjustment, depression, and lower well-being [26–29]. Studies on relocation point out that voluntary relocation to senior care facilities involving choice and control followed by a supportive environment at these settings and compatibility of new residence with the previous one are the primary determinants of well-being [29, 30]. Well-being has long been considered to be an important indicator of general health in older adults and, thus, a significant determinant of successful aging [31, 32]. Many researchers have conceptualized well-being as a multidimensional construct and stressed to measure it along the three dimensions of physical, psychological, and social well-being [33–35]. Nevertheless, very few studies conducted on senior care facilities have focused on all three types of well-being [33, 34].

The published research literature on senior care facilities in Pakistan is scant. Very few studies have been conducted on old age homes, and the exact number of old age homes and the older adults residing in these senior care facilities in Pakistan is not known [1, 2, 5, 6, 11, 36]. Some of the studies conducted on these facilities have reported poor quality of life of older adults due to a lack of financial, structural, and social support [1, 2, 5, 6, 36]. No study was found that empirically investigated the well-being of older adults in these settings. Therefore, this study is the largest of its kind in Pakistan. The aim of this study was both empirical and normative, because it investigates the factors that determine the well-being of older adults in old age homes, a marginalized and neglected setting in Pakistan [1, 2]. It is very important to oversee and evaluate the working of such facilities to ensure the well-being of older adults. The study hypothesized that higher relocation autonomy, lower loneliness, and higher satisfaction with staff and services will be related to higher well-being of older adults living in old age homes of Punjab, Pakistan [22, 33]. Although abuse and neglect were not measured directly in this study, proxy variables such as relocation autonomy, loneliness, satisfaction with staff and services were considered to be reasonable indicators to assess neglect experienced by older adults in old age homes. The study has the potential to expand the current knowledge on (i) neglect of older adults by assessing loneliness, autonomy, and their living conditions, (ii) relocation of older adults to senior care facilities, and (iii) factors determining older adult's well-being in these facilities. Additionally, the study contributes to clinical and research practices by providing psychometric validation of different scales that can be administered to older adults in old age homes and other senior care facilities. By providing

pragmatic interventions, this study is also expected to sensitize policymakers and relevant governments in Pakistan to address the problems of older adults living in old age homes.

Methods

Study design and participants

The study was conducted among 270 residents (out of a total of 350 residents) living in seven public and 11 private senior care residential facilities located across 11 districts of Punjab, Pakistan. Due to the lack of a sampling frame and absence of information related to private old age homes in Punjab, six districts were randomly selected from a total of 35 districts in Punjab. The researchers were able to locate a total of 20 private old age homes in these six districts. Permission to conduct this study was provided by administrations of 11 old age homes. Only those older adults were included in the study who had a minimum age of 60 years, were able to communicate, and did not have any cognitive impairment. Cognitive impairment was not assessed using any tool (for example, mini mental state examination), but residents having some cognitive impairment as identified by administrators and staff of the old age homes were excluded from the study.

Measures

A structured interview questionnaire was administered to the residents by two authors of this study who were assisted by three graduate research assistants. The three research assistants hired for this study were given two trainings and were financially compensated for their assistance. The cross-sectional survey instrument was translated from English to Urdu with the help of three experts using forward and backward translation procedure.

Dependent variable. The dependent variable in this study was well-being which was conceptualized along the three dimensions of physical, psychological, and social well-being. The two sub-scales to measure perceived well-being developed by Reker and Wong [37] were used to measure physical and psychological well-being, respectively. The construct to measure physical well-being had eight items (range: 0–16) and for psychological well-being six items (range: 0–12). Higher scores on both scales indicated higher well-being. Social well-being was measured by the Duke Social Support Index [38] with eleven items (range: 0–22), where higher scores showed higher social well-being. Items comprising these scales along with their psychometric properties are provided in Table 1.

Table 1
Psychometric properties and descriptive statistics of scales used in senior care facilities (n = 270)

<i>Scales</i>	<i>Loadings</i>	<i>Cronbach's alpha</i>	<i>Mean (SD)</i>	<i>Range</i>
Relocation autonomy (AVE = 81.4%)		0.92	4.61 (3.10)	0–8
Was it your decision to live here?	0.82			
How much input you had in the decision?	0.89			
Did you somehow influence the decision?	0.84			
Any consultations about the decision?	0.71			
Loneliness (AVE = 59.4%)		0.93	11.8 (6.83)	0–22
Miss pleasure of company of others?	0.87			
Often feel rejected?	0.73			
Experienced a general sense of emptiness?	0.67			
Plenty of people to lean on in case of problems?	0.66			
Have someone to talk to about daily problems?	0.70			
Circle of friends and acquaintances too limited?	0.77			
Have many people that can be trusted?	0.70			
Miss having people around you?	0.87			
Have enough people to feel close to?	0.61			
Miss having a really close friend?	0.74			
Call on friends whenever needed?	0.51			
Service quality (AVE = 35.2%)		0.90	32.7 (8.36)	4–50
Respectful staff?	0.78			
Polite staff?	0.71			
Questions addressed by staff?	0.65			
Comforted by staff when sad/lonely?	0.61			
Staff has time to talk about what bothers you?	0.62			
Staff knows about personal habits and tastes?	0.72			
Staff less than what is actually required?	0.58			
Staff encourages residents to establish relations?	0.72			
Staff encourages participation in activities?	0.71			
Staff too busy to answer my requests?	0.62			

AVE = Average Variance Extract

<i>Scales</i>	<i>Loadings</i>	<i>Cronbach's alpha</i>	<i>Mean (SD)</i>	<i>Range</i>
Staff meets promise of coming within time?	0.50			
Not have to ask twice before something is done?	0.64			
Immediate response by staff when called?	0.70			
Staff responds instantly in medical emergency?	0.54			
Staff keeps quality of life as high as possible?	0.58			
Staff takes sincere interest in solving problems?	0.55			
Information given by staff about daily activities?	0.57			
Residents involved in decision making?	0.62			
Residents can offer feedback to concerned?	0.61			
Residents can propose suggestions?	0.71			
Residents can decide when they eat?	0.70			
Residents can decide what to eat?	0.71			
Residents decide when they want to go out?	0.79			
Residents decide which clothes they wear?	0.67			
Residents decide when to go to bed and get up?	0.60			
Residents given the privacy they want?	0.57			
Physical well-being (AVE = 62.7%)		0.91	9.58 (4.59)	0–16
Do not have many physical complaints?	0.67			
Do not think that I have a heart condition?	0.65			
Good appetite for food?	0.56			
Have aches and pains?	0.54			
In good shape physically?	0.65			
Is your health deteriorating?	0.58			
Do not get tired very easily?	0.66			
Can stand a fair amount of physical strain?	0.73			
Psychological well-being (AVE = 62.1%)		0.88	7.20 (3.39)	0–12
Is life worth living?	0.67			
Often bored?	0.51			
No one cares if I am dead or alive?	0.50			
Exciting to be alive?	0.60			

AVE = Average Variance Extract

<i>Scales</i>	<i>Loadings</i>	<i>Cronbach's alpha</i>	<i>Mean (SD)</i>	<i>Range</i>
Sometimes I wish I never wake up?	0.74			
Do not seem to care about what happens to me?	0.71			
Social well-being (AVE = 50.3%)		0.89	5.83 (4.94)	0–22
Definite role in family and among friends?	0.58			
Family and friends understand you?	0.68			
Feel useful to family and friends?	0.58			
Feel listened to by family and friends?	0.74			
Can talk about your deepest problems?	0.53			
Know what is happening with family/friends?	0.58			
Satisfied with the relationships you have?	0.71			
Number of family members you can depend on?	0.56			
Number of times spent with someone not living here in past week?	0.65			
Number of times had telephonic conversation with friends/relatives in past week?	0.69			
Number of times attended social meetings/events in past week?	0.64			
AVE = Average Variance Extract				

Independent variables. Relocation autonomy was measured using Perceived Control Measure [29, 39] which comprised of four items (range: 0–8). Higher score on this scale meant higher autonomy. Loneliness was measured through eleven items (range: 0–22) of the de Jong-Gierveld Loneliness Scale (DJGLS-11) [40]. A higher score on DJGLS-11 characterized higher loneliness experienced by the residents. Satisfaction with service quality was measured by the Service Quality Scale (SQS) developed by Lapré [41], which is a 26-items scale (range: 0–52) measuring satisfaction of residents along three dimensions (autonomy, engagement, and relationship with staff). Higher scores on SQS meant higher satisfaction with the services given at the facility. Items comprising these scales along with their psychometric properties are presented in Table 1.

Covariates. The covariates (socio-demographic variables) included in the study were gender, age, marital status, number of children and visitors, time since residence (in years), education, income, social organization (rural/urban), and residents' visiting family/friends (yes/no).

Data analysis

The Statistical Package for Social Sciences (SPSS), version 25, was used to generate descriptive and inferential statistics. Factor analysis using principal components and reliability analysis using Cronbach's alpha were calculated to investigate construct validity and internal consistency of relocation autonomy, loneliness, satisfaction with service quality, and well-being scales. The Kaiser-Meyer-Ohlin (KMO) measure and p-values of Barlett's sphericity were generated to assess sampling adequacy, correlations between items, and linearity. Descriptive statistics such as frequency, percentage, mean, standard deviation (SD), and range of obtained responses were calculated for the variables used in this study. For testing models, Cook's and Leverage distances were used to possibly detect outliers and no outliers were found. The tolerance values for all predictor variables were close to 1 and variance inflation factor (VIF) values were under 5 so multicollinearity

posed no risk [42]. The assumptions of normality, linearity, and homoscedasticity were fulfilled as assessed through histogram, normal p-p plot, and scatter plot, respectively. Three separate multiple regression analyses were carried out to assess the effect of predictor variables on physical, psychological, and social well-being of older adults living in senior care facilities.

Ethics

The study was approved by university's Institutional Review Board of the University of the Punjab, Lahore, Pakistan. Research ethics related to confidentiality, anonymity, informed consent, privacy, and safety were taken care of. Residents were given an overview of the study and those willing to participate signed a consent form. The residents were informed that they would not be compensated for their participation.

Results

Psychometric properties and descriptive statistics of scales administered to older persons

The results showed that KMO values were greater than 0.8 in all scales (relocation autonomy, loneliness, satisfaction with service quality, and well-being scales) which suggested that linearity was present and Barlett's p-values were less than 0.001, which suggested that the items in the relevant scales were not orthogonal. The factor loadings of all items for their relative constructs were greater than or equal to 0.5 and the values for Cronbach's alpha were greater than 0.7 which proved the construct validity and internal consistency of the scales (Table 1). The descriptive statistics (mean, SD, and range of obtained responses) of these scales are also summarized in Table 1. These descriptive statistics show that almost 50% of the residents had higher relocation autonomy, lower loneliness, and higher satisfaction with quality services. Likewise, 50% of the residents had higher physical and psychological well-being. The majority of residents had lower social well-being.

Descriptive statistics of socio-demographic variables and their frequency distribution with respect to loneliness

The frequency distributions and percentages of socio-demographic variables and their breakdown with respect to loneliness, in addition to Chi-Square significance values, are summarized in Table 2. The mean age of older adults was 68.9 (5.79) years. More than half of residents (57.4%) were between the age of 60–69 years, of which 104 (38.5%) experienced loneliness. With respect to gender, 185 (68.5%) were males and 126 (46.6%) of them experienced loneliness in old age homes. The rate of loneliness among females was lower (24.1%). Out of 270 residents, 58.9% were widowed; among them, 43.7% reported loneliness. 238 (88.1%) residents reported that they did not have any source of income, with almost two thirds of them (62.5%) reporting loneliness. 202 (74.8%) residents said that they did not have any visitors and half of them were experienced loneliness. Likewise, 220 (81.5%) of the respondents said that they did not visit their family and friends and 60% of them reported experiencing loneliness.

Table 2: Distribution of socio-demographic variables with respect to loneliness in senior care facilities (n=270)

Variables	n (%)	Loneliness		χ^2 p-value
		Not lonely	Lonely	
		(score ≤ 6)	(score ≥ 7)	
		n (%)	n (%)	
Age (years)				0.085
60–69	155 (57.4)	51 (18.9)	104 (38.5)	
70–79	100 (37.0)	27 (10.0)	73 (27.0)	
80–89	15 (5.6)	1 (0.4)	14 (5.2)	
Gender				0.161
Female	85 (31.5)	20 (7.4)	65 (24.1)	
Male	185 (68.5)	59 (21.9)	126 (46.6)	
Marital status				0.006
Never married	55 (20.4)	25 (9.3)	30 (11.1)	
Currently married	31 (11.5)	4 (1.5)	27 (10.0)	
Widow	159 (58.9)	41 (15.2)	118 (43.7)	
Divorced/separated	25 (9.3)	9 (3.4)	16 (5.9)	
Number of children				<0.001
No children	96 (35.6)	40 (14.8)	56 (20.8)	
1–2	67 (24.8)	26 (9.6)	41 (15.2)	
≥ 3	107 (39.6)	13 (4.8)	94 (34.8)	
Education				0.407
No formal schooling	152 (56.3)	43 (15.9)	109 (40.4)	
Up to five years	38 (14.1)	9 (3.3)	29 (10.8)	
Up to eight years	27 (10.0)	7 (2.6)	20 (7.4)	
\geq Matriculation	53 (19.6)	20 (7.4)	33 (12.2)	
Any source of income				0.792
No	238 (88.1)	69 (25.6)	169 (62.5)	
Yes	32 (11.9)	10 (3.7)	22 (8.2)	
Social organization				0.295
Rural	133 (49.3)	35 (13.0)	98 (36.3)	
Urban	137 (50.7)	44 (16.3)	93 (34.4)	
Duration of residence (years)				0.781
Less than 1 year	43 (15.9)	10 (3.7)	33 (12.2)	

1-Less than 3 years	117 (43.3)	37 (13.7)	80 (29.6)	
3-Less than 5 years	47 (17.4)	14 (5.2)	33 (12.2)	
≥5 years	63 (23.4)	18 (6.7)	45 (16.7)	
Number of visitors				0.072
None	202 (74.8)	62 (23.0)	140 (51.8)	
One	53 (19.6)	9 (3.3)	44 (16.3)	
Two	15 (5.6)	8 (2.9)	7 (2.7)	
Respondent's goes to visit				0.028
No	220 (81.5)	58 (21.5)	162 (60.0)	
Yes	50 (18.5)	21 (7.8)	29 (10.7)	

<i>Predictors</i>	Physical well-being			Psychological well-being			Social well-being		
	B	95% CI	p-value	B	95% CI	p-value	B	95% CI	p-value
Age	-0.16	-0.23–-0.09	< 0.001	-0.01	-0.05–0.04	0.76	0.02	0.05–0.10	0.52
Gender	-0.14	-0.97–0.69	0.74	0.27	-0.29–0.83	0.34	-0.21	-1.06–0.65	0.64
Marital status	0.03	-0.41–0.46	0.91	0.09	-0.20–0.38	0.56	0.29	-0.15–0.74	0.20
Number of children	-0.21	-0.47–0.05	0.11	-0.24	-0.42–-0.07	0.006	0.37	0.10–0.64	0.007
Education	-0.13	-0.44–0.18	0.42	0.02	-0.19–0.22	0.89	-0.06	-0.38–0.26	0.72
Any source of income	-0.15	-1.46–1.17	0.83	0.21	-0.67–1.09	0.64	0.79	-0.56–2.14	0.25
Social organization	1.14	0.33–1.95	0.006	-0.01	-0.54–0.54	0.99	0.77	-0.06–1.60	0.07
Duration of residence	-0.21	-0.37–-0.04	0.01	-0.07	-0.18–0.04	0.24	-0.03	-0.20–0.14	0.70
Number of visitors	0.82	0.19–1.45	0.01	0.80	0.38–1.22	< 0.001	2.40	1.74–3.05	< 0.001
Respondents' visit	0.33	-0.74–1.40	0.54	1.07	0.35–1.78	0.004	2.97	1.87–4.07	< 0.001
Relocation autonomy	0.56	0.34–0.76	< 0.001	0.36	0.21–0.50	< 0.001	0.16	-0.06–0.39	0.16
Loneliness	-0.14	-0.24–-0.04	0.005	-0.19	-0.25–-0.12	< 0.001	-0.36	-0.46–-0.26	< 0.001
Service quality	0.07	0.02–0.12	0.004	0.02	-0.01–0.05	0.23	0.08	0.03–0.13	0.002
Predictors → Physical well-being $F(13,256) = 27.10, p < 0.001, R^2 = 0.579, \text{adj. } R^2 = 0.558$									
Predictors → Psychological well-being $F(13,256) = 37.24, p < 0.001, R^2 = 0.654, \text{adj. } R^2 = 0.637$									
Predictors → Social well-being $F(13,256) = 31.49, p < 0.001, R^2 = 0.615, \text{adj. } R^2 = 0.596$									

Multiple regression analyses estimates for models predicting physical, psychological, and social well-being

The results of multiple regression analyses (Table 3) showed that the model predicting the association of independent variables with physical well-being was statistically significant ($F(13,256) = 27.10, p < 0.001, R^2 = 0.579$). Results show that one unit raise in age decreased physical well-being by 0.16 units ($p < 0.001$). Physical well-being was higher among residents belonging to urban areas ($b = 1.14, p = 0.006$). Duration of stay in the residence was a significant determinant of physical well-being as one unit increase in stay (in years) resulted in 0.21 units decrease in physical well-being ($p = 0.01$).

The number of visitors was also a significant predictor of physical well-being as one unit increase in visitors raised physical well-being by 0.82 units ($p = .01$). Relocation autonomy, loneliness, and satisfaction with service quality were all significant predictors of physical well-being as one unit increase in relocation autonomy increased physical well-being by 0.56 units ($p < 0.001$), one unit increase in loneliness decreased physical well-being by 0.14 units ($p = 0.005$), and one unit increase in satisfaction with service quality increased physical well-being by 0.07 units ($p = 0.004$). The effect of remaining variables on physical well-being was statistically insignificant (Table 3).

The model predicting effect of independent variables on psychological well-being was statistically significant ($F(13,256) = 37.24, p < 0.001, R^2 = 0.654$). Results show that a higher number of children reduces psychological well-being ($b = -0.24, p = 0.006$). Number of visitors and respondents' visit to family and friends were significant predictors of psychological well-being as one unit rise in visitors raised psychological well-being by 0.80 units ($p < 0.001$) and one unit increase in visits resulted in 1.07 units increase in psychological well-being ($p = 0.004$). Relocation autonomy ($b = 0.36, p < 0.001$) and loneliness ($b = -0.19, p < 0.001$) were also significant predictors of psychological well-being. The effect of remaining variables on psychological well-being was statistically insignificant.

Table 3 shows that the model predicting relation of independent variables with social well-being was statistically significant ($F(13,256) = 31.49, p < 0.001, R^2 = 0.615$). Results show that one unit raise in number of children raised social well-being by 0.37 units ($p = 0.007$). Number of visitors ($b = 2.40, p < 0.001$) and respondents' visit to family and friends ($b = 2.97, p < 0.001$) were significant predictors of social well-being. Loneliness and satisfaction with service quality were also significant predictors of social well-being as one unit increase in loneliness decreased social well-being by 0.36 units ($p < 0.001$) and one unit increase in satisfaction with service quality increased social well-being by 0.08 units ($p = 0.002$). The effect of remaining variables on social well-being was statistically insignificant (Table 3).

Discussion

The study found significant associations of various variables with the three dimensions of physical, psychological, and social well-being. Residents who had lower choice in the process of relocation had lower physical and psychological well-being and those who experienced loneliness in the facility had lower physical, psychological, and social well-being. These findings suggest that lower relocation autonomy can result in withdrawal of the residents from their new surroundings which can drive them towards isolation. Some studies which have tried to explore this relation have found that lower well-being can be caused by higher levels of loneliness. One of the causal factors leading to higher levels of loneliness was lower relocation autonomy [29]. These studies highlight that residents who had higher relocation autonomy had higher life satisfaction and lower depression as compared to those who reported lower relocation control [29, 43]. Likewise, another study found that the consequences of lower relocation autonomy were higher depression, anger, and loneliness [44]. Relocating to a senior care facility with low or no control on the process can be traumatic for older persons, which can lead to a loss of interpersonal relations, meanings, identity, and purpose of life [45]. Consistent with previous studies, satisfaction with service quality significantly increased physical and social well-being [46, 47]. Studies have shown that the environment provided in such facilities like autonomy, flexibility, engagement, and respect can increase well-being of older adults [47–49]. A healthy and facilitating environment characterized by quality services can increase confidence, self-respect, and self-worth of the residents which can increase their well-being in the facility [47].

The number of people visiting older adults was a significant predictor of all three types of well-being. As the number of visitors increased, physical, psychological, and social well-being increased. This finding corroborates another finding of the study which suggested that loneliness lowered all three types of well-being in older adults living in old age homes. Researchers pointed out that higher loneliness in such facilities may arise due to a lack of social relationships and support and may thus be brief and reactive [50]. Empirical literature shows that relocation can hamper the development of new social relationships and networks and can, thereby, lower the support available to older adults leading to higher levels

of loneliness and lower well-being [51]. Nevertheless, there is a likelihood that lonely older adults may actively reserve themselves from others which may lead to lower well-being [52]. This finding can be corroborated by another finding of the study which showed that those who had more visitors were less likely to experience loneliness and, therefore, had higher levels of physical, psychological, and social well-being. Similarly, those residents who visited their family and friends had higher psychological and social well-being. This finding is consistent with many studies which confirm that higher social integration and interaction in older adults resulted in higher well-being [53]. Another significant finding of the study was that a higher number of children lowered psychological well-being. This can be explained by the argument that the norms of filial piety in Eastern societies demand that children care for the parents when the latter reach old age. According to this, having more children in these societies is an assurance for the parents that their children will look after them in their late days. Such parents are more likely to face negative psychological consequences and lower well-being when children deny them care in old age, forcing them to seek shelter in old age homes [54–56].

The findings of the study should be interpreted with caution keeping in mind that it was cross-sectional, exploratory, and relational, thus making it difficult to draw causal inferences. Furthermore, the study did not attempt to explore complex interactions such as the role of loneliness and satisfaction quality as mediators of relationship between relocation and well-being of residents of old age homes.

Conclusion

The study proposes pragmatic, evidence-based, and cost-effective interventions for administrators, social workers, and researchers working with older adults in senior care facilities. Interventions such as supporting new residents of senior care residential facilities to become acquainted with the facility and mobilizing staff and adjusted residents to facilitate new residents in structuring relationships can increase the physical and psychological well-being of older adults. Facilitating the residents – particularly new residents – in the adjustment process by offering relocation support programs can help to reduce relocation distress. Mobilizing the staff in these settings to provide social support through discussions, listening to problems, spending time with residents, and making them realize that they are not alone, can reduce well-being problems in senior residential care facilities. Likewise, helping residents to feel comfortable, secure, and engaged in the setting by involving them in various activities taking place in the facility can in turn develop their sense of ownership of the facility which can help to reduce loneliness and increase well-being. Increasing the social support available to the residents by (therapeutic) interventions can increase the social well-being of older adults and may even slow down the cognitive and physical decline. Administrators and managers of senior care facilities should support, facilitate, and encourage volunteers from educational institutions, community, and other welfare institutions to spend time with older adults on regular (e.g. weekly) basis. Likewise, trips may be organized for older residents of senior care facilities so that they can visit places and stay in touch with the outside world. Intergenerational social support can help the residents in the adjustment process. The administrators of old age homes as well as other senior care facilities in Pakistan, South-Asian countries, and other countries with similar contexts, keeping in view the findings of this study, can improve the well-being of older adults residing in these facilities by implementing these interventions.

Abbreviations

DJGLS	de Jong-Gierveld Loneliness Scale
KMO	Kaiser-Meyer-Ohlin
SD	Standard deviation
SPSS	Statistical Package for Social Sciences
SQS	Service Quality Scale

Declarations

Ethics approval and Consent to participate: The study received an ethics approval from the ethics committee at the Advanced Studies and Research Board at the University of the Punjab, Lahore, Pakistan (D.NO/7505/ACAD, dated 07/10/2019). The study was performed in accordance with the Declaration of Helsinki. Written informed consent was taken from all subjects to participate in the study.

Consent for publication: Not applicable.

Availability of data and materials: The dataset for this study is not publicly available due to confidentiality agreement with administrators of senior care facilities. Any individual/organization can make a reasonable request to the corresponding author of this study who will share the dataset after seeking permission from the administrators.

Competing interests: The authors declare that they have no competing interests.

Funding: This study received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

Authors contributions: JT and RZ planned the study and wrote the paper. MVA and AS helped in data collection. RZ and MZZ supervised data analysis and revised the final manuscript. MVA, AS, MZZ and FF contributed to manuscript writing. All authors read and approved the final version of the manuscript.

Acknowledgements: Not applicable.

References

1. Jafree SR, Burhan SK, Khawar A, Mahmood QK, Shahed S. The Impact of Intergenerational Learning on Quality of Life in Older Populations Residing in A Public Sector Old Age Home: A Quasi-experimental Study. *Journal of Intergenerational Relationships*. 2021; doi:10.1080/15350770.2021.1942383.
2. Tariq J, Ali MV, Zakar R, Sajjad A, Tariq H. Institutional Environment of Old Age Homes and Its Relation to Adjustment of Older Adults: Evidence from Pakistan. *Journal of Aging and Environment*. 2020;36(1):40-51. doi:10.1080/26892618.2020.1858385.
3. Global AgeWatch Index: Global Rankings table: Switzerland tops the 2015 index. <https://www.helpage.org/global-agewatch/population-ageing-data/global-rankings-table/> (2015). Accessed 18 Aug 2021.
4. HelpAge International: Ageing population in Pakistan. <https://ageingasia.org/ageing-population-pakistan/> (2019). Accessed 16 Aug 2021.
5. Rizvi Jafree S, Mahmood QK, Burhan SK, Khawar A. Protective Factors for Life Satisfaction in Aging Populations Residing in Public Sector Old Age Homes of Pakistan: Implications for Social Policy. *Journal of Aging and Environment*. 2021;36(2):136-155. doi:10.1080/26892618.2021.1887042.
6. Cassum LA, Cash K, Qidwai W, Vertejee S. Exploring the experiences of the older adults who are brought to live in shelter homes in Karachi, Pakistan: a qualitative study. *BMC Geriatrics*. 2020;20:8. doi:10.1186/s12877-019-1376-8.
7. Ganesh Kumar S, Majumdar A, Pavithra G. Quality of life (QOL) and its associated factors using WHOQOL-BREF among elderly in urban Puducherry, India. *Journal of Clinical and Diagnostic Research*. 2014;8(1):54-57. doi:10.7860/JCDR/2014/6996.3917.
8. Rai S, Khanal P, Chalise HN. Elderly abuse experienced by older adults prior to living in old age homes in Kathmandu. *J Gerontol Geriatr Res*. 2018;7(1): 1000460. doi:10.4172/2167-7182.1000460.

9. Lalan Y. A sociological study of old persons residing in an old age home Delhi, India. *International Research Journal of Social Sciences*. 2014;3(4):21-3.
10. Sethi V, Verma V, Singh U. Impact of ageing on depression and activities of daily livings in normal elderly subjects living in old age homes and communities of Kanpur, UP. *International Journal of Medical Research & Health Sciences*. 2013;2(2):243. doi:10.5958/j.2319-5886.2.2.028.
11. Sabzwari SR, Azhar G. Ageing in Pakistan – a new challenge. *Ageing International*. 2011;36:423-427. doi:10.1007/s12126-010-9082-z.
12. Hashmi, SF, Mubeen B, Tariq MI, Warris SH, Rashid A, Jafar MZ. Prevalence of elder abuse among the patients attending outpatient department of tertiary care hospitals of Lahore, Pakistan. *PJMHS*. 2020;14(1):70-73.
13. Zaidi, A, Stefanoni S, Khalil H. Moving from the margins: Promoting and protecting the rights of older persons in Pakistan. British Council Pakistan. https://www.britishcouncil.pk/sites/default/files/promoting_and_protecting_the_rights_of_older_persons_in_pakistan_british_council_2019.pdf (2019). Accessed 19 Aug 2021.
14. Mushtaq N, Ali R. Challenges to Elderly People in Quetta City (Balochistan) Pakistan. *Elderly Health Journal*. 2020;6(2):91-97. doi:10.18502/ehj.v6i2.5015.
15. Botngård A, Eide AH, Mosqueda L, Blekken L, Malmedal W. Factors associated with staff-to-resident abuse in Norwegian nursing homes: a cross-sectional exploratory study. *BMC Health Services Research*. 2021;21:244. doi:10.1186/s12913-021-06227-4.
16. Yunus RM. Researching institutional elder abuse in Malaysia: challenges and recommendations. *Gerontology*. 2021;67(4):449-452. doi:10.1159/000513932.
17. Withall A, Karystianis G, Duncan D, Hwang YI, Hagos Kidane A, Butler T. Domestic violence in residential care facilities in New South Wales, Australia: a text mining study. *The Gerontologist*. 2021;62(2):223-231. doi:10.1093/geront/gnab068.
18. Lim J. Elder Abuse in Care Facilities: Current State and Suggestions for Preventive Measures. *Research in Brief*. 2021;82:1-8.
19. Andela M, Truchot D, Huguenotte V. Work environment and elderly abuse in nursing homes: the mediating role of burnout. *Journal of Interpersonal Violence*. 2021; 36(11-12):5709-5729. doi:10.1177/0886260518803606.
20. Chen, Z. Knowledge and perceptions: Chinese older adults' willingness to use institutional elder care. Masters' theses, Gerontology Institute, Georgia State University; 2011.
21. Qidwai W. Ageing population: status, challenges and opportunities for health care providers in Pakistan. *Journal of the College of Physicians and Surgeons Pakistan*. 2009;19:399-400.
22. Cesetti G, Vescovelli F, Ruini C. The promotion of well-being in aging individuals living in nursing homes: a controlled pilot intervention with narrative strategies. *Clinical Gerontologist*. 2017;40(5):380-391. doi:10.1080/07317115.2017.1292979.
23. Broadbent J, de Quadros-Wander S, McGillivray JA. Perceived control's influence on wellbeing in residential care versus community dwelling older adults. *Journal of Happiness Studies*. 2014;15(4):845-848. doi:10.1007/s10902-013-9452-9.
24. Street D, Burge S, Quadagno J, Barrett A. The salience of social relationships for resident well-being in assisted living. *The Journals of Gerontology Series B: Psychological Sciences and Social Sciences*. 2007;62(2):S129-S134. doi:10.1093/geronb/62.2.S129.
25. Rohkrämer T, Schulz FR. Space, place and identities. *History Compass*. 2009;7(5):1338-49. doi:10.1111/j.1478-0542.2009.00627.x.

26. Lawton MP. Residential environment and self-directedness among older people. *American Psychologist*. 1990;45(5):638-640. doi:10.1037/0003-066X.45.5.638.
27. Soares E. The care of institutionalized elderly: principles for action of multidisciplinary team. *Gerontology & Geriatric Research*. 2014;3:5. doi:10.4172/2167-7182.1000191.
28. Scocco P, Rapattoni M, Fantoni G. Nursing home institutionalization: a source of eustress or distress for the elderly?. *International Journal of Geriatric Psychiatry*. 2006;21(3):281-287. doi:10.1002/gps.1453.
29. Kim YS. Assessing the effect of relocation control on psychological well-being of assisted living residents. Doctoral dissertation, University of Tennessee, Tennessee; 2012.
30. Rossen EK, Knafl KA. Women's well-being after relocation to independent living communities. *Western Journal of Nursing Research*. 2007;29:2. doi:10.1177/0193945906292539.
31. Borim FS, Neri AL, Francisco PM, Barros MB. Dimensions of self-rated health in older adults. *Revista de Saude Publica*. 2014;48(5):714-722. doi:10.1590/S0034-8910.2014048005243.
32. Fuchs J, Scheidt-Nave C, Hinrichs T, Mergenthaler A, Stein J, Riedel-Heller SG, Grill E. Indicators for healthy ageing – a debate. *International Journal of Environmental Research and Public Health*. 2013;10(12):6630-6644. doi:10.3390/ijerph10126630.
33. Liu WM, Forbat L, Anderson K. Death of a close friend: Short and long-term impacts on physical, psychological and social well-being. *PloS One*. 2019;14(5):e0218026. doi:10.1371/journal.pone.0218026.
34. Baez M, Far IK, Ibarra F, Ferron M, Didino D, Casati F. Effects of online group exercises for older adults on physical, psychological and social wellbeing: a randomized pilot trial. *PeerJ*. 2017;5:e3150. doi:10.7717/peerj.3150.
35. Mansour AMH, Shehadeh JH, Puskar K, El-Hneiti M, M Haourani E. Investigating physical, psychological and social well-being of older persons in Jordan. *Current Aging Science*. 2017;10(3):217-223. doi:10.2174/1874609810666170113093307.
36. Hafeez A, Rafique R. Spirituality and Religiosity as Predictors of Psychological Well-Being in Residents of Old Homes. *Dialogue*. 2013;8(3):285-301.
37. Reker GT, Wong PT. Psychological and physical well-being in the elderly: The Perceived Well-Being Scale (PWB). *Canadian Journal on Aging*. 1984;3(1):23-32. doi:10.1017/S0714980800006437.
38. Wardian J, Robbins D, Wolfersteig W, Johnson T, Dustman P. Validation of the DSSI-10 to measure social support in a general population. *Research on Social Work Practice*. 2013;23:1. doi:10.1177/1049731512464582.
39. Davidson HA, O'Connor BP. Perceived control and acceptance of the decision to enter a nursing home as predictors of adjustment. *The International Journal of Aging and Human Development*. 1990;31(4):307-318.
40. De Jong-Gierveld J, Kamphuis F. The development of a Rasch-type loneliness scale. *Applied psychological measurement*. 1985;9:289-299. doi:10.1177/014662168500900307.
41. Lapré FA. Service quality in nursing homes. A construct, measurement and performance model to increase client focus. Doctoral dissertation, University of Bradford, UK; 2013.
42. Hair JF, Black WC, Babin BJ, Anderson RE, Tatham RL. *Multivariate data analysis*. Upper Saddle River: Prentice Hall; 2010.
43. Chen CK, Zimmerman S, Sloane PD, Barrick AL. Assisted living policies promoting autonomy and their relationship to resident depressive symptoms. *The American Journal of Geriatric Psychiatry*. 2007;15(2):122-129. doi:10.1097/01.JGP.0000247163.49665.5b.
44. Carpenito LJ. *Nursing diagnosis: Application to clinical practice*. Philadelphia: Wolters Kluwer; 2017.
45. Elias SM, Neville C, Scott T. The effectiveness of group reminiscence therapy for loneliness, anxiety and depression in older adults in long-term care: a systematic review. *Geriatric Nursing*. 2015;36(5):372-380. doi:10.1016/j.gerinurse.2015.05.004.

46. Kurup VSK. Life realities of abandoned elderly living under the care of Governmental institutions – a case study. *International Journal of English Language, Literature, & Humanities*. 2014;2(2):293-303.
47. Cheng Y, Rosenberg MW, Wang W, Yang L, Li H. Aging, health and place in residential care facilities in Beijing, China. *Social Science & Medicine*. 2011;72(3):365-372. doi:10.1016/j.socscimed.2010.10.008.
48. Choy KL, Siu KY, Ho TS, Wu CH, Lam HY, Tang V, Tsang YP. An intelligent case-based knowledge management system for quality improvement in nursing homes. *VINE Journal of Information and Knowledge Management Systems*. 2018;48(1):103-121. doi:10.1108/VJIKMS-01-2017-0001.
49. Lee GE. Predictors of adjustment to nursing home life of elderly residents: A cross-sectional survey. *International Journal of Nursing Studies*. 2010;47(8):957-964. doi:10.1016/j.ijnurstu.2009.12.020.
50. Bekhet AK, Zauszniewski JA. Mental health of elders in retirement communities: Is loneliness a key factor?. *Archives of Psychiatric Nursing*. 2012;26(3):214-224. doi:10.1016/j.apnu.2011.09.007.
51. Kudo Y, Saeki K. Reasons for the creation of new social networks by the elderly after relocation. *Health*. 2013;5:31-38. doi:10.4236/health.2013.512A005.
52. Liu L, Gou Z, Zuo J. Social support mediates loneliness and depression in elderly people. *Journal of Health Psychology*. 2016;21(5):750-758. doi:10.1177/1359105314536941.
53. Chen Y, Feeley TH. Social support, social strain, loneliness, and well-being among older adults: An analysis of the Health and Retirement Study. *Journal of Social and Personal Relationships*. 2014;32:2. doi:10.1177/0265407513488728.
54. Cheng ST, Chan AC. Filial piety and psychological well-being in well older Chinese. *The Journals of Gerontology Series B: Psychological Sciences and Social Sciences*. 2006;61(5):262-269. doi:10.1093/geronb/61.5.P262.
55. Cheung CK, Kwan AY. The erosion of filial piety by modernisation in Chinese cities. *Ageing & Society*. 2009;29(2):179-198. doi:10.1017/S0144686X08007836.
56. Dai YT, Dimond MF. Filial Piety. A Cross-Cultural Comparison and Its Implications for the Well-Being of Older Parents. *Journal of Gerontological Nursing*. 1998;24(3):13-18. doi:10.3928/0098-9134-19980301-05.