Grey matter abnormalities in Tourette syndrome: an activation likelihood estimation meta-analysis

Fang Wen

Beijing Children's Hospital

Junjuan Yan

Beijing Children's Hospital

Liping Yu

Beijing Children's Hospital

Fang Wang

Beijing Children's Hospital

Jingran Liu

Beijing Children's Hospital

Ying Li

Beijing Children's Hospital

Yonghua Cui (✉ cuiyonghua@bch.com.cn)

Beijing Children's Hospital, Capital Medical University <https://orcid.org/0000-0002-8244-5884>

Research article

Keywords: Grey Matter, Tourette syndrome, ALE, Thalamus

Posted Date: April 13th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-22072/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Version of Record: A version of this preprint was published at BMC Psychiatry on April 7th, 2021. See the published version at <https://doi.org/10.1186/s12888-021-03187-1>.

Abstract

Background: Tourette syndrome (TS) is a neurodevelopmental disorder defined by the continual presence of primary motor and vocal tics. Grey matter abnormalities have been identified in numerous studies of TS, but conflicting results have been reported. This study was an unbiased statistical meta-analysis of published neuroimaging studies of TS structures.

Methods: A voxel quantitative meta-analysis technique called activation likelihood estimation (ALE) was used. The meta-analysis included six neuroimaging studies involving 247 TS patients and 236 healthy controls. A statistical threshold of $p < 0.05$ was established based on the false discovery rate and a cluster extent threshold of 50 voxels.

Results: We found that grey matter volumes were significantly increased in the bilateral thalamus, right hypothalamus, right precentral gyrus, left postcentral gyrus, left inferior parietal lobule, right lentiform nucleus, and left insula of TS patients compared to those of healthy controls. In contrast, grey matter volumes were significantly decreased in the bilateral postcentral gyrus, bilateral anterior cingulate, bilateral insula, left posterior cingulate, and left postcentral gyrus of TS patients compared to those of healthy controls.

Conclusions: Our present meta-analysis primarily revealed significant increases in grey matter volumes in the thalamus and lentiform nucleus, and decreased grey matter volumes in the anterior cingulate gyrus, of TS patients compared to those in healthy controls. Most of these identified regions are associated with cortico-striato-thalamo-cortical circuits. Further studies with larger sample sizes are needed to confirm these changes in grey matter volumes in TS patients.

1. Background

Tourette syndrome (TS) is a neurodevelopmental disorder defined by the continual presence of primary motor and vocal tics [1]. TS is mainly diagnosed by observing symptoms and by referring to disease history [2]. Magnetic resonance imaging (MRI) has enabled early diagnosis, treatment, evaluation, and pathogenic studies of TS. It has been suggested that TS arises from neurobiological abnormalities, of which imaging studies have revealed many relevant clues [3, 4]. For example, basal ganglia gray matter volume decreased and dorsolateral prefrontal area increased in TS children [5]. Smaller corpus-callosum volumes and thinner sensorimotor cortices have been also reported in structural imaging studies on TS [6, 7]. Some of these regions with grey matter abnormalities may be associated with cortico-striato-thalamo-cortical (CSTC) circuits implicated in TS [5, 8, 9]. However, identification of the specific regions of grey matter abnormalities linked to the CSTC circuit of TS requires further investigation.

Moreover, previous imaging studies on grey matter abnormalities in TS patients have yielded conflicting results. For instance, in previous imaging studies in TS patients, grey matter volumes have been shown to be decreased in the basal ganglia and left hippocampal gyrus, as well as in orbitofrontal, anterior cingulate, and bilateral ventrolateral prefrontal cortices [10], whereas such volumes have been found to be increased in dorsal-lateral prefrontal regions [10, 11], the bilateral ventral putamen [12], posterior thalamus, and hypothalamus. However, decreased grey matter volumes have also been reported in the putamen and some prefrontal regions [13]. Hence, these conflicting findings indicate that grey matter abnormalities in TS patients remain unclear and that further studies are needed.

Therefore, in the present study, we performed a meta-analysis on the results of studies reporting grey matter abnormalities in TS patients. Activation likelihood estimation (ALE), a quantitative meta-analysis for imaging studies, is a statistically-based approach that we used to analyze results of grey matter volumes across studies. Due to most published studies on TS using voxel-based morphometry (VBM) to assess grey matter volumes, we performed VBM-based ALE analysis to assess previous results of grey matter volumes in TS patients.

2. Methods

2.1. Study selection

PubMed, Web of Knowledge, Elsevier, and PsycINFO were searched for articles published between January 31, 2009, and August 31, 2019. The following search terms were used: Tourette syndrome, Tic disorder, TS, MRI, and grey matter. The inclusion criteria for articles were as follows: (1) TS diagnosis based on either the Diagnostic and Statistical Manual of Mental Disorders, 4th Edition, text revision (DSM-IV-TR) or DSM-5 criteria for TS; (2) Grey matter changes were reported by volumes, thicknesses, or densities in structural MRIs; and (3) the foci were reported in either Talairach space or the Montreal Neurological Institute (MNI) space. The exclusion criteria for articles were as follows: (1) duplicated studies; (2) articles that were not original studies; or (3) no coordinates of grey matter were reported.

Ultimately, a total of six studies (247 TS patients and 236 healthy controls) were included in the present meta-analysis. Figure 1 shows a flowchart depicting our steps in the identification of relevant studies. Information on ages, percentages of males, coordinates, measures, Yale Global Tic Severity Scale (YGTSS) values, and the main findings of each study were extracted (Table 1).

Table 1. Demographic characteristics of subjects and Summary of articles included in ALE meta-analysis of TS.

Abbreviations: YGTSS, Yale Global Tic Severity Scale; DOI, Duration Of Illness; HC, Health Control; MNI, Montreal Neurological Institute; GMV, Grey Matter Volume; WMV, White Matter Volume; GMT, Grey Matter Thickness; PSP, Premonitory Sensory P

2.2. Activation likelihood estimation (ALE)

All meta-analysis was accomplished by applying the ALE technique [14], which has been implemented in the Brain Map [15]. ALE analyses were carried out in Talairach space, and a Lancaster transform was conducted if coordinates were instead originally reported in MNI space[16]. Brett's formula [17] and the Lancaster method were used in the transition from MNI space to Talairach space. ALE analysis of TS included six studies comprised 208 foci. Resulting statistical maps were corrected for multiple comparisons using false discovery rates (FDRs) and were then threshold at $p < 0.05$, with a cluster-extent threshold of 50 voxels.

3. Results

ALE results showed that grey matter volumes were significantly increased in the left thalamus, left red nucleus, right thalamus, right hypothalamus, right precentral gyrus, left postcentral gyrus, left inferior parietal lobule, right putamen, and left insula of TS patients compared to those of healthy controls. In contrast, grey matter volumes were significantly decreased in the bilateral postcentral gyrus, bilateral anterior cingulate, bilateral insula, left

Author	Year	YGTSS x±s	N	Age x±s	Males %	DOI	TS, HC	Coordinates	Measure	Main Findings
Greene	2017	18.1±8.3	103	11.9±2.1	78.64	N/A	103, 103	MNI	GMV	The TS group demonstrated lower WM volumes bilaterally in orbital and medial prefrontal cortices, and greater GM volumes in the posterior thalamus, hypothalamus, and midbrain.
Muellner	2015	17.64±7.05		29.5±8.51	55.77	N/A	52, 52	N/A	N/A	Lower depth and reduced thickness of gray matter was found in the pre- and post-central gyri, as well as superior, inferior, and internal frontal sulci.
Draper	2015	N/A	35	14±3.1	91.43	N/A	35, 29	MNI	GMT	PSP was inversely associated with grey matter thickness within insular and sensorimotor cortices. Grey matter thickness was significantly reduced in these areas in individuals with TS. PSP ratings were significantly correlated with tic severity.
Liu	2013	41.71±12.46	21	7.9±1.95	95.24	1.84±0.56	21, 20	MNI	GMV	Brain volume changes were found in the left superior temporal gyrus, bilateral paracentral gyrus, right precuneous cortex, right pre- and post-central gyrus, left temporal-occipital fusiform cortex, right frontal pole, and left lingual gyrus in TS patients. Increases

										were found in the anterior thalamic radiation, right cingulum bundle projecting to the cingulate gurus, and the forceps minor. Decreases in white matter volume (WMV) were found in the right frontal pole. Reductions in grey matter volume were found in orbitofrontal, anterior cingulate, and ventrolateral prefrontal cortices bilaterally. Cortical thinning extended into the limbic mesial temporal lobe..
Draganski	2010	28.7±7.4	40	32.4±11	75	24±11.6	40,40	MNI	GMV& GMT	Decreases in gray matter volumes were found in prefrontal areas, the anterior cingulate gyrus, sensorimotor areas, left caudate nucleus, and left postcentral gyrus. Decreases in white matter volumes were detected in the right inferior frontal gyrus, left superior frontal gyrus, and the anterior corpus callosum. Increases were found in the left middle frontal gyrus and left sensorimotor areas. In MTI, white matter
Müller-Vahl	2009	28.8	19	30.4	100	N/A	19, 20	MNI	GMV	

reductions
were seen in
the right
medial frontal
gyrus and
inferior
frontal gyrus
bilaterally, as
well as in the
right
cingulate
gyrus.

posterior cingulate, and left postcentral gyrus of TS patients compared to those of healthy controls. For more details, see Table 2 and Fig. 2.

Table 2
Results of ALE analyses on grey matter abnormalities in TS patients.

Cluster#	Volume (mm ³)	Peak ALE value	Talairach coordinates(x, y, z)	Brain regions
Groups with increased grey matter volumes				
1	952	0.019318 0.016552817	-12, -30, 10 -14, -28, -2	(L) Thalamus (61.3% Pulvinar, 2.5% Medial Geniculate Body)
2	784	0.01593744 0.015419524	-2, -6, -6 0, -16, -8	(L) Thalamus (Red Nucleus)
3	480	0.019446805	12, -30, 13	(R) Thalamus (100% Pulvinar)
4	424	0.016458694	16, -28, -2	(R) Thalamus (16.7% Pulvinar)
5	384	0.015193136	20, -22, 68	(R) Precentral Gyrus (BA 4)
6	336	0.014851548	8, -4, -14	(R) Hypothalamus
7	80	0.010117263	-56, -18, 33	(L) Postcentral Gyrus (BA 2)
8	80	0.010117263	-36, -34, 45	(L) Inferior Parietal Lobule (BA 40)
9	64	0.008642835	27, -3, 9	(R) Lentiform Nucleus (Putamen)
10	64	0.00938297	-38, 4, 12	(L) Insula (BA 13)
Groups with decreased grey matter volumes				
1	208	0.00997	-56, -18, 34	(L) Postcentral Gyrus (BA 2)
2	96	0.008955	6, 44, 0	(R) Anterior Cingulate (BA 32&24)
3	96	0.009334	-40, -2, 8	(L) Sub-lobar (Insula)
4	80	0.009279	0, 33, -26	(L) Medial Frontal Gyrus (BA 11)
5	80	0.008896	-40, 4, 12	(L) Sub-lobar (Insula)
6	80	0.009279	58, -13, 16	(R) Postcentral Gyrus (BA 43)
7	56	0.009123	-14, 40, 10	(L) Anterior Cingulate (BA 32)
8	56	0.009123	-8, -52, 26	(L) Posterior Cingulate (BA 31)
Abbreviations: ALE, activation likelihood estimation; BA, Brodmann area; L, left; R, right				

4. Discussion

To our knowledge, this is the first ALE analysis of VBM-based MRI studies in TS patients. The main findings of the present study were that grey matter volumes in the thalamus, precentral gyrus, postcentral gyrus, and lentiform nucleus were increased, whereas such volumes were decreased in the anterior cingulate and medial frontal gyrus of TS patients.

4.1 Thalamus

Several imaging studies have found increased total thalamic volumes in children and adults with TS[18]. In the present study, we found that grey matter volume was increased in the thalamus of TS patients, especially within the pulvinar nucleus. The pulvinar nucleus is the largest thalamic nucleus and has robust connectivity with the visual cortex [19]. Previous studies have shown that the most prominent differences in grey matter volumes between TS patients and healthy controls were in the pulvinar nucleus and several other motor nuclei [13]. There is a bidirectional connection between the pulvinar nucleus and the cortex [19]. Medial nucleus pulposus is extensively connected with cortex, including prefrontal cortex, orbital cortex and cingulate cortex [20]. Some regions that connect with the pulvinar nucleus are also associated with TS, such as prefrontal, orbital, and cingulate cortices.

Alternatively, altered thalamic regions (especially the pulvinar nucleus) in TS patients may be due to hyperactive motor circuitry and compensatory mechanisms derived from years of attempting to control tics. In consideration of widespread thalamic afferents and efferents, increased thalamic grey matter volumes in TS patients may be associated with multisensory integration in the thalamus and/or convergence of sensory inputs to cognitive-, motivational-, and /or movement-related cortical areas[13]. It is noteworthy that CSTC circuit dysfunction is thought to cause the occurrence of tics [21]. The pulvinar nucleus is a prototypic association nucleus that takes part in reciprocal cortico-cortical interactions and accelerates synchronized oscillatory activities in functionally related areas of the cortex [19].

4.2 Putamen

Findings from studies on the volume of the putamen in children with TS are inconsistent and have mostly focused on grey matter. For example, some studies have found smaller putamens in children and adults with TS [22, 23], while other studies have shown increased putamen volumes in such patients[13, 24, 25]. In our present meta-analysis, we found that the grey matter volume of the putamen was increased in TS patients. Interestingly, the dorsolateral putamina plays a crucial role in motion planning by projecting the sensorimotor region [26–28]. The possible explanation for our finding is that the volume expansion of the dorsolateral putamina in TS patients might be caused by increased motor activity.

4.3 Other regions associated with TS

In addition, we also found a decreased grey matter volume in the postcentral gyrus (motor-sensory gyrus), which indicates that primary motor pathways may be associated with TS[5]. It has been suggested that sensory-motor pathways may take part in the modulation of tics [29]. In contrast, grey matter volumes in insula have yielded mixed results. Some studies have found smaller insular cortices in children and adults with TS [21], whereas other studies have reported increased grey matter volumes in insular cortices of TS patients[5]. Regardless, these findings suggest that the insular cortex may play an important role in premonitory urges (PUs)[30, 31], which are based on interoceptive awareness [32]. Future studies are needed to further elucidate the role of the insular cortex in PUs.

Additionally, our present meta-analysis revealed decreased grey matter volumes in the medial frontal gyrus, which is consistent with several previous TS studies. Numerous studies have reported reductions in grey matter volumes in prefrontal cortices in TS patients, especially in the orbitofrontal cortex (OFC)[5, 13, 33]. For example,

there is a negative correlation between OFC grey matter volume and tic severity[34]. Additionally, decreased grey matter volumes in the OFC have been reported in children with TS[13]. It was reported that cortical thickness of the right OFC decreased with age compared to the control group [35]. In addition, studies of adult TS patients showed a decrease in OFC gray matter volume, OFC cortical thinning [33, 36]. Changes in OFC activity may involve sensory aspects of twitching. These “uncomfortable feelings or bodily sensations” are referred to as PUs, which have a strong correlation with tic symptoms [37]. Task-specific fMRI studies should be performed in the future to explore the correlation between OFC activity and PUs in TS patients. Additionally, it seems that reduced OFC is found not only in children but also in adults with TS; although the volume of OFC found in this study was small, due to its important role in TS, future studies should follow up on this finding.

In our present meta-analysis, we found that grey matter volumes were reduced in the anterior cingulate gyrus (ACC) in TS patients. Previous studies showed cortical thinning and/or below-normal volumes in the subgenual ACC (sACC) [38]. The severity of the tic was also negatively correlated with tissue change in ACC on the right [34]. From functional imaging studies, tic production may be caused by changes in ACC [30, 39]. ACC activity increased with tic suppression, but ACC hypoperfusion was found in the quiescent period without tic suppression [39]. There was also a positive correlation between tic frequency and ACC activity[38, 40]. ACC abnormalities, however, might appear to play a key role in TS pathology.

In summary, the brain regions with grey matter abnormalities in TS patients that we reported in our present meta-analysis were mainly located in the CSTC circle. Future studies employing other methodologies (e.g., functional imaging and neural networks) will be needed to determine the role of the CSTC circuit in TS.

4.4 Limitations

Three limitations of our study should be acknowledged. First, the results reported in our meta-analysis comprised only six studies; future studies are needed with larger sample sizes. Second, several other factors (e.g., medications and comorbidities) should be considered in future MRI studies; we acknowledge that these factors may have accounted for the structural changes that we found in TS patients. Third, we only used the FDR to correct for our p values, while we did not use the family-wise error (FEW).

5. Conclusion

In the present study, we primarily found that TS patients exhibited grey matter increases in the thalamus and lentiform nucleus, and grey matter decreases in the anterior cingulate gyrus. Most regions identified were associated with CSTC circuitry. Further studies with large sample sizes are needed to confirm these changes in grey matter volumes in TS patients.

Clinical Significance

It found that TS patients exhibited grey matter increases in the thalamus and lentiform nucleus, and grey matter decreases in the anterior cingulate gyrus. Most regions identified were associated with CSTC circuitry.

Abbreviations

TS: Tourette syndrome; ALE: Activation Likelihood Estimation; MRI: Magnetic Resonance Imaging; CSTC: Cortico-Striato-Thalamo-Cortical; VBM: Voxel-Based Morphometry; MNI: Montreal Neurological Institute; YGTSS: Yale Global Tic Severity Scale; OFC: OrbitoFrontal Cortex; ACC: Anterior Cingulate gyrus; FDRs: False Discovery Rates; FEW: Family-Wise Error; PUs: Premonitory Urges; HC: Health Control; GMV: Grey Matter Volume; WMV: White Matter Volume; GMT: Grey Matter Thickness; PSP: Premonitory Sensory Phenomena; DOI: Duration Of Illness; BA: Brodmann area; L: left; R: right

Declarations

Ethics approval and consent to participate Not Applicable.

Consent for publication All authors consent for publishing this article to BMC Psychiatry Journal.

Availability of data and materials All data generated or analyzed during this study are included in this published article and its supplementary information files.

Competing interests The authors declare that they have no competing interests.

Funding This study is supported by the Special Fund of the Pediatric Medical Coordinated Development Center of Beijing Hospitals Authority, No. XTYB201802.

Authors' contributions table

FW did Substantial contributions to the conception and design of the work; and the acquisition, analysis, or interpretation of data for the work and drafted the manuscript. YL did Substantial contributions to the conception and design of the work; and the acquisition, analysis, or interpretation of data for the work, drafted the manuscript and edit the entire English writing. LY and Fang Wang finalized the whole search work and gave suggestions for manuscript. JY contributed in statistical analyses and gave suggestions for manuscript. JL gave suggestions for writing, analysis and revised the manuscript. YC did Substantial contributions to the conception and design of the work; supervise the work from beginning to end, and gave Substantial suggestions to the manuscript. All authors read and approved the final manuscript.

Acknowledgements Not applicable.

Author details

Correspondence should be addressed to:

Yonghua Cui, Beijing Children's Hospital, Capital Medical University, National Center for Children Healthy, 56 Nanlishi Road, Beijing 100101, China; Email: cuiyonghua@bch.com.cn; Tel/Fax number: +86(0)10-59616161.

First Author:

Fang Wen, Beijing Children's Hospital, Capital Medical University, National Center for Children Healthy, 56 Nanlishi Road, Beijing 100101, China; Email: wenfang0812@163.com; Tel/Fax number: +86(0)10-59616161.

Co-Author:

Junjuan Yan, Beijing Children's Hospital, Capital Medical University, National Center for Children Healthy, 56 Nanlishi Road, Beijing 100101, China; Email: Yanzhen0328@163.com; Tel/Fax number: +86(0)10-59616161.

Liping Yu, Beijing Children's Hospital, Capital Medical University, National Center for Children Healthy, 56 Nanlishi Road, Beijing 100101, China; Email: dongyuzui1@163.com; Tel/Fax number: +86(0)10-59616161.

Fang Wang, Beijing Children's Hospital, Capital Medical University, National Center for Children Healthy, 56 Nanlishi Road, Beijing 100101, China; Email: doramon0807@sina.cn; Tel/Fax number: +86(0)10-59616161.

Fang Wang, Beijing Children's Hospital, Capital Medical University, National Center for Children Healthy, 56 Nanlishi Road, Beijing 100101, China; Email: liujingran123@sina.com; Tel/Fax number: +86(0)10-59616161.

Ying Li, Beijing Children's Hospital, Capital Medical University, National Center for Children Healthy, 56 Nanlishi Road, Beijing 100101, China; Email: liying@bch.com.cn; Tel/Fax number: +86(0)10-59616161. Ying Li and Yonghua Cui contributed equally to this study.

References

1. **American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders, 5th edn:** American Psychiatric Association: Arlington, VA; 2013.
2. Wen H LY, Rekik I, Wang S, Chen Z, Zhang J, Zhang Y, Peng Y, He H: **Combining Disrupted and Discriminative Topological Properties of Functional Connectivity Networks as Neuroimaging Biomarkers for Accurate Diagnosis of Early Tourette Syndrome Children.** *Mol Neurobiol* 2018, **55(54)**:3251-3269.
3. Mink: **Neurobiology of basal ganglia and Tourette syndrome: basal ganglia circuits and thalamocortical outputs.** *Adv Neurol* 2006, **99**:89-98.
4. Martino D LJ: **Tourette Syndrome:** Oxford University Press: Oxford; 2013.
5. Liu YM, W.Wang, J. Q.Gao, P. Y.Yin, G. H.Zhang, L. P.Lv, C. K.Ji, Z. Y.Yu, T.Sabel, B. A.He, H. G.Peng, Y.: **Structural Abnormalities in Early Tourette Syndrome Children: A Combined Voxel-Based Morphometry and Tract-Based Spatial Statistics Study.** *Plos One* 2013, **8(9)**.
6. Plessen KJ W-LT, Hugdahl K, Feineigle P, Klein J **Altered interhemispheric connectivity in individuals with Tourette's disorder.** *The American journal of psychiatry* 2004, **161(11)**:2028-2037.
7. Sowell ER KE, Yoshii J, Thompson PM, Bansal R **Thinning of sensorimotor cortices in children with Tourette syndrome.** *Nat Neurosci* 2008, **11(6)**:637-639.
8. Ramkiran S HL, Gaebler A, Shah NJ, Neuner I: **Alterations in Basal Ganglia-Cerebello-Thalamo-Cortical Connectivity and Whole Brain Functional Network Topology in Tourette's Syndrome.** *NeuroImage Clinical* 2019, **24**.
9. Yu Hamamoto YK: **Tourette Syndrome.** *Brain Nerve* 2018, **70(11)**:1237-1245.
10. Peterson K, Scahill, Zhang, Bronen **Basal ganglia volumes in patients with Gilles de la Tourette syndrome.** *Archives of general psychiatry* 2003, **60(4)**:415-424.
11. Peterson S, Scahill, Zhang, Anderson, Leckman et al: **Regional brain and ventricular volumes in Tourette syndrome.** *Archives of general psychiatry* 2001, **58(5)**:427-440.

12. Kassubek J JF, Ludolph AG: **Heterogeneity of voxelbased morphometry findings in Tourette's syndrome: an effect of age?** *Ann Neurol* 2006, **59(5)**:872-873.
13. Greene DJ, Williams Iii AC, Koller JM, Schlaggar BL, Black KJ: **Brain structure in pediatric Tourette syndrome.** *Molecular psychiatry* 2017, **22(7)**(7):972-980.
14. Turkeltaub PE, Eden, G.F., Jones, K.M., & Zeffiro, T.A.: **Meta-analysis of the functional neuroanatomy of single-word reading: Method and validation.** *NeuroImage* 2002, **16(3-part-PA)**:765-780.
15. Laird AR, Fox, P.M., Price, C.J., Glahn, D.C., Uecker, A.M., Lancaster, J.L.: **ALE meta-analysis: Controlling the false discovery rate and performing statistical contrasts.** *Human brain mapping* 2005a, **25(1)**:155-164.
16. Lancaster JL, Tordesillas-Gutierrez D, Martinez M, Salinas F, Evans A, Zille S K, Mazziotta JC, Fox PT: **Bias between MNI and talairach coordinates analyzed using the ICBM-152 brain template.** *Human brain mapping* 2007, **28(11)**(11):1194-1205.
17. Brett: **The MNI Brain and the Talairach Atlas.** Cambridge: MA: MRC Cognition and Brain Sciences Unit; 1999.
18. Miller AM BR, Hao X, Sanchez-Pena JP, Sobel LJ, Liu J **Enlargement of thalamic nuclei in Tourette syndrome.** *Archives of general psychiatry* 2010, **67(9)**:955-964.
19. Benarroch: **Pulvinar: associative role in cortical function and clinical correlations.** *Neurology* 2015, **84(7)**:738-747.
20. Shipp, S.: **The functional logic of cortico-pulvinar connections.** *Philos Trans R Soc Lond B Biol Sci* 2003, **358(1438)**:1605-1624.
21. Amelia Draper GMJ, Paul S. Morgan and Stephen R. Jackson: **Premonitory urges are associated with decreased grey matter thickness within the insula and sensorimotor cortex in young people with Tourette syndrome.** *Journal of Neuropsychology* 2016, **10(1)**:143-153.
22. Bloch MH LJ, Zhu H, Peterson BS: **Caudate volumes in childhood predict symptom severity in adults with Tourette syndrome.** *Neurology* 2005, **65(8)**:1253-1258.
23. Peterson BS, Thomas P, Kane MJ, Scahill L, Zhang H, Bronen R, King RA, Leckman JF, Staib L: **Basal Ganglia Volumes in Patients With Gilles de la Tourette Syndrome.** *Archives of general psychiatry* 2003, **60(4)**(4):415.
24. Ludolph AG JF, Libal G, Ludolph AC, Fegert JM, Kassubek J: **Greymatter abnormalities in boys with Tourette syndrome: magnetic resonance imaging study using optimised voxel-based morphometry.** *Br J Psychiatry* 2006, **188(5)**:484-485.
25. Roessner V OS, Schmidt-Samoa C, Baudewig J, Dechent P, Rothenberger A et al: **Increased putamen and callosal motor subregion in treatment-naïve boys with Tourette syndrome indicates changes in the bihemispheric motor network.** *Child Psychol Psychiatry* 2011, **52(3)**:306-314.
26. Alexander GE CM, DeLong MR: **Basal ganglia-thalamocortical circuits: parallel substrates for motor, oculomotor, "prefrontal" and "limbic" functions.** *Prog Brain Res* 1990, **85**:119-149.
27. Draganski B KF, Kloppel S, Cook PA, Alexander DC, Parker GJ: **Evidence for segregated and integrative connectivity patterns in the human Basal Ganglia.** *J Neurosci* 2008, **28**:7143-7152.
28. Mary Mufford JC, Neda Jahanshad , Celia van der Merwe , Linda Ding , Nynke Groenewold , Nastassja Koen , Emile R Chimusa , Shareefa Dalvie, et al.: **Concordance of Genetic Variation That Increases Risk for Tourette Syndrome and That Influences Its Underlying Neurocircuitry.** 2019, **9(1)**:120.
29. Shprecher D KR: **The Management of Tics.** *Movement disorders clinical practice* 2009, **24**(15-24).

30. Bohlhalter S, Goldfine, A., Matteson, S., Garraux, G., Hanakawa, T., Kansaku, K., Wurzman, R., & Hallett, M.: **Neural correlates of tic generation in Tourette syndrome: An event-related functional MRI study.** *Brain* 2006, **129**:2029-2037.
31. Ganos C, Kassavetis P, Cerdan M, Erro R, Balint B, Price G, Edwards MJ, Bhatia KP: **Revisiting the Syndrome of "Obsessional Slowness".** *Movement disorders clinical practice* 2015, **2**(2):163-169.
32. Craig AD: **How do you feel now? The anterior insula and human awareness.** *Nature Reviews Neuroscience* 2009, **10**(1):59-70.
33. Draganski B, Martino D, Cavanna AE, Hutton C, Orth M, Robertson MM, Critchley HD, Frackowiak RS: **Multispectral brain morphometry in Tourette syndrome persisting into adulthood.** *Brain* 2010, **133**:3661-3675.
34. Muller-Vahl KR, Kaufmann J, Grosskreutz J, Dengler R, Emrich HM, Peschel T: **Prefrontal and anterior cingulate cortex abnormalities in Tourette Syndrome: evidence from voxel-based morphometry and magnetization transfer imaging.** *Bmc Neurosci* 2009, **10**(1).
35. Fahim C YU, Das S, Lyttelton O, Chen J, Arnaoutelis R et al: **Somatosensorymotor bodily representation cortical thinning in Tourette: effects of tic severity,age and gender.** *Cortex* 2010, **46**(6):750-760.
36. Worbe Y GE, Hartmann A, Valabrégue R, Chupin M, Tremblay L: **Distinct structural changes underpin clinical phenotypes in patients with Gilles de la Tourette syndrome.** *Brain* 2010, **133**(12):3649-3660.
37. Li Y WF, Liu J, Wen F, Yan C, Zhang J, Lu X, Cui Y.: **The Correlation Between the Severity of Premonitory Urges and Tic Symptoms: A Meta-Analysis.** *Journal of child and adolescent psychopharmacology* 2019, **29**(9) (9):652-658.
38. O'Neill J PJ, Peterson BS: **Cingulate role in Tourette syndrome.** *Handbook of Clinical Neurology* 2019, **166**.
39. Moriarty J CD, Schmitz B, Trimble MR, Ell PJ, Robertson MM: **Brain perfusion abnormalities in Gilles de la Tourette's syndrome.** *Br J Psychiatry* 1995, **167**(2):249-254.
40. Stern E SD, Chee KY, Holmes A, Robertson MM, Trimble M, et al.: **A functional neuroanatomy of tics in Tourette syndrome.** *Archives of general psychiatry* 2000, **57**(8):741-748.

Figures

Figure 1

The flowchat for the identification of the included study

Figure 2

Results from the ALE meta-analysis on grey matter increased volumes in patients with TS compared with healthy controls

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [PRISMA2009checklist.pdf](#)