Factors associated with maternal overall Quality of Life six months postpartum

Lisbeth Valla (✉ lisbeth.valla@oslomet.no)

OsloMet - storbyuniversitetet <https://orcid.org/0000-0001-6742-7787>

Sølvi Helseth

OsloMet - storbyuniversitetet

Milada Cvancarova Småstuen

OsloMet - storbyuniversitetet

Nina Misvær

OsloMet - storbyuniversitetet

Randi Andenæs

OsloMet - storbyuniversitetet

Research article

Keywords: Quality of Life, Postpartum, the Norwegian Mother, Father and Child Cohort Study

Posted Date: March 19th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-222632/v2>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background: Having good Quality of Life (QoL) is essential, particularly for women after childbirth. However, little is known about the factors associated with maternal QoL after giving birth. We aimed to investigate the relationship between characteristics of the mother (socio-demographic variables), selected symptoms (depression and joy/anger), health perception (perception of birth) and possible characteristics of the environment (infant temperament, colic, sleep, parental relationship), with mothers' overall quality of life when the child is six months of age.

Methods: This study is based on the Norwegian Mother, Father and Child Cohort Study (MoBa), conducted at the Norwegian Institute of Public Health from June 1999 to December 2008, which included a total of 86,724 children. Maternal QoL was assessed by the Satisfaction With Life Scale. Joy and anger were measured using the Differential Emotional Scale, mothers' mental health was assessed using the Edinburgh Postnatal Depression Scale and satisfaction with relationship was measured using the Relationship Satisfaction Scale. Child temperament was measured using the Infant Characteristics Questionnaire and colic, sleep duration and feelings related to childbirth were assessed by mothers' reports. The associations between life satisfaction and selected variables were analysed using stepwise multiple linear regression models.

Results: Maternal feelings of joy of having a baby ($B=0.35$, CI [0.79 to 0.83]), high relationship satisfaction ($B=0.02$, CI [0.28 to 0.60]), as well as having a baby with normal sleep ($B=0.32$, CI [0.21 to 0.22]), are factors associated with higher maternal overall QoL. Postnatal depression was negatively associated with mothers' QoL, and infant colic or child's temperament (fussiness) showed no such association with mothers' QoL.

Conclusions: Health professionals and clinicians should focus on these issues when they develop health interventions or provide counselling to new mothers and their families.

Background

Giving birth and caring for a new infant mark an important transition in life. Although this transition is often exciting and rewarding, it is also a time of vulnerability and many challenges including sleep disruption (1) and heightened psychosocial stress (2), which in turn pose a risk to successful emotional and psychological adjustments that ensure quality of life (3). Theorists have conceptualised quality of life (QoL) in multiple ways, including satisfaction with life (4), subjective well-being (5), and happiness (6). In medical and health science literature, the term 'health-related quality of life' is most commonly used (7). Wilson and Cleary propose a theoretical framework for QoL and suggest that symptoms, functional status, health perception, personal factors and environmental characteristics, in addition to biological and physiological factors, are related to an individual's overall quality of life. In the present study, we use the term QoL throughout this article to characterise a broad representation, and define overall quality of life in accordance with Diener (1984) as subjective well-being, relating to an evaluation

of how people feel and think about their lives (8, 9). QoL constitutes an important goal for individuals and societies with positive consequences for population health (10, 11). However, the overall QoL of mothers may be of particular significance, as it may not only affect the mother herself but also the way she interacts with her child and the environment (12).

Several factors impact maternal QoL postpartum. Issues mentioned in the literature such as infant temperament or fussiness, are likely to affect maternal well-being (13). Infant colic and poor sleep behaviour can give rise to negative emotions in the mother that overshadow the joy of being a mother. Colic is infant fussing and crying that occurs in the infant's second or third week of life, lasts more than three hours a day and gradually resolves in the third or fourth month (14, 15). A study of mothers of newborns (N=78) reported that those with an infant with colic had more symptoms of depression and lower well-being (16). During the postpartum period, most women experience significant disturbances in their sleep patterns due to hormonal changes and responsibility for the care of the newborn (17). Postpartum sleep quality is a critical health index and sleep deprivation is the main source of stress, anxiety and depression for postpartum women (18, 19). In addition, women's QoL after childbirth may also be affected by factors related to pregnancy and delivery (20-22) and their marital relationship (3, 23, 24). Research indicates that women who perceived they were 'very well' cared for during delivery, and women who felt their overall birth experience was 'very positive', were significantly more likely to experience high postnatal functioning (21). On the other hand, women with a negative childbirth experience are at an increased risk of psychological maladjustment in the first five months after giving birth (22).

The majority of previous research focusing on health and quality of life are about the immediate postnatal (21) or postpartum period (25, 26), or risk populations, such as premature birth (27).

There is a paucity of knowledge regarding the role of individual or environmental factors in relation to mothers' QoL after the puerperal period, from large unselected populations. To provide a rich picture of the factors influencing maternal quality of life, the aim of the present study was to investigate the relationship between characteristics of the mother (i.e. socio-demographic variables), selected symptoms (i.e. depression and joy/anger), health perception (i.e. perception of birth), and possible characteristics of the environment (i.e. infant temperament, colic, sleep, parental relationship), with mothers' overall quality of life when the child is about six months.

Methods

Participants and sample

The present study used data from the Norwegian Mother, Father and Child Cohort Study (MoBa), conducted by the Norwegian Institute of Public Health from 1999 through 2008 (28). All pregnant women in Norway were eligible to participate in MoBa if they were able to read Norwegian. The MoBa cohort included 114,500 children and 95,200 mothers yielding a response rate of around 41% (28). Written

informed consent was obtained from all MoBa participants upon recruitment and participants did not receive any financial compensation. The establishment of MoBa and the initial data collection were based on a licence from the Norwegian Data Protection Agency and approval from the Regional Committees for Medical and Health Research Ethics. The MoBa cohort is now based on regulations related to the Norwegian Health Registry Act. The current study was approved by the Regional Committees for Medical and Health Research Ethics (project no; 2017/2199) and is based on version 10 of the MoBa data files. Written informed consent was obtained from all MoBa participants. The sample in the present study constitutes 86,724 mothers and children. Due to possible bias, twins (n=1,480) and triplets (n=14) were excluded.

Measurements

Overall quality of life

Quality of life was measured using the *Satisfaction With Life Scale (SWLS)* (8) six months postpartum. The SWLS includes five items, such as 'The conditions of my life are excellent' and 'I am satisfied with my life', rated on a Likert scale ranging from 1 = Strongly Disagree, to 7 = Strongly Agree. Mean scores of the five items were computed. The scale revealed a satisfactory internal consistency, with a value 0.89. Several studies have confirmed the validity of (8, 29).

Socio-demographic data

Information on child gender, gestation age and maternal age, maternal educational level, marital status, work status and experience of birth was collected from the MoBa study at 16-20 weeks of pregnancy and six months postpartum.

Symptoms

The Edinburgh Postnatal Depression Scale (EPDS)

Mothers' mental health was assessed using the *Edinburgh Postnatal Depression Scale (EPDS)*. The EPDS has a sensitivity of 86% and a specificity of 78% (30). Items are scored on a Likert scale ranging from 0 ('no, never') to 3 ('yes, almost all the time'). A 6-item version was used in the MoBa study. The maximum score is 18, with high scores indicating higher levels of postnatal depression. The internal consistency: Cronbachs α was 0.84 (28).

The Differential Emotional Scale (DES)

The Differential Emotional Scale (31) consists of a series of subscales that capture various emotions. In the present study, items were selected from the Enjoyment and Anger subscales (DES-IV), and included 3 items of Joy and 3 items of Anger. The respondents rated how often they felt the different emotions on a scale from 1 to 5 where 1 = seldom/never, and 5 = very often. Mean scores were computed. The DES has been widely used in research on self-rated affect and has shown good psychometric properties (31, 32).

Feelings related to childbirth

Feelings related to childbirth were assessed six months postpartum based on the single statement '*I felt safe and in good hands*', with the following response options; applies well, applies partly, does not apply.

Environment

Child's sleep

Sleep duration at six months was assessed using the question, 'How many hours does your child sleep per day'. Response categories were 10 hours or less, 11 to 12 hours, 13 to 14 hours, and 15 hours or more. We defined that the normal sleep of a child aged around six months was 13 hours or more a day.

Infant colic

Colic was measured at 6 months using one question in the questionnaire, '*Has your child had the following illness/health problem? Infant colic?*' The response categories were yes or no. Parents thus answered yes if they thought their child had had colic, and no other diagnostic criteria were needed.

Child temperament

Child temperament was measured at six months using the *Infant Characteristics Questionnaire (ICQ)* (33). The original ICQ consists of 24 items describing infant behaviour. Four subscales have been identified; Fussy/Difficult, Unadaptable, Dull and Unpredictable, with varying internal consistency previously reported (.79, .75, .39, .50) (Bates et al 1979). The scale used in MoBa is the Fussy/Difficult scale (8 items). Mothers rank each item on a 7-point Likert scale (1-7), indicating the level of perceived difficulty of handling the described behaviour. Responses to negatively framed questions on the Fussy/Difficult scale were reversed, so that lower scores reflect more difficulty. The total score ranged from 7 to 49 and the internal consistency was 0.69.

Satisfaction with relationship

Satisfaction with relationship was measured using the *Relationship Satisfaction Scale (RS)* (34). RS is a 10-item scale developed for the MoBa study based on previously developed scales. The scale is validated, and showed good psychometric properties (34). The RS refers to a 'partner' instead of being limited to marital spouses, and includes the items 'I am satisfied with the relationship with my partner', 'My partner and I have problems in our relationship', 'I am very happy in my relationship', 'My partner is generally understanding', and 'We agree on how children should be brought up'. Responses were rated on a 6-point Likert scale ranging from 1 = *Strongly Disagree* to 6 = *Strongly Agree*. Negative responses were reversed, and items added to form a sum score with a maximum score of 60.

Statistical Analysis

Data are described with the means and standard deviations (SD) for continuous variables and with frequencies and percentages for categorical data. The associations between life satisfaction and selected variables were analysed using stepwise multiple linear regression models. Variables that were associated with QoL – based on the Wilson & Cleary model and our clinical knowledge – were included in

the multiple regression model. The results are presented as regression coefficients, standard errors (SE) and p -values. The assumptions for linear regression were fulfilled as the residuals were normally distributed. All analyses were considered exploratory, so no correction for multiple testing was performed and p -values <0.05 were considered statistically significant. All statistical analyses were carried out using SPSS, version 25 (IBM Corp, Armonk, NY).

Results

The participants' characteristics are described in Table 1. There were 51.0% boys in the sample, and a large proportion of mothers had a medium to high level of education and were married or co-habiting.

Results from the stepwise linear regression analyses estimating associations between selected factors and mothers' overall life satisfaction are presented in Table 3. As described in the method section, each block of covariates was added step by step. In block 1, lower age of the mother, higher level of education and living alone were all associated with higher life satisfaction scores. Age and educational level remained significant throughout the regression analyses. The second step of the linear regression analysis, adjusted for confounding variables such as age, educational level and marital status, showed that low depression scores, higher feelings of joy, lower feelings of anger and higher feelings of having been taken care of during birth, were all significantly related to higher life satisfaction.

The third step of the analysis, during which the environmental factors were entered, and with control for the variables in step 2, showed that neither infant colic nor the child's temperament were associated with mothers' life satisfaction, while children sleeping 13 hours or more a day, showed significant association. Living alone lost its significant association, while higher partnership satisfaction scores were significantly associated with higher QoL. The final model explained 49% of the variance in mothers' life satisfaction.

Discussion

Our results, from a large Norwegian representative sample six months after birth, suggest that feelings of joy of having a baby, high relationship satisfaction, as well as having a baby with normal sleep, are factors associated with higher maternal overall QoL. Postnatal depression was negatively associated with mothers' QoL, and infant colic or child temperament (fussiness) showed no such association with mothers' QoL.

In accordance with previous studies, our results indicated that positive emotions such as feelings of joy of motherhood are an important variable in maternal overall QoL (3, 35-37). One explanation for these findings may be that joy and positive emotions in motherhood serve as a buffer against stress (38, 39), and are important components of connectedness in the parent-child attachment that enhances psychological health and well-being (40). Wilson and Cleary (1995) theorised that psychological symptoms and emotions influence physical or social functioning, role functioning, mental health and general health perceptions (7). Thus, the feelings of joy mothers experience as a result of having children

are likely to contribute to their global happiness (7, 9). Consequently, children may benefit as well. Indeed, joy and positive emotions in the parent-child interaction have been associated with positive outcomes for the child (12). However, our results showed that postnatal depression was negatively associated with mothers' overall QoL. A majority of studies emphasise the postpartum period as a particularly vulnerable time during which women are at increased risk of mental disorders (41-43), and that postpartum depression has negative consequences on maternal QoL (44).

Maternal QoL was also positively affected by environmental characteristics such as high relationship satisfaction and infant sleep quality (13 hours or more a day at six months). Previous research has demonstrated that relationship dissatisfaction (3, 22-24, 45), and poor infant sleep quality are associated with higher maternal depression scores (46-48) and lower QoL (3). Nelson et al (2014) proposed in a previous review that parents of young children are unhappy to the extent that they encounter sleep disturbance and troubled marriages. Among married/cohabiting couples, the partner is usually the major provider of various types of social support, and living in an emotionally supportive relationship may protect their physical and psychological health in demanding situations. A person's perception of relationship quality may therefore directly affect their wellbeing by eliciting positive emotions or by serving as a buffer against the potentially harmful effects of other external stressors (22).

Notably, neither infant colic nor child temperament (fussiness) showed any association with mothers' QoL. Although factors such as infant colic and fussiness are highly stressful and strongly associated with maternal depression (49, 50) when they are ongoing, our results show that when the child is six months, a history of colic and fussiness has no effect on mothers' QoL. This may indicate that these problems are transient and that the mother's QoL at six months is not affected by such previous challenges. However, higher levels of maternal social support during pregnancy and postpartum have previously been associated with lower rates of maternal reported infant colic (51).

Our findings have important implications for practice and clinical counselling. First, mothers' overall QoL is a significant indicator of positive mental health that may affect the parent-child dyad, which in turn may influence the child's development (10, 12). Our results indicate that healthcare providers should raise awareness about mothers' joy of motherhood, children's sleep habits and encourage parents to look after their relationship. Health professionals should inform both parents of the importance of enhancing partnership relations and partner support. Secondly, our findings highlight the transient nature of the phenomenon of colic, and the short period in which it places a burden on parents. Our findings may be considered in the design of effective health promotion strategies that enable new mothers to maintain optimal health and QoL for themselves and their infants. Identifying health promotion factors that may be particularly positive for life satisfaction in this phase may enable policymakers to prioritise and develop interventions or programmes, or provide social support that will benefit individuals and/or families.

Although the present study has many strengths, including a large national sample, the results must be interpreted in light of several methodological limitations. First, the assessment of feelings related to

childbirth, sleep or colic did not include validated or objective measures. The results may also be impacted by a selection bias because of attrition in the MoBa study, such as higher maternal age, and fewer health-related risks among those who did not participate in the study (28). Hence, the results must be interpreted with some caution.

Conclusions

In conclusion, the present study's results show that joy of motherhood, relationship satisfaction and infant sleep are the most important factors for mothers' overall QoL six months after birth, while a history of infant colic showed no association. Health professionals and clinicians should focus on these issues when they develop health interventions or provide counselling to new mothers and their families.

List Of Abbreviations

MoBa: The Norwegian Mother, Father and Child Cohort Study QoL: Quality of Life

Declarations

Ethics approval and consent to participate: Not applicable

Written informed consent was obtained from all MoBa participants upon recruitment. The study was approved by the Regional Committee for Medical Research Ethics in South-Eastern Norway

Consent for publication: Not applicable

Availability of data and materials: The data that support the findings of this study are available from Norwegian Institute of Public Health, but restrictions apply to the availability of these data, which were used under license for the current study, and so are not publicly available. Data are however available from the author upon reasonable request and with permission of the Norwegian Institute of Public Health

Competing interests: The authors declare that they have no competing interests.

Funding: This study did not receive any specific funding.

Authors' contributions:

LV: Responsibility for the study design, participated in the analytic framework of the study, interpretation and in the writing of the manuscript

SH: Primary responsibility for the study design, Participated in the analytic framework of the study, with the data interpretation, and in the writing of the manuscript

MCS: Was involved in and supervised the statistical analysis.

NM: Has critically revising the manuscript for important intellectual content

RA: Has been involved in the study design, statistical analysis, and in writing of the manuscript

All the authors have given their final approval of the final version of the manuscript.

Acknowledgements: The Norwegian Mother, Father and Child Cohort Study is supported by the Norwegian Ministry of Health and Care Services and the Ministry of Education and Research. We are grateful to all the families in Norway who are taking part in this ongoing follow-up study.

References

1. Hagen EW, Mirer AG, Palta M, Peppard PE. The sleep-time cost of parenting: sleep duration and sleepiness among employed parents in the Wisconsin Sleep Cohort Study. *Am J Epidemiol*. 2013;177(5):394-401.
2. Reid KM, Taylor MG. Social support, stress, and maternal postpartum depression: A comparison of supportive relationships. *Soc Sci Res*. 2015;54:246-62.
3. Nelson SK, Kushlev K, Lyubomirsky S. The pains and pleasures of parenting: when, why, and how is parenthood associated with more or less well-being? *Psychol Bull*. 2014;140(3):846-95.
4. Diener E. Subjective well-being. *Psychological Bulletin*. 1984;95(3):542-75.
5. Diener E, Lucas RE. Subjective well-being: Three decades of progress. *Psychological Bulletin*. 1999;125(2):276.
6. Brülde B. Happiness and the Good Life. Introduction and Conceptual Framework. *An Interdisciplinary Forum on Subjective Well-Being*. 2007;8(1):1-14.
7. Wilson IB, Cleary PD. Linking clinical variables with health-related quality of life. A conceptual model of patient outcomes. *Jama*. 1995;273(1):59-65.
8. Diener E, Emmons RA, Larsen RJ, Griffin S. The Satisfaction With Life Scale. *J Pers Assess*. 1985;49(1):71-5.
9. Diener E, Pressman SD, Hunter J, Delgadillo-Chase D. If, Why, and When Subjective Well-Being Influences Health, and Future Needed Research. *Appl Psychol Health Well Being*. 2017;9(2):133-67.
10. Howell RT, Kern ML, Lyubomirsky S. Health benefits: Meta-analytically determining the impact of well-being on objective health outcomes. *Health Psychology Review*. 2007;1:83–136.
11. Steptoe A, Deaton A, Stone AA. Subjective wellbeing, health, and ageing. *Lancet*. 2015;385(9968):640-8.
12. Sameroff A. The transactional model of development: How children and contexts shape each other. Washington, DC: American Psychological Association; 2009.
13. Brenning K, Soenens B, Mabbe E, Vansteenkiste M. Ups and Downs in the Joy of Motherhood: Maternal Well-Being as a Function of Psychological Needs, Personality, and Infant Temperament. *Journal of Happiness Studies*. 2019;20(1):229-50.
14. Wolke D, Bilgin A, Samara M. Systematic Review and Meta-Analysis: Fussing and Crying Durations and Prevalence of Colic in Infants. *J Pediatr*. 2017;185:55-61.e4.

15. Wessel MA, Cobb JC, Jackson EB, Harris GS, Jr., Detwiler AC. Paroxysmal fussing in infancy, sometimes called colic. *Pediatrics*. 1954;14(5):421-35.
16. Abacı FB, Gökçe S, Tuygun N, Karacan CD, Öner Ö. Psychosocial status and quality of life in mothers of infants with colic. *The Turkish Journal Of Pediatrics*. 2013;55(4):391-5.
17. Hunter LP, Rychnovsky JD, Yount SM. A selective review of maternal sleep characteristics in the postpartum period. *J Obstet Gynecol Neonatal Nurs*. 2009;38(1):60-8.
18. Henderson J, Alderdice F, Redshaw M. Factors associated with maternal postpartum fatigue: an observational study. *BMJ Open*. 2019;9(7):e025927.
19. Giallo R, Gartland D, Woolhouse H, Brown S. "I didn't know it was possible to feel that tired": exploring the complex bidirectional associations between maternal depressive symptoms and fatigue in a prospective pregnancy cohort study. *Arch Womens Ment Health*. 2016;19(1):25-34.
20. Rezaei N, Tavalaei Z, Sayehmiri K, Sharifi N, Daliri S. The relationship between quality of life and methods of delivery: A systematic review and meta-analysis. *Electron Physician*. 2018;10(4):6596-607.
21. Michels A, Kruske S, Thompson R. Women's postnatal psychological functioning: the role of satisfaction with intrapartum care and the birth experience. *Journal of Reproductive and Infant Psychology*. 2013;31(2):172-82.
22. Lemola S, Stadlmayr W, Grob A. Maternal adjustment five months after birth: the impact of the subjective experience of childbirth and emotional support from the partner. *Journal of Reproductive and Infant Psychology*. 2007;25(3):190-202.
23. Delicate A, Ayers S, McMullen S. A systematic review and meta-synthesis of the impact of becoming parents on the couple relationship. *Midwifery*. 2018;61:88-96.
24. Luhmann M, Hofmann W, Eid M, Lucas RE, King L. Subjective Well-Being and Adaptation to Life Events: A Meta-Analysis. *Journal of Personality and Social Psychology*. 2012;102(3):592-615.
25. Van der Woude DAA, Pijnenborg JMA, de Vries J. Health status and quality of life in postpartum women: a systematic review of associated factors. *European Journal Of Obstetrics, Gynecology, And Reproductive Biology*. 2015;185:45-52.
26. Zubaran C, Foresti K. Investigating quality of life and depressive symptoms in the postpartum period. *Women and Birth*. 2011;24(1):10-6.
27. Hill PD, Aldag JC. Maternal perceived quality of life following childbirth. *Journal of Obstetric, Gynecologic, & Neonatal Nursing: Clinical Scholarship for the Care of Women, Childbearing Families, & Newborns*. 2007;36(4):328-34.
28. Magnus P, Birke C, Vejrurp K, Haugan A, Alsaker E, Daltveit AK, et al. Cohort Profile Update: The Norwegian Mother and Child Cohort Study (MoBa). *International Journal Of Epidemiology*. 2016;45(2):382-8.
29. Pavot W, Diener E, Colvin CR, Sandvik E. Further validation of the Satisfaction with Life Scale: evidence for the cross-method convergence of well-being measures. *J Pers Assess*. 1991;57(1):149-61.

30. Cox JL, Holden JM, Sagovsky R. Detection of postnatal depression. Development of the 10-item Edinburgh Postnatal Depression Scale. *The British Journal Of Psychiatry: The Journal Of Mental Science*. 1987;150:782-6.
31. Izard CE, Libero DZ, Putnam P, Haynes OM, Geen R. Stability of Emotion Experiences and Their Relations to Traits of Personality. *Journal of Personality and Social Psychology*. 1993;64(5):847-60.
32. Blumberg SH, Izard CE. Affective and cognitive characteristics of depression in 10- and 11-year-old children. *J Pers Soc Psychol*. 1985;49(1):194-202.
33. Bates JE, Freeland CA, Lounsbury ML. Measurement of infant difficultness. *Child Dev*. 1979;50(3):794-803.
34. Røysamb E, Vittersø J, Tambs K. The Relationship Satisfaction scale - psychometric properties. *Norsk epidemiologi*. 2014;24(1-2):187-94.
35. Nelson SK, Kushlev K, English T, Dunn EW, Lyubomirsky S. In defense of parenthood: children are associated with more joy than misery. *Psychol Sci*. 2013;24(1):3-10.
36. Berntsen D, Rubin DC, Siegler IC. Two versions of life: emotionally negative and positive life events have different roles in the organization of life story and identity. *Emotion*. 2011;11(5):1190-201.
37. Deaton A, Stone AA. Evaluative and hedonic wellbeing among those with and without children at home. *Proc Natl Acad Sci U S A*. 2014;111(4):1328-33.
38. Tugade MM, Fredrickson BL. Resilient individuals use positive emotions to bounce back from negative emotional experiences. *J Pers Soc Psychol*. 2004;86(2):320-33.
39. Cohn MA, Fredrickson BL, Brown SL, Mikels JA, Conway AM. Happiness unpacked: positive emotions increase life satisfaction by building resilience. *Emotion*. 2009;9(3):361-8.
40. Deci EL, Ryan RM. Self-determination theory: A macrotheory of human motivation, development, and health. *Canadian Psychology/Psychologie canadienne*. 2008;49 182–5.
41. O'Hara MW, McCabe JE. Postpartum depression: current status and future directions. *Annu Rev Clin Psychol*. 2013;9:379-407.
42. Vesga-López O, Blanco C, Keyes K, Olfson M, Grant BF, Hasin DS. Psychiatric disorders in pregnant and postpartum women in the United States. *Arch Gen Psychiatry*. 2008;65(7):805-15.
43. Munk-Olsen T, Laursen TM, Pedersen CB, Mors O, Mortensen PB. New parents and mental disorders: a population-based register study. *Jama*. 2006;296(21):2582-9.
44. Slomian J, Honvo G, Emonts P, Reginster JY, Bruyère O. Consequences of maternal postpartum depression: A systematic review of maternal and infant outcomes. *Womens Health (Lond)*. 2019;15:1745506519844044.
45. Doss BD, Rhoades GK, Stanley SM, Markman HJ. The effect of the transition to parenthood on relationship quality: an 8-year prospective study. *J Pers Soc Psychol*. 2009;96(3):601-19.
46. Dørheim SK, Bondevik GT, Eberhard-Gran M, Bjorvatn B. Sleep and depression in postpartum women: a population-based study. *Sleep*. 2009;32(7):847-55.

47. Hall WA, Moynihan M, Bhagat R, Wooldridge J. Relationships between parental sleep quality, fatigue, cognitions about infant sleep, and parental depression pre and post-intervention for infant behavioral sleep problems. BMC Pregnancy Childbirth. 2017;17(1):104.

48. Saxbe DE, Schetter CD, Guardino CM, Ramey SL, Shalowitz MU, Thorp J, et al. Sleep Quality Predicts Persistence of Parental Postpartum Depressive Symptoms and Transmission of Depressive Symptoms from Mothers to Fathers. Ann Behav Med. 2016;50(6):862-75.

49. Vik T, Grote V, Escribano J, Socha J, Verduci E, Fritsch M, et al. Infantile colic, prolonged crying and maternal postnatal depression. Acta Paediatr. 2009;98(8):1344-8.

50. Radesky JS, Zuckerman B, Silverstein M, Rivara FP, Barr M, Taylor JA, et al. Inconsolable infant crying and maternal postpartum depressive symptoms. Pediatrics. 2013;131(6):e1857-64.

51. Alexander CP, Zhu J, Paul IM, Kjerulff KH. Fathers make a difference: positive relationships with mother and baby in relation to infant colic. Child Care Health Dev. 2017;43(5):687-96.

Tables

Table 1 Characteristics of the study population when the child is six months

Child (N= 86,724)	n (%)	mean(SD)	mean (SD)	Median (range)
Male n (%)	40,205 (51.0)			
Missing (n= 7,947)				
Gestation age				
≥ 37 weeks	70,815 (95.0)			
32-36 weeks	3,231 (4.3)			
28-31 weeks	309 (0.4)			
< 28 weeks	192 (0.3)			
Missing (n= 12,177)				
Birth weight in grams	78,816			3,600 (500, 5,960)
Missing (n = 7,908)				
Mother (N= 86,724)				
Maternal age	82,377		29.8 (4.5)	
Missing (n = 4,387)				
Maternal education ^a				
low	26,607 (34.0)			
medium	32,941 (42.0)			
high	18,836 (24.0)			
Missing (n = 8,340)				
Work status ^b				
Education	4,930 (6.0)			
Employed	68,153 (83.1)			
Other	8,938 (10.9)			
Missing (n = 4,703)				
Marital status ^c				
Married/co-habiting	80,333 (97.9)			
Missing (n = 4,640)				

^a Education at 16-20 weeks of pregnancy: low education = completed primary school and or lower and upper secondary school. Medium education = completed college / university 1-3 years. High education = completed college / university 4 years.

^b Work Status at 16-20 weeks of pregnancy: education = student, at home, trainee, military service. Employed = employed in the public or private sector. Other = unemployed.

^c Marital status at 16-20 weeks of pregnancy.

Table 2 Description of study variables

Variable (scoring)	Scale	Range	Mean	SD
Overall quality of life (high scores = higher QoL)	SWLS	5 – 35	28.8	4.9
Missing n = 1,684				
Depression (high scores = more depression)	EPDS	0 – 18	3.1	3.2
Missing n = 1,710				
Joy (high scores = more joy)	DES-J	2 – 15	11.9	2.1
Missing n = 9,109				
Anger (high scores = more anger)	DES-A	2 – 15	6.2	2.2
Missing n = 9,131				
Fussiness (higher scores = more fussy)	ICQ	7 – 49	16.4	5.6
Missing = 4,321				
Relationship satisfaction (higher scores = higher satisfaction)	RS	10 – 60	52.6	7.2
Missing = 6,308				
			%	
Feeling safe during birth				
applies well			84.5	
applies partly			13.3	
does not apply			2.2	
Missing n = 2,333				
Normal sleep of child (13 hours or more a day)			92.6	
Missing = 3,695				

Table 3 Stepwise multivariate linear regression analysis (standardised beta coefficients) with mothers' quality of life as the dependent variable

Independent Variables	B	95% CI	P-value
<i>Block 1</i>			
Age, years	-0.06	-0.07 – -0.06	<0.001
Level of education (1 – 3)	0.13	0.78 – 0.89	<0.001
Marital status (not married/cohabitating as ref.)	-0.02	-3.3 – -1.58	<0.001
<i>Block 2</i>			
Age, years	-0.02	-0.03 – -0.02	<0.001
Level of education (1 – 3)	0.09	0.49 – 0.57	<0.001
Marital status (not married/cohabitating as ref.)	-0.02	-2.35 – -1.00	<0.001
Depression (EPDS 6-item) (high scores = more depression)	-0.15	-0.24 – -0.22	<0.001
Joy (DES-J) (high scores = more joy)	0.48	1.10 – 1.13	<0.001
Anger (DES-A) (high scores = more anger)	-0.13	-0.31 – -0.28	<0.001
Feeling safe during birth (lower scores = more confidence)	-0.03	-0.42 – -0.29	<0.001
<i>Block 3</i>			
Age, years	-0.01	-0.01 – 0.001	0.12
Level of education (1 – 3)	0.08	0.44 – 0.51	<0.001
Marital status (not married/cohabitating as ref.)	-0.001	-0.73 – 0.54	0.76
Depression (EPDS 6-item) (high scores = more depr.)	-0.12	-0.19 – -0.19	<0.001
Joy (DES-J) (high scores = more joy)	0.35	0.79 – 0.83	<0.001
Anger (DES-A) (high scores = more anger)	-0.06	-0.15 – -0.12	<0.001
Feeling safe during birth (lower scores = less confidence)	-0.02	-0.33 – -0.21	<0.001
Colic (no colic as ref.)	-0.004	-0.15 – 0.03	0.18
Child's temperament (ICQ) (lower scores = less fussy)	-0.006	-0.01 – 0.00	0.06
Child sleeping more than 13 hours a day ($\leq$ 13 hours as ref.)	0.02	0.28 – 0.60	<0.001
Relationship satisfaction (RS) (higher scores = more satisfaction)	0.32	0.21 – 0.22	<0.001

$R^2 = 0.49$