

General practitioners' perspectives on barriers to depression care: development and validation of a questionnaire

Arun Senchyna (✉ arun.senchyna@unige.ch)

Universite de Geneve Faculte de Medecine <https://orcid.org/0000-0003-2504-7916>

Milena Abbiati

Universite de Geneve Faculte de Medecine

Juliette Chambe

Universite de Strasbourg

Dagmar Haller

Universite de Geneve Faculte de Medecine

Hubert Maisonneuve

Universite de Geneve Faculte de Medecine

Research article

Keywords: Depression, General practitioners, Primary care, Questionnaire

Posted Date: April 13th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-22270/v1>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published on August 1st, 2020. See the published version at <https://doi.org/10.1186/s12875-020-01224-8>.

Abstract

Background General practitioners (GPs) regularly feel challenged by the care of depressed patients and may encounter several barriers in providing best management. GPs' perspectives on barriers to depression care are a subject of growing interest but there is a lack of validated assessment tools. The aim of this study was to develop and validate a questionnaire assessing barriers to depression care (BDC-Q) encountered by GPs in France and the French-speaking part of Switzerland.

Methods The BDC-Q was constructed in five steps: Item development, content validation, pretesting, testing phase and test-retest reliability. The questionnaire items were generated through a literature search. An expert panel of GPs (n=16) and psychiatrists (n=3) validated the content and 20 GPs pretested the questionnaire to provide response process validity evidence. We then tested the questionnaire among 116 GPs and used principal component analysis and internal consistency testing (Cronbach's alpha) to structure it into consistent dimensions. Test-retest reliability using Pearson correlation coefficient was assessed with 30 GPs who completed the questionnaire twice after an interval of at least 2 weeks.

Results The 25 items BDC-Q was structured in five dimensions: (i) provision of care by the general practitioner, (ii) patient and perspectives about depression, (iii) guidance for care, (iv) collaboration with mental health specialists and (v) access to mental health care.

Conclusions The BDC-Q displays evidence of validity and reliability to meaningfully assess GPs' perspectives on barriers to depression care. It can be used both at a practice level within a quality improvement strategy, and at a broader level, to inform health planners and tailor appropriate strategies to improve depression care in the community.

Introduction

Depression affects more than 300 million people worldwide, accounting now for the first leading cause of disability (WHO). In European countries, the 12 month prevalence of depression is estimated at 6,9%.^{1,2} Along with anxiety disorders, depression is the main mental disorder encountered in primary care (PC) and most depressed patients will be handled in PC settings only.^{3,4} General practitioners (GPs) regularly encounter difficulties to diagnose depression and provide adequate care.^{5,6} Addressing the pitfalls and perspectives for improvements has moved beyond a disease-centred approach to an integrative approach involving the perceptions of patients and care givers.⁷ A growing number of studies analyse GPs' views of their practice and experience to identify barriers to and facilitators of depression care. Authors have described several dimensions of barriers to depression care that involve health care organization, coordination of care, societal influence, patient factors and physician attitudes.⁸⁻¹⁰ To date, the current research on perceived barriers to depression care have mostly focused on explorative studies using qualitative designs. These methodological choices have enabled to gain insight into the complexity of depression care in PC. However, such study designs prevent by nature a broader generalization of the results. Therefore, quantitative approaches are needed to allow wider assessment of the burden of

barriers and explore their relationships and respective weight among communities of GPs. Quantitative assessments of barriers to depression care may be useful to highlight the needs of GPs and in turn, tailor and monitor appropriate improvement programs of care. Therefore, the aim of the current study was to develop and validate a questionnaire assessing barriers to depression care (BDC-Q) from GPs' perspectives in order to provide a structured instrument to use in clinical practices.

Methods

Design and participants

This is a development and validation study conducted within primary care practices (general internists) in France and the French-speaking part of Switzerland between 2014 and 2018.

Figure 1 summarizes the development and validation process of the barriers to depression care questionnaire (BDC-Q) with participants' socio-demographic characteristics.

Step 1 - Item development

We devised 42 items from a literature search conducted in MEDLINE using terms related to depression, family practice, and barriers/facilitators and through bibliographies of retrieved studies. We included qualitative and quantitative studies referring totally or partially to barriers to depression care. We excluded studies that did not involve GPs' views or were focused exclusively on specific aspects of depression care or populations (e.g. depression in youth, end of life care or postpartum depression). We retained 11 studies from the USA^{8,11,12}, UK¹³, France¹⁴, the Netherlands¹⁵, Australia^{16,17}, Hong Kong¹⁸ and 2 international meta-syntheses.^{9,19} We also added the results of a focus group conducted within a local depression improvement program in the French-speaking part of Switzerland (unpublished results). Any elements describing an implicit or explicit barrier were extracted, adapted and translated into French to form an item.²⁰

Step 2 - Content validation

Content validation was undertaken with a French-speaking panel of 19 physicians with expertise in the management of depression in PC. Physicians were GPs, psychiatrists and academics from Lyon (France), Strasbourg (France) and Geneva (Switzerland). Using the content validation index described by Polit, experts rated the relevance and the clarity of each item on a 4-point scale (not relevant, quite relevant, relevant, highly relevant; respectively : not clear, quite clear, clear, very clear).²¹ An item was considered relevant if more than 75% of experts selected it as "quite relevant" or "highly relevant". An item was considered clear if it obtained 80% or more of "quite clear" and "clear" ratings.²¹ Among the 42 items submitted to the binational expert panellists, 16 irrelevant items were dropped. 26 items were retained for pretesting with relevant and unclear items (n = 5) modified according to expert comments or by returning to the literature findings.

Step 3 - Pretesting

Pretesting of the questionnaire was undertaken to provide response process validity evidence defined as the fit between the construct and the detailed nature of the response engaged in by test takers.^{22,23} We used individual semi-directive cognitive interviewing techniques²⁴ among 20 GPs who were recruited using snowball sampling for maximal variation.²⁵ None of the GPs in this step were involved in the content validation step. Two trained investigators (LL and AG) performed the interviews in GPs' practices. Participants read and laudably answered each item before answering standardised question probes.²⁶ The probes explored item understanding (e.g. <<In the item "mental health care professionals are available to take on new patients", what does the word 'available' mean to you?>>), redundancy between items, as well as the response selection process²⁷ (e.g. "How did you decide your answer to this question was strongly agree?"). All the cognitive interviews were recorded, transcribed verbatim and independently coded by LL and AG using a 12-point coding sheet.²⁶ For each interview, items were separately coded as "adequate" or alternatively with a combination of 11 problematic codes (e.g. "Respondent unsure how to answer since experience varies depending on circumstances"; "Respondent asks for clarification of the item"). An item was judged satisfactory if it was coded "adequate" for more than 85% of respondents.²⁶

Among the 26 tested items, 13 were adequate and retained without modification. The 13 problematic items were revised through discussion between the research team members, in accordance with propositions highlighted during the cognitive interviews. In this process, 2 derived items were added leading to a 28 items questionnaire.

For all items, a traditional 5-point Likert scale was suitable, ranging from strongly disagree to strongly agree.²⁰ The neutral position was labelled "no answer" to capture a neutral position or a non-existent opinion. In order to obtain a balanced questionnaire, half of the items were positively worded (e.g. "It is easy to distinguish between simple sadness and a depressive disorder") and the other half negatively (e.g. "Obtaining feedback on patients from mental health care professionals is difficult").

Steps 4 and 5 - Testing phase and Test-Retest reliability

The testing phase of the BDC-Q was carried out among GPs from the Alsace and Rhone-Alpes regions of France. Participants were recruited through regional professional organisation mailing lists. Surveys were web-based and gathered socio-demographic characteristics. The questionnaire items were displayed randomly to avoid response contamination bias.²⁸ We prevented missing data by forcing response to all items. 131 GPs initiated the survey (response rate of 13,3%) with a completion rate of 88,5% (15 incomplete surveys were excluded). Thus, 116 surveys were used for the analyses in the testing phase. We used principal component analysis followed by internal consistency testing to organise the items into descriptive dimensions. Test retest reliability was conducted with a sub-sample of 40 GPs. They were asked to respond to the survey again after a 14 days interval.

Statistical analysis

We calculated descriptive statistics for each item including the mean, standard deviation and range to inspect floor and ceiling effects. Items endorsed by more than 95% of the participants were considered for removal.²⁹ We performed a principal component analysis (PCA) with Promax rotation to aggregate BDC-Q items in factors after using the Kaiser-Meyer-Olkin (KMO) index of sampling adequacy to confirm suitability of the data. We used combined criteria (i.e. eigenvalue > 1 and interpretability) to retain the most relevant factors and order them into consistent dimensions.³⁰ A minimum factor loading of 0.40 was used as the criterion for each retained item.³¹ An item obtaining primary and secondary loading superior to 0.40 was assigned to the dimension with the most theoretical sense. We used classical Cronbach's Alpha coefficient to assess dimensions internal consistency. In doing this our intention was not to create subscales, but just to check that the items were sufficiently related, justifying their grouping under the different dimensions. We chose the following critical values for the dimensions internal consistency : $\alpha > 0.75$ = excellent, α between 0.60 and 0.75 = good, α between 0.40 and < 0.60 = moderate, and $\alpha < 0.40$ = poor.³² We used Pearson correlation coefficient to determine dimensions test-retest reliability between time 1 (T1) and time 2 (T2) with the following critical values for Pearson's r : $r > 0.5$ = high, $r > 0.3 < 0.5$ = moderate, and $r < 0.3$ = low.³³ All statistical analyses were conducted using IBM SPSS Statistics version 24.

Results

Descriptive

Table 1 presents the descriptive statistics for all items. Response means for the 28 items varied from 2.00 to 4.19 with a complete range of responses in 19 out of 25 items (76%) and no floor and ceiling effects observed. Maximum endorsement was 63,8% "disagree" for item 2 ("It is easy to distinguish between simple sadness and a depressive syndrome").

Principal Component analysis

The PCA, shown in Table 2, yielded five factors (KMO 0.57, $p < 0.001$, 41.4% of variance explained) with the first three factors accounting for 31% of the variance. We interpreted and labelled each factor according to the co-varying items. We named Factor 1 "Provision of care by the general practitioner", combining items 5; 14; 15; 16; 19; 28. We named Factor 2 "Patient and perspectives about depression", combining items 2; 6; 9; 10; 11; 12. We named Factor 3 "Guidance for care", combining items 3; 4; 18; 20. We named Factor 4 "Collaboration with mental health specialists", combining items 17; 23; 24; 25; 26; 27. We named Factor 5 "Access to mental health care", combining items 13; 21; 22. We excluded Items 1; 7; 8 since their loadings were < 0.40. Items 15; 17; 24; 28 cross-loaded significantly (> 0.40) on a second Factor but these cross-loadings were always lower than the cross-loadings on their primary Factor except for item 15. We found a weak correlation between Factor 1 and Factor 3 (0.283).

Dimensions internal consistency and test-retest reliability

Table 3 presents the BDC-Q internal consistency (Cronbach's alpha) and the test-retest reliability (Pearson's r) per dimension. Cronbach's alpha ranged from 0.48 to 0.69. Scores were good for dimensions 1 to 4 and moderate for dimension 5. Among the 40 GPs who were asked to respond in both instances, 30 (75%) returned the questionnaire within a median T1-T2 interval of 18 days (range 14–55). Pearson correlation coefficient was high for all dimensions, ranging from 0.54 to 0.65 with all $p < 0.05$.

Discussion

Summary of main findings

The five-step validation study led to the BDC-Q with 25 items, covering five dimensions:

(i) provision of care by the general practitioner, (ii) patient and perspectives about depression, (iii) guidance for care, (iv) collaboration with mental health specialists and (v) access to mental health care. The BDC-Q is available in French (**appendix – BDC-Q French**). We also present an English (unvalidated) translation of the final questionnaire for this publication (**appendix – BDC-Q English**).

Comparison with previous literature

The validation and usage of questionnaires to evaluate GPs' views of depression care has been a subject of growing interest. Haddad et al. recently revised the Depression Attitude Questionnaire (R-DAQ) which explores professional confidence, therapeutic optimism, and views about generalist or specialist perspectives relevant to depression and its care.³⁴ In Norway, Bjertnaes and al. validated a questionnaire on GPs' experience of quality of care in community mental health clinics.³⁵ To the best of our knowledge, this is the first study describing the validation of a questionnaire specifically exploring GPs' views on barriers to depression care. The BDC-Q is structured in five original dimensions, yet embracing previous models.^{8,9} Our findings are also in line with previous studies in France showing that barriers to mental health care involved unsatisfactory co-operation between GPs and mental health services, feelings of stigma and reluctance from patients to consult mental health specialists, as well as high costs of mental health care in the private sector.^{14,36,37}

Strength and Limitations

Preliminary content validation and pretesting steps permitted improvements in terms of relevance and clarity of the items. In turn, the high completion rate during the testing phase suggests that the BDC-Q is acceptable and suitable to administer in everyday practice. Results on the items-level show that there were no remaining floor and ceiling effects, raising the potential to measure cross-sectional differences and responsiveness.

Our study suffers several limitations. First, we devised items through a non-systematic review of the literature. This implies the risk of relevant content loss. However, as we asked for comments, no relevant

item was suggested by participants during the validation steps.

Second, the testing phase may have suffered from selection bias and limited sample size despite a participant/item ratio of 4 which is common practice in this field.³⁸ Factorial analysis rather than PCA is preferred to structure a questionnaire into dimensions, but requires a large sample size.³⁹ Consequently, at this stage we do not suggest subscale scoring of the dimensions presented. The dimensions of the BDC-Q should be considered as indicative and calling for further research, because they may differ in other health care settings or populations of GPs. Finally, the BDC-Q does not account for depression related to specific health or social conditions. Although different barriers to depression care may be encountered depending on the socio-demographic populations of interest⁴⁰, the BDC-Q intends to capture relevant information that is common across different patient groups, and is thus designed for a general use.

Applications and implications for future research

The validation process of the questionnaire encourages its use to meaningfully assess barriers to depression care perceived by GPs. The BDC-Q can be useful to gathered relevant data at a practice level, within a quality improvement effort. Addressing key barriers is crucial to design adequate interventions which have been shown to be effective in improving mental health care both at the clinical and health policy levels.⁴¹ At the physician level, taking the BDC-Q can raise awareness of common pitfalls in current practices, enabling a better recognition of the resources and limitations encountered while dealing with depressed patients. Further longitudinal research should assess the validity of the BDC-Q to measure change, for example to monitor the impact of depression care improvement programs. The BDC-Q may also benefit from translation and cross-cultural validation for use in other settings.

Conclusion

The BDC-Q displays validity and reliability evidence to meaningfully assess GPs' perspectives on barriers to depression care in French-speaking settings. By facilitating broad and reliable assessment of such barriers, the BDC-Q can be a useful tool to target key improvement indicators, inform health planners, and tailor appropriate strategies to improve depression care in the community. Further research steps include the translation and validation of the BDC-Q in other languages to allow use in a broader range of primary care settings.

Declarations

Ethics approval and consent to participate: In accordance with the applicable law in France and Switzerland, since no patients were involved in this study, no ethical approval was required.

Consent for publication: Not applicable

Availability of data and materials: The data that support the findings of this study are available from the authors upon reasonable request.

Competing interests: The authors declare that they have no competing interests. Dagmar M Haller is a member of the Editorial Board (Associate Editor) of BMC.

Funding: None

Authors' contributions: AS designed the study, collected and analyzed the data, interpreted the results and wrote the first draft of this paper. MA analyzed the data, interpreted the results and made critical revisions to the paper. JC made substantial contributions to the collection of the data. HM and DH designed the study, interpreted the results and made critical revisions to the paper. All authors read and approved the final manuscript.

Acknowledgements: The authors thank all the participants for their contribution towards this study. We warmly acknowledge the contribution as data collectors of Alban Glangetas (AG), Carole Champet, Leslie Llorente (LL), Noémie Leuret, as well as Amir Moussa, administrative assistant. We are grateful to all the members of the Primary Care Unit of the University of Geneva for their insights and to Jennifer Hasselgard-Rowe and Joshua Arancino-Gamper for their help in language editing.

References

1. Vos T, Allen C, Arora M, et al. Global, regional, and national incidence, prevalence, and years lived with disability for 310 diseases and injuries, 1990–2015: a systematic analysis for the Global Burden of Disease Study 2015. *The Lancet*. 2016;388(10053):1545–602. doi:.
2. Wittchen HU, Jacobi F, Rehm J, et al. The size and burden of mental disorders and other disorders of the brain in Europe 2010. *Eur Neuropsychopharmacol*. 2011;21(9):655–79. doi:.
3. Rucci P. Subthreshold psychiatric disorders in primary care: prevalence and associated characteristics. *J Affect Disord*. 2003;76(1–3):171–81. doi:.
4. Goldberg D. Psychiatry and primary care. *World Psychiatry*. 2003;5.
5. Mitchell AJ, Vaze A, Rao S. Clinical diagnosis of depression in primary care: a meta-analysis. *The Lancet*. 2009;374(9690):609–19. doi:.
6. Cepoiu M, McCusker J, Cole MG, Sewitch M, Belzile E, Ciampi A. Recognition of Depression by Non-psychiatric Physicians—A Systematic Literature Review and Meta-analysis. *J Gen Intern Med*. 2008;23(1):25–36. doi:.
7. Solberg LI, Rossom RC. Managing mental health disorders requires attending to both primary care and its specialty mental health collaborators. *Fam Pract*. 2019;36(1):1–2. doi:.
8. Whitebird RR, Solberg LI, Margolis KL, Asche SE, Trangle MA, Wineman AP. Barriers to Improving Primary Care of Depression: Perspectives of Medical Group Leaders. *Qual Health Res*. 2013;23(6):805–14. doi:.

9. Schumann I, Schneider A, Kantert C, Lowe B, Linde K. Physicians' attitudes, diagnostic process and barriers regarding depression diagnosis in primary care: a systematic review of qualitative studies. *Fam Pract.* 2012;29(3):255–63. doi:.
10. Mechanic D. Barriers to Help-Seeking, Detection, and Adequate Treatment for Anxiety and Mood Disorders: Implications for Health Care Policy. *J Clin Psychiatry*.:7.
11. Fickel JJ, Parker LE, Yano EM, Kirchner JE. Primary care–mental health collaboration: An example of assessing usual practice and potential barriers. *J Interprof Care.* 2007;21(2):207–16.
12. Nutting PA, Rost K, Dickinson M, et al. Barriers to initiating depression treatment in primary care practice. *J Gen Intern Med.* 2002;17(2):103–11. doi:.
13. Telford R. Obstacles to effective treatment of depression: a general practice perspective. *Fam Pract.* 2002;19(1):45–52. doi:.
14. Mercier A, Kerhuel N, Stalnikiewitz B, et al. Enquête sur la prise en charge des patients dépressifs en soins primaires: les médecins généralistes ont des difficultés et des solutions. *L'Encéphale.* 2010;36:D73–82. doi:.
15. van Rijswijk E, van Hout H, van de Lisdonk E, Zitman F, van Weel C. Barriers in recognising, diagnosing and managing depressive and anxiety disorders as experienced by Family Physicians; a focus group study. *BMC Fam Pract.* 2009;10(1). doi:.
16. Jones KM, Piterman L, Spike N. Difficult-to-Treat-Depression—Perceptions of GPs and GP Trainees. *Open J Psychiatry.* 2014;04(03):228–37. doi:.
17. Richards JC, Ryan P, McCabe MP, Groom G, Hickie IB. Barriers to the effective management of depression in general practice. *Aust N Z J Psychiatry.* 2004;38(10):795–803.
18. Wong SYS, Lee K, Chan K, Lee A. What are the barriers faced by general practitioners in treating depression and anxiety in Hong Kong?: Barriers to Treating Depression and Anxiety in Hong Kong. *Int J Clin Pract.* 2006;60(4):437–41. doi:.
19. Barley EA, Murray J, Walters P, Tylee A. Managing depression in primary care: A meta-synthesis of qualitative and quantitative research from the UK to identify barriers and facilitators. *BMC Fam Pract.* 2011;12(1):47.
20. Passmore C, Dobbie AE, Parchman M, Tysinger J. Guidelines for constructing a survey. *Fam Med-Kans CITY-*. 2002;34(4):281–6.
21. Polit DF, Beck CT, Owen SV. Is the CVI an acceptable indicator of content validity? Appraisal and recommendations. *Res Nurs Health.* 2007;30(4):459–67. doi:.
22. American Educational Research Association, American Psychological Association, & National Council on Measurement in Education [AERA, APA, & NCME]. Standards for educational and psychological testing. Washington, DC: American Educational Research Association; 2014.
23. Artino AR, La Rochelle JS, Dezee KJ, Gehlbach H. Developing questionnaires for educational research: AMEE Guide No. 87. *Med Teach.* 2014;36(6):463–74. doi:.

24. Castillo-Díaz M, Padilla J-L. How Cognitive Interviewing can Provide Validity Evidence of the Response Processes to Scale Items. *Soc Indic Res.* 2013;114(3):963–75. doi:.
25. Noy C. Sampling Knowledge: The Hermeneutics of Snowball Sampling in Qualitative Research. *Int J Soc Res Methodol.* 2008;11(4):327–44. doi:.
26. Napoles-Springer AM, Santoyo-Olsson J, O'Brien H, Stewart AL. Using Cognitive Interviews to Develop Surveys in Diverse Populations. *Med Care.* 2006;44(Suppl 3):21–30. doi:.
27. Beatty PC, Willis GB. Research Synthesis: The Practice of Cognitive Interviewing. *Public Opin Q.* 2007;71(2):287–311. doi:.
28. 10.1093/acprof:oso/9780199231881.001.0001
Streiner DL, Norman GR. *Health Measurement Scales.* Oxford University Press; 2008. doi:.
29. Haller DM, Meynard A, Pejic D, et al. YFHS-WHO + Questionnaire: Validation of a Measure of Youth-Friendly Primary Care Services. *J Adolesc Health.* 2012;51(5):422–30. doi:.
30. Schönrock-Adema J, Heijne-Penninga M, van Hell EA, Cohen-Schotanus J. Necessary steps in factor analysis: Enhancing validation studies of educational instruments. The PHEEM applied to clerks as an example. *Med Teach.* 2009;31(6):e226–32. doi:.
31. Hatcher L. A step-by-step approach to using the SAS system for factor analysis and structural equation modeling. Cary: SAS Institute Inc.; 1994.
32. Kline P. The handbook of psychological testing. London: Routledge; 2000.
33. Cohen J. Statistical Power Analysis for the Behavioral Sciences. 2nd ed. Hillsdale: L. Erlbaum Associates; 1988.
34. Haddad M, Menchetti M, McKeown E, Tylee A, Mann A. The development and psychometric properties of a measure of clinicians' attitudes to depression: the revised Depression Attitude Questionnaire (R-DAQ). *BMC Psychiatry.* 2015;15(1). doi:.
35. Bjertnaes OA, Garratt A, Ruud T, Hunskaar S. The General Practitioner Experiences Questionnaire (GPEQ): validity and reliability following the inclusion of new accessibility items. *Fam Pract.* 2010;27(5):513–9. doi:.
36. Younes N, Gasquet I, Gaudebout P, et al. General Practitioners' opinions on their practice in mental health and their collaboration with mental health professionals. *BMC Fam Pract.* 2005;6(1). doi:.
37. Jegou M, Debaty E, Ouirini L, Carrier H, Beetlestone E. Caring for patients with mental disorders in primary care: a qualitative study on French GPs' views, attitudes and needs. *Fam Pract.* 2019;36(1):72–6. doi:.
38. Anthoine E, Moret L, Regnault A, Sébille V, Hardouin J-B. Sample size used to validate a scale: a review of publications on newly-developed patient reported outcomes measures. *Health Qual Life Outcomes.* 2014;12(1). doi:.
39. Costello AB, Osborne JW. Best Practices in Exploratory Factor Analysis: Four Recommendations for Getting the Most From Your Analysis. *Explor Factor Anal.* 10(7):9.

40. Coventry PA, Hays R, Dickens C, et al. Talking about depression: a qualitative study of barriers to managing depression in people with long term conditions in primary care. *BMC Fam Pract.* 2011;12(1). doi:.
41. Sinnema H, Majo MC, Volker D, et al. Effectiveness of a tailored implementation programme to improve recognition, diagnosis and treatment of anxiety and depression in general practice: a cluster randomised controlled trial. *Implement Sci.* 2015;10(1). doi:.

Tables

Table 1. Descriptive features of the original items submitted to 116 general practitioners during the testing phase (non-validated English translation of the original items in French).

N°	Item	Mean score (SD; Range)
1	The symptoms of depression are specific	3.2 (1.0; 1-5)
2	It is easy to distinguish between simple sadness and a depressive disorder	2.6 (0.9; 1-5)
3	Screening tools for depression, such as the HAD (Hospital Anxiety and Depression scale) for example, lack practical utility	3.3 (0.8; 1-5)
4	Assessment tools for depression, such as the Hamilton scale or the Beck Depression Inventory lack practical utility	2.6 (1.0; 1-5)
5	Best practice recommendations related to depression lack practical applicability	3.4 (0.9; 1-5)
6	The general public is well informed about depression	2.2 (0.8; 1-4)
7	The general public is well informed about the management of depression	3.9 (0.7; 1-5)
8	Patients suffering from depression endure social stigmatization	3.7 (0.8; 1-5)
9	Patients suffering from depression underestimate the severity of their depression	3.6 (0.9; 2-5)
10	Patients suffering from depression easily accept a diagnosis of depression	2.9 (1.1; 1-5)
11	Patients suffering from depression easily accept being referred to a mental health care professional	2.4 (0.9; 1-5)
12	The commitment of patients suffering from depression to the therapeutic project is limited	3.1 (1.0; 1-5)
13	Patients suffering from depression are adequately reimbursed for their mental health care costs	3.4 (1.1; 1-5)
14	Taking care of a patient suffering from depression often takes up more time than I can give him/her	3.9 (1.1; 1-5)
15	I am adequately paid for taking care of patients suffering from depression	2.2 (1.1; 1-5)
16	Working with patients suffering from depression is heavy	3.4 (1.0; 1-5)
17	Mental health care professionals are available to take on new patients	2.0 (0.9; 1-5)
18	I know the specializations of mental health professionals regarding certain pathologies (for example, addiction, bipolar disorders) well.	2.7 (1.1; 1-5)
19	The capacity of specialized mental health care structures is insufficient	4.0 (0.9; 2-5)
20	I know the services offered by mental health care structures well	2.8 (1.0; 1-5)
21	I mistrust mental health care structures	2.7 (0.9; 1-5)
22	I have had bad experiences using structures specialized in mental health	2.9 (1.0; 1-5)
23	Medical information sharing between patients and mental health care professionals is easy	2.1 (1.0; 1-5)
24	Getting advice over the phone from mental health care professionals is easy	2.2 (1.0; 1-5)
25	Obtaining feedback on patients from mental health care professionals is difficult	4.2 (0.8; 2-5)
26	Expectations concerning the communication of information are the same for general practitioners as for mental health care professionals	2.8 (1.0; 1-5)
27	Setting up meetings with mental health care professionals to discuss cases is difficult	4.0 (0.8; 2-5)
28	The clinical situation of a patient suffering from depression is difficult to summarize in writing	3.3 (1.1; 2-5)

Table 2. Principal Components Analysis with Promax Rotation* of the barriers to depression care questionnaire (BDC-Q).

N° Item (Table 1)	Factor loadings				
	F1	F2	F3	F4	F5
14	.811	.121	.244	.173	-.098
16	.680	.103	-.060	.133	-.173
19	.488	.084	.157	.097	.192
28	.437	.415	.257	.298	-.225
5	.402	.072	.212	.130	.094
10 R	.214	.777	.227	.020	.060
12	.159	.601	-.069	.025	-.307
9	.171	.562	-.048	.106	-.125
11 R	-.055	.557	.239	.034	.161
6 R	-.073	.460	.013	-.031	-.010
2 R	.258	.408	.242	.234	-.34
7 D	.052	.371	.248	.153	.078
8 D	.219	.230	.063	.029	-.175
20 R	.235	.022	.755	.205	.017
18 R	.176	-.051	.743	-.065	-.108
4	.144	.282	.645	.108	.082
3	.117	.273	.544	.215	-.050
15 R	.406	.083	.420	-.049	.156
1 RD	.217	.287	.291	.193	-.214
23 R	.007	.139	.038	.776	-.031
25	.386	.044	.088	.635	.151
17 R	.430	-.158	.316	.486	.106
27	.376	.032	.276	.472	-.012
24 R	.151	-.137	.097	.459	.406
26 R	-.098	.036	.099	.454	.242
21	-.041	-.019	.018	.207	.677
22	.031	.024	-.03	.208	.646
13 R	.384	.008	.219	-.196	.554
	% Variance explained				
	F1	F2	F3	F4	F5
	13.906	8.885	6.944	6.255	5.474
	Factor correlations				
	F1	F2	F3	F4	F5
F1	1				
F2	.146	1			
F3	.283	.153	1		
F4	.214	.112	.150	1	
F5	.039	-.138	.087	-.012	1

*Kaiser-Meyer-Olkin index = .57, $p < .001$, 41.4% variance explained

R = reversed; D = dropped

Table 3. Internal consistency (Cronbach's alpha) and Test-retest reliability (Pearson's r) per dimension of the barriers to depression care questionnaire (BDC-Q).

Factor	Dimension	N° Item (Table 1)	Cronbach's alpha	Pearson's r*
			N=116	N=30
F1	Provision of care by the general practitioner	5;14;15;16;19;28	0.610	0.540
F2	Patient and representations about depression	2;6;9;10;11;12	0.639	0.608
F3	Guidance for care	3;4;18;20	0.651	0.631
F4	Collaboration with mental health specialists	17;23;24;25;26;27	0.655	0.653
F5	Access to mental health care	13;21;22	0.477	0.636

*p <0.05

Figures




Figure 1

Development and validation steps of the barriers to depression care questionnaire (BDC-Q) with participants' socio-demographic characteristics.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [supplementarymaterialBDCQfrench.docx](#)
- [supplementarymaterialBDCQenglish.docx](#)