

# Effects of Peak Time of Myocardial Injury Biomarkers On Mid-Term Outcomes of Patients Undergoing OPCABG

**Bo Hu**

Shanghai East Hospital

**Fei Gao**

Huaiyin Hospital of Huai'an city

**Mengwei Lv**

Shanghai east hospital of clinical medical college

**Ban Liu**

Shanghai Tenth People's Hospital

**Yu Shi**

Shanghai East Hospital

**Xi Chen**

Shanghai East Hospital

**Yipeng Feng**

The First Clinical Medical College of Nanjing Medical University

**Xiaoqi Meng**

The Second Clinical Medical College of Nanjing Medical University

**Zhi Li**

Jiangsu Province Hospital

**Yangyang Zhang** (✉ [zhangyangyang\\_wy@vip.sina.com](mailto:zhangyangyang_wy@vip.sina.com))

Shanghai East Hospital

---

## Research Article

**Keywords:** myocardial injury biomarker, coronary artery bypass grafting, off pump, mid-term outcome

**Posted Date:** February 24th, 2021

**DOI:** <https://doi.org/10.21203/rs.3.rs-223381/v1>

**License:**  This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

---

**Version of Record:** A version of this preprint was published at BMC Cardiovascular Disorders on April 24th, 2021. See the published version at <https://doi.org/10.1186/s12872-021-02006-5>.

# Abstract

**Background:** With the development of cardiac surgery technology, myocardial injury is gradually reduced, but cannot be completely avoided. Myocardial injury biomarkers (MIBs) can quickly and specifically reflect the degree of myocardial injury. Due to various reasons, there is no consensus on the specific values of MIBs in evaluating postoperative prognosis. This retrospective study was aimed to investigate the impact of MIBs on the mid-term prognosis of patients undergoing off-pump coronary artery bypass grafting (OPCABG).

**Methods:** Totally 564 patients undergoing OPCABG with normal courses were included. Cardiac troponin T (cTnT) and creatine kinase myocardial band (CK-MB) were assessed within 48 hours before operation and at 6, 12, 24, 48, 72, 96 and 120 hours after operation. Patients were grouped by peak values and peak time courses of MIBs. The profile of MIBs and clinical variables as well as their correlations with mid-term prognosis were analyzed by univariable and multivariable cox regression models.

**Result:** Continuous assessment showed that MIBs increased first (12 hours after surgery) and then decreased. The peak cTnT and peak CK-MB occurred within 24 hours after operation in 76.8% and 67.7% of the patients respectively. No significant correlation was found between CK-MB and mid-term mortality. Delayed cTnT peak (peak cTnT elevated after 24 hours after operation) was correlated with lower Ccr ( $69.36 \pm 21.67$  vs.  $82.18 \pm 25.17$  ml/min/1.73 m<sup>2</sup>), body mass index ( $24.35 \pm 2.58$  vs.  $25.27 \pm 3.26$  kg/m<sup>2</sup>), less arterial grafts ( $1.24 \pm 0.77$  vs.  $1.45 \pm 0.86$ ), higher Euroscore II ( $2.22 \pm 1.12$  vs.  $1.72 \pm 0.91$ ) and mid-term mortality (26.5 vs. 7.9 %). Age (HR: 1.067, CI: 1.005-1.132), left ventricular ejection fraction (HR: 0.950, CI: 0.910-0.992), New York Heart Association score (HR: 1.833, CI: 1.155-2.908), total venous grafting (HR: 2.833, CI: 1.054-7.611) and cTnT peak occurrence within 24 hours (HR: 0.356, CI: 0.731-2.513) were independent predictors of mid-term mortality.

**Conclusion:** cTnT is a better indicator than CK-MB. The peak and peak occurrence of cTnT are related to mid-term mortality in patients undergoing OPCABG, and the peak phases have stronger predictive ability.

## Background

Myocardial injury after cardiac surgery, especially coronary artery bypass grafting (CABG), is inevitable. With the improvement in cardiac surgery techniques, off-pump CABG (OPCABG) has become a favorite choice for surgeons in China.<sup>[1]</sup> This kind of operation can reduce perioperative myocardial injury, but cannot completely eliminate its incidence. Fortunately, most myocardial injuries are asymptomatic and transient accompanied by the increase of cardiac troponin T (cTnT) and creatine kinase myocardial band (CK-MB), and only 1% to 6.4% of patients are diagnosed with postoperative myocardial infarction.<sup>[2-5]</sup>

Serious myocardial injury is a sensible risk factor for short- and long- term outcomes. CABG is a surgical method of myocardial revascularization, and many factors other than surgical injury can cause myocardial injury, such as reperfusion injury, insufficient myocardial protection, and ischemia. Most studies focus on the relationship between cut-off values of myocardial injury biomarkers (MIBs) and short- and long-term postoperative outcomes.<sup>[2-7]</sup> Because of the differences in both assessment approaches and patient situations, results of MIBs cut-off values are inconsistent and lack any unified conclusion. Moreover, in the previous studies, MIBs assessments were mostly limited once or twice at a certain time point, and lacked consequent and dynamic observation. The aim of this study was to describe the release profile of perioperative MIBs (including cTnT and CK-MB) and to investigate effects of these changes on the mid-term prognosis of patients undergoing OPCABG.

## Patients And Methods

## Patients

This was a single-centre, observational, prospective cohort clinical study. Between September 2009 and December 2014, 1386 consecutive patients underwent cardiac operation in the department of cardiovascular surgery. The inclusion criteria were as follows: 1. firstly isolated CABG; 2. No cardiopulmonary bypass (CPB); 3. over 18 years old. The exclusion criteria were as follows: 1. cardiac surgery other than CABG; 2. CABG combined with other procedures; 3. postoperative death; 4. incomplete preoperative MIBs; 5. lack of postoperative follow-up information (Figure 1).

Finally 564 patients with normal postoperative courses were enrolled. All patients were operated by the same group of surgeons and received routine treatment and nursing care after operation. All patients signed written informed consents for surgical procedures and participation in this study, which was approved by the ethics committees of Shanghai East Hospital (ID2019067) with the clinical trial registration number ChiCTR-RRC-19014055.

## Cardiac injury biomarker assay

Blood samples for MIB assay were obtained from venous punctures within 48 hours before operation and at 6, 12, 24, 48, 72, 96 and 120 hours postoperatively. Preoperative MIBs were recorded as baseline values. If a patient recovered well and was discharged within 5 days, MIBs were measured before discharge. MIBs were collected at least until there was a downtrend. Peak values were defined as the highest values measured during postoperative hospitalization. All blood samples were routinely assessed by the same method in the central laboratory of the hospital. The detection limits of cTnT were 0.1-2.0mg/l and the normal range was < 0.1mg/l. The detection limits of CK-MB were 0-400mg/l, and the normal range was 0-40mg/l.

## Follow-up

Patients were followed up in the outpatient clinic or by telephone consultation. The first follow-up was one month after discharge, and the second time was at an interval of 2 months. After that, regular follow-up was conducted every 6 months. All cause mortality was recorded and the median (interquartile range) follow-up time was 76.7(1.0-126.7) months.

## Statistical analysis

Continuous variables were expressed using mean  $\pm$  standard deviation (SD) if they conformed to normal distribution or were expressed using geometric mean with interquartile range (IQR). Categorical variables were summarized as absolute numbers and percentages. Statistical comparisons were performed using t-test, Mann-Whitney U tests for continuous variables, and using Fisher's exact or Chi-square tests for categorical variables.

Mid-term mortality was calculated and plotted according to the Kaplan-Meier survival curves. All variables were tested by univariate cox analysis. Variables with  $P < 0.08$  were entered into the multivariate analysis. Multivariate cox regression analysis by using variables from univariate cox analysis was established to identify the independent risk factors associated with mid-term mortality. Hazard ratios (HR) were expressed with 95% confidence intervals (CI).

Statistical analysis was performed on SPSS 22.0 for windows (IBM, Chicago, USA). Two-tailed  $P < 0.05$  was considered significant for all analysis.

## Results

The 564 patients had 4163 monitoring points (Some patients recovered well and were discharged within 5 days), and 3527 monitoring points were completed (3527/4163, 84.7%). About 557 (557/564, 98.8%) patients had measurements of MIBs at more than 3 postoperative time points. There were 443 males (78.5%) and 121 females (21.5%) who had normal

postoperative courses, with a mean age of  $65.3 \pm 8.2$  years. Moreover, 60 patients died during the 1.0 to 126.7 (geometric mean of 75.7) months of follow-up (**Table 1**).

### **MIBs release**

cTnT and CK-MB were measured at 7 monitoring points postoperatively. The release profile of postoperative MIBs increased first and then decreased. The increase of cTnT was more obvious than that of CK-MB. The median peak values of cTnT and CK-MB were  $0.32 \pm 0.40$  mg/l and  $30.43 \pm 33.22$  mg/l respectively. cTnT peaked within 24 hours after operation in 76.8% of the total cohort (**Figure 2A**). Similarly, CK-MB peaked in 67.7% of patients within 24 hours after surgery (**Figure 2B**).

### **Group classification**

For further evaluation, patients were divided into four groups according to two criteria (peak values and peak time courses of cTnT):

Group I: cTnT peak occurrence within 24 hours, n= 481;

Group II: cTnT peak occurrence after 24 hours, n= 83;

Group III: cTnT peak value  $\leq 0.32$  mg/l, n= 420;

Group IV: cTnT peak value  $> 0.32$  mg/l, n= 144.

### **Perioperative variables**

Patients in group I showed younger age, larger body mass index (BMI), lower serum creatinine, higher Ccr, lower Euroscore II, more arterial grafts, less venous grafts, lower CK-MB and cTnT peak values, lower prevalence of CK-MB peak value  $>30.43$  mg/l, longer follow-up time and lower mid-term mortality compared with group II (**Table 2**).

Patients in group III had lower prevalence of hypertension, total vein bypass grafting, CK-MB peak occurrence within 24 hours, CK-MB peak value  $>30.43$  mg/l and lower serum creatinine, higher Ccr, shorter operation time, less venous grafts, lower CK-MB and cTnT peak values, and lower mid-term mortality compared with group IV (**Table 3**).

### **Follow-up**

All patients were followed up after discharge. The median follow-up time was 76.7 months. The 1-, 5- and 8-year survival rates were  $98.0 \pm 0.6\%$ ,  $96.6 \pm 0.8\%$  and  $86.5 \pm 2.0\%$ , respectively. During the follow-up period, 60 patients died, accounting for 10.6% of the total cohort.

The Kaplan-Meier analyses showed that group I versus group II ( $P < 0.001$ ), and group III versus group IV ( $P = 0.031$ ) had better survival (**Figure 3A and 3B**).

### **Univariate and multivariate cox regression analysis**

All clinical variables were entered into univariate analysis. Age, cerebrovascular disease, combined valvular disease, preoperative Ccr, left ventricular ejection fraction (LVEF), New York Heart Association (NYHA), Euroscore II, number of arterial grafts, number of venous grafts, total vein bypass grafting, cTnT peak occurrence within 24 hours and cTnT peak value  $> 0.32$  mg were significantly associated with mid-term mortality. In multivariate cox regression analysis, age, LVEF, NYHA, total venous grafting and cTnT peak occurrence within 24 hours remained significant, and cTnT peak occurrence within 24 hours was the strongest predictor. Interestingly, in multivariate analysis with the same variables but without

cTnT peak occurrence within 24 hours, cTnT peak value > 0.32 mg/l was a significant independent risk factor (P=0.022). In univariate cox analysis, CK-MB-related variables had no significant effect on mid-term mortality (**Table 4**) (**Figure 4**).

## Discussion

The time course of MIBs increased first and then decreased after OPCABG. The peak value of cTnT in most patients appeared at 24 hours after surgery. We analyzed the risk factors of mid-term outcome in 564 patients undergoing OPCABG with normal postoperative courses and started from both the peak values and time courses of MIBs. It was found that postoperative peak values and peak occurrence time of cTnT were significant predictors of mid-term mortality. Through literature search and review, we know this is the first report in China to clearly put forward that the peak occurrence time is more predictive than absolute value of peak value in OPCABG patients.

Myocardial injury is an inevitable and common phenomenon during and after CABG. Available literature on this aspect has clearly reported, but not limited to, the insufficient myocardial protection, surgical procedure injuries, ischemia and factors related to patient states.<sup>[8-10]</sup> CABG is the most common cardiac performance and the largest cumulative number of cardiac surgery in the world.<sup>[11,12]</sup> Due to the reduction of extracorporeal circulation procedures, OPCABG fundamentally avoids the problem of myocardial protection. The off-pump pattern can reduce postoperative complications and mortality, which is very suitable for the relatively backward cardiac surgery technology in China. Therefore, OPCABG is very popular in China, and up to 60% of patients select CABG<sup>[1]</sup>. It is important and significant to better understand the role of MIBs in mid-term prognosis of patients undergoing OPCABG.

MIBs are the important indicators of postoperative myocardial injury in clinical practice. The cTnT with very high sensitivity and specificity is suitable for early warning of myocardial injury.<sup>[5,6,13-14]</sup> Most of the current articles focus on the relationship between specific peak values of MIBs and short- or long-term prognosis. Due to differences in assessment approaches, time and number of assessment, there is no unified understanding of this issue. Most literature involves only 2-3 measurements at certain time points after operation, including many types of cardiac surgeries and mixed postoperative death cases.<sup>[2-5,7,15-17]</sup> Those factors make it difficult to understand the profile of MIBs after surgical performances. In this study, to focus the dynamic changes of MIBs after OPCABG, we excluded death cases and observed the MIBs in a continuous manner at 7 consecutive points after operation.

Although the assessment approaches were different, the peak occurrence time was not affected in the continuous manner. This method can well unify the differences caused by different assessment approaches. As reported, the peak cTnT occurred between 24 and 48 hours after cardiac surgery,<sup>[4,5,9,18]</sup> which also confirmed our findings. In our cohort study, the cTnT peaked at 24 hours after OPCABG in most cases (76.82%) in the patients with uneventful courses. The proportion of patients whose CK-MB peaked within 24 hours after operation was 67.73%. Patients with maximal peak values had a delayed increase time of peak occurrence, which is similar to another report.<sup>[15]</sup>

Traditional MIBs include cTnT, cTnI and CK-MB. The specificity and sensitivity are higher for both cTnT and cTnI than CK-MB. In this study, although the change trends of CK-MB were similar to those of cTnT, the values and occurrence time of CK-MB peaks were not the risk factors of mid-term mortality (Table 4). This result also reflects that cTnT is superior over CK-MB<sup>[18]</sup>. The possible explanations may be that cTnT is more specific and sensitive than CK-MB in myocardial injury. The skeletal muscle injury can also cause the increase of CK-MB<sup>[19]</sup>. The univariate cox regression model suggested the value and occurrence time of cTnT peak were both risk factors of mid-term mortality. The results of the current study suggest that several factors influence the mid-term prognosis of OPCABG. Age, LVEF, NYHA, total vein bypass grafting and cTnT peak occurrence within 24 hours clearly influence the mid-term mortality. As for the effects of MIBs on the long-term prognosis of CABG patients, Camilla et al.<sup>[7]</sup> also found the cTnT value at 44 hours postoperatively was more predictive than the values at 7 or 20 hours or the peak values. Unfortunately, they did not make further quantitative

analysis or summary. Delayed postoperative elevation of cTnT suggests the persistent injury of cardiomyocytes after surgery, although there were no obvious clinical symptoms and the patients were discharged safely. This asymptomatic myocardial injury will also affect the long-term survival rate of patients undergoing OPCABG.

When other factors in the multivariate cox regression model were unchanged, the value and occurrence time of cTnT peak were included separately (Table 4). It was found that those two factors were independent risk factors of mid-term mortality. Most of studies only included the peak value when evaluating the risk factors of short- and long-term mortality, and ignored the peak occurrence time.<sup>[5-7,20]</sup> Therefore, the peak value was considered as an independent factor of long-term mortality. Interestingly, when the time of peak occurrence was considered, the results changed. The multivariate cox regression model including both peak value and peak occurrence time found only the peak occurrence time was an independent risk factor. This also indicates that the peak occurrence time is more predictive of mid-term mortality than the peak value. Delayed elevation of cTnT more greatly impacts mid-term prognosis than early elevation.

### **Limitation**

There are some limitations in this study. Firstly, the design single-centre retrospective study may cause selection bias. Secondly, the postoperative MIBs were also confected by preoperative myocardial injury (e.g. AMI), which was not considered in the article. Thirdly, the cohort size was small and the incidence of end point events in the follow-up period was also low, which may affect the statistical analysis of the data. Fourthly, the sensitivity and rang of cTnT need to be improved. This limitation will be made up for by the new-generation high-sensitivity cTnT applied in clinic<sup>[6,9]</sup>.

## **Conclusions**

The peak values and peak phases of postoperative MIBs are both risks factors for mid-term prognosis of OPCABG patients. Our data support the view that the peak phases are more powerful than the peak value in predicting mid-term prognosis. In addition, cTnT is superior over CK-MB in predicting mid-term prognosis.

## **Declarations**

### **Ethics approval and consent to participate**

All patients signed written informed consents for surgical procedures and participation in this study, which was approved by the ethics committees of Shanghai East Hospital (ID2019067) with the clinical trial registration number ChiCTR2000033850.

### **Consent for publication**

Written informed consent for publication of their clinical details was obtained from the patient/parent/guardian/ relative of the patient. A copy of the consent form is available for review by the Editor of this journal. The study protocol was carried out in accordance with the Declaration of Helsinki.

### **Availability of data and materials**

The datasets generated and analysed during the current study are not publicly available due because it contains patient personal information but are available from the corresponding author on reasonable request.

### **Competing interests**

The authors declare that they have no competing interests.

## Funding

This research was funded by Shanghai Municipal Natural Science Foundation (NO. 20ZR1446100).

## Authors' contributions

Bo Hu: Data curation, Writing- Original draft preparation. Fei Gao: Data curation, Writing- Original draft preparation. Mengwei Lv: Data curation, Investigation, Formal analysis, Writing- Reviewing and Editing. Ban Liu: Data curation, Validation. Yu Shi: Validation. Xi Chen: Investigation. Yipeng Feng: Investigation. Xiaoqi Meng: Investigation. Zhi Li: Resources, Writing- Reviewing and Editing, Supervision. Yangyang Zhang: Resources, Conceptualization, Project administration, Writing- Reviewing and Editing, Funding acquisition. All authors read and approved the final manuscript.

## Acknowledgements

Not applicable.

## References

1. Hu Shengshou, Gao Runlin, Liu Lisheng, et al. 2017. Report on Cardiovascular Diseases in China 2016. Beijing, China: Encyclopedia of China Publishing House. 210 pp.
2. Carrier M, Pellerin M, Perrault LP, Solymoss BC, Pelletier LC. Troponin levels in patients with myocardial infarction after coronary artery bypass grafting. *Ann Thorac Surg.* 2000;69(2):435-440.
3. Fransen EJ, Diris JH, Maessen JG, Hermens WT, van Dieijen-Visser MP. Evaluation of "new" cardiac markers for ruling out myocardial infarction after coronary artery bypass grafting. *Chest.* 2002;122(4):1316-1321.
4. Lehrke S, Steen H, Sievers HH, et al. Cardiac troponin T for prediction of short- and long-term morbidity and mortality after elective open heart surgery. *Clinical chemistry.* 2004;50(9):1560-1567.
5. Peivandi AA, Dahm M, Opfermann UT, et al. Comparison of cardiac troponin I versus T and creatine kinase MB after coronary artery bypass grafting in patients with and without perioperative myocardial infarction. *Herz.* 2004;29(7):658-664.
6. Mehdiani A, Akhyari P, Kamiya H, et al. Prognostic value of the new high sensitive cardiac troponin T assay (hs-cTnT) after coronary artery bypass grafting. *Acta cardiologica.* 2017;72(3):276-283.
7. Søråas CL, Friis C, Engebretsen KV, Sandvik L, Kjeldsen SE, Tønnessen T. Troponin T is a better predictor than creatine kinase-MB of long-term mortality after coronary artery bypass graft surgery. *American heart journal.* 2012;164(5):779-785.
8. Carlsson AC, Bandstein N, Roos A, Hammarsten O, Holzmann MJ. High-sensitivity cardiac troponin T levels in the emergency department in patients with chest pain but no myocardial infarction. *International journal of cardiology.* 2017;228:253-259.
9. Ge W, Gu C, Chen C, et al. High-sensitivity troponin T release profile in off-pump coronary artery bypass grafting patients with normal postoperative course. *BMC cardiovascular disorders.* 2018;18(1):157.
10. Chung JZ, Dallas Jones GR. Effect of renal function on serum cardiac troponin T—Population and individual effects. *Clinical biochemistry.* 2015;48(12):807-810.
11. Liu J, Zhang W, Wang Q, et al. The Early Diagnostic Value of Procalcitonin in Pneumonia After Off-Pump Coronary Artery Bypass Grafting Surgery. *Medical science monitor : international medical journal of experimental and clinical research.* 2019;25:3077-3089.
12. Arora RC, Singal RK. Is routine use of renal injury biomarkers in cardiac surgery patients putting the cart before the horse? *J Thorac Cardiovasc Surg.* 2017;154(3):938-939.

13. Thygesen K, Alpert JS, Jaffe AS, et al. Fourth Universal Definition of Myocardial Infarction (2018). *J Am Coll Cardiol.* 2018;72(18):2231-2264.
14. Kouvelos GN, Milionis HJ, Arnaoutoglou EM, et al. Postoperative levels of cardiac troponin versus CK-MB and high-sensitivity C-reactive protein for the prediction of 1-year cardiovascular outcome in patients undergoing vascular surgery. *Coronary artery disease.* 2011;22(6):428-434.
15. Tevaearai Stahel HT, Do PD, Klaus JB, et al. Clinical Relevance of Troponin T Profile Following Cardiac Surgery. *Frontiers in cardiovascular medicine.* 2018;5:182.
16. Neshar N, Alghamdi AA, Singh SK, et al. Troponin after cardiac surgery: a predictor or a phenomenon? *Ann Thorac Surg.* 2008;85(4):1348-1354.
17. Lurati Buse GA, Bolliger D, Seeberger E, et al. Troponin T and B-type natriuretic peptide after on-pump cardiac surgery: prognostic impact on 12-month mortality and major cardiac events after adjustment for postoperative complications. *Circulation.* 2014;130(12):948-957.
18. Sadony V, Körber M, Albes G, et al. Cardiac troponin I plasma levels for diagnosis and quantitation of perioperative myocardial damage in patients undergoing coronary artery bypass surgery. *Eur J Cardiothorac Surg.* 1998;13(1):57-65.
19. Tsung JS, Tsung SS. Creatine kinase isoenzymes in extracts of various human skeletal muscles. *Clinical chemistry.* 1986;32(8):1568-1570.
20. Vikenes K, Andersen KS, Melberg T, Farstad M, Nordrehaug JE. Long-term prognostic value of cardiac troponin I and T versus creatine kinase-MB mass after cardiac surgery in low-risk patients with stable symptoms. *Am J Cardiol.* 2010;106(6):780-786.

## Tables

**Table 1.** Clinical characteristics of entire study patients

Variables	Total(n=564)
Male sex (n, %)	443(78.5)
Age(y)	65.30±8.20
Weight (kg)	70.05±10.39
Height (cm)	
BMI (kg/m <sup>2</sup> )	25.13±3.18
BSA (m <sup>2</sup> )	1.76±0.16
Types of CAD	
Unstable angina pectoris (n,%)	332(58.9)
Acute myocardial infarction (n,%)	75(13.3)
Stable angina pectoris (n,%)	157(27.8)
Diabetes mellitus (n, %)	201(35.6)
Hypertension (n, %)	413(73.2)
Cerebrovascular disease	
Stroke (n, %)	22(3.9)
Lacunar infarction (n, %)	141(25.0)
Peripheral vascular disease (n, %)	13(2.3)
Preoperative atrial fibrillation (n, %)	15(2.7)
Pulmonary hypertension (n, %)	12(2.1)
Scrum creatinine (μmol/l)	82.51±24.57
Ccr (mL/min/1.73 m <sup>2</sup> )	80.30±25.09
Renal failure (n,%)	4(0.7)
LVEF (%)	
LVEF >= 45% (n,%)	549(97.3)
LVEF < 45% (n,%)	15(2.7)
Triple vessel disease (n, %)	505(89.5)
COPD (n, %)	9(1.6)
Previous PCI (n, %)	2(0.4)
NYHA	
I (n,%)	30(5.3)
II (n,%)	415(73.6)
III (n,%)	113(20.0)
IV (n,%)	6(1.1)

EuroSCORE II	1.79±0.96
Operating time (min)	267.96±64.04
Number of bypass grafts (n)	3.70±1.08
Number of arterial grafts (n)	1.42±0.85
Number of venous grafts (n)	2.28±1.12
Total arterial grafting (n, %)	42(7.4)
Total venous grafting (n, %)	35(6.2)
Peak of CK-MB (mg/L)	30.45±33.22
Peak of cTnT (mg/L)	0.32±0.40
CK-MB peak occurrence within 24h (n,%)	382(67.7)
CK-MB peak value > 30.43mg/L (n,%)	159(28.2)
Follow-up time (month)	75.65(59.88,93.58)
Mid-term mortality (n,%)	60(10.6)

Abbreviation: BMI, body mass index; BSA, body surface area; CAD, coronary artery disease; Ccr, creatinine clearance rate; LVEF, left ventricular ejection fraction; COPD, chronic obstructive pulmonary disease; PCI, percutaneous coronary intervention; NYHA, New York Heart Association; EuroSCORE II, European system for cardiac operative risk evaluation II; CK-MB, creatine kinase myocardial band; cTnT, cardiac troponin T.

**Table2.** Clinical characteristics of patients with cTnT peak occurrence within 24h after operation vs cTnT peak occurrence more than 24h after operation

Variables	Group I (n=481)	Group II (n=83)	<i>P</i>
Male sex (n, %)	380(79.0)	63(75.9)	0.525
Age(y)	64.77±8.22	68.41±7.39	0.000
Weight (kg)	7.035±10.53	68.30±9.39	0.097
Height (cm)		167.29±7.13	0.487
BMI (kg/m <sup>2</sup> )	25.27±3.26	24.35±2.58	0.005
BSA (m <sup>2</sup> )	1.76±0.16	1.74±0.15	0.231
Types of CAD			0.643
Unstable angina pectoris (n, %)	284(59.0)	48(57.8)	
Acute myocardial infarction (n, %)	66(13.7)	9(10.8)	
Stable angina pectoris (n, %)	131(27.2)	26(31.3)	
Diabetes mellitus (n, %)	164(34.1)	37(44.6)	0.066
Hypertension (n, %)	346(71.9)	67(80.7)	0.095
Cerebrovascular disease			0.301
Stroke (n, %)	18(3.7)	4(4.8)	
Lacunar infarction (n, %)	115(23.9)	26(31.3)	
Peripheral vascular disease (n, %)	10(2.1)	3(3.6)	0.642
Preoperative atrial fibrillation (n, %)	14(2.9)	1(1.2)	0.601
Pulmonary hypertension (n, %)	8(1.7)	4(4.8)	0.153
Scrum creatinine (µmol/l)	80.93±20.31	91.66±40.35	0.020
Ccr (ml/min/1.73 m <sup>2</sup> )	82.18±25.17	69.36±21.67	0.000
Renal failure (n, %)	2(0.4)	2(2.4)	0.197
LVEF (%)			0.878
LVEF >= 45% (n, %)	468(97.3)	81(97.6)	
LVEF < 45% (n, %)	13(2.7)	2(2.4)	
Triple vessel disease (n, %)	426(88.6)	79(95.2)	0.069
COPD (n, %)	7(1.5)	2(2.4)	0.868
Previous PCI (n, %)	2(0.4)	0(0.0)	1.000
NYHA			0.912
I (n, %)	26(5.4)	4(4.8)	
II (n, %)	356(74.0)	59(71.1)	
III (n, %)	94(19.5)	19(22.9)	
IV (n, %)	5(1.0)	1(1.2)	

EuroSCORE II	1.72±0.91	2.22±1.12	0.000
Operating time (min)	266.19±61.80	278.20±75.37	0.115
Number of bypass grafts (n)	3.67±1.09	3.87±1.02	0.119
Number of arterial grafts (n)	1.45±0.86	1.24±0.77	0.028
Number of venous grafts (n)	2.21±1.12	2.67±1.03	0.000
Total arterial grafting (n, %)	40(8.3)	2(2.4)	0.058
Total venous grafting (n, %)	26(5.4)	9(10.8)	0.058
Peak of CK-MB (mg/l)	27.22±27.61	49.06±52.00	0.000
Peak of cTnT (mg/l)	0.28±0.37	0.57±0.45	0.000
CK-MB peak occurrence within 24h (n, %)	327(68.0)	55(66.3)	0.757
CK-MB peak value > 30.43 mg/l (n, %)	114(23.7)	45(54.2)	0.000
Follow-up time (month)	77.10(61.00,94.20)	69.10(34.80,86.60)	0.001
Mid-term mortality (n, %)	38(7.9)	22(26.5)	0.000

Abbreviation: BMI, body mass index; BSA, body surface area; CAD, coronary artery disease; Ccr, creatinine clearance rate; LVEF, left ventricular ejection fraction; COPD, chronic obstructive pulmonary disease; PCI, percutaneous coronary intervention; NYHA, New York Heart Association; EuroSCORE II, European system for cardiac operative risk evaluation; CK-MB, creatine phosphokinase isoenzyme; cTnT, cardiac troponin T.

Group I: cTnT peak occurrence within 24h after operation; Group II: cTnT peak occurrence more than 24h after operation.

**Table3.** Clinical characteristics of patients with cTnT peak value ≤0.32mg/l vs.cTnT peak value >0.32mg/l

Variables	Group III (n=420)	Group IV (n=144)	P
Male sex (n, %)	338(80.5)	105(72.9)	0.057
Age(y)	65.38±8.20	65.08±8.23	0.703
Weight (kg)	70.37±10.51	69.10±10.00	0.205
Height (cm)		166.53±7.06	0.559
BMI (kg/m <sup>2</sup> )	25.22±3.25	24.87±2.96	0.257
BSA (m <sup>2</sup> )	1.77±0.16	1.75±0.16	0.223
Types of CAD			0.222
Unstable angina pectoris (n, %)	249(59.3)	83(57.6)	
Acute myocardial infarction (n, %)	50(11.9)	25(17.4)	
Stable angina pectoris (n, %)	121(28.8)	36(25.0)	
Diabetes mellitus (n, %)	154(36.7)	47(32.6)	0.384
Hypertension (n, %)	298(71.0)	115(79.9)	0.037
Cerebrovascular disease			0.536
Stroke (n, %)	16(3.8)	6(4.2)	
Lacunar infarction (n, %)	110(26.2)	31(21.5)	
Peripheral vascular disease (n,%)	8(1.9)	5(3.5)	0.447
Preoperative atrial fibrillation (n,%)	12(2.9)	3(2.1)	0.843
Pulmonary hypertension (n,%)	7(1.7)	5(3.5)	0.337
Scrum creatinine (μmol/l)	80.73±19.75	87.69±34.59	0.023
Ccr(ml/min/1.73 m <sup>2</sup> )	81.74±24.73	76.09±25.72	0.020
Renal failure (n,%)	2(0.5)	2(1.4)	0.582
LVEF (%)			1.000
LVEF >= 45% (n,%)	409(97.4)	140(97.2)	
LVEF < 45% (n,%)	11(2.6)	4(2.8)	
Triple vessel disease (n,%)	375(89.3)	130(90.3)	0.737
COPD (n,%)	6(1.4)	3(2.1)	0.876
Previous PCI (n,%)	2(0.5)	0(0)	0.986
NYHA			0.600
I (n,%)	25(6.0)	5(3.5)	
II (n,%)	305(72.6)	110(76.4)	
III (n,%)	85(20.2)	28(19.4)	
IV (n,%)	5(1.2)	1(0.7)	

EuroSCORE II	1.75±0.94	1.91±1.01	0.092
Operating time (min)	262.60±59.03	283.58±74.89	0.003
Number of bypass grafts (n)	3.67±1.12	3.76±0.97	0.388
Number of arterial grafts (n)	1.45±0.87	1.33±0.80	0.165
Number of venous grafts (n)	2.21±1.12	2.47±1.09	0.020
Total artery bypass grafting (n, %)	36(8.6)	6(4.2)	0.082
Total vein bypass grafting (n, %)	21(5.0)	14(9.7)	0.043
Peak of CK-MB(mg/l)	20.72±13.47	58.75±52.25	0.000
Peak of cTnT (mg/l)	0.15±0.06	0.84±0.51	0.000
CK-MB peak occurrence within 24h(n,%)	268(63.8)	114(79.2)	0.001
CK-MB peak value > 30.43 (n,%)	61(14.5)	98(68.1)	0.000
Follow-up time (month)	75.90(60.73,93.50)	75.05(45.25,94.68)	0.438
Mid-term mortality(n,%)	37(8.8)	23(16.0)	0.016

Abbreviation: cTnT, cardiac troponin T; BMI, body mass index; BSA, body surface area; Cr, creatinine; Ccr, creatinine clearance rate; COPD, chronic obstructive pulmonary disease; PCI, percutaneous coronary intervention; CAD, coronary artery disease; NYHA, New York Heart Association; LVEF, left ventricular ejection fraction; EuroSCORE II, European system for cardiac operative risk evaluation II.

Group III: Peak of postoperative cTnT ≤ 0.32 mg/l; Group IV: Peak of postoperative cTnT > 0.32 mg/l


**Table 4.** Univariate and multivariate cox regressions analysis for entire study patients.

	Univariate Analysis		Multivariate Analysis excluding (cTnT peak value>0.3219mg/l)		Multivariate Analysis excluding (cTnT peak occurrence within 24h)		Multivariate Analysis(Total)	
	HR (95% CI)	P-value	HR (95% CI)	P-value	HR (95% CI)	P-value	HR (95% CI)	P-value
Sex (n, %)	1.043(0.564-1.928)	0.894	0.922(0.455-1.868)	0.822	1.093(0.549-2.178)	0.800	0.930(0.459-1.885)	0.840
Age (y)	1.097(1.058-1.138)	0.000	1.060(1.000-1.122)	0.048	1.074(1.012-1.139)	0.018	1.067(1.005-1.132)	0.033
BMI (kg/m <sup>2</sup> )	0.925(0.850-1.007)	0.071	0.976(0.880-1.082)	0.976	0.979(0.883-1.084)	0.681	0.972(0.876-1.079)	0.598
BSA (m <sup>2</sup> )	0.350(0.069-1.775)	0.205						
Types of CAD (n, %)	0.717(0.473-1.088)	0.118						
Diabetes mellitus (n, %)	1.392(0.830-2.333)	0.210						
Hypertension (n, %)	1.792(0.927-3.462)	0.083	1.719(0.855-3.455)	0.128	1.734(0.865-3.476)	0.121	1.668(0.827-3.365)	0.153
Cerebrovascular disease (n, %)	1.815(1.251-2.635)	0.002	1.438(0.952-2.172)	0.084	1.508(1.004-2.266)	0.048	1.462(0.970-2.204)	0.069
Peripheral vascular disease (n, %)	0.889(0.123-6.433)	0.907						
Combined valvular disease (n, %)	2.385(1.169-4.864)	0.017	1.057(0.453-2.466)	0.897	1.054(0.450-2.468)	0.904	1.073(0.455-2.529)	0.872
Preoperative atrial fibrillation (n, %)	1.648(0.401-6.765)	0.488						
Pulmonary hypertension (n, %)	3.130(0.756-12.951)	0.115						
Ccr (ml/min/1.73 m <sup>2</sup> )	0.968(0.955-0.981)	0.000	0.982(0.963-1.002)	0.076	0.982(0.963-1.000)	0.056	0.984(0.965-1.003)	0.102
LVEF (%)	0.951(0.918-9.984)	0.004	0.951(0.911-0.993)	0.023	0.951(0.909-0.994)	0.026	0.950(0.910-0.992)	0.021
Number of diseased vessels	0.746(0.458-1.215)	0.239						
COPD (n, %)	1.133(0.157-8.187)	0.901						
Previous PCI (n, %)	0.050(0.000-1.25*10 <sup>9</sup> )	0.806						
NYHA (n,%)	2.160(1.403-3.326)	0.000	1.781(1.125-2.821)	0.014	1.894(1.195-3.003)	0.007	1.833(1.155-2.908)	0.010

EuroSCORE II	1.695(1.383-2.078)	0.000	0.816(0.544-1.225)	0.327	0.777(0.517-1.167)	0.224	0.806(0.536-1.213)	0.302
Operating time (min)	0.999(0.995-1.003)	0.486						
Number of bypass grafts	0.928(0.723-1.191)	0.558						
Number of arterial grafts	0.388(0.239-0.628)	0.000	0.964(0.537-1.732)	0.903	1.057(0.584-1.911)	0.855	0.976(0.540-1.767)	0.937
Number of venous grafts	1.311(1.037-1.659)	0.024	0.901(0.680-1.193)	0.465	0.973(0.734-1.289)	0.848	0.898(0.677-1.190)	0.452
Total artery bypass grafting (n, %)	0.480(0.150-1.538)	0.217						
Total vein bypass grafting (n, %)	6.247(3.077-12.683)	0.000	2.855(1.066-7.647)	0.037	3.040(1.135-8.144)	0.027	2.833(1.054-7.611)	0.039
CK-MB peak(mg/L)	1.001(0.995-1.008)	0.670						
CK-MB peak occurrence within 24h (n, %)	0.773(0.461-1.296)	0.328						
CK-MB peak value>30.43mg/l (n, %)	1.198(0.695-2.065)	0.516						
cTnT peak (mg/l)	1.544(0.936-2.546)	0.089						
cTnT peak occurrence within 24h(n,%)	0.243(0.143-0.413)	0.000	0.318(0.181-0.557)	0.000			0.356(0.194-0.654)	0.001
cTnT peak value>0.32 mg/L(n,%)	1.776(1.048-3.011)	0.033			1.923(1.097-3.369)	0.022	1.355(0.731-2.513)	0.335


Abbreviation: BMI, body mass index; BSA, body surface area; CAD, coronary artery disease; Ccr, creatinine clearance rate; LVEF, left ventricular ejection fraction; COPD, chronic obstructive pulmonary disease; PCI, percutaneous coronary intervention; NYHA, New York Heart Association; EuroSCORE II, European system for cardiac operative risk evaluation II; CK-MB, creatine kinase myocardial band; cTnT, cardiac troponin T.

## Figures





**Figure 1**

Flowchart of patients' election




**Figure 2**

A. Proportion of cTnT peak at each time point after operation B. Proportion of CK-MB peak at each time point after operation



**Figure 3**

A. Kaplan-Meier survival curves for mortality in group I and group II B. Kaplan-Meier survival curves for mortality in group III and group IV



**Figure 4**

Forest plot of the variables for mid-term mortality