Diet quality and its relationship with overweight and laboratory parameters in breast cancer patients undergoing treatment

Krystal Ng LS

Universiti Putra Malaysia

Chan Yoke Mun (✉ yokemun_chan@yahoo.com)

Universiti Putra Malaysia

Zalilah MS

Universiti Putra Malaysia

Research article

Keywords: Breast cancer, Diet quality, Overweight, Serum hemoglobin

Posted Date: September 24th, 2019

DOI: <https://doi.org/10.21203/rs.2.11380/v3>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background Overweight is the common issue existed among breast cancer patients, leading to poor cancer prognosis and survival. However, previous studies have focused merely on single nutrients or food groups. This study aimed to determine whether diet quality was associated with overweight and laboratory parameters among breast cancer patients undergoing treatment. **Methods** A cross sectional study comprises of 163 patients was conducted from January to May 2017. Diet quality and level of physical activity of patients were ascertained using Healthy Eating Index (HEI)-2015 and International Physical Activity Questionnaire (IPAQ)-short form, respectively. Data were available on body mass index and biochemical data, including of serum albumin, serum hemoglobin and neutrophils count. Logistic regression was performed to identify the association between diet quality and overweight, serum albumin, serum hemoglobin and neutrophils count. **Results** The mean total HEI score (standard deviation) of the breast cancer patients was 63.86 (8.75). More than 80% of the subjects did not adhere to the dietary guidelines for whole grains, dairy, refined grains, ratio of poly- and mono-unsaturated fatty acids to saturated fatty acids components. In the multivariate model, those who perceived higher score of total HEI-2015 were less likely to experience overweight (Q4 vs Q1: OR=0.36; 95% CI=0.13, 0.99) while reported higher concentration of serum hemoglobin (Q4 vs Q1: OR=0.36; 95% CI=0.13, 0.98; Q3 vs Q1: OR=0.35; 95% CI=0.13, 0.92). Non-significant associations of diet quality were shown for serum albumin and neutrophils count. **Conclusions** A majority of breast cancer patients require dietary modification, especially for fatty acids, grains and dairy products. Diet quality may associate with overweight. A balanced diet that address the problem of overweight may help to maintain a good nutritional status and to improve cancer care in breast cancer patients during treatment. Future prospective studies are required to confirm this association.

Background

Cancer- and treatment-related factors have resulted in various undesirable health outcomes among the cancer patients. Unlike other cancers such as head and neck, gastrointestinal tract, liver and lung cancers, patients with breast cancer have lower risk of undernutrition^{1,2}. Weight gain is the common issue among breast cancer patients undergoing treatment³, which may have detrimental effect on survival as a result of inflammation^{4,5}. In addition, breast cancer patients often suffer from anemia and neutropenia⁶, leads to poor prognosis. While low serum hemoglobin is related to increased level of fatigue, reduced ability to perform daily tasks and impaired quality of life^{6,7}, an abnormally low neutrophils count predisposes cancer patients to bacterial infection, which could be life-threatening⁸. Such negative impact of altered blood readings on patients' prognosis might indirectly influence treatment compliance, in light of the close linkage between quality of life and treatment compliance⁹.

In addition to tobacco use, physical activity and diet are established modifiable factor of body weight¹⁰ via principle of energy imbalance. A prospective study highlighted that breast cancer patients had low self-reported physical activity level especially during treatment¹¹, which could be in relation to the

treatment-induced fatigue or lymphedema. In regard to dietary assessment, diet quality measure provides a better understanding on the overall picture of diet consumed in accordance with dietary guidelines, rather than depend on single nutrients¹². This tool has been extensively used for epidemiological studies, in regard to the risk prediction of various health outcomes¹². In addition to the reduction in cancer risk¹³, healthier diet was found to associate with reduced inflammation¹⁴ and improved prognosis among breast cancer survivors^{15,16}. The health benefit of diet could be explained by the food synergy concept¹⁷, in which each of the nutrients acts simultaneously to yield an effective biological pathway. Conversely, Wu (2016) highlighted that insufficient protein intake could solely compromise the immune function¹⁸, which may forsake such notion that an evaluation of combined food is more informative in regards to a particular health issue. Nevertheless, how diet quality may associate with overweight and altered biochemical parameters for breast cancer patients, at the critical phase of on-going treatment is relatively sparse.

As poor diet is commonly perceived to be linked with cancer development^{19,20}, cancer patients are likely to consume low fat, high fiber diet upon diagnosis^{21,22}. Nonetheless, few studies highlighted the reduced diet quality during treatment among breast cancer patients, which resulted from the poor adherence to dietary guidelines for the components of fruits and vegetables^{23,24}, grains, solid fats, alcohol and added sugars²³. This could be explained by the altered taste or increased eating pleasure for “heavy taste” food during treatment²⁵. Preference for spicy and salty food was found to associate with overweight in breast cancer patients, in light of its potential impact on increased overall appetite via oral sensory stimulation²⁵. Diet quality covers the evaluation of a combination of food groups and nutrients from the whole food consumed, whereby the diet-disease relationship could be more detectable¹². This is very beneficial for the successful implementation of dietary strategies. The dietary guidelines help to promote healthy eating via the key principles of balanced, moderation, adequacy and variety, and so to maintain a healthy weight during cancer treatment.

To the best of our knowledge, this is the first study investigating the relationships between overall diet consumed and overweight and altered laboratory parameters in breast cancer patients undergoing treatment. This study included only non-terminally ill breast cancer patients, who were susceptible to poor dietary habit as a result of changes in taste perception while receiving treatment²⁵. Evidence is limited on which particular dietary patterns might influence clinical outcomes such as recurrence and mortality among cancer survivors²⁶. In light of the possibilities of epigenetic aberrations occurring in cancer which could be altered by nutrients makes it plausible that dietary changes after successful cancer treatment could improve prognosis²⁷, identification of dietary pattern of non-terminally ill breast cancer patients is needed. A balanced diet could be an impetus for alleviated side effects which resulting from cancer or its treatment. The findings may assist the dieticians to identify patients who need improvement in their dietary intake, thus integrating nutritional care into quality cancer care. However, little is known about the linkage of diet quality with overweight and the common treatment-induced abnormal laboratory values. This study sought to determine the diet quality of breast cancer patients at non-terminal stage as well as

its association with overweight and treatment-related side effects of hypoalbuminemia, low serum hemoglobin and low neutrophils count.

Methods

Design

This was a cross sectional study.

Samples

This study was conducted among breast cancer patients in National Cancer Institute, which is located in Federal Territory of Putrajaya. There was approximately 20 new cases of breast cancer reported each month in NCI, putrajaya. Given that number of predictors = 11, the minimum sample size required based on the formula of Green (1919), $50 + 8k$ was 138. The eligible patients were enrolled using purposive sampling. The inclusion criteria were female, aged 29 – 71 years, receiving treatments (≥ 1 course) and able to provide written consent. Patients were excluded if under hospice care or have life expectancy of ≤ 6 months, pregnant or lactating, diagnosed with severe mental disorders or reduced ability to understand or communicate with others.

Data collection

The data collection took five months (January – May 2017) and was carried out on patients' appointment day of treatment administration. Ethics approval of this study protocol was obtained from the Ministry of Health's Research and Ethics Committee (MREC) and Ethic Committee for Research Involving Human Subject (JKEUPM), Universiti Putra Malaysia, with written informed consents obtained from all patients prior to study enrolment.

Measurements

Measurements of weight and height were taken to the nearest 0.1kg and 0.1cm, respectively, by using Detecto bariatric scale (6857DHR). Overweight was defined according to body mass index (BMI) cut off point of $\geq 25.00 \text{ kg/m}^2$ ²⁸.

Habitual dietary intake of patients over the past one month was assessed using a 165-items semi-quantitative Malaysians' Food Frequency Questionnaires (FFQ)²⁹. Dietary data were analysed using Nutritionist Pro version 4.0.0 (Axxya system, 2017) based on Malaysian food composition database as the primary source, and supplemented by Singaporean and USDA database (if there's lacking of food items in Malaysian database). Food consumption data was transformed into Healthy eating index-2015 (HEI-2015) scoring according to nine adequacy and four moderation components³⁰. Each food component was scored proportionately between the minimum and maximum standard as shown in Table 1. The summation of individual components yields a total HEI-2015 score, with higher score indicates better adherence to Dietary Guidelines for Americans 2015-2020³⁰. The total HEI-2015 score was divided

into quartiles. In view of the lack of well-validated diet quality tool for Malaysians³¹, the decision of utilizing HEI-2015 tool was made to assess patients' diet quality as it covers different food groups or subgroups which enables the result to be more discussable.

International Physical Activity Questionnaire (IPAQ)-short form was used to assess level of physical activity level of patients. Total intensity level of physical activity (MET-min/week) was identified by taking consideration into frequency and duration of each activity type at one time. The higher reading of METs score indicates more intense physical activity. Patients' physical activity level was categorized into low and moderate or high according to the total frequency, duration and intensity level of physical activities that have been carried out over the past one week³².

Demographic information on age, marital status, educational level, employment status and monthly income were collected. The medical characteristics included time since diagnosis, cancer stage, treatment modalities and comorbidities were also collected. Serum albumin, serum hemoglobin and neutrophils count after receiving at least one course of treatment were retrieved as secondary data. Hypoalbuminemia, low serum hemoglobin and low neutrophils count were categorised using the cut-off points of below 35 g/L, 12.00 g/dL and $2.00 \times 10^9/L$, respectively.

Statistical analysis

All statistical analyses were performed using Statistical Software Package (IBM SPSS statistics version 22.0), with the statistical significance set at $p < 0.05$ (two-sided). Patients' characteristics were described as means $\pm$ standard deviations for continuous variables or counts with percentages for categorical variables. The differences in total mean HEI-2015 score were compared according to patients' characteristics, using analysis of variance (ANOVA) for age and month since diagnosis or chi-square test for categorical variables. Associations of HEI-2015 score with overweight, levels of albumin, hemoglobin and neutrophils count were analyzed using binary logistic regression, controlling for demographic and medical characteristics (Model 1). Additional adjustments of moderate-to-vigorous physical activity and/or BMI were performed in Model 2, to examine any changes of observed association in the first model.

Results

Of 227 breast cancer patients received treatments in hospital, 48 patients were excluded as result of unwillingness to participate ($n=41$), unable to understand the questions being addressed ($n=1$), engaged with medical procedures ($n=1$) and feeling unwell upon recruitment ($n=5$). A total of 163 patients with completed data was included in the final analysis. Table 2 illustrated the patients' characteristics by quartiles of total HEI-2015 scores. Mean age of the patients was 50.34 years old, with approximately 40% of them had tertiary education. In this study, the mean duration of treatment was 8.67 months with chemotherapy (57.7%) as the most common modality received by the patients, followed by a combination of chemotherapy and radiotherapy (37.4%). Physical inactivity was prevalent with

approximately two-third of the patients had low level of physical activity, which was accompanied with more than 50% of the patients were overweight. There was no significant difference between the HEI-2015 quartiles with respect to demographic factors of age and education level. The HEI-2015 score was poorer with longer duration of diagnosis, in which post-hoc test revealed that quartile 1 had significantly longer duration of diagnosis than quartile 4. Lower HEI-2015 score was associated with the presence of comorbidity and overweight. None of the subjects was receiving concurrent treatment, while approximately half of them were receiving chemotherapy alone. The overall mean (SD) of HEI score was 63.86 (8.75). A total of 89.6% subjects (n=146) need dietary improvement, while only 4.3% subjects (n=7) adhered to dietary guidelines at overall. In term of HEI components (Figure 1), the median scores for adequacy components of Total Fruit, Whole Fruit, Total Vegetables, Greens and Beans, Total Protein Food, Seafood and Plant Proteins, as well as moderation components of Sodium, Added Sugars and Saturated Fats achieved the maximum scores of 100%. More than half of the subjects met the recommendation for above components. On the other hand, Whole Grains, Dairy, Fatty Acids and Refined Grains scored less than 25% of the maximum score, with only 20% of patients comply with the recommended intake of these four components.

As shown in Table 3, patients in quartile 4 were less likely to be overweight as compared to patients in quartile 1 after adjusting for demographic and medical characteristics [Model 1: OR=0.36; 95%CI=0.13,0.99]. An additional variable of moderate-to-vigorous physical activity did not substantially change this significant association. In model 1, higher diet quality was associated with lower odds of low serum hemoglobin in quartile 3. The significance value was reduced after adjusting for moderate-to-vigorous physical activity and overweight [Model 2: OR=0.35; 95%CI=0.13, 0.92]. In quartile 4, the better diet quality was significantly showed with improved serum hemoglobin only after adjusting for behavioural factors [Model 2: OR=0.36; 95% CI=0.13, 0.98].

Discussion

This study investigated the associations of diet quality with overweight and treatment-related side effects among breast cancer patients undergoing treatment, with most of them underwent chemotherapy alone with a mean duration of therapy was less than 9 months. In this study, the consumption of whole grains, dairy, fatty acids and refined grains were low. This finding was consistent with earlier studies^{23,24}. Avoidance and abstinence of certain foods are common among cancer patients, owing to belief that animal products is the cause of cancer development^{19,20} or treatment-related side effects that alter the taste perception²⁵. On the other hand, dairy products are not habitual intakes of Malaysian³³, which may further attribute to the low consumption pattern of dairy products among the patients.

Despite dairy products are good sources for protein, multivitamins and minerals which are essential for bodily function, breast cancer patients may limit their dairy intake attributed to the concern about the safety of growth hormone usage in animal products²⁰. World Cancer Research Fund (WCRF) reviewed that the strength and direction of the effect of dairy products on cancer risk is varied across the cancer

sites, including of prostate, breast and colorectal cancers^{34,35}. More studies are needed to confirm the impact of dairy products especially on breast cancer. Contrary to popular belief, the intake of whole grain products that commonly known as healthy food was reduced in breast cancer patients. During treatment, cancer patients experience various side effects which could interfere with the ability of consuming a balanced diet^{25,36,37}, in spite of the high awareness of practising healthy eating. The compliance towards refined grain guideline could be deduced from the assessment of whole grains, where both of these components yield identical interpretation. As compared to whole grains, breast cancer patients prefer refined grains products with softer texture such as white rice, white bread or any food made with white flour. With regards to fatty acid component, the low score is originated from inadequate consumption of mono- and poly-unsaturated fatty acids. The individual component of saturated fat reflected that patients limited the food that are high in saturated fat, however, they did not adhere to the guidelines for unsaturated fatty acids. Although there is limited information regarding the fatty acid ratio, a consistent finding of the good compliance with saturated fatty acids intake was documented^{23,24}. Polyunsaturated fatty acids from marine sources are suggested to ameliorate treatment-related side effects and improve treatment efficacy via antineoplastic effect³⁸. In addition to saturated fatty acid, it is essential to assess the patients' intake of unsaturated fatty acids particularly during treatment. The biggest barrier to adherence to healthy fats recommendation could be high food prices³⁹, for instance, margarines with low saturated fat are more expensive than butter that high in saturated fat. This finding may suggest an enhanced accessibility to healthy food by offering affordable price or lower rate than less healthier options. Furthermore, Malaysians hardly consume raw vegetables or fruits accompanied with oil-based dressings, which is not their ordinary food habit. This elucidates that an unhealthy eating habit in general could persist over cancer diagnosis, which may influence patients' prognosis especially during the critical treatment period.

Current finding demonstrated that a better diet quality has protective effect against overweight among breast cancer patients, as supported by Custódio et al. (2016)²⁴. Similar relationship between diet quality and obesity has been widely reported by public health studies^{40–42}. On the other hand, the finding of the relationship between diet quality and cancer risk or mortality was not consistent^{13,43}. As cancer is a multi-site disease with different treatment, the relationship of diet with cancer prognosis should be studied according to various cancer types, whereby the development of cancer site-specific diet quality tools is deemed necessary. As previous studies indicated the negative impact of obesity on quality of life⁴⁴ and survival^{4,5} of cancer patients, being overweight is not a desirable outcome for cancer patients who are usually considered to be malnourished. Adopting a healthy eating habit should be a bedrock of lifestyle strategy to achieve a normal weight range as well as to maintain an overall good health. Drenowatz et al. (2014) highlighted that high consumption of protein, sodium and empty calories were the contributors of overweight or obesity among healthy adults⁴¹, indicating the necessity of achieving an adequate, balanced, moderate and varied diet. In view of the significant role of healthy eating in disease prognosis, the utilization of diet quality indices is relatively pragmatic for breast cancer patients. Despite the unhealthy dietary behaviour is strongly attributed to poor education in public health area¹⁰, current

finding failed to show the significant relationship between education and diet quality. This could be because that the distribution of studied population according to education level was not widely covered, which may reduce the ability to determine the significant result.

Iron status of cancer patients should be closely monitored as the incidence of anemia could compromise greatly their quality of life⁴⁵. Current finding showed that higher quality diet was associated with reduced risk of low serum hemoglobin only after adjusting for physical activity and BMI (Model 2), suggests that diet quality might indirectly improve anemia by reducing excess body weight. It is noteworthy that the low serum hemoglobin could be owing to multiple nutrient deficiencies^{46,47} rather than a single nutrient. Vegetables and soy food that rich in iron or components enhancing iron absorption were found to associate positively with serum hemoglobin among middle-aged adults and elderly population^{47,48}. Vitamin C, in addition to iron, was found to be essential to prevent iron deficiency for obese patients who undergone bariatric surgery⁴⁹. Vegetables and soy groups are mainly included in Healthy Eating Index assessment, elucidating the necessity of consuming these components in line with the dietary guidelines. Likewise, Mardas et al. (2017) showed that a whole food rather than a single nutrient was related with chemotherapy-induced gastrointestinal symptoms⁵⁰. Iron rich food or dietary enhancers of iron absorption could be somehow beneficial to anemic cancer patients, despite the low hemoglobin level is generally attributable to cancer or treatment-related factors. With regards to the incidence of anemia, Thomson et al. (2011) recommended to keep body weight within a healthy range⁴⁶. Inadequate diet intake could predispose patients to iron deficiency while obese women are likely to experience interrupted iron absorption^{46,48}. In view of the potential inter-relationship between health indicators, investigation on the mediating effect of body mass index in respect to the association between diet quality and low serum hemoglobin are warranted. Despite the potential linkage of diet quality with serum hemoglobin, this marker should be interpreted carefully as it may be affected by cancer prognosis related inflammation⁵¹.

Breast cancer patients were shown to be more likely to adhere to dietary guidelines within the shorter duration of diagnosis. Despite earlier studies demonstrated non-significant relationship of diagnosis time with diet quality in mixed cancer population or survivors^{15,52}, such relationship was implicitly supported by the study of Custódio et al. (2016)²⁴. Over the cancer treatment, a decline in diet quality was evident at the intermediate cycle of chemotherapy²⁴. In general, poor diet quality is attributed to lack of food varieties or an imbalanced diet. As a result of cancer treatment, nutrition impact symptoms, for instance, taste changes and nausea²⁵ could act as barriers to consuming a good quality diet. In making a decision for treatment plan, comorbidity is always a major issue for cancer patients in light of its potential dual burden on quality of life and survival⁵³. In the present study, patients with comorbidities were shown to have poorer diet quality. Noting of the non-prospective study design, this finding does not preclude the notion of substantial impact of poor diet quality on the burden of multiple health outcomes. Despite demographic factors are considered as the risk factors of unhealthy diet in the causation pathway for non-communicable diseases¹⁰, none of the variables was shown to associate with diet quality in current

study. This could be explained by the varying extent of demographic influences on diet behaviour across the population groups.

Present study yields new insights into how a quality diet was linked to the clinical outcomes of patients with breast cancer, with the evaluation of the degree of conformance in consumption of various food groups with dietary guidelines. Achieving a healthy weight range has been set as a primary goal to prevent nutrient deficiencies and to ensure an optimal health for cancer population⁵⁴. The assessment of diet quality in breast cancer patients, particularly those receiving treatment, therefore, provides valuable information for the identification of patients' nutritional status during this critical phase. Moreover, the association between diet quality and treatment-related side effects was performed using multivariate analysis, together with the adjustment of multiple confounders. This minimizes the bias that resulted from the factors that are out of study interest.

The limitations of this study included cross sectional design which determined both exposure and outcome simultaneously, whereby, the causal effect of poor diet towards treatment-related side effects could not be assured. The limited sample size of this study may result in the issue of data sparsity, which may not be representative of the overall findings. In light of the need for increased power of study and strength of evidence, a larger sample size should be considered in the future studies. The significant associations between diet quality and overweight or low haemoglobin should be interpreted with cautious as the confidence interval for odds ratio were closed to null value of 1 indicates weak implication of results. As the patient-reported outcomes rely on cancer types and treatment factors, these findings may not be generalized to other cancer groups or trajectory phases. For instance, the study population were mainly characterized by non-terminal stage, in which the study finding could not be extrapolated to breast cancer patients under hospice care with poor prognosis. The measurement periods of blood serum or body weight were inconsistent across phases of cancer treatment, which does not allow parallel comparison with 1 month of diet history. This implies that the dietary assessment over the past one month is insufficient to reflect the overall dietary pattern during treatment and the assessment period should be extended considering the data reliability without compensating patients' burden. Besides, short assessment period of laboratory measures and body weight may also be insufficient to reflect the nutritional status of patients throughout the treatment, which may undermine the study findings.

Despite Healthy Eating Index was found to be a well-established tool to predict obesity⁵⁵, it does not consider the key principle of balance, as referring that each food groups should be taken in right amount or proportion. In regard to this limitation, an underestimation of over-eating might be existed. For instance, though protein is important for muscle building, over-consumption of proteins can contribute significant calorie which could lead to overweight. Lack of nutrition information, particularly for saturated, mono- and polyunsaturated fatty acids, may underscore the component of fatty acid in overall diet quality evaluation. However, additional food databases with higher availability of food items or nutrients were utilized in order to minimize such bias.

Conclusions

In conclusion, breast cancer patients reported that their overall diet quality during treatments need improvement. A better diet quality was statistically associated with lower likelihood of overweight, however, the strength of such association was weak. On the other hand, non-significant associations were shown between diet quality and altered laboratory values. Findings from this study may serve as fundamental data regarding the impact of diet quality on overweight, which enable healthcare practitioners to implement effective strategies in the improvement of patients’ nutritional status and cancer care. More investigations are deemed necessary to identify the exact mechanism of how diet links to overweight among breast cancer patients.

List Of Abbreviations

Abbreviations	
MREC	Medical Research and Ethics Committee
IPAQ-SF	International Physical Activity Questionnaire-short form
FFQ	food frequencies questionnaires
HEI	Healthy eating index
BMI	Body mass index

Declarations

Ethics approval and consent to participate

Ethics approval was provided by Medical Research and Ethics Committee (MREC), Ministry of Health Malaysia and Ethics Committee for Research Involving Human Subject Universiti Putra Malaysia. Patients’ written informed consents were obtained prior to study enrolment.

Consent for publication

The authors give the consent for this manuscript titled Nutrition Indicators, Physical Functions and Health-related quality of life to be published in BMC Nutrition.

Availability of data and materials

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

Competing interests

The authors declare no conflict of interest regarding the publication of this articles.

Funding

This work was supported by the University Putra Malaysia (grant number GP-IPS 9507400).

Authors’ contributions

All the authors planned and designed the study. The data were collected and computed by Krystal NLS. Chan YM and Zalilah MS were involved in data analysis and interpretation. Krystal NLS prepared the manuscript and this manuscript was editing or reviewed by Chan YM and Zalilah MS.

Acknowledgements

Special thanks are dedicated to all patients and staffs of National Cancer Institute at Putrajaya, for the enthusiasm in data collection. The authors thank Dr Subashini Elangkovan from the Clinical Research Centre; Norshariza Jamhuri, Betti Sharina Mohd Haniff, Siti Nuraini Mohd Samwil, and Ng Wai Han from the Department of Dietetics, for assistance along the data collection.

References

1. Hébuterne X, Lemarié E, Michallet M, De Montreuil CB, Schneider SM, Goldwasser F. Prevalence of malnutrition and current use of nutrition support in patients with cancer. *J Parenter Enter Nutr*. 2014;38(2):196-204.
2. Pressoir M, Desné S, Berchery D, Rossignol G, Poiree B, Meslier M, Traversier S, Vittot M, Simon M, Gekiere JP et al. Prevalence, risk factors and clinical implications of malnutrition in french comprehensive cancer centres. *Br J Cancer*. 2010;102(6):966-971.
3. van den Berg MMGA, Winkels RM, de Kruif JT, van Laarhoven HW, Visser M, de Vries JHM, de Vries YC, Kampman E. Weight change during chemotherapy in breast cancer patients: a meta-analysis. *BMC Cancer*. 2017;17(1):259.
4. Jiralerspong S, Goodwin PJ. Obesity and breast cancer prognosis: Evidence, challenges, and opportunities. *J Clin Oncol*. 2016;34(35):4203-4216.
5. Picon-ruiz M, Morata-tarifa C, Valle-goffin JJ, Friedman ER, Slingerland JM. Obesity and Adverse Breast Cancer Risk and Outcome: Mechanistic Insights and Strategies for Intervention. *A Cancer Journals Clin*. 2017;67(5).
6. Tachi T, Teramachi H, Tanaka K, Asano S, Osawa T, Kawashima A, Yasuda M, Mizui T, Nakada T, Noguchi Y et al. The impact of outpatient chemotherapy-related adverse events on the quality of life of breast cancer patients. *PLoS One*. 2015;10(4):1-15.
7. Bower JE. Cancer-related fatigue: Mechanisms, risk factors and treatments. *Nat Rev Clin Oncol*. 2014;11(10):597-609.
8. Lyman GH, Michels SL, Reynolds MW, Barron R, Tomic KS, Yu J. Risk of mortality in patients with cancer who experience febrile neutropenia. *Cancer*. 2010;116(23):5555-5563.
9. Kane HL, Halpern MT, Squiers LB, Treiman KA, McCormack LA. Implementing and Evaluating Shared Decision Making in Oncology Practice. *CA Cancer J Clin*. 2014;64(6):377-388.
10. Dans A, Ng N, Varghese C, Tai ES, Firestone R, Bonita R. The rise of chronic non-communicable diseases in southeast Asia: Time for action. *Lancet*. 2011;377(9766):680-689.

11. Gal R, Monninkhof EM, Peeters PHM, van Gils CH, van den Bongard DHJG, Wendel-Vos GCW, Zuithoff NPA, Verkooijen HM, May AM. Physical activity levels of women with breast cancer during and after treatment, a comparison with the Dutch female population. *Acta Oncol.* 2019;58(5):673-681.
12. Wirt A, Collins CE. Diet quality - What is it and does it matter? *Public Health Nutr.* 2009;12(12):2473-2492.
13. Du M, Liu SH, Mitchell C, Fung TT. Associations between diet quality scores and risk of postmenopausal estrogen receptor-negative breast cancer: A systematic review. *J Nutr.* 2018;148(1):100-108.
14. George SM, Neuhouster ML, Mayne ST, Irwin ML, Albanes D, Gail MH, Alfano CM, Bernstein L, McTiernan A, Reedy J et al. Postdiagnosis diet quality is inversely related to a biomarker of inflammation among breast cancer survivors. *Cancer Epidemiol Biomarkers Prev.* 2010;19(9):2220-2228.
15. George SM, Ballard-Barbash R, Shikany JM, Caan BJ, Freudenheim JL, Kroenke CH, Vitolins MZ, Beresford SA, Neuhouster ML. Better Postdiagnosis Diet Quality Is Associated with Reduced Risk of Death among Postmenopausal Women with Invasive Breast Cancer in the Women's Health Initiative. *Cancer Epidemiol Biomarkers Prev.* 2014;23(4):575-583.
16. Terranova CO, Protani MM, Reeves MM. Overall Dietary Intake and Prognosis after Breast Cancer: A Systematic Review. *Nutr Cancer.* 2018;70(2):153-163.
17. Jacobs DR, Gross MD, Tapsell LC. Food synergy: an operational concept for understanding nutrition. *Am J Clin Nutr.* 2009;89(5):1543-1548.
18. Wu G. Dietary protein intake and human health. *Food Funct.* 2016;7(3):1251-1265.
19. Beeken RJ, Williams K, Wardle J, Croker H. "What about diet?" A qualitative study of cancer survivors' views on diet and cancer and their sources of information. *Eur J Cancer Care (Engl).* 2016;25(5):774-783.
20. Thomson AK, Heyworth JS, Girschik J, Slevin T, Saunders C, Fritschi L. Beliefs and perceptions about the causes of breast cancer: a case-control study. *BMC Res Notes.* 2014;7:558.
21. Shaharudin SH, Sulaiman S, Shahril MR, Emran NA, Akmal SN. Dietary changes among breast cancer patients in Malaysia. *Cancer Nurs.* 2013;36(2):131-8.
22. Yong HY, Shariff ZM, Kandiah M, Yong HW, Saibul N, Sariman S, Hashim Z. Diet and physical activity in relation to weight change among breast cancer patients. *Asian Pac J Cancer Prev.* 2014;15(1):39-44.
23. Ceccatto V, Faria Di Pietro P, Nogueira Previdelli A, Kunradi Vieira FG, Cesa Schiavon C, Engel R, Lizane Cardoso A, Altenburg de Assis MA, Gilberto Crippa C, Gonzalez Chica DA. Brazilian healthy eating index revised (BHEI-R) of women before and during adjuvant treatment for breast cancer. *Nutr Hosp.* 2014;30(5):1101-1109.
24. Custódio IDD, Marinho EDC, Gontijo CA, Pereira TSS, Paiva CE, Maia YC de P. Impact of chemotherapy on diet and nutritional status of women with breast cancer: A prospective study. *PLoS*

- One. 2016;11(6):1-21.
25. Marinho E da C, Custódio IDD, Ferreira IB, Crispim CA, Paiva CE, Maia YC de P. Impact of chemotherapy on perceptions related to food intake in women with breast cancer: A prospective study. PLoS One. 2017;12(11):1-14.
26. Jochems SHJ, Van Osch FHM, Bryan RT, Wesselius A, van Schooten FJ, Cheng KK, Zeegers MP. Impact of dietary patterns and the main food groups on mortality and recurrence in cancer survivors: A systematic review of current epidemiological literature. BMJ Open. 2017;8(2):1-12.
27. Leenders M, Sluijs I, Ros MM, Boshuizen HC, Siersema PD, Ferrari P, Weikert C, Tjønneland A, Oisen A, Boutron-Ruault M et al. Fruit and vegetable consumption and mortality. Am J Epidemiol. 2013;178(4):590-602.
28. World Health Organization. Regional Office for the Western Pacific. The Asia-Pacific Perspective: Redefining Obesity and Its Treatment. Sydney: Health Communications Australia. 2000. Available from <http://www.wpro.who.int/nutrition/documents/docs/Redefiningobesity.pdf>
29. Institute for Public Health. National Health and Morbidity Survey 2014: Malaysian Adult Nutrition Survey Volume I: Methodology and General Findings. 2014. Available from <http://iku.moh.gov.my/images/IKU/Document/REPORT/NHMS2014-MANS-VOLUME-1-MethodologyandGeneralFind.pdf>
30. National Cancer Institute; National Institutes of Health. Overview and Background of The Healthy Eating Index. [Internet]. 2018. Available from <https://epi.grants.cancer.gov/hei/>. Cited 19 July 2018.
31. Fokeena WB, Jamaluddin R, Khaza'ai H. Development and Assessment of the Reliability and Validity of a Diet Quality Index in a Sample of Malaysian University Students. J Food Nutr Res. 2016;4(4):251-257.
32. IPAQ (2005). Guidelines for data processing and analysis of the International Physical Activity Questionnaire (IPAQ) - short and long forms. 2005. Available from <https://sites.google.com/site/theipaq/scoring-protocol>
33. Shahril MR, Sulaiman S, Shaharudin SH, Akmal SN. Healthy eating index and breast cancer risk among Malaysian women. Eur J Cancer Prev. 2013;22(4):342-347.
34. World Cancer Research Fund, American Institute for Cancer Research. Diet, Nutrition, Physical Activity and Cancer: A Global Perspective. Meat, Fish and Dairy Products and the Risk of Cancer. 2018. Available from <https://www.wcrf.org/sites/default/files/Meat-Fish-and-Dairy-products.pdf>.
35. World Cancer Research Fund/American Institute for Cancer Research. Diet, Nutrition, Physical Activity and Cancer: A Global Perspective. [Internet]. 2018. Available from <https://www.wcrf.org/dietandcancer>. Cited 31 July 2019.
36. Boltong A, Aranda S, Keast R, Wynne R, Francis PA, Chirgwin J, Gough K. A prospective cohort study of the effects of adjuvant breast cancer chemotherapy on taste function, food liking, appetite and associated nutritional outcomes. PLoS One. 2014;9(7):1-9.
37. de Vries YC, van den Berg MMGA, de Vries JHM, Boesveldt S, DE Kruif JTCM, Buist N, Haringhuizen A, Los M, Sommeijer DW, Timmer-Bonte JHN et al. Differences in dietary intake during chemotherapy

- in breast cancer patients compared to women without cancer. *Support Care Cancer*. 2017;25(8):2581-2591.
38. D'Eliseo D, Velotti F. Omega-3 Fatty Acids and Cancer Cell Cytotoxicity: Implications for Multi-Targeted Cancer Therapy. *J Clin Med*. 2016;5(2):15.
 39. Rao M, Afshin A, Singh G, Mozaffarian D. Do healthier foods and diet patterns cost more than less healthy options? A systematic review and meta-analysis. *BMJ Open*. 2013;3(12).
 40. Livingstone KM, McNaughton SA. Diet quality is associated with obesity and hypertension in Australian adults: a cross sectional study. *BMC Public Health*. 2016;16(1):1-10.
 41. Drenowatz C, Shook RP, Hand GA, Hébert JR, Blair SN. The independent association between diet quality and body composition. *Sci Rep*. 2014;4:1-6.
 42. Drewnowski A, Aggarwal A, Tang W, Hurvitz PM, Scully J, Stewart O, Moudon AV. Obesity, diet quality, physical activity, and the built environment: the need for behavioral pathways. *BMC Public Health*. 2016;16(1):1-12.
 43. Potter J, Brown L, Williams RL, Byles J, Collins CE. Diet quality and cancer outcomes in adults: A systematic review of epidemiological studies. *Int J Mol Sci*. 2016;17(7):1-30.
 44. Fang P, Tan KS, Troxel AB, Rengan R, Freedman G, Lin LL. High body mass index is associated with worse quality of life in breast cancer patients receiving radiotherapy. *Breast Cancer Res Treat*. 2013;141(1):125-133.
 45. Kanuri G, Sawhney R, Varghese J, Britto M, Shet A. Iron deficiency anemia coexists with cancer related anemia and adversely impacts quality of life. *PLoS One*. 2016;11(9):1-11.
 46. Thomson CA, Stanaway J, Neuhauser ML, Snetselaar LG, Stefanick ML, Arendell L, Chen Z. Nutrient Intake and Anemia Risk in the WHI Observational Study. *J Am Diet Assoc*. 2011;111(4):532-541.
 47. Zhang Q, Qin G, Liu Z, Li Z, Li J, Varma DS, Wan Q, Zhao J, Min X, Han X et al. Dietary balance index-07 and the risk of anemia in middle aged and elderly people in southwest China: A cross sectional study. *Nutrients*. 2018;10(2):1-12.
 48. Song P, Li L, Man Q, Wang C, Meng L, Zhang J. Case-control study of anaemia among middle-aged and elderly women in three rural areas of China. *BMJ Open*. 2014;4(8).
 49. Mischler RA, Armah SM, Wright BN, Mattar SG, Rosen AD, Gletsu-Miller N. Influence of diet and supplements on iron status after gastric bypass surgery. *Surg Obes Relat Dis*. 2016;12(3):651-658.
 50. Mardas M, Madry R, Stelmach-Mardas M. Link between diet and chemotherapy related gastrointestinal side effects. *Wspolczesna Onkol*. 2017;21(2):162-167.
 51. Fankhauser CD, Sander S, Roth L, Goss O, Eberli D, Sulser T, Seifert B, Beyer J, Hermanns T. Systemic inflammatory markers have independent prognostic value in patients with metastatic testicular germ cell tumours undergoing first-line chemotherapy. *Br J Cancer*. 2018;118(6):825-830.
 52. Kane K, Ilic S, Paden H, Lustberg M, Grenade C, Bhatt A, Diaz D, Beery A, Hatsu I. An evaluation of factors predicting diet quality among cancer patients. *Nutrients*. 2018;10(8):1-10.

53. Sarfati D, Koczwara B, Jackson C. The impact of comorbidity on cancer and its treatments. CA Cancer J Clin. 2016;00(0):1-13.
54. Rock CL, Doyle C, Demark-Wahnefried W, Meyerhardt J, Courneya KS, Schwartz AL, Bandera EV, Hamilton KK, Grant B, McCullough M et al. Nutrition and Physical Activity Guidelines for Cancer Survivors. CA Cancer J Clin. 2012;00:000-000.
55. Asghari G, Mirmiran P, Yuzbashian E, Azizi F. A systematic review of diet quality indices in relation to obesity. Br J Nutr. 2017;117(8):1055-1065.

Tables

Table 1: Standard scores of HEI-2015 components

HEI-2015 components	Maximum scores	Standard for maximum score	Standard for minimum score
Adequacy			
Total fruits	5	≥ 0.8 cup	No Fruits
Whole fruits	5	≥ 0.4 cup	No Whole Fruits
Total vegetables	5	≥ 1.1 cup	No Vegetables
Greens and beans	5	≥ 0.2 cup	No Greens and Beans
Whole grains	10	≥ 1.5 oz	No Whole Grains
Dairy	10	≥ 1.3 cup	No Dairy
Total Protein Food	5	≥ 2.5 oz	No Protein Foods
Seafood and Plant Proteins	5	≥ 0.8 cup	No Seafood and Protein food
Fatty Acids	10	$(\text{PUFAs} + \text{MUFAs})/\text{SFAs} \geq 2.5$	$(\text{PUFAs} + \text{MUFAs})/\text{SFAs} \leq 1.2$
Moderation			
Refined Grains	10	≤ 1.8 oz	≥ 4.3 oz
Sodium	10	≤ 1.1 g	≥ 2.0 g
Added sugars	10	$\leq 6.5\%$ of energy	$\geq 26\%$ of energy
Saturated fats	10	$\leq 8\%$ of energy	$\geq 16\%$ of energy

Table 2: Patients' characteristics by total HEI-2015 score

Parameters	Quartiles of HEI-2015 score				<i>p</i>
	1 (41 to <58.7)	2 (58.7 to <62.5)	3 (62.5 to <69.5)	4 (69.5 to 89)	
Number of patients	41	41	41	40	
Age (years), mean±SD	50.95±11.40	48.83±10.28	51.41±10.09	50.15±9.13	0.684
Educational level					0.567
Secondary level and below	25 (61.0)	27 (65.9)	21 (51.2)	25 (62.5)	
Higher education	16 (39.0)	14 (34.1)	20 (48.8)	15 (37.5)	
Month since diagnosis, mean±SD	11.01±6.69*	8.28±5.96	8.00±5.63	7.31±4.92*	0.027
Cancer stage					0.380
0/I/II	12 (29.3)	16 (39.0)	16 (39.0)	16 (40.0)	
III	19 (46.3)	15 (36.6)	18 (43.9)	10 (25.0)	
IV	10 (24.4)	10 (14.4)	7 (17.1)	14 (35.0)	
Chemotherapy (ongoing)					0.164
Yes	21 (51.2)	25 (61.0)	29 (70.7)	29 (72.5)	
No	20 (48.8)	16 (39.0)	12 (29.3)	11 (27.5)	
Radiotherapy (ongoing)					0.061
Yes	13 (31.7)	11 (26.8)	5 (12.2)	5 (12.5)	
No	28 (68.3)	30 (73.2)	36 (87.8)	35 (87.5)	
Hormonal therapy (ongoing)					0.316
Yes	13 (31.7)	7 (17.1)	7 (17.1)	8 (20.0)	
No	28 (68.3)	34 (82.9)	34 (82.9)	32 (80.0)	
Targeted therapy (ongoing)					0.904
Yes	7 (17.1)	5 (12.2)	6 (14.6)	7 (17.5)	
No	34 (82.9)	36 (87.8)	35 (85.4)	33 (82.5)	
Combined & single treatment modality					0.062†
Chemotherapy alone	19 (46.3)	24 (58.5)	27 (65.9)	24 (60.0)	
Radiotherapy alone	3 (7.3)	5 (12.2)	0 (0.0)	0 (0.0)	
Chemotherapy + Radiotherapy	19 (46.3)	12 (29.3)	14 (34.1)	16 (40.0)	
Presence of comorbid condition					0.021
Yes	24 (58.5)	13 (31.7)	22 (53.7)	13 (32.5)	
No	17 (41.5)	28 (68.3)	19 (46.3)	27 (67.5)	
Meets moderate/vigorous PA					0.779
Yes	12 (29.3)	14 (34.1)	15 (36.6)	16 (40.0)	
No	29 (70.7)	27 (65.9)	26 (63.4)	24 (60.0)	
Overweight (≥24kg/m ²)					0.021
Yes	28 (68.3)	19 (46.3)	23 (56.1)	14 (35.0)	
No	13 (31.7)	22 (53.7)	18 (43.9)	26 (65.0)	
Hypoalbuminemia					0.691
Yes	3 (7.3)	3 (7.3)	4 (9.8)	1 (9.1)	
No	38 (92.7)	38 (92.7)	37 (90.2)	39 (97.5)	
Low haemoglobin					0.186†
Yes	28 (68.3)	25 (62.5)	19 (46.3)	21 (52.5)	
No	13 (31.7)	15 (37.5)	22 (53.7)	19 (47.5)	
Low neutrophils count					0.700
Yes	3 (7.3)	5 (12.2)	6 (14.6)	6 (15.0)	

No	38 (92.7)	36 (87.8)	35 (85.4)	34 (85.0)	
----	-----------	-----------	-----------	-----------	--

Data were presented as mean±SD and number of frequency (%)

Figure in bold indicate statistical significance based on ANOVA and chi-square test

*Post-hoc test revealed that Q1 had significantly higher score than Q4 (p=0.025)

†Statistical significance based on fisher's exact test

Table 3: Multivariate associations of total HEI-2015 score with overweight status and treatment-related side effects

	HEI-2015 score (ref: Quartile 1= 41 to <58.7)					
	2 (58.7 to <62.5)		3 (62.5 to <69.5)		4 (69.5 to 89)	
	OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p
Overweight ^a						
Model 1	0.58 (0.22, 1.54)	0.270	0.70 (0.26, 1.83)	0.463	0.36 (0.13, 0.99)	0.047
Model 2	0.58 (0.22, 1.54)	0.270	0.70 (0.26, 1.83)	0.462	0.36 (0.13, 0.99)	0.047
Hypoalbuminemia ^b						
Model 1	0.94 (0.13, 6.74)	0.951	1.21 (0.20, 7.18)	0.832	0.17 (0.01, 2.15)	0.170
Model 2	0.87 (0.11, 6.73)	0.898	1.20 (0.19, 7.52)	0.847	0.13 (0.10, 1.96)	0.142
Low hemoglobin ^b						
Model 1	0.65 (0.24, 1.76)	0.398	0.37 (0.14, 0.95)	0.039	0.39 (0.14, 1.05)	0.062
Model 2	0.64 (0.24, 1.72)	0.374	0.35 (0.13, 0.92)	0.033	0.36 (0.13, 0.98)	0.046
Low neutrophils count ^b						
Model 1	1.47 (0.26, 8.32)	0.662	2.85 (0.55, 14.89)	0.214	2.36 (0.44, 12.63)	0.317
Model 2	1.15 (0.17, 7.65)	0.887	2.63 (0.45, 15.59)	0.286	1.88 (0.31, 11.34)	0.492

n=163 for each model fitted

^aModel 1: adjusted for age (continuous), higher education, time since diagnosis (continuous), cancer stage, hormonal therapy, comorbidity; Model 2: Model 1 + moderate-to-vigorous physical activity;

^b Model 1: adjusted for age (continuous), higher education, time since diagnosis (continuous), cancer stage, hormonal therapy, comorbidity; Model 2: Model 1 + moderate-to-vigorous physical activity + overweight

Figures

Figure 1: Radar graph of median components score in HEI-2015 and percentage of meeting dietary recommendations

Median scores of component were expressed in percentage (%)
 Higher % indicate better conformance towards dietary guidelines
 *Fatty Acids: ratio of poly- and mono-unsaturated fatty acids to saturated fatty acids

Figure 1

Radar graph of median components score in HEI-2015 and percentage of meeting dietary recommendations

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Tables.pdf](#)