

Association Between Plasma Homocysteine Levels and Cognitive Deficits in Han Chinese Patients with Schizophrenia across Age Groups

Sumiao Zhou

The Affiliated Brain Hospital of Guangzhou Medical University

Yuanyuan Huang

The Affiliated Brain Hospital of Guangzhou Medical University

Yangdong Feng

The Affiliated Brain Hospital of Guangzhou Medical University

Hehua Li

The Affiliated Brain Hospital of Guangzhou Medical University

Kai Wu

School of Materials Science and Engineering, South China University of Technology (SCUT)

Mingzhe Yang

The Affiliated Brain Hospital of Guangzhou Medical University

Fengchun Wu

The Affiliated Brain Hospital of Guangzhou Medical University

Xingbing Huang (✉ hxbing2002@163.com)

The Affiliated Brain Hospital of Guangzhou Medical University

Research Article

Keywords: Homocysteine, Cognitive deficits, Schizophrenia, Age

Posted Date: February 23rd, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-226210/v1>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background: It was still unclear how homocysteine (Hcy) levels and cognitive deficits change in patients with schizophrenia of various ages. The present article attempts to assess the relationship between Hcy levels and cognitive deficits in patients with schizophrenia across age groups, especially in young people.

Methods: Totals of 103 patients and 122 healthy controls were included. All participants were stratified into four groups according to their age: 18-29 years, 30-39 years, 40-49 years, and 50-59 years. Clinical data, plasma Hcy levels, and cognitive function score were collected. Cognitive function was evaluated using the MATRICS Consensus Cognitive Battery of tests assessing speed of processing, verbal learning and memory, visual learning and memory, working memory, and attention/vigilance.

Results: Compared with the healthy group, Hcy levels increased significantly, and all the measured cognitive function score were significantly lower in all age groups of patients with schizophrenia ($p < 0.001$). Hcy levels were negatively associated with speed of processing (SoP), working memory (WM), and visual learning and memory (Vis Lrng) score in 18-29 years. Further multiple regression analysis showed that SoP were independently associated with Hcy levels in patients with schizophrenia aged 18-29 years ($B = 0.61, t = 3.17, p = 0.013$).

Conclusions: Based on our results, patients with schizophrenia performed worse on cognitive assessments and Hcy levels were more closely related to cognition in young patients.

Introduction

Schizophrenia is a severe psychiatric syndrome characterized by psychotic symptoms, negative symptoms and cognitive deficits. It is among the top 10 global causes of disability¹. Biomarkers, such as homocysteine (Hcy), kynurenic acid, C-reactive protein, and triiodothyronine, have been shown to be potential risk factors for schizophrenia²⁻⁵. Hcy interacts with NMDA receptors, induces apoptosis, triggers oxidative stress, injures mitochondrial function and leads to vascular damage, which played an essential role in the occurrence and development of schizophrenia⁶. Elevated plasma Hcy levels might also contribute to cognitive deficits that are common in patients with schizophrenia, even when the analyses are corrected for the relevant covariates⁷. However, several epidemiological studies have shown that elevated Hcy levels do not appear to be correlated with cognitive deficits in patients with schizophrenia⁸. Therefore, the association between Hcy levels and cognitive deficits in patients with schizophrenia continues to be debated.

Abnormal accumulation of Hcy in plasma is a high-risk factor for neurodegeneration in individuals aged 65 years⁹. Moreover, a previous study showed that plasma Hcy levels increased significantly with age in a Northeast Chinese population¹⁰. In addition, Chen PH et al⁶ showed a significant association between higher plasma Hcy concentration and worse cognitive performance in patients with late-onset bipolar disorder. Taken together, age may play an important role in determining Hcy levels in both the healthy

population and patients with psychiatric disorders. However, we do not know the association between Hcy levels and cognitive deficits in patients with schizophrenia across of various ages.

Based on the increased plasma Hcy levels and cognitive deficits in patients with schizophrenia, as well as the close relationship between Hcy levels and age, we were very interesting in exploring their possible association in patients with schizophrenia. To the best of our knowledge, no study has reported the association between Hcy levels and cognitive deficits in patients with schizophrenia across age groups. Therefore, there are two purposes for the present study: (1) to determine whether differences in plasma Hcy levels exist in patients with schizophrenia across age groups by performing a statistical analysis and (2) to determine whether plasma Hcy levels and cognitive deficits are correlated in patients with schizophrenia from different age groups.

Methods

Subjects and Procedures

Totals of 103 patients who met DSM-V diagnosis for schizophrenia were recruited from the inpatient department of the Affiliated Brain Hospital of Guangzhou Medical University. All patients were conducted from December 2018 to May 2019. The study was approved by the ethics committee of the Affiliated Brain Hospital of Guangzhou Medical University. All individuals signed the informed consent form. The patients also met the following criteria: (a) Han Chinese, age 18–59 years, (b) with a duration of illness of at least 5 years and at least 5 years of education, (c) a diagnosis conducted by an independent experienced psychiatrist on the basis of the Structured Clinical Interview for DSM-V (SCID), and (d) take a stable dose of antipsychotic drugs for ≥ 4 weeks before the start of the study.

The exclusion criteria were as follows: (a) use of immunosuppressive agents; (b) past history of autoimmune disorders; (c) substance dependence/abuse; (d) breastfeeding or pregnancy; and (e) major medical abnormalities, including central nervous system diseases and cancer.

Totals of 122 healthy individuals were recruited from community. All subjects were recruited during the same period from Guangzhou. None of them had serious physical diseases or drug abuse/dependence, excluding nicotine.

Psychotic Symptoms

The Positive and Negative Syndrome Scales (PANSS) were performed to assess the clinical symptoms of all patients by two psychiatrists who participated in a training course. Repeated assessment found that the interrater correlation coefficient remained ≥ 0.80 for the PANSS total score.

Cognitive Assessment

Cognitive function was evaluated using the MATRICS Consensus Cognitive Battery (MCCB) of tests assessing speed of processing (SoP), working memory (WM), verbal learning and memory (Vrbl Lrng),

visual learning and memory (Vis Lmg), and attention/vigilance (AV). The two researchers participated in a training course in the use of MCCB to ensure the consistency and reliability of ratings. They maintained an interrater correlation coefficient of 0.85 on the MCCB in repeated assessments.

Hcy Analysis

Plasma samples were obtained to analyze plasma Hcy levels on the same day as the cognitive function assessments. Peripheral venous blood was collected from the participants after at least 8 hours of fasting. Plasma Hcy levels were measured using the enzyme cycling assay and analyzed by a technician from Affiliated Brain Hospital of Guangzhou Medical University.

Statistical Analysis

Clinical and demographic data were analyzed using analysis of variance (ANOVA) for continuous variables and the chi-square test for categorical variables. If the ANOVA results were significant, we then performed an analysis of covariance (ANCOVA) of cognitive deficits and controlled for the confounding effects of age, BMI, and education. Second, we separately determined associations between plasma Hcy levels and cognitive deficits in stable patients with schizophrenia across age groups. Finally, multiple regression analyses was used to qualify the correlations between plasma Hcy levels and cognitive deficits in patients with schizophrenia after adjustment for age, BMI, and education. SPSS version 22.0 were used to do all data, and statistical significance was defined as $p \leq 0.05$ (two-sided).

Results

Differences in demographics, cognitive function and Hcy levels between patients and healthy controls.

Totals of 103 patients and 122 healthy controls were analyzed. Table 1 shows significant differences in age, BMI, and education between patients with schizophrenia and healthy controls ($p < 0.001$). Thus, we controlled for age, BMI, and education in subsequent analyses.

In the patient group, all individuals were stratified into four groups according to age: ≤ 18 -29 years ($n = 17$), 30–39 years ($n = 19$), 40–49 years ($n = 32$), and 50-59 years ($n = 35$). Significant differences were observed in education and duration of illness ($p < 0.05$). However, no significant differences in positive symptoms, negative symptoms, general psychopathology syndrome, or PANSS total score were observed across age groups.

Changes in cognitive deficits and Hcy levels in schizophrenia and healthy controls across age groups.

Plasma Hcy levels were significantly lower in patients with schizophrenia than in healthy controls ($p < 0.001$). As shown in Figure 1, the highest plasma Hcy levels were detected in the 50–59 years group and the lowest levels were detected in the 30-39 years group. Compared with healthy controls, significantly higher Hcy levels were observed in patients with schizophrenia across all age groups, which remained significant after adjustment for age, BMI, and education (all $p < 0.05$).

Figure 2 shows significant differences in cognitive test score for the five indexes of MCCB between patients with schizophrenia and healthy controls among the four groups, which remained significantly after adjustment for age, BMI, and education, except for 40-49 years in WM score ($p = 0.103$), 30-39 years in Vrbl Lrng score ($p = 0.199$), 30-39 years and 40-49 years in Vis Lrng score ($p = 0.087$, $p = 0.084$), and 18-29 years in AV score ($p = 0.195$). The lowest SoP, WM, Vis Lrng, and AV score were recorded by patients aged 50–59 years, and the lowest Vrbl Lrng score were recorded by patients aged 18–29 years.

Age-related differences in the relationships between Hcy levels and clinical symptoms and cognitive deficits in schizophrenia.

Table 2 shows the correlations between Hcy levels, clinical symptoms and cognitive deficits in patients. As shown, plasma Hcy levels showed negatively correlated with SoP, WM, and Vis Lrng score in schizophrenia aged 18-29 years ($r = -0.60$, $p = 0.011$; $r = -0.62$, $p = 0.008$; and $r = -0.50$, $p = 0.039$, respectively). In addition, Hcy levels were negatively correlated with Vrbl Lrng score in patients aged 40-49 years ($r = -0.48$, $p = 0.015$).

After adjusting for confounding factors (including education and duration of illness), a further multiple regression analysis performed in the subgroup of patients with schizophrenia showed independent associations between SoP and Hcy levels ($B = 0.61$, $t = 3.17$, $p = 0.013$) in patients with schizophrenia aged 18-29 years. However, no significant relationships were observed between Hcy levels and Vrbl Lrng score in patients aged 40-49 years.

Discussion

To our best knowledge, the report is firstly explore the age-specific difference in Hcy levels and the correlation with cognitive deficits in schizophrenia. Our study reported several major findings: (1) Higher plasma Hcy levels were detected in patients with schizophrenia than in healthy controls across age groups. (2) Significant differences in MCCB test score were observed for schizophrenia compared to healthy controls among the four age groups. (3) With increasing age, the incidences of increased Hcy levels were not significantly increased. (4) More interestingly, we observed age-related differences in the relationships between increased plasma Hcy levels and cognitive deficits in patients with schizophrenia. Significant relationships between SoP, WM, and Hcy levels were only observed in the 18-29 year age group but not in the other subgroups.

Regland et al. first detected the Hcy levels, and high levels were detected in 45% of patients with schizophrenia¹¹. Similar to our results, the Hcy levels were increased in schizophrenia compared to healthy controls, even across age groups. This result is consistent with findings supporting that hypothesis that Hcy levels might be confounding factors in schizophrenia¹².

Haidemenos et al. defined schizophrenia as a disease characterized by generalized cognitive deficits¹³;¹⁴. Consistent with previous research, our study showed that patients with schizophrenia exhibited

significantly worse performance on neurocognitive tests than healthy controls in most age groups¹⁵, which suggests the occurrence of cognitive deficits in the early course of the disease¹⁶.

As shown in the present study, the Hcy concentration first decreased and then increased, as the lowest levels were at 30-39 years of age and significantly increased after 40 years of age. In several longitudinal studies, the progression of changes in Hcy levels associated with age was not confirmed. Additionally, a recent large-sample study including 7872 subjects evaluated the age-related differences in Hcy levels in a general Chinese population and showed that the Hcy levels first decreased, then increased, and significantly increased after 50 years of age¹⁷, similar to the results of our study. However, some previous studies did not observe a statistically significant association between Hcy levels and age¹⁸, whereas other studies reported that plasma Hcy levels increased significantly with age in the general population and patients with schizophrenia^{19; 20}. The inconsistent results for this association are unclear. One possible reason may be due to the differences in subjects who had suffered from different diseases, BMIs, regions of residence, races, dietary habits, genetic backgrounds or courses of illness²¹⁻²⁴; for example, the subjects in the present study were inpatients, and most had suffered from schizophrenia for a long time.

Regarding cohort studies, some recent studies in humans displayed that an increase in the Hcy concentration was negatively associated with the evolution of cognitive deficits in the general population and patients with Alzheimer's disease and schizophrenia, particularly in studies of elderly individuals²⁵⁻²⁷. Interestingly, we found that Hcy levels were negatively associated with SoP score, but not score for other cognitive domains, in patients with schizophrenia aged 18-29 years after adjustment for age, BMI, and education. Additionally, with increasing age, no relationship was observed between these factors, in contrast most previous studies²⁸⁻³⁰. The inconsistent results may be because plasma Hcy levels result from a complex process mediated by the interaction between genetic and environmental factors, and the independent factors associated with Hcy levels and cognitive deficits differed according to age³¹. Moreover, the negative effects of various risk factors on cognitive deficits increase with aging¹⁶. Although the mechanism remains unclear, these findings indicated that age-related changes in cognitive deficits in patients with chronic schizophrenia are subtle and not uniform across multiple cognitive domains. These age-related differences must be considered when applying strategies to prevent cognitive deficits.

Limitations

Several limitations of the present study should be mentioned. One significant drawback of our study is that it was a cross-sectional study and was unable to determine the direct causal relationship between cognitive function and Hcy levels in patients with schizophrenia across age groups. Second, all patients in this study were diagnosed with chronic schizophrenia, limiting the generalizability in other disease states, including first-episode and drug-naive schizophrenia. Furthermore, Hcy levels and cognitive deficits are related to many confounding factors, such as dietary patterns and lifestyle. Unfortunately, we did not collect these data in our current study, and further investigation is warranted.

Conclusions

Based on our results, patients with schizophrenia exhibited cognitive deficits and Hcy levels were more closely related to cognition in young patients than in other subgroups. Therefore, researchers should focus on the complex age-Hcy interaction, and further research is warranted to investigate the mechanisms underlying the effect of Hcy levels on cognitive deficits in different age groups.

Declarations

Conflict of interest

None declared.

Compliance with ethical standards

All clinical studies described in the manuscript were carried out in accordance with Declaration of Helsinki promulgated by the National Institute of Health and all methods were carried out in accordance with relevant guidelines and regulations.

Funding: This research was supported by the National Natural Science Foundation of China (Grant No. 31771074), the Science and Technology Plan Project of Guangdong Province (Grant No. 2019B030316001), the Science and Technology Plan Project of Guangzhou (Grant Nos. 201804010259, 201807010064, 201704020168, and 201804010259), and the Science and Technology Plan Project of Liwan District, Guangzhou (Grant No. 201804011).

References

1. Marder, S. R. & Cannon, T. D. Schizophrenia. *N Engl J Med.* **381**, 1753-1761 (2019).
2. Huang, X. et al. Increased Plasma Kynurenic Acid Levels are Associated with Impaired Attention/Vigilance and Social Cognition in Patients with Schizophrenia. *Neuropsychiatr Dis Treat.* **16**, 263-271 (2020).
3. Jiang, B. et al. Effects of Differences in Serum Total Homocysteine, Folate, and Vitamin B12 On Cognitive Impairment In Stroke Patients. *BMC Neurol.* **14**, 217 (2014).
4. Bunevicius, A., Kazlauskas, H., Raskauskiene, N., Janusonis, V. & Bunevicius, R. Ischemic Stroke Functional Outcomes are Independently Associated with C-reactive Protein Concentrations and Cognitive Outcomes with Triiodothyronine Concentrations: A Pilot Study. *Endocrine.* **45**, 213-220 (2014).
5. Balcioglu, Y. H. & Kirlioglu, S. S. C-Reactive Protein/Albumin and Neutrophil/Albumin Ratios as Novel Inflammatory Markers in Patients With Schizophrenia. *Psychiatry Investig.* **17**, 902-910 (2020).
6. Chen, P. H. et al. Homocysteine, Rather than Age of Onset, is a Better Predictor for Cognitive Function in Older Adults with Bipolar Disorder. *Int J Geriatr Psychiatry.* **34**, 1473-1480 (2019).

7. Setién-Suero, E., Suárez-Pinilla, M., Suárez-Pinilla, P., Crespo-Facorro, B. & Ayesa-Arriola, R. Homocysteine and Cognition: A Systematic Review of 111 Studies. *Neurosci Biobehav Rev.* **69**, 280-298 (2016).
8. Di Lorenzo, R. et al. Homocysteine Levels in Schizophrenia Patients Newly Admitted to an Acute Psychiatric Ward. *Acta Neuropsychiatr.* **27**, 336-344 (2015).
9. Ostrakhovitch, E. A. & Tabibzadeh, S. Homocysteine and Age-Associated Disorders. *Ageing Res. Rev.* **49**, 144-164 (2019).
10. Wang, F. et al. The Relationship Between Plasma Homocysteine Levels and MTHFR Gene Variation, Age, and Sex in Northeast China. *Niger. J. Clin. Pract.* **22**, 380-385 (2019).
11. Regland, B., Johansson, B. V., Grenfeldt, B., Hjelmgren, L. T. & Medhus, M. Homocysteinemia is a Common Feature of Schizophrenia. *J Neural Transm Gen Sect.* **100**, 165-169 (1995).
12. García-Miss, M. R. et al. Folate, Homocysteine, Interleukin-6, and Tumor Necrosis Factor Alfa Levels, but Not the Methylenetetrahydrofolate Reductase C677T Polymorphism, are Risk Factors for Schizophrenia. *J. Psychiatr. Res.* **44**, 441-446 (2010).
13. Haidemenos, A. et al. Plasma Homocysteine, Folate and B12 in Chronic Schizophrenia. *Prog Neuropsychopharmacol Biol Psychiatry.* **31**, 1289-1296 (2007).
14. Bagney, A. et al. Negative Symptoms and Executive Function in Schizophrenia: Does their Relationship Change with Illness Duration? *Psychopathology.* **46**, 241-248 (2013).
15. Vinogradov, S., Fisher, M. & Nagarajan, S. Cognitive Training in Schizophrenia: Golden Age Or Wild West? *Biol Psychiatry.* **73**, 935-937 (2013).
16. Mosio Ek, A., Gierus, J., Koweszko, T. & Szulc, A. Cognitive Impairment in Schizophrenia Across Age Groups: A Case-Control Study. *BMC Psychiatry.* **16**, 37 (2016).
17. Xu, R. et al. Gender- and Age-Related Differences in Homocysteine Concentration: A Cross-Sectional Study of the General Population of China. *Sci Rep.* **10**, 17401 (2020).
18. Omer Sultan, M. et al. Correlation of Homocysteine Level and Age in Patients with Ischemic Stroke. *Cureus.* **12**, e7785 (2020).
19. Wang, F. et al. The Relationship Between Plasma Homocysteine Levels and MTHFR Gene Variation, Age, and Sex in Northeast China. *Niger. J. Clin. Pract.* **22**, 380-385 (2019).
20. Di Lorenzo, R. et al. Homocysteine Levels in Schizophrenia Patients Newly Admitted to an Acute Psychiatric Ward. *Acta Neuropsychiatr.* **27**, 336-344 (2015).
21. Huang, Y. et al. Homocysteine Level, Body Mass Index and Clinical Correlates in Chinese Han Patients with Schizophrenia. *Sci Rep.* **10**, 16119 (2020).
22. Yazici, E. et al. Klotho, Vitamin D and Homocysteine Levels During Acute Episode and Remission Periods in Schizophrenia Patients. *Nord J Psychiatry.* **73**, 178-184 (2019).
23. Numata, S. et al. Evaluation of an Association Between Plasma Total Homocysteine and Schizophrenia by a Mendelian Randomization Analysis. *BMC Med. Genet.* **16**, 54 (2015).

24. Moustafa, A. A., Hewedi, D. H., Eissa, A. M., Frydecka, D. & Misiak, B. Homocysteine Levels in Schizophrenia and Affective Disorders-Focus On Cognition. *Front. Behav. Neurosci.* **8**, 343 (2014).
25. Ayesa-Arriola, R. et al. Homocysteine and Cognition in First-Episode Psychosis Patients. *Eur Arch Psychiatry Clin Neurosci.* **262**, 557-564 (2012).
26. Agrawal, A. et al. Age Dependent Levels of Plasma Homocysteine and Cognitive Performance. *Behav. Brain Res.* **283**, 139-144 (2015).
27. Setién-Suero, E., Suárez-Pinilla, M., Suárez-Pinilla, P., Crespo-Facorro, B. & Ayesa-Arriola, R. Homocysteine and Cognition: A Systematic Review of 111 Studies. *Neurosci Biobehav Rev.* **69**, 280-298 (2016).
28. Trze Niowska-Druka A, B. et al. Evaluation of Hyperhomocysteinemia Prevalence and its Influence On the Selected Cognitive Functions I N Patients with Schizophrenia. *Prog Neuropsychopharmacol Biol Psychiatry.* **95**, 109679 (2019).
29. Gonzalez-Liencre, C. et al. Oxidative Stress in Schizophrenia: A Case-Control Study On the Effects On Social Cognition and Neuroc Ognition. *BMC Psychiatry.* **14**, 268 (2014).
30. Xu, R. et al. Gender- and Age-Related Differences in Homocysteine Concentration: A Cross-Sectional Study of the Gen Eral Population of China. *Sci Rep.* **10**, 17401 (2020).
31. Legdeur, N. et al. Age Dependency of Risk Factors for Cognitive Decline. *BMC Geriatr.* **18**, 187 (2018).

Tables

Table 1. Sociodemographic data and clinical profiles of patients with schizophrenia and healthy controls.

Variables	schizophrenia n=103	healthy controls n=122	F/X ²	<i>p</i>
Male	56 (54.4%)	58 (47.9%)	0.92	0.337
Age, years	42.89 (11.83)	35.02 (12.87)	12.87	<0.001
BMI, kg/m ²	24.56 (4.35)	22.44 (3.44)	3.44	<0.001
Education, years	10.89 (3.21)	14.83 (3.05)	3.05	<0.001
Hcy, mmol/L	16.28 (10.66)	9.53 (4.63)	4.63	<0.001
SoP score	33.36 (14.80)	49.85 (9.15)	9.15	<0.001
WM score	37.64 (13.44)	47.97 (9.75)	9.75	<0.001
Vrbl Lrng score	33.77 (12.34)	44.27 (10.23)	10.23	<0.001
Vis Lrng score	36.14 (11.88)	48.07 (10.35)	10.35	<0.001
AV score	37.73 (10.38)	48.67 (10.30)	10.30	<0.001

Abbreviations: Hcy=homocysteine, SoP=speed of processing, WM=working memory, Vrbl Lrng=verbal learning and memory, Vis Lrng=visual learning and memory, AV=attention/vigilance.

Table 2. The correlations between the plasma Hcy levels and clinical symptoms or cognitive deficits in patients with schizophrenia across age groups.

Figures

Group	SoP score	WM score	Vrbl Lrng score	Vis Lrng score	AV score	P subscore	N subscore	G subscore	Total PANSS score
18-29 y	r -0.60 p 0.011	-0.62 0.008	-0.09 0.722	-0.50 0.039	0.44 0.08	0.03 0.913	0.14 0.592	0.14 0.595	0.15 0.576
30-39 y	r -0.04 p 0.869	-0.11 0.662	-0.04 0.864	-0.25 0.301	-0.06 0.819	0.22 0.366	0.33 0.168	0.30 0.216	0.40 0.094
40-49 y	r -0.14 p 0.440	0.05 0.779	-0.48 0.005	-0.30 0.099	-0.19 0.299	0.33 0.070	-0.01 0.950	0.16 0.375	0.13 0.48
50-59 y	r -0.06 p 0.726	0.07 0.694	-0.12 0.498	-0.11 0.534	-0.19 0.276	0.01 0.964	-0.17 0.345	-0.17 0.327	-0.17 0.324

Abbreviations: Hcy=homocysteine, SoP=speed of processing, Vrbl Lrng=verbal learning and memory, Vis Lrng=visual learning and memory, WM=working memory, AV=attention/vigilance, P=positive symptom, N=negative symptom, G=general psychopathology syndrome, r=Pearson's correlation coefficient.




Figure 1

Differences in the plasma Hcy levels between patients with schizophrenia and healthy controls across age groups. *Significant differences in Hcy levels was observed between patients with schizophrenia and healthy controls ($p < 0.05$). Abbreviations: Hcy=homocysteine.




Figure 2

Differences in cognitive test score between patients with schizophrenia and healthy controls across age groups. *Significant differences in cognitive function score was observed between patients with schizophrenia and healthy controls ($p < 0.05$). Abbreviations: SoP=speed of processing, WM=working memory, VrbLrng=verbal learning and memory, Vis Lrng=visual learning and memory, AV=attention/vigilance.