Using the Medical Research Council Framework for the development of a communication partner training intervention for people with primary progressive aphasia (PPA): Better Conversations with PPA

Anna Volkmer (✉ a.volkmer.15@ucl.ac.uk)

University College London

Aimee Spector

University College London

Kate Swinburn

University College London

Jason D. Warren

University College London

Suzanne Beeke

University College London

Research Article

Keywords: Primary Progressive Aphasia, Speech and Language Therapy, Intervention, conversation, co-production, consensus

Posted Date: February 24th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-228838/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background: Primary Progressive Aphasia (PPA) is a language-led dementia resulting in a gradual dissolution of language. PPA has a significant psychosocial impact on both the person and their families. Speech and language therapy (SLT) is one of the only available management options, and communication partner training (CPT) interventions offer a practical approach to identify strategies to support conversation. The aim of this study was to define and refine a manual and an online training resource for speech and language therapists to deliver CPT to people with PPA and their communication partners (CPs) called Better Conversations with PPA (BCPPA).

Methods: The BCPPA manual and training program were developed using the Medical Research Council framework for developing complex interventions. The six-stage development process included 1. Review of existing literature, 2. Consultation and co-production 3. Development of an initial draft 4. Review by a group of speech and language therapists, using consensus methods 5. Two focus groups to gather opinions of people with PPA and their families 6. Refinement.

Results: The BCPPA manual and training program comprises seven modules, and four CPT intervention sessions. Eight important components of CPT were identified in the consensus work undertaken with speech and language therapists, including use of video feedback to focus on strengths as well as areas of conversation breakdown. Three themes arose from the focus groups with people with PPA and their families 1) Timing of intervention, 2) speech and language therapists' understanding of types of dementia, and 3) Knowing what helps. Refinements to the manual included additional practice activities and useful strategies for the future.

Conclusions: Using the MRC framework to develop an intervention that is underpinned by a theoretical rationale based on a review of the literature, increases the likelihood of behaviour change. Co-development of the manual and training materials ensures the intervention will meet the needs of people with PPA and their CPs. Gathering further data from speech and language therapists and people living with PPA and their families to refine the manual and the training materials enhances the feasibility of delivering this in preparation for a phase II NHS-based randomised controlled pilot-feasibility study, currently underway.

Background

The number of people living with dementia worldwide continues to rise, estimated at around 50 million at present with nearly 10 million new cases each year¹. Of these, perhaps a half a million people worldwide and several thousand in the United Kingdom have primary progressive aphasia (PPA): a group of language-led dementias associated with Frontotemporal Dementia and Alzheimer's disease². PPA presents as an insidious dissolution of language skills with relative sparing of other cognitive functions². At present there are three internationally recognised PPA variants; people with semantic variant (svPPA) experience a gradual loss of word meanings affecting both comprehension and naming, people with

logopenic variant PPA (lvPPA) present with difficulties in word retrieval and processing of complex sentences, and people with non-fluent agrammatic variant PPA (nfvPPA) demonstrate effortful, distorted articulation of speech sounds (apraxia) and/or an agrammatism³. Each variant presents with a distinct neuroanatomical distribution of atrophy and underlying neuropathology^{2,3}. Though it constitutes only a small proportion of the total dementia burden, PPA is of disproportionate clinical importance because it tends to strike people in older midlife with devastating impact on occupational and social functioning and because it presents a number of unique challenges not well met by conventional models of aphasia and dementia management.

People with PPA report increasing social isolation and reduced confidence as a result of their worsening communication difficulties⁴. More than one third of people with PPA experience depression and symptoms of anxiety are not uncommon. These likely impact directly on reports of reduced quality of life amongst people with PPA⁵. Spouses of people with PPA report a long trajectory of change, even prior to diagnosis. This results in feelings of loss of relationship and meaningful social interaction, increasing dependency of their spouse with PPA on them for communication, and overwhelming responsibility⁶.

The research literature on speech and language treatment approaches for people with PPA is developing. The majority of research has focused on impairment-focused interventions that aim to maintain or improve the person's ability to use words^{7,8}. Many people with PPA disengage from such naming therapies due to the frustration of practising individual words they will inevitably lose as the disease progresses⁹. More recently there has been a growing focus on functional communication interventions for PPA, which aim to support a person to execute an activity or participate in a life situation¹⁰. A systematic review of these diverse interventions identified two key shared components; building on existing strategies, and practising strategies with a communication partner¹⁰.

Despite barriers to therapy access, such as a lack of awareness of the role of the speech and language therapist in PPA, and restrictive service criteria, the number of people with this condition being referred to SLT is increasing¹¹. In contrast to a research focus on naming therapies, in clinical practice speech and language therapists prioritise communication partner training (CPT) interventions for people with PPA and their communication partners (CPs)^{11,12}.

CPT interventions for stroke and dementia have arisen from studies of conversation between people with communication disorders and their CPs. This research demonstrates that both people with dementia and aphasia draw on areas of retained strength, such as gesture, to maintain interactional flow^{13,14,15}. Some CPs are seen to facilitate conversational interaction, for example through giving time, but can equally expose their partners' difficulties by using barrier behaviours, for example, test questions (to which they already know the answer, a pedagogic behaviour used with children). CPT interventions aim to change conversation behaviours, enhancing conversational skill and confidence, and reducing barriers to facilitate the flow of natural conversation¹⁶. CPT interventions result in improved quality of life and wellbeing for people with dementia, and improved competency in their CPs¹⁷.

Many speech and language therapists report delivering CPT to people with PPA and describe using resources developed for stroke aphasia or brain injury related communication difficulties¹². CPT has a growing evidence base in stroke aphasia^{16,18} and delivers positive changes in the conversation skills of people with aphasia as well as their CPs^{19,20}. However, CPT approaches in stroke aphasia are not designed to meet the needs of people with progressive communication difficulties. Currently there are only case study reports of CPT for people with PPA^{21,22}. There is some suggestion of increased communicative effectiveness as a result, however, it is difficult to attribute these gains to CPT due to the fact that individuals were concurrently participating in additional interventions. Thus, there is a clinical need to develop a CPT intervention designed to meet the needs of people with PPA and their families^{6,23,24}.

To our knowledge there has been no specific research undertaken asking people with PPA and their families what interventions are important or need to be developed. People with PPA have written about their general experiences of SLT and the value of developing “a wide range of personalized strategies that continually evolve as the disease progresses”²⁵. Spouses report a need to develop practical approaches to deal with communication difficulties and maintain a close bond with their loved ones⁶. These issues are more likely to be met by tailored interventions, that build capacity by helping them to adjust and reframe their communication over time⁶. Speech and language therapists themselves have identified a need to engage family who are motivated to understand how they can best support their loved ones²⁶. Therefore, gathering ideas and contributions of people living with PPA, often described as Public Involvement (PI), is important to ensuring an intervention will meet their needs. PI is defined by the UK Standards for PI as research that is carried out with members of the public rather than to them. These standards include ensuring that people are involved as early as possible and that participation is made accessible. The aim of this study was to work with people with PPA and their families, from the beginning, to co-produce a CPT intervention to meet their needs.

The Medical Research Council provide a framework for developing and evaluating complex interventions²⁷. The guidance outlines the importance of preliminary development and testing of an intervention’s procedures prior to piloting and evaluation. This paper therefore describes how the MRC framework was used to develop Better Conversations with PPA (BCPPA), a 4-session, manualised, CPT intervention to help people with PPA and their CPs to identify and practice strategies to reduce barriers (such as interjecting when a person may not have finished) and increase facilitators in conversations (such as giving more time). A manual and an online training resource for speech and language therapists, hosted on a life-learning platform at UCL, were developed to enable speech and language therapists to deliver the intervention. In line with stages 1 and 2 of the MRC Framework the underlying theory and proposed mechanisms of change for the BCPPA program will be described as well as primary research which informed the co-production of the manual and online training resource.

Aim

To use the MRC framework for developing complex interventions to define and refine a manual and an online training resource for speech and language therapists to deliver BCPPA to people with PPA and their CPs.

Methods

Intervention development activities were based on phases one and two in the MRC framework for development of complex interventions²⁷. This comprised six stages including 1. examination of existing literature, 2. consultation and co-production work, 3. development of an initial draft, 4. consensus work with speech and language therapists, 5. focus groups with people with PPA and their families, 6. Refinement of the BCPPA intervention and manual in preparation for the randomised controlled pilot-feasibility study. Figure 1 demonstrates how these activities map onto the MRC guidance. Intervention development also followed the GUIDED guidelines for reporting for intervention development studies²⁸. Further patient and public involvement work undertaken to finalise outstanding training modules identified as supplementary to the RCT will not be discussed here. The first author, A.V., an experienced speech and language therapist, led all stages.

Recruitment

Consultation and co-production work (Stage 2): An opportunistic sample of people with PPA and their families, specialist speech and language therapists and neuropsychologists were invited to join the project steering group. A.V. emailed people who were known to her through clinical work, and asked the facilitator of the PPA branch of the Rare Dementias Support Group based at UCL (<https://www.raredementiasupport.org>) to forward an invitation email to individuals in the support group, inviting them to participate.

Consensus work (Nominal Group Technique, NGT) with speech and language therapists (Stage 4): speech and language therapists were recruited to participate in the NGT consensus study through the RCSLT Dementia and Mental Health Clinical Excellence Network, of which A.V. was a committee member. The NGT was carried out at one of the Clinical Excellence Network meetings and advertised in the RCSLT clinical practice magazine (Bulletin) and via emails circulated to members two months prior to the study day taking place.

Focus groups with people with PPA and their families (Stage 5): People with PPA and their families who attend the PPA branch of the Rare Dementias Support Group at UCL were invited to participate in one of two focus group meetings held at an accessible venue on the university campus. The aim was to recruit eight people to each focus group, totaling 16 participants. To optimize opportunities for individuals with communication difficulties to contribute to discussion²⁹, group numbers were capped at eight participants. Potential participants who responded to the advert were contacted by A.V. on the telephone to judge if they met the inclusion criteria of a) a diagnosis or possible diagnosis of PPA/relative with such a diagnosis, b) the ability to communicate to participate in a focus group c) see and hear well enough to

participate d) English as their language of daily use. Potential participants were excluded if they had a) a history of brain lesion or major head trauma, b) major physical illness or disability which could impact on participation. criteria required.

Examination of existing literature (Stage 1)

Existing models of dementia, principles of applied Conversation Analysis (CA), behaviour change theory and frameworks for chronic disease self- management were explored. This informed the preliminary contents and focus of the intervention.

Consultation and co-production work (Stage 2)

Four people with PPA and their spouses, two expert speech and language therapists, a neuropsychologist and the group facilitator (A.V.) took part in 12 formal BCPPA public involvement (PI) steering group meetings. Work to co-produce the BCPPA intervention materials and training modules was informed by feedback from people with PPA who had previously received CPT³⁰, research undertaken by A.V.^{10,11,12} and research into the BCA program for people with stroke aphasia³¹. In order to support communication, steering group members were informed of the topic for discussion in advance of each meeting and invited to contribute in advance, during or after meetings using verbal, written or visual means, e.g. bringing photos, drawing pictures or re-assembling draft materials.

First draft of the manual (Stage 3)

A draft of the BCPPA manual was developed using PowerPoint software. In order to upload these to the UCLeXtend website an online software package called Articulate was used to adapt the PowerPoint slides to an appropriate format. The work was undertaken with assistance from speech and language therapist researchers and four postgraduate researchers in speech and language sciences who were paid for their time.

Consensus work (Nominal Group Technique) with speech and language therapists (Stage 4)

Draft one of the manual was made available to a group of speech and language therapists, who were attending a Clinical Excellence Network study day. In order to gain an understanding of the clinical experiences and reality of speech and language therapists a qualitative research method was identified as appropriate. Speech and language therapists were encouraged to review the resource and pilot it with their clients. To ensure the BCPPA intervention reflected a consensus view of the most important components to include in a CPT intervention for people with PPA and their families a Nominal Group Technique (NGT) method was chosen. Given that many of the speech and language therapists participating in the study day had pre-existing professional relationships that could result in certain voices being represented over others in discussions, the NGT method was also chosen to provide opportunities to consider ideas and experiences equally, yet allowing for clarification and discussion prior to rating³².

Six weeks prior to attending the study day speech and language therapists were sent an email inviting them to anonymously complete a 12-item feedback survey comprising all open questions (supplementary document 1), hosted online on the Google Forms platform. Survey questions were developed by A.V. in consultation with the steering group and included questions about speech and language therapists' experiences and views on the content and format of the manual.

The NGT meeting itself comprised a two-stage ranking process commencing with a 90-minute group session (stage one), followed by email consultation (stage two). Meeting facilitators (AV and SB) agreed the session plan and central question for discussion in advance (see supplementary document 2), in line with guidelines for conducting NGT meetings¹⁴. At stage two, results of the group session were circulated via email to all participants, providing information on scores and mean rankings for each item. As per guidelines for conducting NGT meetings³², items describing the same ideas from the two groups were merged, following discussion and agreement between A.V. and S.B. Participants were asked to reply via email identifying and ranking their top eight items from this list (by placing a number from 1-8 to reflect which is most important - 8 and least important - 1). Following NGT guidelines³², scores were tallied and mean rankings calculated to identify the top eight ranked items overall.

Focus groups with people with PPA and their families (Stage 5)

Two focus groups took place, to provide people with PPA and their families the choice of attending with or without partners. Discussion was guided by the question 'How can speech and language therapists support people with PPA to live well and maintain relationships?'. The focus groups were jointly facilitated by A.V., alongside volunteer student speech and language therapists from UCL (one per focus group). A topic guide was co-produced with the BCPPA steering group and attendees of the PPA branch of the Rare Dementia Support Group at UCL (see supplementary document 3).

Focus group discussions were video recorded and transcribed by UCL student speech and language therapists (using transcription guidance³³). Given the researchers objectives to understand the lived experiences of people with PPA and their families, and gather opinions from them, qualitative methods employing a realist approach to reflexive thematic analysis was undertaken^{34,35}. Initial codes were generated by systematically coding interesting features (phase 2), collating these into potential themes (phase 3) and reviewing them in relation to the coded extracts (phase 4). Potential themes were refined to generate definitions and names (phase 5), further inspected to identify and report any additional key elements (phase 6). In addition, to improve reliability of analysis, four speech and language therapist researchers with experience of thematic analysis independently extracted data from a randomly selected section of transcript, discussed and reached agreement on the coding of themes arising from the data.

Refinement of the BCPPA manual (Stage 6)

The BCPPA manual was refined based on work undertaken in stage 4 and 5 (Figure 1).

Results

Examination of existing literature (Stage 1)

Existing literature comprising the bio-psychosocial model of dementia, applied CA, behaviour change theory and self-management and self-efficacy theory was examined.

Bio-psychosocial model of dementia

The bio-psychosocial model³⁶ proposes that there are factors other than the organic causes of dementia that influence the nature and speed of deterioration in daily functioning. These include some factors that are fixed, such as PPA variant, that cannot be changed. The BCPPA manual therefore provides practice tasks, to maximise generalisation for people with svPPA, for whom this is more difficult than those with nfvPPA (insert ref to cadario). Tractable factors, such as the way a CP interacts with a person with PPA, may be amenable to change and are directly targeted in the BCPPA intervention. Adaptive mechanisms used by the CP, such as multiple questions or test questions, may result in the person with PPA feeling incompetent³⁷. On the other hand, the use of gesture and enactment (whole body gesture and pantomime) by a person with PPA when they are having difficulty retrieving a spoken word³⁸ could be described as an effective coping strategy. The BCPPA intervention seeks to take account of fixed factors whilst targeting tractable factors to support the dyad (person with PPA and their CP) achieve their potential function.

Applied Conversation Analysis (CA)

CA is an approach to the study of human social interaction through the analysis of spontaneous, naturally occurring talk³⁹. A number of CA informed stroke aphasia intervention studies and clinical resources have been developed⁴⁰ such as SCA¹⁸, SPPARC⁴¹ and BCA⁴². These have in common the analysis of video recordings of natural conversations between the person with aphasia and their CP, and providing these as video feedback, as a foundation for targeting therapy⁴⁰. The speech and language therapist (who typically delivers such an intervention) analyses 10-15 minute video-recorded interaction to identify behaviours resulting in conversational breakdown, known as barriers, and ways in which members of a dyad successfully resolve or circumvent troubles to maintain interaction, known as facilitators. The aim of video feedback is to increase awareness in one or both members of the dyad of the impact of their behaviours, and jointly agree on goals for therapy. Once the goals of therapy are agreed upon, a process of practice, through supported conversations, role play and reflection, is commonly employed⁴⁰. The BCPPA intervention is informed⁴³ by this well-described⁴³, CA-underpinned approach to CPT.

Behaviour change theory

Recognising conversational barrier behaviours in video recordings of oneself and setting a goal to cease these, or adopt facilitative strategies instead, does not guarantee that a change in behaviour will occur⁴⁴.

Behaviour change theory, specifically the COM-B model⁴⁵ accounts for an individual's behaviour change as the product of three equally weighted components namely Capability, Opportunity and Motivation. Researchers examined video recordings of CA-underpinned CPT being delivered to people with stroke aphasia and their CPs⁴⁶ and used the COM-B model⁴⁵ to identify the essential change processes and the core procedures that serve them⁴⁷. The BCPPA intervention incorporates the seven core mechanisms that have been identified as essential to behaviour change in a CPT⁴⁶, specifically the processes to motivate change and those that embed changes (See supplementary material 4).

Self-management and self-efficacy

Central to self-management is the concept of the client as an active participant whose current status is influenced not only by diagnosis but by psychological responses and experiences. This implies interventions should address the ability to self-manage daily activities and the emotional journey, not just medical symptoms^{48,49}. Taking action to accomplish a plan to self-manage their condition is more likely to succeed if a person has the confidence or self-efficacy to achieve it⁵⁰. Self-efficacy is a mechanism that directs behaviour change, for if one feels in control of a behaviour it becomes easier to make a change to it⁵⁰. Five core self-management skills and four key self-efficacy mechanisms have been highlighted for inclusion in speech and language therapist interventions with people with progressive communication difficulties⁴⁹ and these have been considered in the development of the BCPPA intervention (see supplementary file 4).

Consultation and co-production work with the steering group (Stage 2)

Decisions made included:

1. Identification of seven subjects to form distinct training modules within the BCPPA program. Table 1 provides an overview of the learning objectives and how these were co-produced. The three modules required for the phase II NHS based randomised controlled pilot-feasibility study (Module 3: How to make a video, Module 4: What to target in therapy and Module 5: the BCPPA therapy) were prioritised for development over the four only needed for the future general release of the online BCPPA program. Table 2 provides an overview of the content of these three modules.
2. Development of a topic list, for Module 3: How to make a video, to support participants when making video recordings of their own conversations.
3. Distillation of the components of the eight BCA sessions into four 1- hour BCPPA sessions (the duration agreed-upon by speech and language therapists as feasible^{11,12})

Table 1. Learning objectives and timeline for development of the BCPPA training modules including the therapy program

BCPPA training modules	Learning objectives for speech and language therapists accessing the module	Co-produced module components	Development timeline
Module 1: What is PPA?	To explain what PPA is according to: People with PPA and their relatives who have worked on this module, Speech and language therapists working in the area The research literature in this area	Co- produced with steering group. References selected by steering group.	Prior to launch of online BCPPA program
Module 2: What is conversation training?	To explain what conversation training is to clients, based on video recorded interviews with: Speech and language therapists working in the area People with PPA and their relatives who have worked on this module.	Co- produced with steering group Video clips planned, filmed and selected by steering group	Prior to launch of online BCPPA program
Module 3: How to make a video	<ul style="list-style-type: none"> • To have an appropriate tool available to gain consent for the purpose of videoing of a couple in conversation with one another to be used in the conversation training intervention, BCPPA. • To be aware of the Mental Capacity Act (2005) and how this will impact on consent. • For speech and language therapists to be supported to make and store videos, in line with the data management guidance and policies of their local organisation, of conversation between a client and their conversation partner for the purpose of the BCPPA intervention. • To be able to set up an optimal environment for the purposes of making a video for the BCPPA intervention 	Co-produced work with the steering group included: a topic sheet to support participants in identifying what to discuss during video recording, example consent forms, video samples and formatting of module.	Prior to Phase II RCT Feasibility Pilot Study
Module 4: What to target in therapy	To understand the three stages of the goal setting process: 1) Identification of facilitators and barriers from pre-therapy videos 2) Selection of suitable video clips of appropriate length and focus to show clients, and 3) Negotiation of goals with a person with PPA and their conversation partner	Co- produced work with people with PPA included: video samples and formatting of module.	Prior to Phase II RCT Feasibility Pilot Study
Module 5: BCPPA therapy	To deliver the four synchronous BCPPA therapy sessions, supporting people with PPA and their communication partners (as a dyad) to: - Understand concept of barriers and facilitators in conversation and consider this briefly in relation to their own conversation - Identify barriers and facilitators in their own conversation - Set goals for therapy based on this discussion - Practice conversation using the strategies identified during goal setting	Co- produced work with steering group included: Therapy handouts for sessions 1 and 4, therapy activities for session 3, video samples and formatting of module.	Prior to Phase II RCT Feasibility Pilot Study

	<ul style="list-style-type: none"> - Problem solve any issues that have arisen in using identified strategies in conversations outside of therapy sessions - Consider planning for future changes in communication 		
Module 6: Measuring it	<ul style="list-style-type: none"> • To consider what options are available for measuring outcomes for BCPPA; • To think about the pros and cons of different outcome measures; • To consider how to use outcome measures in clinical practice. 	Co-produced with speech and language therapists working with people with PPA (local collaborators who participated in the Phase II RCT Feasibility Study)	Prior to launch of online BCPPA program
Module 7: Useful Resources	<ul style="list-style-type: none"> • To learn about some activities people with PPA enjoy; • To find out about some useful websites and resources; • To have thought about what has been useful in your therapy. 	Co-produced with steering group Online resources selected by steering group	Prior to launch of online BCPPA program

NB: The language used for module titles and learning objectives reflects vocabulary selected by the steering group during co-production and was felt appropriate and accessible for the target audience (clinical speech and language therapists).

Table 2. Overview of content for the first draft of the BCPPA manual (Modules 3, 4 & 5).

Module	Overview of content
<p data-bbox="87 134 500 161">Module 3: How to make a video</p> <div data-bbox="162 197 597 674"> <p data-bbox="186 275 570 289">The following diagram summarises the key steps that have discussed in this module.</p> <div data-bbox="201 310 558 428"> </div> <div data-bbox="201 470 570 569"> <p>Part 1: Develop a tool to gain consent for video recording clients and their conversation partners for conversation therapy.</p> <p>Part 2: Make and store a video of a client and their conversation partner in your work environment</p> <p>Part 3: Set up the optimal environment for the purposes of making a video for the BCPPA intervention.</p> </div> </div>	<p data-bbox="704 134 1289 161">Discussion of barriers to using consent forms</p> <p data-bbox="704 197 1344 224">Activity for user to identify local policy re consent</p> <p data-bbox="704 260 1487 287">Practical task to evaluate the accessibility of a consent form</p> <p data-bbox="704 323 1192 350">Provision of an example consent form</p> <p data-bbox="704 386 1523 449">Basic overview of key aspects of the MCA (2005) and the MCA Code of Practice (2007)</p> <p data-bbox="704 485 1507 548">Tips on creating accessible information and practical exercise to create a consent form</p> <p data-bbox="704 583 1463 646">Overview of functional capacity assessment from the MCA (2005) and case study of how to gain consent</p> <p data-bbox="704 682 1403 709">Common barriers to making videos in clinical practice</p> <p data-bbox="704 745 1159 772">Practical task to overcome barriers</p> <p data-bbox="704 808 1446 835">Tips on making a good quality video- using video samples</p> <p data-bbox="704 871 1523 934">Practical tasks on what to consider when making a good video- using video samples</p> <p data-bbox="704 970 1474 1033">Tips and ideas to choose the topic of conversation for video recording</p>
<p data-bbox="87 1031 558 1058">Module 4: What to target in therapy</p> <div data-bbox="162 1094 597 1499"> <p data-bbox="186 1100 576 1127">Summary: What to target in therapy</p> <div data-bbox="175 1157 586 1241"> </div> <div data-bbox="180 1251 558 1304"> <p>1. Video record 1-2 conversations of the person with aphasia and their partner.</p> <p>2. Watch the videos and identify the facilitators and barriers.</p> <p>3. Identify which facilitators and barriers to work on in therapy.</p> </div> <div data-bbox="175 1314 586 1377"> </div> <div data-bbox="186 1388 570 1419"> <p>4. Select the best example.</p> <p>5. Preparing to show a clip.</p> <p>6. Ready for agreeing goals</p> </div> </div>	<p data-bbox="704 1031 1333 1058">An overview of what facilitators and barriers are</p> <p data-bbox="704 1094 1523 1157">Examples of facilitators and barriers in conversations between a person with PPA and their partners</p> <p data-bbox="704 1192 1507 1276">Video examples of PwPPA and their CPs and the barriers and facilitators that may arise and practical exercise to identify these</p> <p data-bbox="704 1312 1523 1375">Tips on how to link barriers and facilitators to what to work on in therapy</p> <p data-bbox="704 1411 1490 1474">Practical tasks on selecting and presenting the video clips to show PwPPA and their families</p> <p data-bbox="704 1509 1203 1537">Things to consider when setting a goal</p> <p data-bbox="704 1572 1507 1635">Practical tasks using example goals from therapists who have shared real goals that they set for people in therapy</p>
<p data-bbox="87 1640 428 1667">Module 5: BCPPA therapy</p>	<p data-bbox="704 1640 1263 1667">Prompt / reminder to look at Modules 3 &4</p> <p data-bbox="704 1703 1511 1766">Sessions 1: provision of aims, sessions plan, therapy handouts and home-based tasks for PwPPA and CP</p> <p data-bbox="704 1801 1484 1864">Session 2: provision of aims, session plan, therapy handouts and home-based tasks for PwPPA and CP.</p> <p data-bbox="704 1900 1484 1963">Session 3: provision of aims, session plan, therapy handouts and home-based tasks for PwPPA and CP.</p>

Summary of BCPPA therapy.

The following diagram summarises the four sessions in BCPPA.

Session 1: What is conversation?	Session 2: Goal setting	Session 3: Practice the strategies identified during the goal setting session	Session 4: Problem solving and planning for future changes in communication
Introduce the therapy and conversation	View the dyads videos and set goals together		

Module Overview

1. Learning Objectives

2. Before you start

3. BCPPA Session 1

4. BCPPA Session 2

5. BCPPA Session 3

6. BCPPA Session 4

Module Summary

Session 4: provision of aims, session plan, therapy handouts and home-based tasks for PwPPA and CP.

First draft of BCPPA manual (Stage 3)

Module 5: the BCPPA therapy, hosted the BCPPA manual comprising session plans, session handouts and home-based tasks for each of the four BCPPA intervention sessions. The session plans identified intervention components as either core or non-essential components that can be tailored to an individual’s needs.

The draft manual was evaluated by the steering group to ensure information was presented in an accessible way. This included decisions on images and formatting.

The first draft of the manual was uploaded to a secure area on the UCLeXtend website and made available to speech and language therapists participating in the stage 4 consensus work via a bespoke URL. It was not publicly accessible.

Consensus work (Nominal Group Technique) with speech and language therapists (Stage 4)

Demographics and characteristics of speech and language therapist participants

Thirty-six speech and language therapists took part. Of these, 17 had completed the pre- NGT meeting survey, 22 had viewed the first draft of the BCPPA manual and training program prior to attending, and two had been able to use the BCPPA manual with a client with PPA. Table 3 presents speech and language therapist participant demographics and their familiarity with the BCPPA manual and training program. Following the meeting, 20 of the 36 participants completed the final NGT ranking task by email.

Table 3: Demographics of SLTs who participated in the NGT meeting and their familiarity with the BCPPA program

	SLT participants (n=36)
Gender (m:f)	2:34
Years practicing as an SLT (mean and range)	12.5 (0-21)
Number of clients with PPA seen in clinical career (mean and range)	9 (0-20)
BCPPA modules viewed online prior to meeting:	
None but knows of BCA	
None	1
Module 3 How to make a video Module	11
4 What to target in therapy	22
Module 5 BCPPA therapy	21
	22

m: male, f: female, SLT: speech and language therapist, PPA: primary progressive aphasia, BCA: Better Conversations with Aphasia program, BCPPA: Better Conversations with PPA program.

Pre-NGT meeting survey

When asked what surprised them when they first accessed the online BCPPA program five of 17 respondents (29%) commented on there being a lot of detail. Five respondents (29%) described the program as clear, easy to use and accessible; one person highlighted the comprehensive and detailed step by step guidance. A further four respondents (24%) stated that they were unsurprised by the BCPPA program, given their familiarity with the BCA program on which BCPPA is based. Respondents provided feedback on the BCPPA program including the most useful aspects (17, 100%, respondents), formatting (16, 94%, respondents), additions or changes (14, 82%, respondents) and the least useful aspects of the program (10, 60%, of respondents). Five themes arose from these data: 1. General usefulness; 2. Specific 'helpful' tasks or sections; 3. Access issues, 'I had trouble with'; 4. 'Could you add'; 5. 'Not a fan'. These

themes are illustrated with quotes in Figure 2. Notably, access issues were generally related to glitches in the program, though some local NHS browser systems posed restrictions.

Nominal Group Technique

After two iterations of consensus work with speech and language therapists, focused on the question “*What components of the BCPPA therapy sessions are important for people with PPA and their conversation partners?*”, eight components were identified, and ranked in order of importance, see Table 4.

Table 4: Final eight ranked components identified as important for the BCPPA program, from two stage NGT consensus work

1	Use of video feedback to identify facilitators versus barriers in conversation when focusing on people’s strengths as well as areas of potential breakdown
2	Tailored and person centred: <ul style="list-style-type: none">- goals,- conversational topics,- strategies- practice opportunities
3	Emphasising a focus on getting message across rather than a perfect interaction
4	Focusing individual attention on non-verbal communication strategies such as body language, gesture, facial expression and other methods of total communication.
5	Recognising and building on current communication strengths.
6	Working with both the person with PPA and the CP together.
7	Providing opportunities to practice strategies and get feedback from the SLT.
8	Providing an opportunity to discuss their communication difficulties

Focus groups with people with PPA and their families (Stage 5)

Demographics of participants

Thirteen participants, six people with PPA and seven family members, responded to the advertisement. All were eligible and agreed to participate but one couple withdrew the day before the focus group due to a conflicting commitment. The remaining 11 participants attended two focus groups (NB: these were mixed groups, whereby people with PPA and their CPs attended together, alongside some CPs and people with PPA who attended independently, group 1: seven participants; group 2: four participants). Participants

with PPA represented all three variants, and atypical mixed variants. Demographic information is outlined in Table 5.

Table 5: Demographic information for focus group participants

	Person with PPA (PwPPA) and communication partner (CP)	PPA variant	Time since symptom onset	Time since diagnosis
Focus Group 1:	PwPPA (m) + CP (f)	lvPPA	4 years,	2 years
	PwPPA (f) + CP (m)	Mixed	3 years	2 years
	CP (f)	(Mixed)	(9 years)	(4 years)
	PwPPA (f) + CP (m)	nfvPPA	5 years	4 years
Focus Group 2:	PwPPA (m)	lvPPA	4 years,	1 year
	PwPPA (f) + CP (m)	svPPA	5 years	4 years
	CP (m)	(Mixed)	(8 years)	(5 years)

PwPPA: person with primary progressive aphasia, CP: communication partner, lvPPA: logopenic variant primary progressive aphasia, svPPA: semantic variant primary progressive aphasia, nfvPPA: non-fluent agrammatic variant primary progressive aphasia

Themes arising from the focus groups

Three overarching themes emerged: 1) Timing of intervention, 2) speech and language therapists' understanding of types of dementia, and 3) Knowing what helps. Theme 3 encompassed five further subthemes: 'No one size fits all', 'I've discovered that', 'who's targeted', 'therapy approaches' and 'toolkit'. All themes and subthemes are presented in relation to illustrative units of data in Figure 3.

Refinement of BCPPA manual (Stage 6)

Refinements that arose from stages 4 and 5 of intervention development were decided on in conjunction with the steering group and are presented in Table 6. The refined BCPPA program was consequently made available to participating local speech and language therapist collaborators on UCLeXtend as part of their training in preparation for delivering the intervention during the randomised controlled pilot-feasibility study. The final intervention is described in detail, using the template for Intervention Description and Replication (TiDIER), in the authors PhD thesis which this paper is based on⁵¹, and a published protocol for study which remains currently underway⁵². Further to this, the project steering

group made plans to continue working to co-produce the remaining four modules, in anticipation of a future launch of the BCPPA program. This paper is based on work from the authors PhD thesis.

Table 6: Refinements for BCPPA manual and intervention

Decisions made	Examples of refinements made
Provide more options on strategies and practice activities in the intervention materials.	Addition of Home based task 2: Strategies to help turntaking and expansion of session plan 3 to include a list of 11 optional additional strategy practice ideas based on ideas collated from SLTs, people with PPA and their families and a review of manuals for stroke aphasia CPT manuals.
Provide more information on resources and other services.	Expansion of session plan 4 to include a list of resources and other services for SLTs making recommendations for the future.
Develop video examples of the intervention being delivered.	<p>Addition of video recordings of conversation breakdown and intervention being delivered inserted to Module 5: The BCPPA therapy. These included:</p> <p>Session 1:</p> <p>Video examples of Keith and Rose watching videos of themselves and the SLT facilitating them to identify barriers and facilitators.</p> <p>Session 2:</p> <p>Video examples of Keith and Rose goal setting with the SLT.</p> <p>Session 4:</p> <p>Video example of Keith and the SLT discussing a difficult subject around future planning.</p>
Include more testimonies from people with PPA in Module 1: What is PPA and Module 2: What is communication partner training?	Use of quotes to illustrate experience of communication facilitators and barriers in Module 4: What to target in therapy.
Provide more information on how PPA impacts on daily communication.	Refinement of Session 1. Handout 1. How does conversation work? And addition of Session 1. Handout 2. What can go wrong in conversations? in co- production with project steering group.
Provide a	Addition of summary handout for session 4:

summary sheet including suggestions for future changes on one handout at the end of the intervention.	Handout 6: Your strategies
---	----------------------------

BCPPA= Better Conversations with Primary Progressive Aphasia, SLT= speech and language therapist, PPA=primary progressive aphasia

Discussion

The BCPPA manual and training program were developed using the framework described in the MRC guidelines for development of complex interventions²⁷. The intervention content is underpinned by the bio-psychosocial model of dementia, applied CA, behaviour change theory, and self-management and self-efficacy literature. Consultation and co-production work with a project steering group made up of people with PPA and their family members provided the first draft of the BCPPA manual and training program. Consensus work using a Nominal Group Technique with practicing speech and language therapists, and focus groups with people with PPA and their families, identified further refinements. These included additions to the manual, and modifications to improve access to and use of the materials within the modules.

Speech and language therapists report seeing people with PPA in their clinics who feel incompetent in conversations, whilst their CPs feel helpless to support them in these situations⁵³. Addressing this by exploring meaningful strategies to maintain conversation via CPT that involves both a person with PPA and their CP has been recommended by expert speech and language therapists²⁶. Currently, speech and language therapists delivering CPT to people with PPA and their CPs report using tools designed for people with stroke aphasia because there are no PPA-specific materials^{11,12}. The BCPPA manual and training program address this gap in the speech and language therapists’ “toolkit” (described as such by participants in the focus groups) of interventions for PPA, and provides an evidence based, manualised training resource designed by and for people with PPA and their CPs.

Strengths And Limitations

Drawing on the best available evidence and appropriate theory to develop the BCPPA manual, in accordance with MRC guidance²⁷, should increase the likelihood that components of the intervention result in behaviour change. Extensive use of theory has been associated with larger effect sizes in a review of online behaviour change interventions⁵⁴. This work has involved new research with those targeted by the intervention as well as those delivering it.

There are, however, some methodological limitations. NGT does not allow for anonymisation in the way that other consensus methods such as Delphi do, and can thus bias the responses of participants.

Though, the NGT did provide a method of involving large participant numbers and incorporating mathematical voting techniques to aggregate group judgements equally³². Making the intervention manual available for scrutiny of its practicality for clinical practice in this way has allowed for prioritisation of key components in anticipation of the phase II NHS based randomised controlled pilot-feasibility study. Notably, only two males were recruited to the NGT, though this is generally representative of the current SLT community⁵⁵. Despite being a useful method for eliciting participant's genuine and honest opinions, a focus group can be a challenging communication environment⁵⁶. The role of the speech and language therapist facilitator and the student speech and language therapist co-facilitators was to mitigate this by enabling participants to contribute to discussion. The option to attend with CPs to support communication was also provided, but instead participants prioritised the convenience of meeting dates and times. Given the steering group was established a number of years prior to the recently published practice standards for PI⁵⁷ it is likely that the methods employed may have limited the effectiveness of the co-produced work. Some have criticised the steering group model for consulting with only a small number of individuals. There were only three couples with PPA in this group and that may have limited its value. PPA is, however, a relatively rare condition and people were approached to reflect the known diversity within the condition. Additionally, new members were sought when others withdrew due to disease progression, and the author sought to gather perspectives of other people and their families through individual telephone contact. Despite approaching professionals from other disciplines, including medicine and social work, interested individuals were not able to attend steering group meetings. The author was able to consult with the research team, including neurology colleagues, to gather feedback and ideas.

A manualised approach enables standardised delivery of the intervention for a future trial. Given that speech and language therapists in clinical practice may have limited experience of working with people with PPA^{10,11}, this helps to maximise ease and fidelity of delivery for future implementation. However, a manualised intervention may limit the potential to tailor an intervention to individual clients, for example by deciding not to use video recording or by delivering the intervention to a person accompanied by two CPs. Person-centred components have been identified as important for functional communication interventions for people with PPA, and have been highlighted as important for behaviour change^{46,49}. The development of this intervention took behaviour change theory into account and embedded the core processes and mechanisms that had been identified in previous CPT research as essential components. These were clearly signposted in the manual and distinguished from non-essential components that were amendable to tailoring. Furthermore, expecting four 1-hour therapy sessions to result in a change may seem ambitious. However, the decision on dosage was made based on the average number of sessions that speech and language therapists reported having available to deliver functional communication interventions for PPA¹¹. Developing an intervention that meets this requirement increases the chance of implementation.

Conclusions

The six-stage process of development included a review of existing literature, and consultation and co-production with the project steering group to develop an initial draft. Consensus work undertaken with speech and language therapists and focus groups with people with PPA and their families identified further refinements. The BCPPA manual was refined in preparation for a phase II NHS based randomised controlled pilot-feasibility study which is currently underway⁵².

Abbreviations

PPA: Primary Progressive Aphasia; svPPA: semantic variant PPA; lvPPA: Logopenic variant PPA; nfvPPA: non-fluent variant PPA; SLT: Speech and Language Therapy; CP: Communication Partner; CPT: Communication Partner Training; BCPPA: Better Conversations for Primary Progressive Aphasia; PI: Public Involvement; NGT: Nominal Group Technique; NHS: National Health Service; MRC: Medical Research Council; CA: Conversation Analysis.

Declarations

Ethics approval and consent to participant

All work undertaken in this study was conducted in accordance with the Declaration of Helsinki.

Consultation and co-production work (Stage 2): Ethical approval is not required when involving individuals in the planning or design of research, for example when they are members of a research steering or advisory group (Health Research Authority, 2019). In order to equalize participation and power issues specific strategies were used in and outside of the steering group meetings. All communication, written and spoken, was made accessible to ensure no individuals were disadvantaged. All steering group members were required to use the same methods to contribute (raising a card to indicate they had a question or comment). People with communication difficulties were invited to contribute before others, and the author made contact with individuals with communication difficulties prior to the meeting to gather initial thoughts, to support facilitation during group discussions.

Consensus work with speech and language therapists (Stage 4): The UCL Research Ethics Committee confirmed the NGT consensus work (Stage 4) with speech and language therapist participants to be service evaluation. Participants were informed that all responses would be anonymous and at the start of the event they provided written informed consent to participate.

Focus groups with people with PPA and their families (Stage 5): Camden and Kings Cross Research Ethics Committee (IRAS ID: 202353, Rec Ref: 17/LO/0357) approved the focus group study. Informed consent was obtained by A.V. following the current guidance from the Mental Capacity Act (Department of Health, 2005) and Royal College of Speech and Language Therapy (RCSLT), regarding gaining consent

from people with communication difficulties. A caregiver (a friend or relative) was asked to witness the informed consent process whenever possible.

Participant information sheets, consultee information sheets, consent forms and consultee declaration forms were designed to be accessible to support the process of gaining informed consent. They were designed using a resource for researchers in communication disability “Engaging people who have aphasia” (Pearl, 2014) and modified with advice from the project steering group. Transcriptions of focus group data were anonymised via the allocation to each participant of a unique research number, used at all times. All names, places and personal information mentioned in the discussions were pseudonymised.

The addition of video recordings demonstrating delivery of the intervention had ethical implications, requiring a minor amendment to HRA ethical approval. Having received this approval, a separate dyad were recruited through an email advert to members of the PPA branch of the UCL Rare Dementia Support Group. During the consent process it was made clear to the dyad there would be a risk that their faces and voices may be recognized from their video recordings. Information was provided regarding the course registration process and expected registrants, such as health professionals and people with PPA and their families. After consenting to participate, the dyad made four pre-intervention video recordings of their conversations, and received BCPPA therapy from the author, an experienced speech and language therapist. All four therapy sessions were video recorded. The author then identified a selection of short video clips that illustrated key components of the intervention such as the process of supporting dyads to identify barriers and facilitators in their conversation sample, goal setting, and discussion about planning for the future. The dyad viewed these clips prior to giving final consent for their inclusion in Module 5.

Consent for Publication

Not applicable

Availability of data and materials

Not applicable

Competing Interests

The authors declare that they have no competing interests

Funding

AV was funded by a National Institute for Health Research Doctoral Research Fellowship 2015-08-182 JDW receives grant support from the Alzheimer’s Society, Alzheimer’s Research UK and the NIHR UCLH Biomedical Research Centre

This paper presents independent research funded by the National Institute for Health Research (NIHR). The views expressed are those of the authors and not necessarily those of the NHS, the NIHR or the

Department of Health.

Components of work described in this paper were presented at the Academy of Aphasia 57th Annual Meeting, Macau, Macau, 27 Oct - 29 Oct, 2019 as a conference talk with interim findings. The abstract was published in "Aphasia in neurodegenerative conditions." in *Frontiers in Human Neurosciences*: Hyperlink with DOI: https://www.frontiersin.org/10.3389%2Fconf.fnhum.2019.01.00106/event_abstract

Authors' contributions

This paper is based on work from the first authors PhD thesis. AV conceived and designed the study, collected, analysed and interpreted data with supervision and support from SB, AS, KS and JW. AV drafted the article with guidance from SB, AS, KS and JW, all authors critically reviewed the article and are accountable for all aspects of the work.

Acknowledgments

Dr Abi Roper, Dr Claudia Bruns, Dr Gwen Breklemans and Anna Sowerbutts all contributed to the development of the electronic version of the BCPPA modules in preparation for the upload to the UCLeXtend website. Sara Bunker and Claire Tryfonidis were the SLT students, now qualified therapists, who aided in the facilitation of the focus groups with people with PPA and their families. Nicola Sirman and Firl Beckley acted as SLT researchers working on the thematic analysis of the focus group data.

References

1. World Health Organisation. Dementia factsheet. 2019. <https://www.who.int/news-room/factsheets/detail/dementia>. Accessed June 2020.
2. Marshall CR, Hardy CJ, Volkmer A, Russell LL, Bond RL, Fletcher PD, Clark CN, Mummery CJ, Schott JM, Rossor MN, Fox NC. Primary progressive aphasia: a clinical approach. *Journal of Neurology*. 2018;265(6):1474-90.

<https://doi.org/10.1007/s00415-018-8762-6>

3. Gorno-Tempini ML, Hillis AE, Weintraub S, Kertesz A, Mendez M, Cappa SF, Ogar JM, Rohrer JD, Black S, Boeve BF, Manes F. Classification of primary progressive aphasia and its variants. *Neurology*. 2011 Mar 15;76(11):1006-14. <https://doi.org/10.1212/WNL.0b013e31821103e6>
4. Gould Mike's lifeline. *RCSLT Bulletin*, 2015;760:17. Retrieved from https://www.rcslt.org/docs/bulletin/2015/aug_2015
5. Ruggero L, Nickels L, Croot K. Quality of life in primary progressive aphasia: What do we know and what can we do next?. *Asphasiology*. 2019;33(5):498-519.

<https://doi.org/10.1080/02687038.2019.1568135>

6. Pozzebon M, Douglas J, Ames D. Spousal recollections of early signs of primary progressive aphasia. *International journal of language & communication disorders*. 2018;53(2):282-93.
<https://doi.org/10.1111/1460-6984.12347>
7. Jokel R, Graham NL, Rochon E, Leonard C. Word retrieval therapies in primary progressive aphasia. *Aphasiology*. 2014;28(8-9):1038-68.
<https://doi.org/10.1080/02687038.2014.899306>
8. Carthery-Goulart MT, Silveira AD, Machado TH, Mansur LL, Parente MA, Senaha ML, Brucki SM, Nitrini R. Nonpharmacological interventions for cognitive impairments following primary progressive aphasia: a systematic review of the literature. *Dementia & Neuropsychologia*. 2013;7(1):122-31.
<https://doi.org/10.1590/S1980-57642013DN70100018>
9. Croot K. Progressive language impairments: Definitions, diagnoses, and prognoses. *Aphasiology*. 2009;23(2):302-26. <https://doi.org/10.1080/02687030801942981>
10. Volkmer A, Spector A, Meitanis V, Warren JD, Beeke S. Effects of functional communication interventions for people with primary progressive aphasia and their caregivers: a systematic review. *Aging & mental health*. 2020;24(9):1381-93.
Doi:10.1080/13607863.2019.1617246
11. Volkmer A, Spector A, Warren JD, Beeke S. Speech and language therapy for primary progressive aphasia: Referral patterns and barriers to service provision across the UK. *Dementia*. 2020;19(5):1349-63. Doi: 10.1177/1471301218797240
12. Volkmer A, Spector A, Warren JD, Beeke S. Speech and language therapy for primary progressive aphasia across the UK: A survey of current practice. *International journal of language & communication disorders*. 2019;54(6):914-26. Doi:10.1111/1460-6984.12495
13. Kindell J, Keady J, Sage K, Wilkinson R. Everyday conversation in dementia: a review of the literature to inform research and practice. *International journal of language & communication disorders*. 2017;52(4):392-406. <https://doi.org/10.1111/1460-6984.12298>
14. Perkins L, Whitworth A, Lesser R. Conversing in dementia: A conversation analytic approach. *Journal of Neurolinguistics*. 1998;11(1-2):33-53. [https://doi.org/10.1016/S0911-6044\(98\)00004-9](https://doi.org/10.1016/S0911-6044(98)00004-9)
15. Wilkinson R. Conversation and aphasia: Advances in analysis and intervention. 2015;29(3):257–268. <https://doi.org/10.1080/02687038.2014.974138>
16. Simmons-Mackie N, Savage MC, Worrall L. Conversation therapy for aphasia: a qualitative review of the literature. *International Journal of Language & Communication Disorders*. 2014;49(5):511-26. <https://doi.org/10.1111/1460-6984.12097>
17. Eggenberger E, Heimerl K, Bennett MI. Communication skills training in dementia care: a systematic review of effectiveness, training content, and didactic methods in different care settings. *Database of*

- Abstracts of Reviews of Effects (DARE): Quality-assessed Reviews [Internet]. 2013.
<https://doi.org/10.1017/S1041610212001664>
18. Kagan A. Supported conversation for adults with aphasia: Methods and resources for training conversation partners. *Aphasiology*. 1998;12(9):816-30.
<https://doi.org/10.1080/02687039808249575>
 19. Beeke S, Johnson F, Beckley F, Heilemann C, Edwards S, Maxim J, Best W. Enabling better conversations between a man with aphasia and his conversation partner: Incorporating writing into turn taking. *Research on Language and Social Interaction*. 2014;47(3):292-305.
<https://doi.org/10.1080/08351813.2014.925667>
 20. Best W, Maxim J, Heilemann C, Beckley F, Johnson F, Edwards SI, Howard D, Beeke S. Conversation therapy with people with aphasia and conversation partners using video feedback: a group and case series investigation of changes in interaction. *Frontiers in human neuroscience*. 2016;10:562.
<https://doi.org/10.3389/fnhum.2016.00562>
 21. Murray LL. Longitudinal treatment of primary progressive aphasia: A case study. *Aphasiology*. 1998;12(7-8):651-72. <https://doi.org/10.1080/02687039808249564>
 22. Wong SB, Anand R, Chapman SB, Rackley A, Zientz J. When nouns and verbs degrade: Facilitating communication in semantic dementia. *Aphasiology*. 2009;23(2):286-301.
<https://doi.org/10.1080/02687030801943112>
 23. Taylor-Rubin C, Croot K, Power E, Savage SA, Hodges JR, Togher L. Communication behaviors associated with successful conversation in semantic variant primary progressive aphasia. *International Psychogeriatrics*, 2017;10:619–1632. <https://doi.org/10.1017/S1041610217000813>
 24. Jokel R, Meltzer J. Group intervention for individuals with primary progressive aphasia and their spouses: Who comes first?. *Journal of communication disorders*. 2017;66:51-64.
<https://doi.org/10.1016/j.jcomdis.2017.04.002>
 25. Douglas JT. Adaptation to early-stage nonfluent/agrammatic variant primary progressive aphasia: A first-person account. *American Journal of Alzheimer's Disease & Other Dementias®*. 2014;29(4):289-92. <https://doi.org/10.1177%2F1533317514523669>
 26. Kindell J, Sage K, Cruice M. Supporting communication in semantic dementia: clinical consensus from expert practitioners. *Quality in ageing and older adults*. 2015;3:153–164.
<https://doi.org/10.1108/QAOA-08-2014-0016>
 27. Craig P, Dieppe P, Macintyre S, Michie S, Nazareth I, Petticrew M. Developing and evaluating complex interventions: the new Medical Research Council guidance. *Bmj*. 2008;29:337.
<https://doi.org/10.1136/bmj.a1937>
 28. Duncan E, O'Cathain A, Rousseau N, Croot L, Sworn K, Turner KM, Yardley L, Hoddinott P. Guidance for reporting intervention development studies in health research (GUIDED): an evidence-based consensus study. *BMJ open*. 2020;10(4):e033516. <http://dx.doi.org/10.1136/bmjopen-2019-033516>
 29. Barrett J, Kirk S. Running focus groups with elderly and disabled elderly participants. *Applied ergonomics*. 2000;31(6):621-9. [https://doi.org/10.1016/S0003-6870\(00\)00031-4](https://doi.org/10.1016/S0003-6870(00)00031-4)

30. Volkmer A, Beeke S. How to help couples have better conversations. *Journal of Dementia Care*. 2015;23(5):22-4.
31. Beckley F, Best W, Johnson F, Edwards S, Maxim J, Beeke S. Conversation therapy for agrammatism: Exploring the therapeutic process of engagement and learning by a person with aphasia. *International journal of language & communication disorders*. 2013;48(2):220-39.
<https://doi.org/10.1111/j.1460-6984.2012.00204.x>
32. Delbecq A L, Van de Ven A H, Gustafson D H. Group techniques for program planning : a guide to nominal group and Delphi processes (S. F. and Company., Ed.). Glenview:Illinois; 1975.
33. McLellan E, MacQueen KM, Neidig JL. Beyond the qualitative interview: Data preparation and transcription. *Field methods*. 2003 Feb;15(1):63-84. <https://doi.org/10.1177/1525822X02239573>
34. Braun V, Clarke V. Using thematic analysis in psychology. *Qualitative research in psychology*. 2006 Jan 1;3(2):77-101. <https://doi.org/10.1191/1478088706qp063oa>
35. Braun V, Clarke V. One size fits all? What counts as quality practice in (reflexive) thematic analysis?. *Qualitative research in psychology*. 2020 Aug 14:1-25.
<https://doi.org/10.1080/14780887.2020.1769238>
36. Spector A, Orrell M. Using a biopsychosocial model of dementia as a tool to guide clinical practice. *International Psychogeriatrics*. 2010 Sep 1;22(6):957. <https://doi.org/10.1017/S1041610210000840>
37. Kindell J, Keady J, Sage K, Wilkinson R. Everyday conversation in dementia: a review of the literature to inform research and practice. *International journal of language & communication disorders*. 2017 Jul;52(4):392-406. <https://doi.org/10.1111/1460-6984.12298>
38. Kindell J, Sage K, Keady J, Wilkinson R. Adapting to conversation with semantic dementia: using enactment as a compensatory strategy in everyday social interaction. *International Journal of Language & Communication Disorders*. 2013 Sep;48(5):497-507.

<https://doi.org/10.1111/1460-6984.12023>

39. Sidnell J, Stivers T, editors. *The handbook of conversation analysis*. John Wiley & Sons; 2012 Aug 3.
<https://doi.org/10.1002/9781118325001>
40. Wilkinson R. Intervening with conversation analysis in speech and language therapy: Improving aphasic conversation. *Research on Language and Social Interaction*. 2014 Jul 3;47(3):219-38.
<https://doi.org/10.1080/08351813.2014.925659>
41. Wilkinson R. Intervening with conversation analysis in speech and language therapy: Improving aphasic conversation. *Research on Language and Social Interaction*. 2014 Jul 3;47(3):219-38.
<https://doi.org/10.1080/08351813.2014.925659>
42. Lock S, Wilkinson R, Bryan K, Maxim J, Edmundson A, Bruce C, Moir D. Supporting partners of people with aphasia in relationships and conversation (SPPARC). *International Journal of Language & Communication Disorders*. 2001 Jan 1;36(sup1):25-30. DOI:[3109/13682820109177853](https://doi.org/10.1080/13682820109177853)
43. Beeke S, Sirman N, Beckley F, Maxim J, Edwards S, Swinburn K, Best W. Better Conversations with Aphasia: an e-learning resource. 2013 <https://extend.ucl.ac.uk/> Access June 2020

44. Cruice M, Blom Johansson M, Isaksen J, Horton S. Reporting interventions in communication partner training: A critical review and narrative synthesis of the literature. *Aphasiology*. 2018 Oct 3;32(10):1135-66. <https://doi.org/10.1080/02687038.2018.1482406>
45. Beckley F, Best W, Johnson F, Edwards S, Maxim J, Beeke S. Conversation therapy for agrammatism: Exploring the therapeutic process of engagement and learning by a person with aphasia. *International journal of language & communication disorders*. 2013 Mar;48(2):220-39.<https://doi.org/10.1111/j.1460-6984.2012.00204.x>
46. Michie S, Van Stralen MM, West R. The behaviour change wheel: a new method for characterising and designing behaviour change interventions. *Implementation science*. 2011 Dec;6(1):1-2. <https://doi.org/10.1186/1748-5908-6-42>
47. Johnson FM, Best W, Beckley FC, Maxim J, Beeke S. Identifying mechanisms of change in a conversation therapy for aphasia using behaviour change theory and qualitative methods. *International journal of language & communication disorders*. 2017 May;52(3):374-87. <https://doi.org/10.1111/1460-6984.12279>
48. Johnson F. What works in conversation therapy for aphasia, and how? Searching for mechanisms of change and active ingredients using tools and theory from behaviour change research. Unpublished PhD thesis: University College London. 2015
[https://discovery.ucl.ac.uk/id/eprint/1472090/1/Johnson_Oct 2015 SUBMISSION Fiona Johnson Phd THESIS.pdf](https://discovery.ucl.ac.uk/id/eprint/1472090/1/Johnson_Oct%202015%20SUBMISSION%20Fiona%20Johnson%20Phd%20THESIS.pdf) Accessed June 2020.
49. Turner A, Anderson JK, Wallace LM, Bourne C. An evaluation of a self-management program for patients with long-term conditions. *Patient education and counseling*. 2015 Feb 1;98(2):213-9. <https://doi.org/10.1016/j.pec.2014.08.022>
50. Yorkston K, Baylor C, Britton D. Incorporating the principles of self-management into treatment of dysarthria associated with Parkinson's disease. In *Seminars in speech and language* 2017 Jul (Vol. 38, No. 3, p. 210). NIH Public Access. <https://doi.org/10.1055/s-0037-1602840>
51. Volkmer A. Better Conversations with Primary Progressive Aphasia (BCPPA): Developing and piloting a communication partner training intervention for people with primary progressive aphasia and their significant others. Unpublished PhD thesis: University College London. 2020.
<https://discovery.ucl.ac.uk/id/eprint/10100280/> Accessed June 2020.
52. Volkmer A, Spector A, Warren JD, Beeke S. The 'Better Conversations with Primary Progressive Aphasia (BCPPA)' program for people with PPA (Primary Progressive Aphasia): protocol for a randomised controlled pilot study. *Pilot and feasibility studies*. 2018 Dec;4(1):1-0. <https://doi.org/10.1186/s40814-018-0349-6>
53. Taylor C, Croot K, Power E, Savage SA, Hodges JR, Togher L. Trouble and repair during conversations of people with primary progressive aphasia. *Aphasiology*. 2014 Sep 2;28(8-9):1069-91. <https://doi.org/10.1080/02687038.2014.930411>
54. Webb T, Joseph J, Yardley L, Michie S. Using the internet to promote health behavior change: a systematic review and meta-analysis of the impact of theoretical basis, use of behavior change

techniques, and mode of delivery on efficacy. Journal of medical Internet research. 2010;12(1):e4. <https://doi.org/10.2196/jmir.1376>

55. Litosseliti L, Leadbeater C. Speech and language therapy/pathology: perspectives on a gendered profession. International Journal of Language & Communication Disorders. 2013 Jan;48(1):90-101. <https://doi.org/10.1111/j.1460-6984.2012.00188.x>
56. Wilkinson S. Focus groups in health research: Exploring the meanings of health and illness. Journal of Health Psychology. 1998 Jul;3(3):329-48. <https://doi.org/10.1177/135910539800300304>
57. Staniszewska S, Denegri S, Matthews R, Minogue V. Reviewing progress in public involvement in NIHR research: developing and implementing a new vision for the future. BMJ open. 2018 Jul 1;8(7):e017124. <https://doi.org/10.1136/bmjopen-2017-017124>

Figures

Figure 1

The six stages in the development BCPPA intervention and manual drafting, mapped on to the MRC framework for development of complex interventions.

Figure 2

Themes identified from survey responses in Stage 4 consensus work.

Figure 3

Themes and subthemes arising from focus groups with people with PPA and their CPs

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupplementaryMaterials.docx](#)