

Chronic Anal Fissure in Female Patients: Is Fissurectomy and V-Y Anoplasty an Effective Procedure?

BEATRICE D'ORAZIO

University of Palermo

Girolamo Geraci

University of Palermo

Sebastiano Bonventre

University of Palermo

Dario Calì

University of Palermo

Gaetano Di Vita (✉ divitagaetano@libero.it)

University of Palermo

Research Article

Keywords: proctology, fissurectomy, anoplasty, anal fissure, female patients

Posted Date: February 24th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-228865/v1>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

INTRODUCTION: Lateral internal sphincterotomy (LIS) is still the approach of choice for the treatment of chronic anal fissure (CAF) regardless to the internal anal sphincter tone but it is burdened by high risk post-operative faecal incontinence (FI). In female patient there are some anatomical and functional differences of the sphincteric system which make them more at risk of FI and vaginal birth could cause sphincteric lesions affecting the anal continence function. The aim of our study is to evaluate the results of saving sphincter procedure as treatment for female patients affected by CAF.

METHODS: We studied 110 female patients affected by CAF undergone fissurectomy and anoplasty with V-Y cutaneous flap advancement associating pharmacological sphincterotomy in patients with hypertonic IAS. The follow up was at least for 2 years. The goals were patient's complete healing, the evaluation of FI, recurrence rate and manometry parameters.

RESULTS: All wounds healed within 40 days after surgery. We recorded 8 cases of recurrences 6 healed with medical therapy and 2 with dilatation. We recorded 2 "de novo" temporary and low grade post-operative cases of FI. Post-operative value of MRP were unmodified in patient with normotonic IAS but significantly lower at 12 months follow up as compared with the pre-operative ones in patients with hypertonic IAS; after 24 months from surgery MRP values were within the normal range.

CONCLUSION: The fissurectomy and anoplasty with V-Y cutaneous flap alone or in association with a pharmacological sphincterotomy in patients with hypertonic IAS may represent an effective approach for the treatment of CAF in female patients.

Introduction

Chronic anal fissure (CAF) is a frequently occurring proctology lesion, which presents itself as a painful superficial tear in the cutaneous- mucosal transition zone of the anal canal. First line therapy is medical treatment with a dietary-higenyc and pharmacological approach. Nevertheless, in the event of CAFs, which are refractory to dietary and pharmacological therapy, lateral internal sphincterotomy (LIS) is still suitable as the treatment of choice independently from IAS tone. In fact, this latter surgical procedure is characterised by a low rate of post-operative complications and a high healing rate. The greatest disadvantage of this latter procedure is represented by the high rate of faecal incontinence occurrence (FI), which accounts for up to 30-40%. A meta-analysis from 2013, which evaluated long-term incidence of FI after LIS, showed an overall continence alteration risk of 14%[1]; however, on severity analysis, flatus incontinence and soilage/seepage were much commoner than frank incontinence to liquid or solid stool [2]. FI has a strong impact on the quality of life of patients and it can be more disabling than CAF itself ; actually, patients tend to better bear a recurrence than FI [3]. From the guidelines for CAF treatment from different countries [4-9] we understand that the risk factors for FI occurrence are female gender, old age and the absence of an hypertonic IAS. In female patient there are some anatomical and functional [10] of the sphincteric system as compared to male patients, which make them more at risk of FI after

proctological procedures; moreover, with vaginal birth sphincteric lesions may occur, which could affect the anal continence function. To reduce the risk of post-operative FI, the mostly used surgical procedure is fissurectomy alone or in association with pharmacological sphincterotomy and/or cutaneous or mucosal flap advancement.

The aim of our study is to evaluate the results of fissurectomy and anoplasty with V-Y cutaneous flap advancement in female patients affected by CAF.

Materials And Methods

This study involves 110 female patients, all affected by idiopathic and non-recurrent CAF who underwent surgical procedure, from January 2010 to January 2019. Exclusion criteria for the study were: the presence of multiple fissures, fistulas in ano, syphilis, inflammatory bowel disease, anal abscess, malignant disease and previous anorectal surgery. All patients were followed up for at least 2 years after the surgical procedure. The patients' outcome data were retrieved from a prospectively monitored database.

We conduct this study in compliance with the principles of declaration of Helsinki, the protocol for this study has been submitted to the Ethical Committee of our institution, which did not consider necessary to approve it. Written informed consent was obtained from all the study participants.

Preoperative manometric evaluation was performed after a reasonable period of suspension of all medical therapy influencing IAS tone. The manometric evaluation was carried out by a manometric sensor using the station pull-through as described in our previous work [11]. A manometric evaluation has been undertaken at 12 and 24 months after surgery.

Data collected on healthy subjects by our anorectal pathophysiological laboratory showed that normal values of maximum resting pressure (MRP) and maximum squeeze pressure (MSP) were respectively $68,1 \pm 12,3$ mmHg and $112 \pm 36,2$ mmHg [11]. The normal range of MRP, according to Jones et al. [12], were 45-85 mmHg; so that CAAF without hypertonic IAS were defined as those with MRP values < 85 mmHg.

All patients underwent fissurectomy and anoplasty with V-Y skin flap advancement lying in a gynecological position under spinal or general anesthesia.

The sentinel skin tags and hypertrophied anal papilla located at the dentate line were excised, if present; the tissue at the base of the fissure was curetted until clean IAS muscle fibers were reached. Technical details concerning the surgical procedure have already been widely explained in a previous work from our group [13].

The patients with hypertonic IAS were treated with intraoperative local injection of 30 U.I. of botulinum toxin A (Botox, Allergan Westport, Ireland) [14] or with local administration of post-operative nifedipine and lidocaine for 15 days after surgery (Antrolin®) [15]. Before surgery, all patients received a small

volume of phosphate-saline enema. Metronidazole was administered intravenously in a dose of 500mg 1h before surgery, subsequently, it was administered per os at the dosage of 250 mg for 7 days, three times daily. During the first two weeks after the surgery, patients took variable doses of psyllium fibers. A laxative preparation (sennosides) was given orally to subjects who had not yet passed stools 3 days after surgery. Immediately after surgery, all patients received 100 mg of diclofenac intramuscularly for analgesia and were instructed to take only 100mg of nimesulide tablets as requires.

A complete healing was defined as a complete epithelialization of the advancement skin flap. Recurrent CAFs were defined as those who occurred after the complete healing of the previous wound. Both duration and intensity of post-defecatory pain have been evaluated; the intensity was evaluated with a visual analogue scale (VAS).

FI was assessed preoperatively and 6,12 and 24 months after surgery according to Pescatori's grading system[16] : A incontinence for flatus and mucus; B for liquid stool; C for solid stool; 1 for occasional; 2 for weekly and 3 for daily. Patients were discharged within 24 hours after surgery, afterwards they were examined until they were completely healed and they were also followed up until 24 months following the surgical procedure. Independently of the scheduled appointments, patients have been seen on request.

Results

Demographic, clinical characteristics of CAF of the patients object to study are reported in Table 1. We observed an anterior location of the CAF in the 34,5% of patients, while we recorded an hypertonic IAS in the 65,5 % of them (Table 2). Among the 72 patients with hypertonic IAS, we treated 40 of them with the injection of Botulinum Toxin A, while 32 of them with topical administration of nifedipine and lidocaine ointment. At 40 days after the surgical procedure we achieved a complete wound healing a resolution of clinical symptoms. The duration and intensity of post-defecatory pain have significantly reduced as compared with the pre-operative values since the first defecation, data not shown.

Manometry findings

At 12 and 24 months after surgery MRP and MSP values were substantially unmodified in the patients with normotonic IAS. In those patients with hypertonic IAS MSP values were immodified as compared to the values at 12 and 24 months after surgery. Whereas MRP ones were significantly lower at 12 months follow-up as compared to the pre-operative ones but still significantly higher than those measured in healthy controls. At 24 months follow up after surgery MRP values were within normal range.

Faecal incontinence

We recorded 15 cases of pre-operative FI (13,63%), in 7 of them CAF was located at the anterior commissure (18,42%), in 8 of them at the posterior one (11,11%). FI cases and rate recorded pre-operatively related to childbirths are reported in Table 3 and those related to IAS tone and location in Table 4..

According to Pescatori's grading system [19] 10 were classified as A1, 4 as A2 and 1 as A3. We observed only 2 "de novo" case of post-operative FI both classified as A1 and temporaries. We recorded a temporary worsening of A1 pre-operative grading in 2 cases: in the first one it occurred in patients treated with Botulinum Toxin A injection and the CAF was located at the posterior commissure; in the other one the IAS tone was normal and the CAF located at the anterior commissure.

Recurrence

At 24 months after the surgical procedure follow-up we recorded 8 recurrences (7,3%). In 6 of them MRP values were normal and they were successfully treated with conservative therapy (pharmacological and hygienic-dietary approach). In 2 cases there was a mild hypertonic IAS, these latter have been treated with dilatation with complete healing.

Complications and follow-up

We did not observe any anal stenosis nor keyhole deformity. At the same time, we did not record any necrosis of the transposed flap. The post-operative complications we were able to observe were of slight entity and they didn't require, in any case, further surgical procedures; in particular we observed 1 case of infection located at the donor site as well as 8 case of partial break down.

Discussion

The results of this study show that fissurectomy and anoplasty associated with a pharmacological therapy in case of hypertonic IAS is feasible and effective; this procedure allows a perfect post-operative faecal continence and a low rate of recurrence.

LIS is the treatment of choice for CAFs, but in female patient both of childbearing age or elderly it is not appropriate to undertake this procedure, nor in the traditional nor in the new proposed variant [17-23];, due to the high risk of immediate or long term post-operative FI, caused by some anatomical, clinical and functional features typical of these patients as compared to male ones, which expose female patients to a higher risk of FI [24].

Murad-Regadas et al. [25] , through a 3-D anal endosonography have demonstrated an asymmetrical configuration of anal canal and a difference between the two genders. In women patients both external anal sphincter (EAS) and IAS are shorter at the anterior level and they are characterized by a longer gap, which may justify the higher incidence of pelvic dysfunction . It is still not known whether some women have slightly different pressure distribution in the distal anal canal predisposing to a higher risk than man for the development of anterior CAF [26]. Ellis [27] reported that patients affected by anterior CAF are frequently affected also by rectocele at physical examination with a typical manometric profile (high pressure zone shortened with low to normal resting pressure) .

We shall consider that female patients of childbearing age coul give birth by vaginal delivery during which the lesions of the anal sphincteric system may occur. Sultan et al [28], report in their study that an

isolated dysfunction of the IAS might occur after vaginal delivery in the 13% as well as a dysfunction of EAS in the 59% of cases, while a dysfunction of both sphincters arrives in the 10% of cases. Richter et al. [29] observed a sphincteric dysfunction after vaginal delivery in the 35% of cases.

As for the differences in manometric parameters in various research, different authors [30-32] report that MSP values are significantly lower in female patients, as they represent the strength of the EAS, which is made by striated muscle. As a matter of fact, men have a greater striated muscle due to the well-known effect of testosterone of muscle growth [33]. MRP values, on the contrary, are similar in both sexes[30-32,34].

Available research focusing on LIS outcomes in female patients are few. Regadas et al [35] and Fernandes et al. [36] analyzed respectively a series on 31 and 36 female patients affected by CAF hypertonic IAS, without FI or previous proctological procedure, who underwent LIS extended to the apex of the fissure; they observed, in the 52-53% of cases. The authors highlighted, with a post-operative evaluation endoanal ultrasound (EAUS), that if LIS extension was greater than 20-25% of the IAS length (corresponding to less than 1 cm), the incidence of FI was significantly higher. It appears difficult, if not impossible, to divide exactly a quarter of the IAS length, as there is not any reference landmark neither with EAUS.

In their recent research Brillantino et al. [37] performed a sphincterotomy limited to the 20% of the IAS length as treatment for female patient affected by CAF with hypertonic IAS and without any clinical sign of FI, with the help of a new device introduced by themselves. This latter procedure requires a pre-operative evaluation with an EAUS and the researchers achieved very good results with a rate of healing of the 93,3% and a rate of recurrence of 6,2% without alteration of the anal continence. Nevertheless, this proctological surgery appears complex and requires equipment, which are not available in every hospital.

Conclusion

Even if a LIS limited to the 20% of the IAS length allows to achieve good results both in terms of healing and FI rate, in female patients affected by CAF with hypertonic IAS without signs of FI or previous proctological procedure, this procedure appears difficult and requires further technical development.

So that, up to date sphincter saving surgical procedure represent the treatment of choice for female patients affected by CAF regardless of the IAS tone, age, previous proctological procedure or presence of FI [38-42]. The saving sphincter procedure we undertook, fissurectomy with V-Y advancement flap associated with chemical- pharmacological sphincterotomy for IAS tone reduction, in case of hypertonic IAS may represent an effective approach for the treatment of CAF in female patients. Only in case of failure of these latter LIS could be taken into account as an alternative approach.

Bibliography

1. Garg P, Garg M, Menon GR. Long-term continence disturbance after lateral internal sphincterotomy for chronic anal fissure: a systematic review and meta-analysis. *Colorectal Dis.* 2013;15:e104-17.
2. Griffin N, Acheson AG, Tung P, Sheard C, Glazebrook C, Scholefield JH. Quality of life in patients with chronic anal fissure. *Colorectal Dis.* 2004; 6:39-44.
3. Sobrado Júnior CW, Hora JAB, Sobrado LF, Guzela VR, Nahas SC, Ceconello I. Anoplasty with skin tag flap for the treatment of chronic anal fissure. *Rev Col Bras Cir.* 2019; 46: e20192181.
4. Marti L, Post S, Herold A, Scwandner O, Hetzer F, Strittmatter B, Iesalnieks I, Huth M, Schmit-Lauber M, Weyandt G, Ommer A, Ebinger SM. S3-Leitlinie: Analfissur. AWMF-Registriernummer: 081-010. *Coloproctology* 2020 ; 42:90-6
5. Altomare DF, Binda GA, Canuti S, Landolfi V, Trompetto M, Villani R. The management of patients with primary chronic anal fissure: a position paper. *Tech Coloproctol.* 2011; 15:135-41.
6. Alonso-Coello P, Marzo-Castillejo M, Mascort JJ, Hervás AJ, Viña LM, Ferrús JA, Ferrándiz J, López-Rivas L, Rigau D, Solà I, Bonfill X, Piqué JM. [Clinical practice guideline on the treatment of hemorrhoids and anal fissure (update 2007)]. *Gastroenterol Hepatol.* 2008; 31:668-81
7. Holzgang M., Jayne D. Lateral Internal sphincterotomy (LIS)- still top gun in chronic anal fissure treatment? *Coloproctology* 2020; 42:478-84
8. Cross KL, Massey EJ, Fowler AL, Monson JR; ACPGBI. The management of anal fissure: ACPGBI position statement. *Colorectal Dis.* 2008; 3:1-7
9. Emile SH. Indications and Technical Aspects of Internal Anal Sphincterotomy: Highlighting the Controversies. *Dis Colon Rectum.* 2017; 60:128-32.
10. Rociu E, Stoker J, Eijkemans MJ, Laméris JS. Normal anal sphincter anatomy and age- and sex-related variations at high-spatial-resolution endoanal MR imaging. 2000 ;217:395-401.
11. Patti R, Territo V, Aiello P, Angelo GL, Di Vita G. Manometric evaluation of internal anal sphincter after fissurectomy and anoplasty for chronic anal fissure: a prospective study. *Am Surg.* 2012; 78:523-7.
12. Jones OM, Ramalingam T, Lindsey I, Cunningham C, George BD, Mortensen NJ. Digital rectal examination of sphincter pressures in chronic anal fissure is unreliable. *Dis Colon Re* 2005; 48:349-52.
13. Patti R, Guercio G, Territo V, Aiello P, Angelo GL, Di Vita G. Advancement flap in the management of chronic anal fissure: a prospective study. *Updates Surg.* 2012; 64:101-6.
14. D'Orazio B., Geraci G., Martorana G., Sciumé C., Corbo G., Di Vita G. Fissurectomy and anoplasty with botulinum toxin injection in patients with chronic anal posterior fissure with hypertonia: a long-term evaluation. *Updates Surg.* 2020. Ahead of print. DOI: 10.1007/s13304-020-00846-y
15. D'Orazio B., Sciumé C., Famà F, Bonventre S., Martorana G., Cali D., Terranova G., Di Vita G., Geraci G. Surgical Sphincter Saving Approach and Topical Nifedipine for Chronic Anal Fissure with Hypertonic Internal Anal Sphincter. *Chirurgia*, 2020; 115:585-94

16. Pescatori M, Anastasio G, Bottini C, Mentasti A. [New grading and scoring for anal incontinence. Evaluation of 335 patients.](#) *Dis Colon Rectum.*1992; 35:482-7.
17. Garcia-Aguilar J, Belmonte C, Wong WD, Lowry AC, Madoff RD. [Open vs. closed sphincterotomy for chronic anal fissure: long-term results.](#) *Dis Colon Rectum.* 1996;39(4):440-3.
18. Alawady M, Emile SH, Abdelnaby M, Elbanna H, Farid M. [Posterolateral versus lateral internal anal sphincterotomy in the treatment of chronic anal fissure: a randomized controlled trial.](#) *Int J Colorectal Dis.* 2018;33:1461-7.
19. Elsebae MM. [Elsebae MM. A study of fecal incontinence in patients with chronic anal fissure: prospective, randomized, controlled trial of the extent of internal anal sphincter division during lateral sphincterotomy.](#) *World J Surg.* 2007;31:2052-7.
20. Ho KS, Ho YH. [Randomized clinical trial comparing oral nifedipine with lateral anal sphincterotomy and tailored sphincterotomy in the treatment of chronic anal fissure.](#) *Br J Surg.* 2005;92:403-8.
21. Menteş BB, Güner MK, Leventoglu S, Akyürek N. [Fine-tuning of the extent of lateral internal sphincterotomy: spasm-controlled vs. up to the fissure apex.](#) *Dis Colon Rectum.* 2008;51:128-33
22. Brown CJ, Dubreuil D, Santoro L, Liu M, O'Connor BI, McLeod R. [Lateral internal sphincterotomy is superior to topical nitroglycerin for healing chronic anal fissure and does not compromise long-term fecal continence: six-year follow-up of a multicenter, randomized, controlled trial.](#) *Dis Colon Rectum.* 2007;50:442-8.
23. Nessar G, Topbas M. [Lateral Internal Partial Sphincterotomy Technique for Chronic Anal Fissure.](#) *Indian J Surg.* 2017;79:185-7.
24. D’Orazio B., Bonventre S , Sciumé C, Cudia B., Terranova G., Di Vita G., Geraci G. [Lateral Internal Sphincterotomy is still crucial to heal hypertonic chronic anal fissure and normalize the internal anal sphincter tone? Is fissurectomy and V en Y plastic a valuable alternative?](#) *Ital. Chir* 2021; Vol 92. Accepted for publication
25. [Regadas FS, Murad-Regadas SM, Lima DM, Silva FR, Barreto RG, Souza MH, Regadas Filho FS.](#) [Anal canal anatomy showed by three-dimensional anorectal ultrasonography.](#) *Surg Endosc.*2007; 21:2207-11.
26. D’Orazio B, Geraci G, Famà F, Cudia B., Martorana G., Sciumé C., Corbo G., Terannova G., Bonventre S., Di Vita G. [Long term outcomes of fissurectomy and anoplasty for chronic anterior anal fissure without hypertonia: low recurrences and continence conservation.](#) *Ann Ital Chir,* 2020; 9. pii: S0003469X20034727. Online ahead of print
27. [Ellis CN.](#) [Anterior levatorplasty for the treatment of chronic anal fissures in females with a rectocele: a randomized, controlled trial.](#) *Dis Colon Rectum.*2004; 47:1170-3.
28. Sultan AH, Kamm MA, Hudson CN, Thomas JM, Bartram CI. [Anal-sphincter disruption during vaginal delivery.](#) . *N Engl J Med.* 1993;329:1905-11.
29. Richter HE, Fielding JR, Bradley CS, Handa VL, Fine P, FitzGerald MP, Visco A, Wald A, Hakim C, Wei JT, Weber AM; Pelvic Floor Disorders Network. [Endoanal ultrasound findings and fecal incontinence](#)

- symptoms in women with and without recognized anal sphincter tears. . *Obstet Gynecol.* 2006;108:1394-401.
30. Lee HR, Lim SB, Park JY. Anorectal manometric parameters are influenced by gender and age in subjects with normal bowel function. *Int J Colorectal Dis.* 2014;29:1393-9
 31. Williams AB, Cheetham MJ, Bartram CI, Halligan S, Kamm MA, Nicholls RJ, Kmiot WA. Gender differences in the longitudinal pressure profile of the anal canal related to anatomical structure as demonstrated on three-dimensional anal endosonography. *Br J Surg.* 2000 ;87(:1674-9.
 32. Gundling F, Seidl H, Scalercio N, Schmidt T, Schepp W, Pehl C. Influence of gender and age on anorectal function: normal values from anorectal manometry in a large caucasian population. 2010;81:207-13.
 33. Storer TW, Woodhouse L, Magliano L, Singh AB, Dzekov C, Dzekov J, Bhasin S.. Changes in muscle mass, muscle strength, and power but not physical function are related to testosterone dose in healthy older men. *J Am Geriatr Soc.* 2008 ;56:1991-9.
 34. Felt-Bersma RJ, Gort G, Meuwissen SG Normal values in anal manometry and rectal sensation: a problem of range.. 1991;38(5):444-9.
 35. Murad-Regadas SM, Fernandes GO, Regadas FS, Rodrigues LV, Pereira Jde J, Regadas Filho FS, Dealcanfreitas ID, Holanda Ede C How much of the internal sphincter may be divided during lateral sphincterotomy for chronic anal fissure in women? Morphologic and functional evaluation after sphincterotomy. . *Dis Colon Rectum.* 2013;56:645-51.
 36. Fernandes GO, Murad-Regadas SM, Regadas FS, Rodrigues LV, Dealcanfreitas ID, Pereira Jde J, de Carvalho E, Regadas Filho FS. Clinical, functional and morphologic evaluation of patients undergoing lateral sphincterotomy for chronic anal fissure treatment. Identification of factors that can interfere with faecal continence. *J Coloproctol (Rio J).* 2014; 34: 174-80.
 37. Brilliantino A, Izzo D, Iacobellis F, Maglio M, Grillo M, Vincenzo L, Bottino V, Renzi A. Safety and effectiveness of minimal sphincterotomy in the treatment of female patients with chronic anal fissure. *Updates SurgOnline* ahead of print. <https://doi.org/10.1007/s13304-020-00874-8>
 38. Hancke E, Rikas E, Suchan K, Völke K. Dermal flap coverage for chronic anal fissure: lower incidence of anal incontinence compared to lateral internal sphincterotomy after long-term follow-up. *Dis Colon Rectum.* 2010;53:1563-8.
 39. Iesalnieks I. Kombinationen therapie Fissurektomie und Botulinum toxin in der Behandlung der chronischen Analfissur. *Coloproctology* 2020; 42: 464-8
 40. Chambers W, Sajal R, Dixon A.. V-Y advancement flap as first-line treatment for all chronic anal fissures. *Int J Colorectal Dis.* 2010;25:645-8.
 41. Giordano P, Gravante G, Grondona P, Ruggiero B, Porrett T, Lunniss PJ. Simple cutaneous advancement flap anoplasty for resistant chronic anal fissure: a prospective study. *World J Surg.* 2009; 33:1058-63.
 42. Hancke E, Suchan K, Völke K. Anokutaner Advancement-Flap zur sphinkterchonenden chirurgischen Therapie der chronischen Analfissur. *Coloproctology.* 2020; 42: 270-6

Declarations

AUTHORS AND ACKNOWLEDGEMENT

All authors contributed to the study.

D’Orazio B., Geraci G. and Di Vita G. conceived, devised and designed the manuscript.

Cali D. collected the case and iconography.

D’Orazio B. and Geraci G. wrote the manuscript and participated in the sequence alignment.

Di Vita G. and Bonventre S. revised and approved the final manuscript and its conclusions.

No acknowledgements are required.

FUNDING

No financial support has been obtained in the preparation of this study.

DISCLOSURE AND COMPETING INTEREST

The Authors report no conflict of interest in this work.

The Authors declare that they have no competing interests.

This article has not been presented nor published elsewhere, and no financial support has been obtained in its preparation.

CONSENT SECTION

Informed written consent was obtained from all individuals participants included in this study.

The consent for publication has been obtained by all participants to the work.

ETHICS APPROVAL

All procedures performed in studies involving human participants were in accordance with the ethical standards of the 1964 Helsinki Declaration and its later amendments or comparable ethical standards.

The protocol for this study has been submitted to the Ethical Committee of our institution “Ethical Committee Palermo 1-AUOP P.Giaccone” on the 1st of May 2020, which did not consider necessary to approve it and waved off, in that occasion, the requirement for ethical approval.

AVAILABILITY OF DATA

The datasets generated during and analysed during the current study are not publicly available because we did not consider it necessary to publish them in a public repository but are available from the corresponding author on reasonable request.

References

1. Garg P, Garg M, Menon GR. Long-term continence disturbance after lateral internal sphincterotomy for chronic anal fissure: a systematic review and meta-analysis. *Colorectal Dis.* 2013;15:e104-17.
2. Griffin N, Acheson AG, Tung P, Sheard C, Glazebrook C, Scholefield JH. Quality of life in patients with chronic anal fissure. *Colorectal Dis.* 2004; 6:39–44.
3. Sobrado Júnior CW, Hora JAB, Sobrado LF, Guzela VR, Nahas SC, Cecconello I. Anoplasty with skin tag flap for the treatment of chronic anal fissure. *Rev Col Bras Cir.* 2019; 46: e20192181.
4. Marti L, Post S, Herold A, Scwandner O, Hetzer F, Strittmatter B, Iesalnieks I, Huth M, Schmit-Lauber M, Weyandt G, Ommer A, Ebinger SM. S3-Leitlinie: Analfissur. AWMF-Registriernummer: 081 – 010. *Coloproctology* 2020; 42:90 – 6
5. Altomare DF, Binda GA, Canuti S, Landolfi V, Trompetto M, Villani R. The management of patients with primary chronic anal fissure: a position paper. *Tech Coloproctol.* 2011; 15:135–41.
6. Alonso-Coello P, Marzo-Castillejo M, Mascort JJ, Hervás AJ, Viña LM, Ferrús JA, Ferrándiz J, López-Rivas L, Rigau D, Solà I, Bonfill X, Piqué JM. [Clinical practice guideline on the treatment of hemorrhoids and anal fissure (update 2007)]. *Gastroenterol Hepatol.* 2008; 31:668–81
7. Holzgang M., Jayne D. Lateral Internal sphincterotomy (LIS)- still top gun in chronic anal fissure treatment? *Coloproctology* 2020; 42:478 – 84
8. Cross KL, Massey EJ, Fowler AL, Monson JR; ACPGBI. The management of anal fissure: ACPGBI position statement. *Colorectal Dis.* 2008; 3:1–7
9. Emile SH. Indications and Technical Aspects of Internal Anal Sphincterotomy: Highlighting the Controversies. *Dis Colon Rectum.* 2017; 60:128–32.
10. Rociu E, Stoker J, Eijkemans MJ, Laméris JS Normal anal sphincter anatomy and age- and sex-related variations at high-spatial-resolution endoanal MR imaging. *Radiology.* 2000;217:395–401.
11. Patti R, Territo V, Aiello P, Angelo GL, Di Vita G. Manometric evaluation of internal anal sphincter after fissurectomy and anoplasty for chronic anal fissure: a prospective study. *Am Surg.* 2012; 78:523–7.
12. Jones OM, Ramalingam T, Lindsey I, Cunningham C, George BD, Mortensen NJ. Digital rectal examination of sphincter pressures in chronic anal fissure is unreliable. *Dis Colon Rectum.* 2005; 48:349–52.
13. Patti R, Guercio G, Territo V, Aiello P, Angelo GL, Di Vita G. Advancement flap in the management of chronic anal fissure: a prospective study. *Updates Surg.* 2012; 64:101–6.
14. D’Orazio B., Geraci G., Martorana G., Sciumé C., Corbo G., Di Vita G. Fisurectomy and anoplasty with botulinum toxin injection in patients with chronic anal posterior fissure with hypertonia: a long-term evaluation. *Updates Surg.* 2020. Ahead of print. DOI: 10.1007/s13304-020-00846-y

15. D’Orazio B, Sciumé C, Famà F, Bonventre S., Martorana G., Cali D., Terranova G., Di Vita G., Geraci G. Surgical Sphincter Saving Approach and Topical Nifedipine for Chronic Anal Fissure with Hypertonic Internal Anal Sphincter. *Chirurgia*, 2020; 115:585 – 94
16. Pescatori M, Anastasio G, Bottini C, Mentasti A. New grading and scoring for anal incontinence. Evaluation of 335 patients. *Dis Colon Rectum*.1992; 35:482–7.
17. Garcia-Aguilar J, Belmonte C, Wong WD, Lowry AC, Madoff RD. Open vs. closed sphincterotomy for chronic anal fissure: long-term results. *Dis Colon Rectum*. 1996;39(4):440–3.
18. Alawady M, Emile SH, Abdelnaby M, Elbanna H, Farid M. Posterolateral versus lateral internal anal sphincterotomy in the treatment of chronic anal fissure: a randomized controlled trial. *Int J Colorectal Dis*. 2018;33:1461–7.
19. Elsebae MM. Elsebae MM. A study of fecal incontinence in patients with chronic anal fissure: prospective, randomized, controlled trial of the extent of internal anal sphincter division during lateral sphincterotomy. *World J Surg*. 2007;31:2052–7.
20. Ho KS, Ho YH. Randomized clinical trial comparing oral nifedipine with lateral anal sphincterotomy and tailored sphincterotomy in the treatment of chronic anal fissure. *Br J Surg*. 2005;92:403–8.
21. Menteş BB, Güner MK, Leventoglu S, Akyürek N. Fine-tuning of the extent of lateral internal sphincterotomy: spasm-controlled vs. up to the fissure apex. *Dis Colon Rectum*. 2008;51:128–33
22. Brown CJ, Dubreuil D, Santoro L, Liu M, O’Connor BI, McLeod RS Lateral internal sphincterotomy is superior to topical nitroglycerin for healing chronic anal fissure and does not compromise long-term fecal continence: six-year follow-up of a multicenter, randomized, controlled trial. *Dis Colon Rectum*. 2007;50:442–8.
23. Nessar G, Topbas M. Lateral Internal Partial Sphincterotomy Technique for Chronic Anal Fissure. *Indian J Surg*. 2017;79:185–7.
24. D’Orazio B., Bonventre S, Sciumé C, Cudia B., Terranova G., Di Vita G., Geraci G. Lateral Internal Sphincterotomy is still crucial to heal hypertonic chronic anal fissure and normalize the internal anal sphincter tone? Is fissurectomy and V en Y plastic a valuable alternative? *Ann. Ital. Chir* 2021; Vol 92. Accepted for publication
25. Regadas FS, Murad-Regadas SM, Lima DM, Silva FR, Barreto RG, Souza MH, Regadas Filho FS. Anal canal anatomy showed by three-dimensional anorectal ultrasonography. *Surg Endosc*. 2007; 21:2207–11.
26. D’Orazio B, Geraci G, Famà F, Cudia B., Martorana G., Sciumé C., Corbo G., Terannova G., Bonventre S., Di Vita G. Long term outcomes of fissurectomy and anoplasty for chronic anterior anal fissure without hypertonia: low recurrences and continence conservation. *Ann Ital Chir*, 2020; 9. pii: S0003469X20034727. Online ahead of print
27. Ellis CN. Anterior levatorplasty for the treatment of chronic anal fissures in females with a rectocele: a randomized, controlled trial. *Dis Colon Rectum*. 2004; 47:1170–3.
28. Sultan AH, Kamm MA, Hudson CN, Thomas JM, Bartram CI Anal-sphincter disruption during vaginal delivery.. *N Engl J Med*. 1993;329:1905–11.

29. Richter HE, Fielding JR, Bradley CS, Handa VL, Fine P, FitzGerald MP, Visco A, Wald A, Hakim C, Wei JT, Weber AM; Pelvic Floor Disorders Network Endoanal ultrasound findings and fecal incontinence symptoms in women with and without recognized anal sphincter tears.. *Obstet Gynecol.* 2006;108:1394–401.
30. Lee HR, Lim SB, Park JY. Anorectal manometric parameters are influenced by gender and age in subjects with normal bowel function. *Int J Colorectal Dis.* 2014;29:1393–9
31. Williams AB, Cheetham MJ, Bartram CI, Halligan S, Kamm MA, Nicholls RJ, Kmiot WA. Gender differences in the longitudinal pressure profile of the anal canal related to anatomical structure as demonstrated on three-dimensional anal endosonography. *Br J Surg.* 2000;87(16):1674–9.
32. Gundling F, Seidl H, Scalercio N, Schmidt T, Schepp W, Pehl C. Influence of gender and age on anorectal function: normal values from anorectal manometry in a large caucasian population. *Digestion.* 2010;81:207–13.
33. Storer TW, Woodhouse L, Magliano L, Singh AB, Dzekov C, Dzekov J, Bhasin S.. Changes in muscle mass, muscle strength, and power but not physical function are related to testosterone dose in healthy older men. *J Am Geriatr Soc.* 2008;56:1991–9.
34. Felt-Bersma RJ, Gort G, Meuwissen SG Normal values in anal manometry and rectal sensation: a problem of range.. *Hepatogastroenterology.* 1991;38(5):444–9.
35. Murad-Regadas SM, Fernandes GO, Regadas FS, Rodrigues LV, Pereira Jde J, Regadas Filho FS, Dealcanfreitas ID, Holanda Ede C How much of the internal sphincter may be divided during lateral sphincterotomy for chronic anal fissure in women? Morphologic and functional evaluation after sphincterotomy.. *Dis Colon Rectum.* 2013;56:645–51.
36. Fernandes GO, Murad-Regadas SM, Regadas FS, Rodrigues LV, Dealcanfreitas ID, Pereira Jde J, de Carvalho E, Regadas Filho FS. Clinical, functional and morphologic evaluation of patients undergoing lateral sphincterotomy for chronic anal fissure treatment. Identification of factors that can interfere with faecal continence. *J Coloproctol (Rio J).* 2014; 34: 174–80.
37. Brillantino A, Izzo D, Iacobellis F, Maglio M, Grillo M, Vincenzo L, Bottino V, Renzi A. Safety and effectiveness of minimal sphincterotomy in the treatment of female patients with chronic anal fissure. *Updates Surg 2020.* Online ahead of print. <https://doi.org/10.1007/s13304-020-00874-8>
38. Hancke E, Rikas E, Suchan K, Völke K. Dermal flap coverage for chronic anal fissure: lower incidence of anal incontinence compared to lateral internal sphincterotomy after long-term follow-up. *Dis Colon Rectum.* 2010;53:1563–8.
39. Iesalnieks I. Kombinationen therapie Fissurektomie und Botulinum toxin in der Behandlung der chronischen Analfissur. *Coloproctology 2020;* 42: 464–8
40. Chambers W, Sajal R, Dixon A.. V-Y advancement flap as first-line treatment for all chronic anal fissures. *Int J Colorectal Dis.* 2010;25:645–8.
41. Giordano P, Gravante G, Grondona P, Ruggiero B, Porrett T, Lunniss PJ. Simple cutaneous advancement flap anoplasty for resistant chronic anal fissure: a prospective study. *World J Surg.* 2009; 33:1058–63.

Tables

TABLE 1 Clinical and demographic characteristics of CAF

	No.	%
Gender		
Female	110	100
Age (mean ± standard deviation)	36,5±15,8	
Hypertrophied anal papilla	88	80
Skin tags	82	75
Symptoms		
Pain	110	100
Bleeding	88	80
Pruritus	66	60
Duration of Symptoms		
Months (mean ± standard deviation)	12,1±7,3	

TABLE 2 Location and IAS tone in the patients object to study

Location	Normotonic IAS		Hypertonic IAS		Tot.	
	N°	%	N°	%	N°	%
Anterior	21	55,3	17	44,7	38	34,5
Posterior	17	23,6	55	76,4	72	65,5
Tot.	38	34,5	72	65,5	110	100

TABLE 3 Faecal Incontinence assessed preoperatively related to childbirths

	Childbirth		Faecal Incontinence	
	N°	%	N°	%
Nulliparous	38	34,5	3	7,8
Multiparous	72	65,6	12	16,6
C-section	27	24,5	3	11,1
Vaginal birth	45	40,5	9	20

TABLE 4 Faecal Incontinence assessed preoperatively related to IAS tone and CAF's location.

Location of the CAF	Patients		Faecal Incontinence			
	N°	%	IAS tone		Total	
Normal			High	N°	%	
Anterior	38	34,5	4	3	7	18,42
Posterior	72	65,6	3	5	8	11,1