

WDPCP modulates cilia beating through MAPK/ERK pathway in chronic rhinosinusitis

Yun Ma

Sun Yat-sen Memorial Hospital <https://orcid.org/0000-0002-4672-3584>

Peng Tian

Sun Yat-sen Memorial Hospital

Hua Zhong

First Affiliated Hospital of Sun Yat-sen University

Fan Wu

Sun Yat-sen Memorial Hospital

Qining Zhang

Sun Yat-sen Memorial Hospital

Xiang Liu

Sun Yat-sen Memorial Hospital

Hua Dang

Sun Yat-sen Memorial Hospital

Qiujian Chen

Sun Yat-sen Memorial Hospital

Hua Zou

Sun Yat-sen Memorial Hospital

Yiqing Zheng (✉ zhengyiq@mail.sysu.edu.cn)

Research

Keywords: chronic rhinosinusitis, WDPCP, cilia beating, mitochondria

Posted Date: May 3rd, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-22917/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background: Cilia loss and dysfunction is one of the typical pathological features of chronic rhinosinusitis. Tryptophan-aspartic acid (W-D) repeat containing planar cell polarity effector (WDPCP) has been proven to be an essential element for ciliogenesis in human nasal epithelium, but its role in the beating of cilia remains unclear. Cilia beating requires energy from the mitochondria, which is regulated by the MAPK/ERK pathway. In this study, we sought to investigate the role of WDPCP and its underlying mechanism behind the dysfunction in the beating of cilia in chronic rhinosinusitis.

Methods: We demonstrated WDPCP expression in the epithelium of nasal polyps. We also investigated the MAPK/ERK pathway in primary human sinonasal epithelial cells to explore the function of WDPCP. The air-liquid interface culture system was used as a model to verify the role of WDPCP and the MAPK/ERK pathway in the beating of cilia.

Results: With the dysfunction of cilia beating, we observed a low expression of WDPCP in the epithelium of nasal polyps. Within the *in vitro* study, we found that WDPCP was critical for mitochondrial biogenesis and mitochondrial function in human sinonasal epithelial cells, possibly due to the activation of the MAPK/ERK pathway. The mitochondrial dysfunction caused by U0126 or lacking WDPCP could be partially recovered by dexamethasone.

Conclusion: The low expression of WDPCP in nasal epithelium could affect mitochondria via the MAPK/ERK pathway, which may contribute to the dysfunction in the beating of cilia in chronic rhinosinusitis.

Background

Chronic rhinosinusitis (CRS) is a common disease which affects 5.7%-16% of the world population[1–4]. CRS is divided into CRS with nasal polyps (CRSwNP) and CRS without nasal polyps (CRSsNP), depending on the presence or absence of nasal polyps (NP). CRS has been identified as an inflammatory disease. Cilia cells play an important role in the nasal epithelium. The beating of cilia occurs in a coordinated manner to generate flow for the clearance of debris and pathogens in mucus. The beating of cilia is affected by a variety of factors such as bacterial pathogens[5], inflammatory cytokines[6], tobacco smoke[7], temperature[8] and pH value[9]. Cilia loss and dysfunction is one of the predominant pathological changes present in CRS[10]. However, the mechanism of poor ciliation and cilia dysfunction in CRS remains unclear.

The planar cell polarity (PCP) pathway, which is also the non-canonical Wnt pathway, regulates the convergence and extensive movement of cells during embryogenesis and gastrulation of vertebrates[11, 12]. Proteins in the PCP pathway control the morphogenesis and motility of multiciliated epithelial cells[13–15]. WD repeat containing planar cell polarity effector (WDPCP) was critical for ciliogenesis in *Xenopus* and controlled cell polarity in *Drosophila*[16, 17]. In our previous study, we have discovered that

WDPCP was essential for ciliogenesis in human nasal epithelium, but it was still unclear whether WDPCP regulated cilia motility in the human airway[18].

The beating of cilia depends on ATP provided by mitochondria[19]. Decreased energy levels caused by mitochondrial damage is an important reason for ciliary dysfunction in airway diseases[20, 21]. The mitogen activated protein kinase (MAPK) and extracellular signal-regulated kinase (ERK) signaling pathway is a protein-serine/threonine kinase cascade which includes dual-specificity mitogen-activated protein kinases kinases 1/2 (MEK1/2) that activates the effector kinases ERK1/2. This pathway participates in a variety of nuclear transcription factors and affects cell proliferation, apoptosis, differentiation, inflammatory response, and other biological processes[22]. Moreover, mitochondrial membrane potential, mitochondrial oxidative stress and mitochondrial energy metabolism were also modulated by the MAPK/ERK pathway[23]. However, studies regarding cilia disorders related to mitochondrial damage in human nasal epithelium remain lacking.

Therefore, the aim of this study was to investigate whether WDPCP plays a role in the function of cilia in human nasal epithelium, and to unveil corresponding molecular mechanisms of mitochondrial damage regulated by the MAPK/ERK pathway.

Methods

Human tissue procurement

Our study was approved by the Ethics Committee of the Sun Yat-sen Memorial Hospital of Sun Yat-sen University (2018-212), and written consent was obtained from each participant. Both control subjects and patients with CRSwNP which were aged from 18 to 65 were recruited from the Department of Otorhinolaryngology at the Sun Yat-sen Memorial Hospital of Sun Yat-sen University from September 2018 to July 2019. Their clinical characteristics were listed in Table 1. The diagnosis of CRSwNP was based on a European position paper on rhinosinusitis and nasal polyps published in 2012[24]. Patients with genetic defects such as primary ciliary dyskinesia or any systemic diseases were excluded from this study. Control subjects had no evidence of sinonasal mucosal inflammation but have underwent sinonasal surgery for the repair of cerebrospinal fluid rhinorrhea and optic nerve decompression. Mucosa was removed from the ethmoid sinuses (ES), which can be easily accessed during surgery. Mucosa of ethmoid sinuses from control subjects and polyp tissues from patients with CRSwNP were harvested during surgery. The samples were used for various further analyses as followed.

Air-liquid interface (ALI) cultures

The complete method for the preparation of primary sinonasal epithelial cultures can be found in our previous study[18]. Human sinonasal epithelial cells (HSECs) were transferred to Transwell inserts

(Corning, 0.4µm) to initiate the ALI culture. 14 days were required to obtain the differentiated cilia via the PneumaCult™-ALI Medium (Stem cell). The ALI membranes were collected on Day-0, -7 and -14.

Quantitative RT-PCR

Quantitative real-time PCR (RT-PCR) was performed on human sinonasal epithelial cells and clinical samples. β-actin was used as a reference for normalization. Nuclear respiratory factor 1 (NRF1), nuclear respiratory factor 2 (NRF2), transcription factor A, mitochondrial (TFAM) and cytochrome c oxidase subunit 4 (COX4) PCR was performed with the Roche LightCycler 480 Real-Time PCR System using SYBR Premix Ex Taq (Takara). Relative gene expression was calculated using the $2^{-\Delta\Delta CT}$ method.

Histological staining

Immunohistochemical staining was performed on the paraffin sections. After dehydration, antigen retrieval, quenching of endogenous peroxides, and blocking, sections were incubated overnight at 4 °C with monoclonal mouse anti-NRF1 at 1:100 (Abcam, ab175932), monoclonal mouse anti-NRF2 at 1:100 (Abcam, ab62352), monoclonal mouse anti-TFAM at 1:50 (Abcam, ab131607) and polyclonal rabbit anti-COX4 at 1:200 (Cell signaling, 4850S). DAB and hematoxylin were used for staining.

Immunofluorescence staining was performed for air-liquid interface (ALI) membranes and HSECs. In brief, HSECs were incubated overnight at 4 °C with monoclonal mouse anti-beta tubulin α at 1:100 (Abcam, ab11315), and subsequently with Alexa flour 488 secondary antibodies (Invitrogen) at 1:200 at room temperature for 1 hour, and 40,6-diamidino-2-phenylindole (DAPI) (10 mg/mL, Sigma-Aldrich) for 10 minutes.

Mitochondrial structure was detected via fluorescent MitoTracker (Invitrogen), which was cultured with HSECs at 1:500 dilution at 37 °C for 1 hour. Then the cells were fixed and stained with DAPI for 10 minutes. HSECs were observed and imaged with a Zeiss LSM 780 confocal microscope.

Western blot analysis

Total protein from nasal mucosa and HSECs were extracted in RIPA lysis buffer. Protein concentrations were determined by bicinchoninic acid assay (BCA). Samples which contained 30 µg of protein were resolved via SDS-PAGE in 10%~12% Tris-glycine gels and transferred onto a polyvinylidene fluoride membrane (Milipore) and blocked with 5% bovine serum albumin (BSA). The membranes were incubated overnight with monoclonal mouse anti-NRF1 (Abcam, ab175932), monoclonal mouse anti-NRF2 (Abcam, ab62352), monoclonal mouse anti-TFAM (Abcam, ab131607), polyclonal rabbit anti-COX4 (Cell signaling,

4850S) at 1:1000, then incubated with an appropriate secondary antibody (1: 10,000) for 1 hour. The relative protein levels were quantified by densitometric image analysis of bands using Fiji (National Institutes of Health) and were normalized against β -actin.

Scanning electron microscopy

The membranes which contained HSECs from the Transwell inserts were fixed with 2.5% glutaraldehyde for 4 hours at room temperature. Subsequently, the cells were progressively dehydrated in 30%, 50%, 70% and 90% ethanol for one time per concentration at 15-minutes intervals, 3 times in 100% ethanol for 15 minutes, and 3 times in 100% tert-butyl alcohol for 15 minutes. The filters were then glued onto scanning electron microscopic (SEM) stubs and sputter coated to a thickness of 12 nm by gold palladium. The surface of the Transwell membrane was examined at an accelerating voltage of 10 kV using a Quanta-400 scanning electron microscope (FEI).

Cell stimulation and transfection

Human sinonasal epithelial cells were transfected with siRNA targeted to WDPCP using Lipofectamine 3000 (Invitrogen) following the manufacturer's instructions. HSECs were stimulated by U0126 (10 μ M) or dexamethasone (0.01 mg/mL). After 24 hours of incubation, the protein was collected for further experiments.

Mitochondrial DNA copy number analysis

Total DNA was extracted from HSECs using the QIAamp DNA Mini Kit (Qiagen). The mitochondrial DNA (mtDNA) copy number was determined by amplifying genes encoding genomic DNA and mitochondrial DNA. The mtDNA levels were quantified by quantitative real-time polymerase chain reaction on a Roche Light Cycler 96 (Roche) using HV1 primers (forward: 5'-TTGCACGGTACCATAAATACTTGAC-3', reverse: 5'-GAGTTGCAGTTGATGTGTGATAGTTG-3'). Nuclear gene β -globin primers (forward: 5'-ACACAAGTGTTCCTACTAGC-3', reverse: 5'-CAACTTCATCCACGTTCCACC-3') were used as a nuclear control. Relative quantification of mitochondrial DNA copy number was calculated after using the $2^{-\Delta\Delta CT}$ method to obtain the expression fold change.

Mitochondrial membrane potential measurement

The mitochondrial membrane potential was evaluated with JC-1 fluorescent dye (Beyotime, China). JC-1 displayed red or green fluorescence depending on the mitochondrial potential. Normal mitochondrial

membrane potential exhibited red fluorescence, whereas damaged mitochondria exhibited green fluorescence. The HSECs were stained with JC-1 in culture media for 20 minutes at 37 °C and were then washed for 3 times with washing buffer. The mitochondrial membrane potential was measured by flow cytometry (BD Biosciences).

Measurement of ATP levels

An ATP Assay Kit (Beyotime, China) was used to conduct the ATP assay. Briefly, HSECs were collected in lysis buffer and centrifuged at 12,000 × g for 5 minutes at 4 °C. ATP detection reagent was added into 96-wells, then the standards and samples were added into the wells and the detection solution was mixed. Chemiluminescence was detected by a Synergy H1 Hybrid Multi-Mode Reader (BioTek, USA). The levels of ATP were calculated based on the standard curve and were normalized to the protein content.

Cilia beating frequency

Both ciliated HSECs in ALI culture and clinical samples collected within 30 minutes were used for cilia beating frequency (CBF) analysis. The room temperature was controlled at 25 °C, and the humidity was maintained at 70-80%, which would not significantly affect the CBF[25]. A high-speed digital video camera (Basler AG) captured images at 100 frames per second and Sisson-Ammons Video Analysis (SAVA) software (National Instruments) was used for video analysis. Each measurement was obtained 3 times and recorded 15 seconds each time.

Statistical analysis

Data were presented as mean ± standard error of the mean (SEM), and analyzed using GraphPad PRISM 7 (GraphPad Software). One-way ANOVA, Student's t-test, one sample t-tests and Fisher's exact tests were performed for statistical analysis. A value of $p < 0.05$ was considered statistically significant.

Results

Expression of WDPCP and the cilia beating function were decreased in nasal polyps

In our previous study, we have previously discovered that the expression of WDPCP was reduced in the epithelial layer of mucosa from patients with CRS compared with control subjects, accompanied with cilia loss[18]. According to immunohistochemical staining, we discovered that WDPCP was mainly expressed in the cytoplasm of nasal mucosa epithelium and the staining (brown color) was reduced in

nasal polyps (Figure 1A). Western blot also showed decreased protein expression of WDPCP in nasal polyp tissue compared with ethmoid sinus tissue from control subjects (Figure 1B-C). Consistent with the protein levels, RT-PCR demonstrated a decrease of WDPCP mRNA levels in nasal polyps (Figure 1D). Moreover, to explore the function of cilia, we measured the cilia beating frequency (CBF), and found that the cilia from nasal polyps beat at a slower rate compared with control mucosa (Figure 1E). Additional movie files show the cilia beating in more detail (see Additional file 1-2).

WDPCP potentially acted through the MAPK/ERK pathway to regulate ciliogenesis and cilia function

Evidence showed that WDPCP participated in ciliogenesis and could possibly affect the beating of cilia[16]. To investigate the potential mechanism, we detected the changes of expression in the MAPK/ERK pathway. We silenced the expression of WDPCP in HSECs by transferring small interfering RNAs (Si-WDPCP). Twenty-four hours post-transfection, the expression of phosphorylated ERK1/2 (p-ERK1/2) was significantly decreased in HSECs compared to the control. However, the expression of MEK1/2, p-MEK1/2 and ERK1/2 remained constant after WDPCP silencing (Figure 2A-B).

HSECs could differentiate into respiratory epithelium when they were grown in an air-liquid interface (ALI). It took approximately 14 days to obtain fully mature cilia. This ALI culture was also used to explore the role of WDPCP and the MAPK/ERK pathway in ciliogenesis and cilia beating. The cilium in HSEC ALI cultures was detected on Day-14 by scanning electron microscopy and immunofluorescence staining. HSECs which lacked WDPCP at the beginning of ciliogenesis, differentiated into fewer and shorter cilia compared with the controls (Figure 2C-D). Similarly, HSECs cultured with MAPK/ERK inhibitor U0126 also developed fewer and shorter cilia (Figure 2C-D). We also examined the cilia beating frequency (CBF) on Day-14. HSECs treated with Si-WDPCP and U0126 displayed slower cilia beating frequencies compared with controls (Figure 2E). Additional movie files show the cilia beating of HSECs in more detail (see Additional file 3-5).

WDPCP regulated mitochondrial biogenesis through the MAPK/ERK pathway

Substantial evidence suggested that ATP from mitochondria is essential for the beating of cilia[26-28]. Thus, we hypothesized that WDPCP acts on mitochondria through the MAPK/ERK pathway to affect cilia beating in CRS. First, we measured the mitochondrial DNA (mtDNA) copy number to observe the number of mitochondria in HSECs. We found that the mtDNA copy numbers were decreased in HSECs after WDPCP silencing or after being cultured with MAPK/ERK pathway inhibitor U0126. Glucocorticoids are known to be the mainstay of medical treatment for CRS. Based on previous research, we explored the

effects of glucocorticoids on the mitochondria. mtDNA copy numbers were not significantly decreased after treatment with dexamethasone in Si-WDPCP HSECs and HSECs inhibited by U0126 (Figure 3A).

Decreased mitochondrial biogenesis or increased apoptosis can lead to the reduction in the number of mitochondria. To understand the specific reasons of mitochondrial reduction, we examined the expression of biomarkers for both mitochondrial biogenesis and apoptosis. Western blot results revealed that the apoptosis markers, Cleaved Caspase 3 and Cleaved Caspase 7 were not activated in HSECs which lacked WDPCP nor HSECs inhibited by U0126 (Figure 3B). However, we found the low expression of NRF1, NRF2, TFAM and COX4 in Si-WDPCP HSECs compared with controls (Figure 3C-D). Interestingly, NRF1 protein levels were increased after dexamethasone treatment in Si-WDPCP HSECs (Figure 3C-D). Additionally, the protein expression of NRF1, NRF2 and TFAM could be inhibited by U0126 (Figure 3E-F). Negative control siRNAs (Si-NC) were used to rule out the effect of transfection on mitochondria. Moreover, the decreased expression of NRF1, NRF2, TFAM and COX4 was also observed in the nasal epithelium and nasal polyps by immunohistochemical staining (Figure 3G).

The temporal mitochondrial biogenesis in the ALI culture of human sinonasal epithelial cells

The temporal mitochondrial biogenesis in the ALI cultures of HSECs was established to explore the changes in mitochondrial biogenesis during ciliogenesis. Western blot results showed that NRF1 and NRF2 mRNA and protein expression increased on Day-7 compared with Day-0, and then decreased after 14 days in the control ALI culture. In contrast, Si-WDPCP HSECs and U0126-treated HSECs demonstrated a decreased expression of NRF1 and NRF2 mRNA and protein on Day-7. TFAM protein levels continued to rise on Day-7 and Day-14 in the control ALI culture. Although similar TFAM expression trends were observed in the ALI culture of Si-WDPCP HSECs and U0126-treated HSECs, there was a decrease in the mRNA level in U0126-cultured HSECs on Day-14 and the protein level in Si-WDPCP HSECs on Day-7 compared with controls. In the control group, COX4 expression was significantly increased on Day-7. However, the COX4 gene and protein levels were reduced in Si-WDPCP HSECs on Day-7 and Day-14, and COX4 protein expression was also decreased in U0126-treated HSECs on Day-7 (Figure 4).

WDPCP stabilized mitochondrial function through the MAPK/ERK pathway

In addition to the number of mitochondria, the function of mitochondria is also critical for the beating of cilia[29]. Flow cytometry analysis revealed a depolarization of the mitochondrial membrane in Si-WDPCP HSECs and U0126-treated HSECs (Figure 5A-B). We also measured mitochondrial function by analyzing ATP production, and we found that ATP levels were reduced in Si-WDPCP HSECs and U0126-treated HSECs (Figure 5C). Interestingly, dexamethasone could repair the damage of mitochondrial function caused by the lack of WDPCP as well as inhibition by U0126 (Figure 5A-C). To monitor mitophagy in the HSECs, mitochondria were visualized with fluorescent MitoTracker. From the MitoTracker staining, rod-

shaped mitochondria can be detected in the control and Si-NC group, but not in the Si-WDPCP or U0126 group. Despite the use of dexamethasone, the mitochondrial structure was difficult to recover (Figure 5D).

Discussion

Patients with CRS were subjected to repeated cycles of inflammation and infection, which resulted in severe cilia loss and increased mucus secretion. In addition to direct ciliary loss, the surviving cilia that experienced inflammatory and/or microbial damage appeared to be dysfunctional. However, the reason behind poor ciliation and cilia dysfunction is still unclear. PCP pathway proteins, such as Intu, Vangl2 and WDPCP were important units which controlled ciliogenesis and cilia function. Studies showed that WDPCP governs ciliogenesis in *Xenopus* embryos[14, 17] and human nasal epithelium[18]. Although short cilia might contribute to cilia beating dysfunction[30], it is still unclear whether WDPCP directly regulated the beating of cilia. To our knowledge, this is the first study of WDPCP on cilia function in human sinonasal epithelium. Our data showed that the low expression of WDPCP in nasal polyps was accompanied with a decreased frequency of cilia beating. Following WDPCP silencing in ALI cultures of HSECs, there was a decrease in cilia beating frequency. These data suggest that in addition to its role in the ciliogenesis of HSECs, WDPCP is also critical to cilia beating function.

Mitochondria are highly dynamic organelles that are essential for energy production and cell homeostasis. Mitochondrial function modulates cell survival, metabolism and health status. The impairment of mitochondrial function has been involved in a variety of pathological conditions and diseases. For example, mitochondrial dysfunction could provoke heterotaxy via aberrant ciliogenesis and reduced the beating of cilia[29]. Our finding showed that the expression of mitochondrial biogenesis markers such as NRF1, NRF2, TFAM and COX4 were decreased in the epithelium of nasal polyp. HSECs which lacked the expression of WDPCP had decreased mtDNA copy number, low expression of NRF1, NRF2, TFAM and COX4, and poor mitochondrial function and structure. However, the mitochondrial apoptosis markers were not activated. Overall, these findings support the notion that WDPCP regulated cilia function from its effects on the mitochondria.

The MAPK/ERK pathway regulates a series of cellular biological behaviors. Although the importance of MAPK/ERK activity in morphogenesis and metabolism is emerging[31], the role of the MAPK/ERK pathway in ciliogenesis and the function of human airway epithelium is still unknown. Our results showed that after WDPCP silencing, the expression of p-ERK in HSECs was decreased. Also, when MAPK/ERK activity was inhibited by U0126, HSECs developed fewer and shorter cilia in ALI cultures, a phenotype reminiscent of WDPCP deficiency. Moreover, in the absence of MAPK/ERK activity, the mtDNA copy number and the expression of NRF1, NRF2, and TFAM were decreased in HSECs, and the structure of mitochondria was altered. Collectively, these results demonstrate that WDPCP acts through the MAPK/ERK pathway to regulate ciliogenesis and mitochondrial biogenesis. Consistent with our findings, the role of MAPK/ERK activity in cellular differentiation has already been investigated. For example, MAPK/ERK activation was involved in the differentiation of nephron progenitors by the Wnt and Notch

pathways[32], in addition to the crucial role of the MAPK/ERK pathway in airway development[31]. Our findings highlight the importance of MAPK/ERK activity in airway ciliogenesis and cilia function.

Glucocorticoids and their receptors could modulate mitochondrial function. Glucocorticoids activated nuclear-encoded genes, which resulted in enhanced mitochondrial biogenesis[33, 34]. Additionally, glucocorticoids could directly regulate mitochondrial function by interpolating into the mitochondrial membrane[35, 36]. Hence, we speculated that glucocorticoids could also repair mitochondrial damage in airway diseases. In order to address this hypothesis, we examined the expression of mitochondrial biogenesis markers and mitochondrial function after treatment with dexamethasone in Si-WDPCP HSECs and U0126-treated HSECs. Interestingly, we found that the decreased NRF1 in HSECs which lacked WDPCP expression could be repaired by dexamethasone. However, dexamethasone could not repair the mitochondrial structure. Taken together, dexamethasone could partially recover the damage on mitochondrial function in HSECs which lacked WDPCP expression or HSECs with an inactivated MAPK/ERK pathway. These findings provide a new insight into the mechanism underlying glucocorticoid treatment of CRS.

Conclusions

In summary, WDPCP regulates the mitochondrial biogenesis and mitochondrial function through the MAPK/ERK pathway to affect cilia beating function. Dexamethasone could partially repair the mitochondrial damage. Further investigation of PCP proteins may offer a deeper understanding of the pathophysiology and treatment of CRS.

Abbreviations

CRS

chronic rhinosinusitis

CBF

cilia beating frequency

NP

nasal polyps

HSECs

human sinonasal epithelial cells

PCP

planar cell polarity

WDPCP

tryptophan-aspartic acid (W-D) repeat containing planar cell polarity effector

MAPK

mitogen activated protein kinase

ERK

signal-regulated kinase

MEK1/2

dual-specificity mitogen-activated protein kinases kinases 1/2

NRF1

nuclear respiratory factor 1

NRF2

nuclear respiratory factor 2

TFAM

transcription factor A, mitochondrial

COX4

cytochrome c oxidase subunit 4

Declarations

Ethics approval and consent to participate

This study was approved by the Sun Yat-sen Memorial Hospital Institutional Review Board. Written consent has been obtained from all participants.

Consent for publication

All participants signed their written consent for publication.

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Competing interests

The authors declare that there is no potential conflict of interest.

Funding

Yat-sen Scientific Research Startup Foundation, the Young Investigator Award, No. YXQH201914. For Yun Ma

National Natural Science Foundation of China, the Young Investigator Award, No. 81500773. For Peng Tian

Authors' Contributions

Conceived and designed the experiments: Yun Ma, Hua Zou, Yiqing Zheng

Performed the experiments: Yun Ma and Peng Tian did experiments on Western blot, real-time PCR and cell culture; Hua Zhong performed tests on cilia beating frequency and immunostaining; Fan Wu and Qining Zhang were responsible for mitochondrial related experiments.

Analyzed the data and wrote the manuscript: Yun Ma, Hua Zhong

Recruited patients and collected specimen: Xiang Liu, Dang Hua, Qiu Jian Chen and Hua Zou

Acknowledgements

We thank Prof. Yinyan Lai for her help in measurement of cilia beating.

References

1. Hastan D, Fokkens WJ, Bachert C, Newson RB, Bislimovska J, Bockelbrink A, et al. Chronic rhinosinusitis in Europe—an underestimated disease. A GA(2)LEN study. *Allergy*. 2011;66:1216–23.
2. Blackwell DL, Lucas JW, Clarke TC. Summary health statistics for U.S. adults: national health interview survey, 2012. *Vital Health Stat*. 2014;10:1–161.
3. Shi JB, Fu QL, Zhang H, Cheng L, Wang YJ, Zhu DD, et al. Epidemiology of chronic rhinosinusitis: results from a cross-sectional survey in seven Chinese cities. *Allergy*. 2015;70:533–9.
4. Kim JH, Cho C, Lee EJ, Suh YS, Choi BI, Kim KS. Prevalence and risk factors of chronic rhinosinusitis in South Korea according to diagnostic criteria. *Rhinology*. 2016;54:329–35.
5. Ferguson JL, McCaffrey TV, Kern EB, Martin WJ 2. The effects of sinus bacteria on human ciliated nasal epithelium in vitro. *Otolaryngol Head Neck Surg*. 1988;98:299–304. nd .
6. Lennard CM, Mann EA, Sun LL, Chang AS, Bolger WE. Interleukin-1 beta, interleukin-5, interleukin-6, interleukin-8, and tumor necrosis factor-alpha in chronic sinusitis: response to systemic corticosteroids. *Am J Rhinol*. 2000;14:367–73.
7. Elliott MK, Sisson JH, West WW, Wyatt TA. Differential in vivo effects of whole cigarette smoke exposure versus cigarette smoke extract on mouse ciliated tracheal epithelium. *Exp Lung Res*. 2006;32:99–118.

8. Schipor I, Palmer JN, Cohen AS, Cohen NA. Quantification of ciliary beat frequency in sinonasal epithelial cells using differential interference contrast microscopy and high-speed digital video imaging. *Am J Rhinol.* 2006;20:124–7.
9. Sutto Z, Conner GE, Salathe M. Regulation of human airway ciliary beat frequency by intracellular pH. *J Physiol.* 2004;560:519–32.
10. Gudis D, Zhao KQ, Cohen NA. Acquired cilia dysfunction in chronic rhinosinusitis. *Am J Rhinol Allergy.* 2012;26:1–6.
11. Wallingford JB, Rowning BA, Vogeli KM, Rothbacher U, Fraser SE, Harland RM. Dishevelled controls cell polarity during *Xenopus* gastrulation. *Nature.* 2000;405:81–5.
12. Jessen JR, Topczewski J, Bingham S, Sepich DS, Marlow F, Chandrasekhar A, et al. Zebrafish trilobite identifies new roles for Strabismus in gastrulation and neuronal movements. *Nat Cell Biol.* 2002;4:610–5.
13. Gray RS, Abitua PB, Wlodarczyk BJ, Szabo-Rogers HL, Blanchard O, Lee I, et al. The planar cell polarity effector Fuz is essential for targeted membrane trafficking, ciliogenesis and mouse embryonic development. *Nat Cell Biol.* 2009;11:1225–32.
14. Park TJ, Kim SK, Wallingford JB. The planar cell polarity effector protein Wdpcp (Fritz) controls epithelial cell cortex dynamics via septins and actomyosin. *Biochem Biophys Res Commun.* 2015;456:562–6.
15. Sittaramane V, Pan X, Glasco DM, Huang P, Gurung S, Bock A, et al. The PCP protein Vangl2 regulates migration of hindbrain motor neurons by acting in floor plate cells, and independently of cilia function. *Dev Biol.* 2013;382:400–12.
16. Kim SK, Shindo A, Park TJ, Oh EC, Ghosh S, Gray RS, et al. Planar cell polarity acts through septins to control collective cell movement and ciliogenesis. *Science.* 2010;329:1337–40.
17. Cui C, Chatterjee B, Lozito TP, Zhang Z, Francis RJ, Yagi H, et al. Wdpcp, a PCP protein required for ciliogenesis, regulates directional cell migration and cell polarity by direct modulation of the actin cytoskeleton. *PLoS Biol.* 2013;11:e1001720.
18. Ma Y, Sun Y, Jiang L, Zuo K, Chen H, Guo J, et al. WDPCP regulates the ciliogenesis of human sinonasal epithelial cells in chronic rhinosinusitis. *Cytoskeleton (Hoboken).* 2017;74:82–90.
19. Seminario-Vidal L, Okada SF, Sesma JI, Kreda SM, van Heusden CA, Zhu Y, et al. Rho signaling regulates pannexin 1-mediated ATP release from airway epithelia. *J Biol Chem.* 2011;286:26277–86.
20. Cloonan SM, Glass K, Laucho-Contreras ME, Bhashyam AR, Cervo M, Pabon MA, et al. Mitochondrial iron chelation ameliorates cigarette smoke-induced bronchitis and emphysema in mice. *Nat Med.* 2016;22:163–74.
21. Liu G, Summer R. Cellular Metabolism in Lung Health and Disease. *Annu Rev Physiol.* 2019;81:403–28.
22. Liu X, Wang X, Chen L, Shi Y, Wei Y. Effects of Erythromycin on the Proliferation and Apoptosis of Cultured Nasal Polyp-Derived Cells and the Extracellular Signal-Regulated Kinase (ERK)/Mitogen-Activated Protein Kinase (MAPK) Signaling Pathway. *Med Sci Monit.* 2018;24:8048–55.

23. Zhang J, Wang L, Xie W, Hu S, Zhou H, Zhu P, et al. Melatonin attenuates ER stress and mitochondrial damage in septic cardiomyopathy: A new mechanism involving BAP31 upregulation and MAPK-ERK pathway. *J Cell Physiol.* 2019.
24. Fokkens WJ, Lund VJ, Mullol J, Bachert C, Alobid I, Baroody F, et al. European Position Paper on Rhinosinusitis and Nasal Polyps 2012. *Rhinol Suppl.* 2012;23:3 p preceding table of contents, 1-298.
25. Kempeneers C, Seaton C, Garcia Espinosa B, Chilvers MA. Ciliary functional analysis: Beating a path towards standardization. *Pediatr Pulmonol.* 2019;54:1627–38.
26. Fliegauf M, Benzing T, Omran H. When cilia go bad: cilia defects and ciliopathies. *Nat Rev Mol Cell Biol.* 2007;8:880–93.
27. Auguste Y, Delague V, Desvignes JP, Longepied G, Gnisci A, Besnier P, et al. Loss of Calmodulin- and Radial-Spoke-Associated Complex Protein CFAP251 Leads to Immotile Spermatozoa Lacking Mitochondria and Infertility in Men. *Am J Hum Genet.* 2018;103:413–20.
28. Workman AD, Carey RM, Chen B, Saunders CJ, Marambaud P, Mitchell CH, et al. CALHM1-Mediated ATP Release and Ciliary Beat Frequency Modulation in Nasal Epithelial Cells. *Sci Rep.* 2017;7:6687.
29. Burkhalter MD, Sridhar A, Sampaio P, Jacinto R, Burczyk MS, Donow C, et al. Imbalanced mitochondrial function provokes heterotaxy via aberrant ciliogenesis. *J Clin Invest.* 2019;129:2841–55.
30. Bottier M, Thomas KA, Dutcher SK, Bayly PV. How Does Cilium Length Affect Beating? *Biophys J.* 2019;116:1292–304.
31. Boucherat O, Nadeau V, Berube-Simard FA, Charron J, Jeannotte L. Crucial requirement of ERK/MAPK signaling in respiratory tract development. *Development.* 2014;141:3197–211.
32. Ihermann-Hella A, Hirashima T, Kupari J, Kurtzeborn K, Li H, Kwon HN, et al. Dynamic MAPK/ERK Activity Sustains Nephron Progenitors through Niche Regulation and Primes Precursors for Differentiation. *Stem Cell Reports.* 2018;11:912–28.
33. Goffart S, Wiesner RJ. Regulation and co-ordination of nuclear gene expression during mitochondrial biogenesis. *Exp Physiol.* 2003;88:33–40.
34. Psarra AM, Sekeris CE. Glucocorticoids induce mitochondrial gene transcription in HepG2 cells: role of the mitochondrial glucocorticoid receptor. *Biochim Biophys Acta.* 2011;1813:1814–21.
35. Sionov RV, Kfir S, Zafir E, Cohen O, Zilberman Y, Yefenof E. Glucocorticoid-induced apoptosis revisited: a novel role for glucocorticoid receptor translocation to the mitochondria. *Cell Cycle.* 2006;5:1017–26.
36. Lee SR, Kim HK, Song IS, Youm J, Dizon LA, Jeong SH, et al. Glucocorticoids and their receptors: Insights into specific roles in mitochondria. *Prog Biophys Mol Bio.* 2013;112:44–54.

Table 1

Table 1 Patients' characteristics

	Control subjects (n=10)	CRSwNPs (n=15)	p Value
Gender(M/F)	6/4	6/9	NS
Mean age (Y)	30.8±2.53	45.67±3.63	0.0061
Smoker	3	8	NS
Allergy(a)	0	5	NS
Asthma(b)	0	3	NS
Inflammatory cell count from peripheral blood			
Neutrophilic (×10 ⁹ /L)	8.20±0.57	7.06±0.51	NS
Eosinophilic (×10 ⁹ /L)	0.19±0.05	0.39±0.07	0.044

M, male; F, Female; Y, Years; NS, not significant.

(a) Allergy was diagnosed by specific IgE level test from peripheral blood.

(b) Physician-diagnosed asthma with treatment of β_2 -agonist (n=2) or β_2 -agonist plus glucocorticoid inhaler (n=1).

Figures




Figure 1

Epithelium of nasal polyps demonstrated a low expression of WDPCP and had decreased cilia beating frequency compared with control subjects. (A) The expression of WDPCP in the cytoplasm of epithelial cells from nasal polyps (NP) was decreased as determined by immunohistochemical staining (brown color). Scale bar: 50 μ m. (B, C) Western blot images and densitometry showed decreased WDPCP protein levels in NP compared with nasal mucosa from control subjects (n=3). (D) The copy number of WDPCP

mRNA was decreased in NP (n=3). (E) Cilia beating frequency (CBF) was significantly reduced in NP compared with control mucosa (n=5).




Figure 2

WDPCP regulated ciliogenesis and cilia beating through the MAPK/ERK pathway in HSECs. (A-B) The expression of p-ERK1/2 was decreased in Si-WDPCP HSECs (n=3) by Western blot and densitometry analysis. (C) Immunofluorescence images showed the decrease of cilia length and quantity in HSECs

which lacked WDPCP and U0126-treated HSECs. Scale bar: 20 μ m. (D) SEM images also showed that the cilia length and quantity was decreased in HSECs which lacked WDPCP and U0126-treated HSECs. Scale bar: 2 μ m. (E) Cilia beating frequency was reduced on Day-14 in HSECs which lacked WDPCP and U0126-treated HSECs (n=4).




Figure 3

WDPCP silencing or inhibition of the MAPK/ERK pathway could reduce mitochondrial biogenesis. (A) mtDNA copy number was decreased in Si-WDPCP HSECs and U0126-treated HSECs (n=3). (B) WDPCP silencing or the inhibition of the MAPK/ERK pathway did not activate mitochondrial apoptosis. (C-D) NRF1, NRF2, TFAM and COX4 protein levels were reduced after WDPCP silencing in HSECs by Western blot and densitometry analysis. The reduction of NRF1 could be recovered by dexamethasone (n=3). (E-F) NRF1, NRF2 and TFAM protein levels were reduced in U0126-treated HSECs (n=3). (G) Mitochondrial biogenesis markers NRF1, NRF2, TFAM and COX4 had low expressions in the epithelium of nasal polyps by immunohistochemical staining. Scale bar: 50 μ m.




Figure 4

Expression of mitochondrial biogenesis makers in ALI cultures of HSECs. (A) NRF1, NRF2 and COX4 mRNA levels increased over 7 days in ALI cultures, but this increase was lesser in Si-WDPCP HSECs and U0126-treated HSECs. TFAM mRNA levels continued to increase over 14 days in ALI cultures, whereas the mRNA expression was reduced in U0126-treated HSECs on Day-14 (n=3). (B-C) NRF1, NRF2 and COX4 protein levels also increased over 7 days in control ALI cultures, and the expression was decreased in Si-

WDPCP HSECs and U0126-treated HSECs on Day-7. TFAM protein levels continued to increase over 14 days in ALI cultures, and its protein expression was decreased in Si-WDPCP HSECs on Day-14 (n=3). (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$)




Figure 5

WDPCP impaired mitochondrial function in HSECs by the MAPK/ERK pathway. (A-B) The mitochondrial membrane potential was decreased in Si-WDPCP HSECs and U0126-treated HSECs as observed by JC-1

fluorescent dye analysis, and the mitochondrial membrane potential could be partially restored after adding dexamethasone (n=5). (C) ATP levels were decreased in Si-WDPCP HSECs and U0126-treated HSECs, after adding dexamethasone, the ATP levels increased (n=5). (D) MitoTracker staining showed that rod-shaped mitochondria were decreased in Si-WDPCP HSECs and U0126-treated HSECs, and the mitochondrial morphology cannot be restored by dexamethasone. Scale bar: 8 μ m.




Figure 6

Schematic diagram of WDPCP regulating ciliogenesis and cilia function in nasal epithelium. Our results showed that WDPCP could work through MAPK/ERK pathway to regulate mitochondrial biogenesis and function. In the case of CRS, HSECs low expressed WDPCP, and phosphorylation of ERK1/2 was partially blocked, further affecting the number and function of mitochondria. This mechanism might lead to the poor ciliation and cilia dysfunction of nasal epithelium in CRS.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Video5U0126.mp4](#)
- [Video1controltissue.mp4](#)
- [Video4SiWDPCP.mp4](#)
- [Video3ControlALI.mp4](#)

- [Video2NP.mp4](#)