Incomplete Staging in Ovarian Cancer

Jogchum Beltman (✉ j.j.beltman@lumc.nl)

Leids Universitair Medisch Centrum <https://orcid.org/0000-0002-0048-7360>

P Laven

Maastricht Universitair Medisch Centrum+

JE Bense

Leids Universitair Medisch Centrum

Maaïke van der Aa

National Institute of Cancer Prevention and Research

MC Vos

Elisabeth-TweeSteden Ziekenhuis

D Boll

Catharina Ziekenhuis

HGJ Arts

Universitair Medisch Centrum Groningen

N Reesink

Medisch Spectrum Twente

JB Trimpos

Leids Universitair Medisch Centrum

RFPM Kruitwagen

Maastricht Universitair Medisch Centrum+

Research

Keywords: epithelial ovarian cancer, surgical oncology, lymph nodes, biopsy

Posted Date: April 23rd, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-23321/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Objective Incomplete surgical staging of patients with early stage epithelial ovarian cancer (EOC) has been reported in up to 85% of cases, when based on the International Federation of Obstetrics and Gynaecology (FIGO) staging procedure. The aim of the present retrospective study was to clarify the reasons for incomplete staging. **Methods** The PRISMA (Prevention Recovery Information System for Monitoring and Analysis) technique was used to evaluate cases with FIGO I-IIa EOC based on incomplete staging from five gynecologic oncologic center hospitals in the Netherlands in the period 2010-2014. **Results** Fifty cases with an incomplete surgical staging of EOC according to national guidelines were included. The most common reasons for incomplete staging were insufficient random biopsies of the peritoneum (n=34, 68%), and less than ten lymph nodes being resected and/or found at pathology (n=16, 32%). The most mentioned reasons for not performing biopsies were forgetting to do so, and believing that after careful inspection and palpation, taking biopsies is irrelevant and/or already are being taken while performing a hysterectomy (cul-de-sac, bladder). The value of contralateral pelvic lymph node dissection in case of a unilateral ovarian malignancy was also doubted, influencing the number of lymph nodes resected. **Conclusion** The most important reasons for incomplete staging in EOC are, besides omitting elements by accident, questioning the importance of obligatory elements of the staging procedure. A structured list of staging steps during surgery and more evidence based consensus concerning these obligatory elements might increase the number of complete staging procedures in EOC.

Introduction

In newly diagnosed clinical early stage epithelial ovarian cancer (EOC), the aim of surgery is to resect the tumor combined with an adequate staging procedure. This will provide prognostic information by means of the definite stage and will define whether adjuvant chemotherapy is needed or can be safely omitted [1]. According to the International Federation of Obstetrics and Gynaecology (FIGO) a staging procedure should include a peritoneal washing or a sample of ascites, hysterectomy and bilateral oophorectomy, infracolic omentectomy (complete, partial or biopsies), numerous peritoneal biopsies from several locations at risk for tumor implantation, pelvic and para-aortic lymphadenectomy and biopsies of suspected lesions and adhesions [2]. The Dutch guideline has incorporated these staging elements with the adjustment that overall at least 10 lymph nodes should be resected from specific locations in the pelvic and paraaortic area .

Despite the FIGO recommendation, incomplete surgical staging has been reported in up to 85% of cases in different studies [3–6]. Studies examining the causes of incomplete staging are scarce and often based on assumptions such as lack of specialized skills and/or knowledge of the gynecologist [3,5,6].

The Prevention Recovery Information System for Monitoring and Analysis (PRISMA) was originally developed to manage human error in the chemical process industry, but has also been applied in healthcare (PRISMA-Medical) [7,8]. The main goal of PRISMA is to build a quantitative database of

incidents and process deviations, in order to facilitate the development and evaluation of system-based preventive strategies.

In this study, we used PRISMA to study the causes of incomplete staging in EOC.

Materials And Methods

Five hospitals participated in the study and all hospitals fulfilled the Dutch criteria to be a gynecologic oncology center within their region and performing 20 or more debulking procedures each year. Patients with EOC FIGO stage I – IIa undergoing a staging procedure in the period from 2010 until 2014 were included if one of the following incidences had occurred: no sampling of ascites or peritoneal washing, less than five peritoneal biopsies resection of less than ten lymph nodes; no high para-aortal, ipsi- and/or contralateral pelvic lymph nodes; no omentum biopsy or resection.

Cases were selected retrospectively from the Netherlands Cancer Registry (NCR). The NCR is maintained by the Netherlands Comprehensive Cancer Organization and contains data of all cancer patients in the Netherlands. The NCR, which reached full national coverage in 1989, relies on notification of all newly diagnosed malignancies via the automated nationwide pathology archive. Trained registrars use standardized forms to collect patient information from medical records, and regular consistency checks are performed to ensure the quality of the data held in the NCR. The following data were examined in this study: age; histological subtype; tumor stage; differentiation grade; inspection and palpation of all peritoneal surfaces; biopsies of any suspect lesions for metastases; biopsies or removal of any adhesions surrounding the primary tumor; peritoneal washing or ascites; infracolic omentectomy or biopsy; biopsies of peritoneum of the bladder, cul-de-sac, pelvic sidewalls, ovarian fossa, right and left paracolic gutter, and right hemi-diaphragm; number of lymph nodes removed; regions of lymph nodes removed (high para-aortal / paracaval, ipsilateral pelvic, contralateral pelvic). All surgery was performed by or under supervision of qualified gynecologic oncologists.

The PRISMA commission consisted of eight gynecologists, a member of the NCR (MA), and a medical researcher. All included cases were analyzed and discussed with the gynecologic oncologists of each participating hospital. All members participating in the commission signed a non-disclosure agreement.

Results

Study population

Figure 1 shows the study population and reasons for exclusion. Fifty-eight patients were selected by the NCR data managers. One selected patient had advanced stage disease at pathology (IIIA1 based on a lymph node metastasis) and was excluded. After analyzing and discussing the patients with the gynecologists of each participating hospital, seven patients (12%) appeared to have had a complete staging after all, but were interpreted differently during selection by the NCR data managers. In two of these cases the location of the aortocaval lymph node dissection was not well described, and in five

cases the gynecologist used anatomical terms describing the location, that were unfamiliar to the data managers. Therewith, 50 cases remained for inclusion in this study.

Most patients (n = 34, 68%) underwent one operation. In 15 patients (30%) a second operation was performed, and in one patient (2%) despite three surgical procedures the staging procedure still was incomplete.

Nineteen patients (38%) received adjuvant chemotherapy. In two cases, it was unknown whether adjuvant chemotherapy was given (4%) and one patient refused chemotherapy. For the remaining 29 patients (58%), according to the gynecologic oncologist, there was no indication for adjuvant chemotherapy.

Incomplete staging

The number of missing elements was one in 12 patients (24%), two in 17 patients (34%), three in eight patients (16%), four in four patients (8%), five in four patients (8%), and six or more in the remaining five patients (10%).

Table 1 shows incompleteness of the staging procedure with respect to the different elements. The most frequent reason for incomplete staging was insufficient peritoneal biopsies of the different locations (n = 34; 68%). Reasons for not performing certain biopsies were 1) forgetting to do so (n = 11), 2) presumption that the biopsy (cul-de-sac and bladder peritoneum) is already part of the hysterectomy (n = 5), 3) the opinion of the surgeon that after careful inspection taking biopsies of macroscopically normal peritoneum is irrelevant (n = 6), 4) chemotherapy already being indicated, hence complete staging would not be contributory (n = 6), 5) technical problems (n = 3), 6) comorbidity or malignancy elsewhere (n = 2), and 7) the wish of the patient to not receive adjuvant therapy (n = 1) (Fig. 2).

In eight patients (16%) no lymph nodes were resected, either because it was technically impossible (n = 4) or because the outcome of the lymph nodes, according to the gynecologist, had no therapeutic consequences (n = 4, Fig. 2). In 25 patients (50%) no contralateral pelvic lymph nodes were resected. In eight patients (16%) the number of resected lymph nodes was less than ten. This could be partly explained by the fact that in five of these patients only lymph nodes were resected on the ipsilateral pelvic region. Furthermore, although according to the gynecologist enough tissue was resected from the para-aortic region, the pathologist did not identify any lymph nodes in one patient (2%). Taking ascites or peritoneal washing was omitted by accident in 12 patients (24%), taking biopsies from adhesions surrounding the tumor was omitted in 14%, a omentum biopsy or omentectomy in 2%, and a biopsy of suspicious lesions in 2% of cases. A hysterectomy or contralateral ovariectomy was not performed in five cases (10%) to preserve fertility or because of the wish of the patient to not receive adjuvant chemotherapy, and therefore a complete staging was not performed.

Discussion

In this retrospective analysis we found that incomplete staging in EOC, performed by gynecologic oncologists, is most often due to insufficient biopsies of different locations of the peritoneum (n = 34;

68%), and less than ten lymph nodes being resected and/or found at pathology (n = 16; 32%). The most mentioned reasons for not performing biopsies were forgetting to do so, and believing that, after careful inspection and palpation, taking biopsies is irrelevant and/or biopsies already are being taken while performing a hysterectomy (peritoneum of the cul-de-sac, broad ligament and bladder). Forgetting to take blind biopsies was registered in 11 cases and there can be little excuse to justify such negligence.

Previous studies, mainly focused on patients treated before 2000, report incidences of incomplete staging up to 85% [3,5,6,9]. In the majority of these patients no or inadequate lymph node resection was performed [3]. But also no or an insufficient number of random peritoneal biopsies was frequently found. These studies conclude that lack of specialized skills and/or knowledge of the gynaecologist might be a possible explanation for incomplete staging [3,5,6,10]. However, in the present study only gynaecologic oncological center hospitals were included, (of which one can assume that they do have the skills and knowledge that is required), taking blind biopsies was simply forgotten in about one third (11/34) of the incomplete biopsy cases. A standardized surgical report, and check lists with all parts of the staging procedure according to the FIGO guidelines available in the operation rooms, might be a solution to forgetting parts of the procedure.

The assumption that taking blind biopsies is unnecessary is a different matter. In a recent review on surgical staging of EOC the average yield of positive blind biopsies of the peritoneum in a number of studies was around 8% [11]. It has also been argued that upstaging of EOC on the basis of positive blind biopsies is lower but this would be true for all the different staging steps and data on this specific issue are scarce [11]. Ayhan et al. found microscopic disease in random peritoneal biopsies or appendectomy in 7.1% of 127 EOC patients [12]. Shroff et al. had similar results in that microscopically positive biopsies from normal-appearing tissue were found in 6/122 patients (5%) [13]. Five of these six patients were upstaged by random biopsies alone. Taking a more general perspective over the debate of percentages it has been argued that random peritoneal biopsies remain indicated in early-stage disease owing to the low morbidity of the procedure and the present possibility of upstaging and altered management. Furthermore, systematic peritoneal biopsies ensure careful palpation and examination of all surfaces. The latter notion is the more important in view of the recent trend to perform staging of EOC laparoscopically, therewith losing the contribution of palpating subperitoneal abnormalities.

Adequate lymph node sampling appears to be an essential part of the staging procedure in clinical early stage EOC. In a review on lymph node metastases, the overall incidence varied from 6.1–29.6% (mean 14.2%) [14]. Despite its importance, the reported incidence of lymph node staging worldwide varies between 10% and 30% [15]. In a recent report on lymph node staging in The Netherlands, the incidence of lymph node dissection improved from 26% in 2000 to 67% in 2012. Moreover, the percentage of patients from whom 10 or more lymph nodes were removed also increased during the study period (from 2.3–47.6%) [16].

In the present study, in 16 patients with an incomplete staging procedure (32%), either no lymph nodes were resected or the number of lymph nodes identified was below ten. Various reasons for this were

mentioned by the gynecologists. First of all, in five patients no lymph nodes were resected because it had no therapeutic consequences according to the gynecologist (already indication for adjuvant chemotherapy). In another four patients it was mentioned that the procedure was technically impossible, mainly related to intense intra-abdominal adhesions, extensive bloodloss and/or obesity. Secondly, although according to the gynecologist enough tissue at the different pelvic and paraaortal locations were sampled, the number of lymph nodes identified by the pathologist were less than ten, hence insufficient according to the Dutch guideline. This could be partly explained by the fact that in five patients pelvic lymph nodes were obtained only at the ipsilateral side. In 25 patients (50%), including these five patients, the value of a contralateral pelvic lymphadenectomy in case of a unilateral ovarian tumor was questioned and therefore omitted. In the already mentioned review examining the incidence and locations of lymph node metastases in early stage EOC, 20% of all lymph node metastases were located at the pelvic regions only. Only contralateral lymph node metastases of a unilateral tumour was found in 16% of positive pelvic nodes and 11% of positive paraaortic nodes [14]. These findings provide a strong argument against an ipsilateral node dissection only.

Besides the resection of not enough fatty tissue containing the lymph nodes by the gynecologist, the pathologist examining the tissue resected may influence the definite number of lymph nodes recognized. In a study that observed the impact of nodal retrieval after educating the pathologist, found that extra education about the importance of nodal count significantly raised the nodal count at pathology [17]. New pathological methods are being investigated and have shown promising results. Svec et al. showed that re-fixing in a lymph node revealing solution containing ethanol, diethyl ether, glacial acetic acid and formalin, increased the number of revealed lymph nodes in colorectal resection specimens [18]. In another study comparing standard formalin fixation with fat-clearing by acetone in specimens after gastrectomy, more lymph nodes were found using acetone, though this was not statistically significant. More high quality trials are needed to study the optimal technique for retrieving lymph nodes at pathology.

Complete staging has been proven to be an important prognostic factor in EOC [4,16]. There is evidence that, in contrast to incompletely staged patients, adjuvant chemotherapy in completely staged patients does not improve survival and should be omitted [4,16,19,20]. This argument makes a comprehensive and complete surgical staging the more important. Our study demonstrates that the staging quality in daily life practice is still too low. We have analyzed the reasons for this and these data might serve to improve the staging performance of EOC. It should be said that the low percentages of complete surgical staging of EOC according to the existing guidelines might be flattered. In our study we found some reasons of the gynecologic oncologists for deliberately omitting certain staging steps in the individual clinical situation, perfectly plausible. In 34 patients the taking of blind peritoneal biopsies was incomplete. Of these, taking the biopsies was considered to have no therapeutic consequences in six patients, there were technical problems in three patients and comorbidity problems or a second malignancy in two. Together they represent one third of the incomplete biopsy group.

In 16 patients the yield of lymph nodes was incomplete with technical problems in four and no therapeutic consequences also in four. To convict these cases as victims of inadequate surgical performance would be unfair. For this reason a plea has been made by Bense et al (2019, in press) to consider cases such as these separately from the categories complete and incomplete staging and the term 'considerate adjustment of staging procedure (CASP)' has been proposed. It might be worthwhile to include this category in future studies on staging of EOC.

Conclusions

The most common reasons for incomplete staging were insufficient random biopsies of the peritoneum, and less than ten lymph nodes being resected and/or found at pathology. The most mentioned reasons for not performing biopsies were forgetting to do so, and believing that, after careful inspection and palpation, taking biopsies is irrelevant and/or already are being taken while performing a hysterectomy (cul-de-sac, bladder). The value of contralateral pelvic lymph node dissection in case of a unilateral ovarian malignancy was also doubted, influencing the number of lymph nodes resected. Introducing a third staging category (considered adjustment of staging procedure) might give a more realistic picture of the actual staging performance. A structured list of staging steps to be used during surgery and evidence based consensus of obligatory staging steps might help in achieving a higher rate of complete staging procedures.

Declarations

Ethics approval and consent to participate. The PRISMA commission consisted of eight gynecologists, a member of the NCR (MA), and a medical researcher. All included cases were analyzed and discussed with the gynecologic oncologists of each of the 5 participating hospitals. All members participating in the commission signed a non-disclosure agreement. Ethical approval was obtained in all hospitals.

Consent for publication: All authors consented for publication

Availability of data and material: Cases were selected retrospectively from the Netherlands Cancer Registry (NCR). The NCR is maintained by the Netherlands Comprehensive Cancer Organization and contains data of all cancer patients in the Netherlands. The NCR, which reached full national coverage in 1989, relies on notification of all newly diagnosed malignancies via the automated nationwide pathology archive. Trained registrars use standardized forms to collect patient information from medical records, and regular consistency checks are performed to ensure the quality of the data held in the NCR.

Conflict of Interest. The authors declare that they have no conflict of interest.

Funding: there was no funding for this study

Author's contributions: JJB,PL, JB, MA, MV, DB, HA, NR and JBT analyzing, reviewing and co-writing. MA selection of cases, analyzing and reviewing, RK design of study, analyzing, co-writing and reviewing.

References

1. Ledermann JA, Raja FA, Fotopoulou C, Gonzalez-Martin A, Colombo N, Sessa C. Newly diagnosed and relapsed epithelial ovarian carcinoma: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Ann Oncol Off J Eur Soc Med Oncol*. 2013 Oct;24 Suppl 6:vi24-32.
2. Benedet JL, Bender H, Jones III H, Ngan HYS, Pecorelli S. FIGO staging classifications and clinical practiceguidelines in the management of gynecologic cancers. *FIGO Comm Gynecol Oncol Int J Gynecol Obstet [Internet]*. 2000;70:209–62. Available from: <https://obgyn.onlinelibrary.wiley.com/doi/pdf/10.1002/ijgo.12611>
3. Timmers PJ, Zwinderman AH, Coens C, Vergote I, Trimbos JB. Understanding the problem of inadequately staging early ovarian cancer. *Eur J Cancer [Internet]*. 2010;46(5):880–4. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/20074933>
4. Zanetta G, Rota S, Chiari S, Bonazzi C, Bratina G, Torri V, et al. The accuracy of staging: an important prognostic determinator in stage I ovarian carcinoma. A multivariate analysis. *Ann Oncol Off J Eur Soc Med Oncol*. 1998 Oct;9(10):1097–101.
5. McGowan L, Leshner LP, Norris HJ, Barnett M. Misstaging of ovarian cancer. *Obstet Gynecol*. 1985 Apr;65(4):568–72.
6. Trimbos JB, Schueler JA, van Lent M, Hermans J, Fleuren GJ. Reasons for incomplete surgical staging in early ovarian carcinoma. *Gynecol Oncol*. 1990 Jun;37(3):374–7.
7. Kaplan HS, Battles JB, Van der Schaaf TW, Shea CE, Mercer SQ. Identification and classification of the causes of events in transfusion medicine. *Transfusion*. 1998;38(11–12):1071–81.
8. Snijders C, van der Schaaf TW, Klip H, van Lingen RA, Fetter WPF, Molendijk A. Feasibility and reliability of PRISMA-medical for specialty-based incident analysis. *Qual Saf Health Care*. 2009 Dec;18(6):486–91.
9. Sijmons EA, van Lankveld MAL, Witteveen PO, Peeters PHM, Koot VCM, van Leeuwen JS. Compliance to clinical guidelines for early-stage epithelial ovarian cancer in relation to patient outcome. *Eur J Obstet Gynecol Reprod Biol*. 2007 Apr;131(2):203–8.
10. Soper JT, Johnson P, Johnson V, Berchuck A, Clarke-Pearson DL. Comprehensive restaging laparotomy in women with apparent early ovarian carcinoma. *Obstet Gynecol*. 1992 Dec;80(6):949–53.
11. Trimbos JB. Surgical treatment of early-stage ovarian cancer. *Best Pr Res Clin Obs Gynaecol [Internet]*. 2016/11/30. 2017;41:60–70. Available from: https://ac.els-cdn.com/S1521693416300943/1-s2.0-S1521693416300943-main.pdf?_tid=6a674bca-ff1c-487b-93e3-2f087d157c8f&acdnat=1541062495_8cce1f6fc1277687a0d39abdfdf2ee38
12. Ayhan A, Gultekin M, Celik NY, Dursun P, Taskiran C, Aksan G, et al. Occult metastasis in early ovarian cancers: risk factors and associated prognosis. *Am J Obs Gynecol [Internet]*. 2007/01/24.

- 2007;196(1):81.e1-6. Available from:
<https://www.sciencedirect.com/science/article/pii/S0002937806011835?via%3Dihub>
13. Shroff R, Brooks RA, Zigelboim I, Powell MA, Thaker PH, Mutch DG, et al. The utility of peritoneal biopsy and omentectomy in the upstaging of apparent early ovarian cancer. *Int J Gynecol Cancer* [Internet]. 2011/06/03. 2011;21(7):1208–12. Available from:
<https://www.ncbi.nlm.nih.gov/pubmed/21633295>
 14. Kleppe M, Wang T, Van Gorp T, Slangen BFM, Kruse AJ, Kruitwagen RFPM. Lymph node metastasis in stages I and II ovarian cancer: A review. *Gynecol Oncol* [Internet]. 2011 Dec 1 [cited 2019 May 1];123(3):610–4. Available from:
<https://www.sciencedirect.com/science/article/pii/S0090825811007839>
 15. Trope C, Kaern J. Adjuvant chemotherapy for early-stage ovarian cancer: review of the literature. *J Clin Oncol*. 2007/07/10. 2007;25(20):2909–20.
 16. Kleppe M, van der Aa MA, Van Gorp T, Slangen BFM, Kruitwagen RFPM. The impact of lymph node dissection and adjuvant chemotherapy on survival: A nationwide cohort study of patients with clinical early-stage ovarian cancer. *Eur J Cancer*. 2016 Oct;66:83–90.
 17. Jeyarajah DR, Khithani A, Siripurapu V, Liu E, Thomas A, Saad AJ. Lymph node retrieval in pancreaticoduodenectomy specimens: does educating the pathologist matter? *HPB (Oxford)*. 2014 Mar;16(3):263–6.
 18. Svec A, Horak L, Novotny J, Lysy P. Re-fixation in a lymph node revealing solution is a powerful method for identifying lymph nodes in colorectal resection specimens. *Eur J Surg Oncol*. 2006 May;32(4):426–9.
 19. Trimbos JB, Vergote I, Bolis G, Vermorken JB, Mangioni C, Madronal C, et al. Impact of adjuvant chemotherapy and surgical staging in early-stage ovarian carcinoma: European Organisation for Research and Treatment of Cancer-Adjuvant ChemoTherapy in Ovarian Neoplasm trial. *J Natl Cancer Inst*. 2003 Jan;95(2):113–25.
 20. Trimbos B, Timmers P, Pecorelli S, Coens C, Ven K, van der Burg M, et al. Surgical staging and treatment of early ovarian cancer: long-term analysis from a randomized trial. *J Natl Cancer Inst* [Internet]. 2010;102(13):982–7. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/20445161>

Tables

Table 1. Incompleteness of the staging procedure per element

Part of staging procedure	n (%) Complete	n (%) Incomplete	Inapplicable
Inspection	50 (100%)	0 (0%)	
Palpation	43 (86%)	7 (14%)	
Ascites	39 (76%)	12 (24%)	
Ovaria and uterus	45 (90%)	5 (10%)	
Omentum	48 (96%)	2 (4%)	
Biopsies (> 4)	16 (32%)	34 (68%)	
Douglas	23 (46%)	27 (54%)	
Bladder peritoneum	32 (64%)	18 (36%)	
Fossa ovarica right	8 (16%)	42 (84%)	
Fossa ovarica left	9 (18%)	41 (82%)	
Paracolic right	37 (74%)	13 (26%)	
Paracolic left	37 (74%)	13 (26%)	
diaphragm	25 (50%)	25 (50%)	
Lymph nodes (≥ 10)	34 (68%)	16 (32%)	
Lymph nodes location			
Aortocaval high	25 (50%)	25 (50%)	
Pelvic ipsilateral	41 (82%)	9 (18%)	
Pelvic contralateral	25 (50%)	25 (50%)	
Suspicious lesions	30 (60%)	1 (2%)	19 (38%)

Figures

Figure 1

Flowchart. Fifty-eight patients with incomplete staging of early epithelial ovarian cancer were included. One patient was excluded due to pathological confirmed advanced stage ovarian cancer. Complete staging was the reason for exclusion in seven other patients.

Figure 2. Reasons for incomplete staging

Figure 2

Reasons for incomplete staging. Per staging element reasons for omitting the particular element of the staging procedure are described and counted.