Statistical characteristics of the spatial distribution of wind and snow in the Xinjiang Uygur Autonomous Region

Xuetao Yu (✉ yuxuetaoxiaoping@163.com)

Shijiazhuang Tiedao University <https://orcid.org/0000-0002-8466-5653>

Mengke Liu

Shijiazhuang Tiedao University

Junyu Wang

Shijiazhuang Tiedao University

Qingkuan Liu

Shijiazhuang Tiedao University

Jianqing Bu

Shijiazhuang Tiedao University

Research Article

Keywords: the Xinjiang Uygur Autonomous Region, snowdrift, spatial distribution, daily data sets of climate data for the China international surface exchange station

Posted Date: February 17th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-233920/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Natural Hazards on January 10th, 2022.
See the published version at <https://doi.org/10.1007/s11069-021-05127-4>.

Statistical characteristics of the spatial distribution of wind and snow in the Xinjiang Uygur Autonomous Region

Xuetao Yu¹⁻³ Mengke Liu³ Junyu Wang³ Qingkuan Liu^{*1, 4, 5} Jianqing Bu^{*1-3}

1-(State Key Laboratory of Mechanical Behavior and System Safety of Traffic Engineering Structures, Shijiazhuang Tiedao University, Shijiazhuang 050043, P. R. China)

2-(Key Laboratory of Traffic Safety and Control of Hebei Province, Shijiazhuang 050043, P. R. China)

3-(School of Traffic and Transportation, Shijiazhuang Tiedao University, Shijiazhuang 050043, P. R. China)

4-(Wind Engineering Research Center, School of Civil Engineering, Shijiazhuang Tiedao University, Shijiazhuang 050043, P. R. China)

5-(Innovation Center for Wind Engineering and Wind Energy Technology of Hebei Province, Shijiazhuang 050043, P. R. China)

Abstract: At present, the wind and snow loads are calculated independently when determining the design specifications for building structures. Yet, when snow drifting occurs, the basic wind and snow pressures exist almost simultaneously. Therefore, building specifications based on the independent calculations of the wind and snow loads cannot be used effectively in areas that are severely impacted by snow drifting. Some parts of the Xinjiang Uygur Autonomous Region of China have suffered particularly severe snow drifting natural disasters. In this paper, we offer an analysis of the statistical characteristics of the spatial distribution of wind and snow in this region. In particular, we extract the values of the combined distribution of the wind and snowfall in the Xinjiang Uygur Autonomous Region of China by using parameters such as snowfall intensity, snowfall amount, wind speed, and wind direction as indicators that show the snow drifting disasters. This study found that, after heavy snowfall events, the accumulative wind scale is high and the accumulative snowfall is large in the Altay, the Bortala Mongol Autonomous, and the Tacheng Prefectures of northern Xinjiang, and the Kizilsu Kirghiz Autonomous Prefecture of western Xinjiang. It has an important practical significance for the design of building structures and the treatment of snow drifting disasters.

Keywords: the Xinjiang Uygur Autonomous Region; snowdrift; spatial distribution; daily data sets of climate data for the China international surface exchange station

1. Introduction

A snow drifting disaster is one of many types of snow damage that occurs frequently in northern provinces of China like the Xinjiang Uygur Autonomous Region, the Inner Mongolia Autonomous Region, the Jilin Province and so on. This type of disaster consists of deep drifts of snow which may close roads or simply block traffic. In severe cases, it can result in the structural damage to buildings, the collapse of roofs and houses, and other problems which disrupt the daily lives of the inhabitants, as well as industrial and agricultural production. For example, the biggest snowstorm on record occurred in Liaoning Province of China in March 2007. For the most part, it halted traffic; it forced airports and expressways to close, and stopped several of the trains—and this caused huge economic losses to the Liaoning Province (Li, Zhu et al. 2007). Since January 10 of 2008, southern

**Corresponding author 1 address:* Qingkuan Liu, Wind Engineering Research Center, School of Civil Engineering, Shijiazhuang Tiedao University, 17 Northeast, Second Inner Ring, Shijiazhuang, Hebei, P.R.C. 050043. E-mail: lqk@stdu.edu.cn

Corresponding author 2 address: Jianqing Bu, School of Traffic and Transportation, Shijiazhuang Tiedao University, 17 Northeast, Second Inner Ring, Shijiazhuang, Hebei, P.R.C. 050043. E-mail: buiq2004@163.com

China suffered a heavy snow and sleet natural disaster which was the worst in 50 years. More than 20 of China's provinces were affected. The snow and ice storms cut off travel by road and rail, and cost 151.65 billion yuan in direct economic losses(Chen and Fan 2009). In January 2010, the Altay district, the Tacheng district and other places nearby in the Xinjiang Uygur Autonomous Region suffered a once-in-60-year blizzard disaster that closed off the main access roads of the city of Tacheng, and caused direct economic losses of 650 million yuan. The snowfall intensity, duration, and amount of rainfall (or snowfall) for this event have rarely been seen in modern Chinese history(Liu, Liu et al. 2010).

Generally, snowdrift formation is a joint probability event of wind and snow variables, with the two primary variables being snowfall intensity and wind strength. Only when heavy snowfall and strong winds appear together will snowdrifts be formed(Zha, Yu et al. 2012). In the *Load Code for the Design of Building Structure*, the wind and snow loads are calculated separately. So, the calculated load based on this load standard cannot be used to design safe architectural structures for those areas that are affected by snow drifting disasters(Research, University et al. 2012).

Snow drifting disasters have a serious impact on the inhabitants of colder regions. Aiming at this problem, many scholars have studied the formation and prevention of snow drifting disasters. Guoping Su has studied the causes of snow drifting disaster, and the factors that influence it. More particularly, Su has analyzed the load status of the snow particles and proposed the design parameters of a retaining wall(Su 2017). Chengliang Ying has studied the technology of the prevention and cure of a snow drifting disaster on highways(Ying 2007). Having studied the causes of snow drifting disasters on the highways in the Xinjiang Uygur Autonomous Region, Jian Liu et al. have summarized the main types of snow deposition on the region's highways, including windward snow cover, leeward snow cover, curve flow snow, cutting snow, and high embankment snow types(Liu, Yao et al. 2008). Tingliang Wang implemented the numerical simulation of snow disasters to sum up the drifting snow disaster prevention technology of the railway. He pointed out that, in general, three conditions are required for the formation of a drifting snow disaster: the source of sufficient snow, the wind which can sweep the snow grains, and the landform which can help to decrease the accumulation of snow grains(Wang 2009). Therefore, in this paper, the authors studied the joint distribution of wind and snow in the Xinjiang Uygur Autonomous Region by using a mass of historical meteorological data, and summarized the distribution rules of wind and snow in Xinjiang, thereby providing a guide for the prevention and control of future snow drifting disasters in the Xinjiang Uygur Autonomous Region which, in turn, provides significant guidance on the construction and maintenance of the transportation infrastructure in the region.

2. Meteorological data

The meteorological data used in this study is the "Daily Data Sets of Climate Data for China International Surface Exchange Station (V3.0)". It includes eight major items, including the daily air pressure values, air temperature, precipitation, evaporation, relative humidity, wind, sunshine time, and 0 cm ground temperature. Beginning in January 1951, the data was acquired daily from 824 basic meteorological stations located in China. After effective quality control, the availability rate of the meteorological factor data was, in general, more than 99%, and the accuracy rate was nearly 100%.

In this study, the precipitation and wind data were selected from the above data sets. In the precipitation data, the accumulated precipitation from 20:00 of the previous day to 20:00 was used. In the wind data, the average wind speed, the maximum wind speed, and the direction of maximum

wind speed were used. Under normal circumstances, the daily average wind speed was obtained by calculating the average of four values observed at 02:00, 08:00, 14:00, and 20:00. However, when the observation station was not equipped with a self-recording instrument, the daily average wind speed is the average of the three wind speed values observed at 08:00, 14:00, and 20:00. Maximum wind speed refers to the average maximum wind speed over 10 minutes during a given period. The wind direction was expressed relative to 16 azimuths. The representation directions were N, NNE, NE, ENE, E, ESE, SE, SSE, S, SSW, SW, WSW, W, WNW, NW, and NNW. The absence of wind was also recorded.

As there was no distinction between rainfall and snowfall in the data sets, the precipitation over the whole winter period was used to represent snowfall in our analysis of the statistical characteristics of snow and wind. The winter period is defined as the months of January, February, March, November, and December. After importing the above data into a database, the effective data from 64 observation stations in the Xinjiang Uygur Autonomous Region were selected for this analysis. The location and number of the 64 stations are shown in Figure 1. The selected period was from 1968 to 2016 and included a total of 7412 days. There was a lack of observation data for a portion of this time due to force majeure. However, the number of missing days at some sites was very small. Station 51704 had no data for 25 days, which comprised the greatest number of days with no data among all the stations. For this situation, the missing data were inferred based on the mean of the data measured before and after the gap.

Figure 1. Meteorological station distribution map in the Xinjiang Uygur Autonomous Region.

3. Wind scale distribution in winter

In the field of meteorological forecasting, the wind scale of a certain day is divided into 18 classes according to the average wind speed. The basis of the classification is the “Wind Scale”(Centre 2012). The specific classification method is shown in Table 1(Centre 2012). According to this

method, the appearance frequency of different daily wind scales at 64 stations in the Xinjiang Uygur Autonomous Region was extracted over the periods of 1st, 2nd, 3rd, 11th, and 12th months from 1968 to 2016. The specific results are shown in Table 2.

Table 1. Wind scale classification.

Wind scale	Wind speed (m/s)	Wind scale	Wind speed (m/s)	Wind scale	Wind speed (m/s)
0	0.0~0.2	6	10.8~13.8	12	32.7~36.9
1	0.3~1.5	7	13.9~17.1	13	37.0~41.4
2	1.6~3.3	8	17.2~20.7	14	41.5~46.1
3	3.4~5.4	9	20.8~24.4	15	46.2~50.9
4	5.5~7.9	10	24.5~28.4	16	51.0~56.0
5	8.0~10.7	11	28.5~32.6	17	≥56.1

Table 2. Appearance frequency of different daily wind scales in the Xinjiang Uygur Autonomous Region during the period of 1st, 2nd, 3rd, 11th, and 12th months from 1968 to 2016.

Station ID	Wind scale 0	Wind scale 1	Wind scale 2	Wind scale 3	Wind scale 4	Wind scale 5	Wind scale 6	Wind scale 7	Wind scale 8	Wind scale 9	Wind scale 10	Wind scale 11
51358	952	4673	1650	107	7	0	0	0	0	0	0	0
51367	65	3020	3808	488	29	1	0	0	0	0	0	0
51059	56	2360	3366	1175	316	104	27	5	0	0	0	0
52112	193	1591	3189	1362	640	263	120	45	6	1	0	0
51060	118	714	2242	2270	1415	574	74	4	0	0	0	0
51720	166	5604	1522	104	15	1	0	0	0	0	0	0
51639	732	4594	1894	165	25	2	0	0	0	0	0	0
51655	446	4492	2215	212	31	6	1	0	0	0	0	0
51704	663	5655	869	153	40	5	2	0	0	0	0	0
51818	211	4957	2119	116	9	0	0	0	0	0	0	0
51730	332	5346	1570	145	17	1	0	0	0	0	0	0
51433	180	3804	2765	537	97	18	1	0	0	0	0	0
51526	12	2335	4725	309	30	1	0	0	0	0	0	0
52118	32	1289	4075	1505	405	87	18	1	0	0	0	0
51633	975	6104	313	13	2	2	0	0	0	0	0	0
51701	158	1877	3165	1634	497	70	9	0	0	0	0	0
51642	1273	4477	1557	98	4	1	0	0	0	0	0	0
51365	669	4538	2070	129	6	0	0	0	0	0	0	0
51470	34	966	5338	1008	57	3	1	1	0	0	0	0
51656	204	3420	2858	686	196	41	7	0	0	0	0	0
51330	341	4569	2252	214	27	6	1	0	0	0	0	0
51238	673	4948	1661	122	4	0	0	0	0	0	0	0
51068	253	3004	3109	866	153	18	9	0	0	0	0	0
51053	61	968	1959	2066	1551	642	157	7	0	0	0	0
51542	455	3614	2182	673	344	122	22	0	0	0	0	0
51467	214	2987	3864	325	20	2	0	0	0	0	0	0
51186	2822	4063	466	38	2	1	0	0	0	0	0	0

51288	88	2628	3908	695	87	4	1	0	0	0	0	0
51810	899	4727	1587	163	20	2	0	0	0	0	0	0
51804	1109	3730	1939	536	88	7	1	0	0	0	0	0
51765	368	4470	2083	374	97	17	3	0	0	0	0	0
51711	6	1316	4624	1373	91	1	0	0	0	0	0	0
51839	121	4933	2218	128	8	0	0	0	0	0	0	0
52101	834	2994	2623	749	201	10	1	0	0	0	0	0
51243	506	4127	2070	446	159	77	19	4	2	1	0	0
51241	100	2785	3771	630	118	6	0	0	0	0	0	0
51232	47	1077	2696	1774	852	454	332	141	36	3	0	0
51477	0	241	1273	2110	2060	1140	483	86	19	0	0	0
51931	126	5321	1909	54	2	0	0	0	0	0	0	0
51705	837	3967	1501	643	348	93	22	1	0	0	0	0
51087	2546	3907	673	215	60	7	2	0	0	0	0	0
51855	343	3982	2646	385	53	3	0	0	0	0	0	0
51437	161	4284	2517	369	77	4	0	0	0	0	0	0
51076	687	4068	2131	443	71	11	1	0	0	0	0	0
51156	632	4156	1907	425	214	56	19	3	0	0	0	0
51811	367	5213	1723	99	10	0	0	0	0	0	0	0
51628	106	5373	1850	72	9	1	0	0	0	0	0	0
51133	66	3370	3165	612	154	37	6	1	0	0	0	0
51346	189	4755	2233	221	13	0	0	0	0	0	0	0
51777	68	3126	3404	607	153	38	12	1	0	0	0	0
51828	73	3719	3479	134	6	1	0	0	0	0	0	0
51716	881	4453	1739	300	38	1	0	0	0	0	0	0
51334	495	5117	1539	207	45	4	0	0	0	0	0	0
51495	357	1568	2011	1174	965	712	315	167	97	40	5	1
52313	2	214	2615	3396	888	233	51	9	2	0	0	0
51431	222	4704	2178	273	28	4	1	0	0	0	0	0
51573	1328	5422	636	22	2	2	0	0	0	0	0	0
51567	695	4364	2049	268	31	4	0	0	0	0	0	0
51379	31	1207	4923	1169	78	3	0	0	0	0	0	0
51463	94	3325	3670	275	31	12	5	0	0	0	0	0
52203	211	4161	2757	248	28	7	0	0	0	0	0	0
51709	283	4984	2008	110	24	3	0	0	0	0	0	0
51644	134	3818	3297	149	14	0	0	0	0	0	0	0
51581	592	5322	1431	63	2	0	0	0	0	0	0	0

From Table 2, we know that strong winds appeared frequently at stations 51059, 52112, 51060, 51053, 51232, 51477, 51495 and 52313. As shown in Table 1, the wind speed at the different grades of wind scale was approximately equal to a mathematical progression. For the convenience of making calculations, we multiplied the different wind scales and their corresponding appearance frequency for every station, and then carried the accumulation. Based on this method, the accumulative wind scale distribution map can be obtained if the contemporaneous data from the

nearby regional sites are added during the calculation process. Taking station 52112 as an example, the total number of days with valid data were 7412. According to the valid data, the nominal accumulative wind grade was $0 \times 193 + 1 \times 1591 + 2 \times 3189 + 3 \times 1362 + 4 \times 640 + 5 \times 263 + 6 \times 120 + 7 \times 45 + 8 \times 6 + 9 \times 1 = 17022$, and its average wind scale was about 2.29717. Due to the absence of data for two days, the weighted cumulative wind scale calculated in this study was $17022 + 2.29717 \times 2 = 17026.5943$. After this calculation, the final distribution map representing the weighted accumulative wind scale was generated, as shown in Figure 2.

Figure 2. Distribution map of the weighted accumulative wind scale in the Xinjiang Uygur Autonomous Region.

It can be seen from Figure 2 that the weighted accumulated wind scale is large in the northwest of Altay, the east and west of Hami, and at the junction of Urumqi and Turpan. However, it is relatively small in other regions. The Altay region is known as one of the nine strong wind regions in Xinjiang, and it is a typical terrain named “a valley between two mountains”, which is composed of the Altay and Saur Mountains, and the Irtysh River Valley. For this reason, the weighted accumulated wind scale in this region is large. The junction of Urumqi and Turpan lies between the middle and the southern Tianshan Mountains. This place is called the Daban Town wind region and is also well known as one of the nine strong wind regions in Xinjiang. It’s the key airflow channel between north and south Xinjiang. The development and construction conditions regarding wind energy in this area is the best among the nine strong wind regions. The special terrain in the Hami area is called the “four mountains lock three basins,” where cold air activity is frequent. So, the wind scale is high in this area.

Meanwhile, the authors extracted the maximum wind direction data for all the stations in Xinjiang during this period, and drew the wind rose maps of the stations with high wind scales (including

51053, 51059, 51060, 51232, 51477, 51495, 52112 and 52313). They are shown in Figure 3. The results show that the east wind prevails in the northwest of Altay. The southeast and west winds prevail at the junction of Urumqi and Turpan, and in the southwest corner of Tacheng city. The north wind prevails in the west of Hami city, while the west and east winds prevail in the east.

Figure 3. Maximum wind direction rose diagrams for some stations in the Xinjiang Uygur Autonomous Region.

4. Basic wind speed in winter

In China's design specifications for buildings that deal with wind load, the reoccurrence period of wind load for general structures is set at 50 years. Therefore, it is important to calculate the reoccurrence period of wind and snow accurately. Consequently, the annual maximum wind speed in winter during the 49-year period was extracted. For historical reasons, this data for the years 1968, 1969, and 1970 for all the stations in this region are missing. In addition, the data for station 54618 in the year 1971 and for station 54606 for the years 1971 and 1972 are also absent.

Based on the above available data, the basic wind speed in the reoccurrence period of 50 and 100 years was calculated using a Gumbel curve with an extreme type I distribution. The calculation steps are as follows.

First, the average maximum wind speed $\bar{v}_i$ and its root variance σ_i within a specified period of each station were calculated. The calculation formula $\bar{v}_i$ for station number i is shown in

equation (1). The formula σ_i for station number i is shown in equation (2). v_{ij} is the maximum wind speed for station number i in the winter of the year j , and n is the year from 1971 to 2016 for most stations.

$$\bar{v}_i = \sum_{j=1}^n v_{ij} \quad (1)$$

$$\sigma_i = \sqrt{\frac{\sum_{j=1}^n (v_{ij} - \bar{v}_i)^2}{n-1}} \quad (2)$$

Second, the assurance rate P (according to the recurrence period T) was obtained. The calculation formula for P is shown in equation (3). The value of T was set to 50 and 100 years.

$$P = 1 - \frac{1}{T} \quad (3)$$

Based on the assurance rate P , the assurance coefficient ϕ can be further obtained according to the reference table presenting the assurance coefficient of the Gumbel curve with an extreme type I distribution, which is shown in Table 3.

Table 3. Reference table of the assurance coefficient of the Gumbel curve with an extreme type I distribution.

P	0.60	0.70	0.80	0.90	0.95	0.97	0.98	0.99	0.995	0.999	0.9995	0.9999
ϕ	0.07	0.35	0.72	1.30	1.87	2.27	2.59	3.14	3.68	4.94	5.48	6.73

Finally, formula (4) was adopted to calculate the basic wind speed at each station during the different reoccurrence periods. The calculated results are shown in Table 4.

$$V_i = \bar{v}_i + \sigma_i \cdot \phi \quad (4)$$

Table 4. Basic wind speed of each station in the Xinjiang Uygur Autonomous Region at different reoccurrence periods.

Station ID	Valid data count	Basic wind speed in 50 year	Corresponding wind scale in 50 years	Basic wind speed in 100 years	Corresponding wind scale in 100 years
51358	20	13.77	6	14.84	7
51367	23	13.85	6	14.89	7
51059	31	24.87	10	26.7	10
52112	12	23.04	9	24.2	9
51060	13	15.15	7	16.03	7
51720	12	14.71	7	15.98	7
51639	10	10.41	5	11	6
51655	37	18.33	8	19.91	8
51704	39	20.85	9	22.62	9
51818	46	14.33	7	15.63	7
51730	12	12.96	6	14.02	7

51433	30	16.14	7	17.21	8
51526	32	15.07	7	15.92	7
52118	8	22.59	9	23.88	9
51633	12	15	7	16.6	7
51701	16	21.59	9	22.84	9
51642	12	17.7	8	19.39	8
51365	12	14.52	7	15.69	7
51470	11	22.97	9	24.74	10
51656	46	22.98	9	25.03	10
51330	32	23.99	9	26.02	10
51238	32	12.79	6	13.78	6
51068	33	19.33	8	20.5	8
51053	32	22	9	23.48	9
51542	45	22.39	9	23.49	9
51467	12	12.86	6	13.77	6
51186	30	12.25	6	13.13	6
51288	11	20.5	8	21.66	9
51810	25	15.18	7	16.53	7
51804	31	22.88	9	24.75	10
51765	12	12.69	6	13.38	6
51711	46	19.22	8	20.97	9
51839	46	14.91	7	16.01	7
52101	12	18.48	8	19.72	8
51243	46	31.96	11	34.46	12
51241	12	12.75	6	13.17	6
51232	46	35.31	12	37.34	12
51477	46	30.57	11	32.6	11
51931	46	13.45	6	14.59	7
51705	46	21.62	9	22.76	9
51087	32	19.72	8	21.3	9
51855	12	11.73	6	12.3	6
51437	46	20.46	8	22.14	9
51076	49	18.18	8	19.48	8
51156	12	25.91	10	28.41	10
51811	20	10.69	5	11.41	6
51628	46	19.39	8	21.37	9
51133	37	21.17	9	22.57	9
51346	32	10.73	5	11.32	6
51777	37	19.98	8	21.48	9
51828	49	14.36	7	15.59	7
51716	34	15.55	7	16.78	7
51334	32	17.51	8	18.77	8
51495	12	37.15	12	38.59	12

52313	12	19.68	8	20.58	8
51431	48	18.72	8	20.11	8
51573	46	19.98	8	22.15	9
51567	25	13.73	6	14.45	7
51379	32	23.92	9	26.42	10
51463	49	26.8	10	29.26	11
52203	47	19.35	8	21.03	9
51709	45	24.4	9	26.91	10
51644	46	17.87	8	19.47	8
51581	32	11.72	6	12.61	6

Due to historical reasons, only partial annual data is available for some stations, but the amount of time for which there is valid data is over 12 years for most stations. This data can reflect the realities and be used in the above calculation. Based on the results presented in Table 4, a distribution map of the basic wind speed in winter at the 50-year reoccurrence period in the Xinjiang Uygur Autonomous Region was generated, as shown in Figure 4. This analysis indicated that the wind speed was high in the west of the Changji Hui Autonomous Prefecture, the west of Turfan, the west of the Bayingolin Mongol Autonomous Prefecture, Hami, Tacheng, and the Bortala Mongol Autonomous Prefecture. The wind speed was low in Aksu, Khotan, Kashgar, and the east of the Bayingolin Mongol Autonomous Prefecture. Consequently, these results can be referred to the code of building structure design in these regions.

Figure 4. Distribution map of the basic wind speed in winter at the 50-year reoccurrence period in the Xinjiang Uygur Autonomous Region.

5. Snowfall distribution in winter

In winter, precipitation is usually represented by snowfall. China has a strict regulation about the snowfall grade standard. Snowfall is the depth of an equal amount of water that originated as snow. Like rainfall, snowfall refers to the amount of snow that falls within a certain period, and this is generally measured over 24 hours. According to the standard *Short-Range Weather Forecast*(Centre 2017), snowfall is divided into seven grades: sporadic light snow, light snow, moderate snow, heavy snow, blizzard, storm snow, and super-big blizzard(Centre 2017). If the snowfall is much more than 10 mm, the grade super-big blizzard can be sub-divided into two levels: big blizzard and super-big blizzard. The defined snowfall occurring at each grade within 24 hours is shown in Table 5.

As the occurrence of sporadic light snow is very small, the sporadic light snow and light snow data can be merged. Subsequently, the times of the different snowfall grades at the 64 stations in the Xinjiang Uygur Autonomous region within 1st, 2nd, 3rd, 11th, and 12th months from 1968 to 2016 were extracted. The results are shown in Table 6.

Table 5. Snowfall grade classification.

Snowfall Grade	24 hours of precipitation (mm)
sporadic light snow	≤ 0.1
light snow	0.1~2.4
moderate snow	2.5~4.9
heavy snow	5.0~9.9
blizzard	10.0~19.9
storm snow	20.0~30.0
super-big blizzard	> 30.0

Table 6. Occurrence times of different snowfall grades in the Xinjiang Uygur Autonomous Region within 1st, 2nd, 3rd, 11th, and 12th months from 1968 to 2016.

Station ID	Times of light snow	Times of moderate snow	Times of heavy snow	Times of blizzard	Times of storm snow	Times of super-big blizzard
51358	1243	201	104	30	2	0
51367	927	154	95	22	0	0
51059	1391	230	105	16	0	0
52112	142	14	3	3	0	0
51060	1287	172	57	10	1	0
51720	260	29	5	7	1	0
51639	227	20	7	2	0	0
51655	165	16	4	0	0	0
51704	253	52	35	12	4	1
51818	210	26	15	3	0	0
51730	148	9	3	2	0	0
51433	1084	327	195	61	4	0
51526	176	8	5	2	0	0
52118	360	25	7	2	1	0
51633	549	55	34	9	0	0
51701	1113	107	32	10	0	0

51642	213	23	12	6	0	0
51365	1074	150	57	4	0	0
51470	898	170	117	26	2	0
51656	242	19	5	2	1	0
51330	1212	94	44	12	1	0
51238	1279	98	55	15	3	0
51068	1156	120	25	3	0	0
51053	1354	219	90	27	2	0
51542	1118	57	17	3	0	0
51467	121	15	2	0	0	0
51186	825	167	91	21	3	0
51288	981	125	39	5	0	0
51810	154	22	7	5	0	0
51804	292	32	12	3	0	0
51765	68	8	3	1	0	0
51711	420	55	29	15	4	0
51839	161	14	2	0	0	0
52101	709	73	21	3	0	0
51243	1107	62	17	3	0	0
51241	1394	153	38	10	0	0
51232	1400	57	16	5	2	1
51477	336	14	1	0	0	0
51931	176	12	6	1	0	0
51705	566	70	42	10	4	0
51087	1090	210	115	35	5	1
51855	94	6	0	0	0	0
51437	1165	183	74	9	1	0
51076	1043	238	143	38	4	0
51156	856	75	23	2	0	0
51811	180	30	16	3	1	0
51628	341	24	15	6	1	0
51133	1193	344	181	70	10	2
51346	1260	118	68	16	1	1
51777	103	15	5	0	0	0
51828	199	16	10	4	0	0
51716	151	16	8	2	0	0
51334	1465	58	29	4	0	0
51495	68	3	1	0	0	0
52313	347	12	4	1	0	0
51431	918	331	223	78	16	1
51573	96	9	4	3	0	0
51567	253	17	10	3	0	0
51379	971	159	68	5	0	0

51463	1113	206	130	45	4	1
52203	229	23	11	1	0	0
51709	308	42	34	8	1	1
51644	305	33	15	4	0	0
51581	124	11	6	0	1	0

As can be seen in Table 6, the times of high snowfall grade at stations 51433, 51133, and 51431 in the west of the Xinjiang Uygur Autonomous Region are relatively more than that of the other stations. According to the snowfall amount at each snowfall grade over 24 hours, the grades were standardized: light snow, 1; moderate snow, 2; heavy snow, 4; blizzard, 8; storm snow, 12; and super-big blizzard, 16.

As an example, at station 51431, the defined weighted cumulative snowfall grade was $918 \times 1 + 331 \times 2 + 223 \times 4 + 78 \times 8 + 16 \times 12 + 1 \times 16 = 3304$. Based on this method, the weighted accumulative snowfall grade for all of the stations in this region, as well as those nearby, was calculated, and the distribution map of the weighted cumulative snowfall grade in the Xinjiang Uygur Autonomous region was generated, as shown in Figure 5.

Figure 5. Distribution map of weighted accumulative snowfall grade in the Xinjiang Uygur Autonomous region.

According to Figure 5, we know that the snowfall areas of the Xinjiang Uygur Autonomous Region are mainly in the north, and the heaviest snowfalls are in the Yili Kazak Autonomous and the northwestern Tacheng Prefectures. The snowfall is medium in Altay, the Changji Hui Autonomous Prefecture, Shihezi City, Urumqi, and the Tacheng Prefectures except for the northwest. Snowfall in other areas is very slight.

6. Accumulative snowfall and wind scale for the top ten maximum snowfall days in winter

As most snow drift disasters occur after a heavy snowfall, the authors extracted the wind scales of six days (including the day of the snowfall) after the top ten maximum snowfall days in winter in the Xinjiang Uygur Autonomous Region by using the above data, and then the wind grades on these 60 days are classified. The numbers of occurrences of different wind scales at all stations during these 60 days are shown in Table 7. In this data of 60 days at all stations, one day's data at stations 51431 and 51470 and two days' data at stations 51433 and 51704 are missing. To analyze the wind power during these 60 days, the authors multiplied the different wind scales and their corresponding appearance frequency for each station, and then calculated the accumulation. Based on this method, the cumulative wind scale distribution map can be obtained if the contemporaneous data from the nearby regional sites are added during the calculation process. The distribution map is shown in Figure 6. For the missing data at different stations, we used the average value on the other days.

Table 7. The numbers of occurrences of different wind scales for six days (including the day of snowfall) after the top ten maximum snowfall days in winter at all stations in the Xinjiang Uygur Autonomous Region.

Station ID	Times of Wind scale 0	Times of Wind scale 1	Times of Wind scale 2	Times of Wind scale 3	Times of Wind scale 4	Times of Wind scale 5	Times of Wind scale 6	Times of Wind scale 7	Times of Wind scale 8	Count of missing data	Accumulative wind scale
51053	0	6	18	19	17	0	0	0	0	0	167
51059	0	24	16	12	5	1	2	0	0	0	129
51060	0	4	23	27	5	1	0	0	0	0	156
51068	0	14	35	9	2	0	0	0	0	0	119
51076	3	33	20	3	1	0	0	0	0	0	86
51087	13	32	10	5	0	0	0	0	0	0	67
51133	1	29	21	8	1	0	0	0	0	0	99
51156	6	19	25	7	3	0	0	0	0	0	102
51186	18	36	6	0	0	0	0	0	0	0	48
51232	0	6	18	20	6	0	4	2	4	0	196
51238	0	31	28	1	0	0	0	0	0	0	90
51241	1	24	32	3	0	0	0	0	0	0	97
51243	8	21	20	7	4	0	0	0	0	0	98
51288	2	15	33	6	3	1	0	0	0	0	116
51330	3	38	19	0	0	0	0	0	0	0	76
51334	4	38	15	3	0	0	0	0	0	0	77
51346	1	29	28	2	0	0	0	0	0	0	91
51358	11	34	15	0	0	0	0	0	0	0	64
51365	1	27	30	2	0	0	0	0	0	0	93
51367	0	19	40	1	0	0	0	0	0	0	102
51379	0	5	46	8	1	0	0	0	0	0	125

51431	0	28	28	3	0	0	0	0	0	1	94.4
51433	8	35	9	5	1	0	0	0	0	2	73.6
51437	0	30	27	3	0	0	0	0	0	0	93
51463	0	21	36	3	0	0	0	0	0	0	102
51467	5	23	30	2	0	0	0	0	0	0	89
51470	0	1	36	21	1	0	0	0	0	1	142
51477	0	4	18	12	13	9	4	0	0	0	197
51495	2	11	17	13	2	9	2	3	1	0	178
51526	0	34	26	0	0	0	0	0	0	0	86
51542	0	27	30	3	0	0	0	0	0	0	96
51567	5	35	16	4	0	0	0	0	0	0	79
51573	11	42	7	0	0	0	0	0	0	0	56
51581	2	48	10	0	0	0	0	0	0	0	68
51628	0	42	18	0	0	0	0	0	0	0	78
51633	5	48	6	1	0	0	0	0	0	0	63
51639	5	35	20	0	0	0	0	0	0	0	75
51642	12	30	18	0	0	0	0	0	0	0	66
51644	0	30	29	1	0	0	0	0	0	0	91
51655	3	46	9	2	0	0	0	0	0	0	70
51656	1	30	25	2	2	0	0	0	0	0	94
51701	0	7	33	15	3	2	0	0	0	0	140
51704	4	43	11	0	0	0	0	0	0	2	67.5
51705	3	35	15	3	1	3	0	0	0	0	93
51709	0	36	24	0	0	0	0	0	0	0	84
51711	0	12	33	14	1	0	0	0	0	0	124
51716	1	36	21	2	0	0	0	0	0	0	84
51720	6	37	17	0	0	0	0	0	0	0	71
51730	1	44	14	1	0	0	0	0	0	0	75
51765	0	30	26	4	0	0	0	0	0	0	94
51777	1	20	27	7	5	0	0	0	0	0	115
51804	12	35	11	2	0	0	0	0	0	0	63
51810	8	41	10	1	0	0	0	0	0	0	64
51811	1	38	16	3	2	0	0	0	0	0	87
51818	0	26	27	4	3	0	0	0	0	0	104
51828	0	23	34	3	0	0	0	0	0	0	100
51839	6	35	18	1	0	0	0	0	0	0	74
51855	5	36	15	4	0	0	0	0	0	0	78
51931	2	45	12	1	0	0	0	0	0	0	72
52101	2	18	29	8	3	0	0	0	0	0	112
52112	1	12	27	12	8	0	0	0	0	0	134
52118	3	10	20	20	7	0	0	0	0	0	138
52203	1	41	17	1	0	0	0	0	0	0	78
52313	0	2	28	24	5	1	0	0	0	0	155

Figure 6. Distribution map of the accumulative wind scale from the top ten maximum snowfall days at all stations in the Xinjiang Uygur Autonomous Region.

From Figure 6, we can know that the wind is very strong in northern Altay, western Hami city, western Urumqi, and at the junction of the Bortala Mongol Autonomous and Tacheng Prefectures after heavy snowfalls. The wind in eastern Hami city and the northern Kizilsukirgiz Autonomous Prefecture is weaker than the above areas. While in the other areas, the wind is weakest.

The above is the analysis of the cumulative wind scale for six days (including the day of snowfall) after the top ten maximum snowfall days in winter at all stations in the Xinjiang Uygur Autonomous Region, without considering the total snowfall on the top ten maximum snowfall days. Therefore, the authors analyzed the joint distribution of the total snowfall on the top ten maximum snowfall days and the accumulated wind scale on six days (including the day of the snowfall) after the top ten maximum snowfall days. First, the authors calculated the total snowfall for the top ten maximum snowfall days, and then calculated the cumulative wind scale for six days (including the day of snowfall) after the top ten maximum snowfall days. In order to compare the joint distribution of the total snowfall and the accumulated wind scale, these data are normalized according to Formulas 1 and 2. After these calculations, the total snowfall, the accumulated wind scale, average wind scale, normalized total snowfall, and normalized accumulated wind scale are shown in Table 8.

$$Wind^* = \frac{Wind}{Wind_{max}} \quad (1)$$

$$Snow^* = \frac{Snow}{Snow_{max}} \quad (2)$$

Finally, the authors display the normalized total snowfall and accumulated wind scale on the map

of the Xinjiang Uygur Autonomous Region, which is shown in Figure 7.

Table 8. The normalized total snowfall and the normalized accumulated wind scale of all stations in the Xinjiang Uygur Autonomous Region.

Station ID	Accumulative amount of snowfall (mm)	Accumulative wind scale	Average daily wind scale	Normalized accumulative snowfall	Normalized accumulative wind scale
51053	181.8	167.000	2.783	0.697	0.848
51059	131.8	129.000	2.150	0.505	0.655
51060	135.8	156.000	2.600	0.521	0.792
51068	91.4	119.000	1.983	0.350	0.604
51076	189.6	86.000	1.433	0.727	0.437
51087	224.6	67.000	1.117	0.861	0.340
51133	255.6	99.000	1.650	0.980	0.503
51156	84.1	102.000	1.700	0.322	0.518
51186	193.5	48.000	0.800	0.742	0.244
51232	170.4	196.000	3.267	0.653	0.995
51238	180.3	90.000	1.500	0.691	0.457
51241	129	97.000	1.617	0.495	0.492
51243	90.4	98.000	1.633	0.347	0.497
51288	110.8	116.000	1.933	0.425	0.589
51330	145	76.000	1.267	0.556	0.386
51334	99.9	77.000	1.283	0.383	0.391
51346	175.2	91.000	1.517	0.672	0.462
51358	182.5	64.000	1.067	0.700	0.325
51365	106.3	93.000	1.550	0.408	0.472
51367	128.6	102.000	1.700	0.493	0.518
51379	112.2	125.000	2.083	0.430	0.635
51431	260.8	94.576	1.576	1.000	0.480
51433	198.6	74.483	1.241	0.762	0.378
51437	130.8	93.000	1.550	0.502	0.472
51463	214.6	102.000	1.700	0.823	0.518
51467	42.2	89.000	1.483	0.162	0.452
51470	177.5	142.373	2.373	0.681	0.723
51477	41.3	197.000	3.283	0.158	1.000
51495	25.4	178.000	2.967	0.097	0.904
51526	63.2	86.000	1.433	0.242	0.437
51542	97	96.000	1.600	0.372	0.487
51567	94.3	79.000	1.317	0.362	0.401
51573	71.4	56.000	0.933	0.274	0.284
51581	72.2	68.000	1.133	0.277	0.345
51628	125.3	78.000	1.300	0.480	0.396
51633	126.5	63.000	1.050	0.485	0.320
51639	80.8	75.000	1.250	0.310	0.381

51642	106.9	66.000	1.100	0.410	0.335
51644	99	91.000	1.517	0.380	0.462
51655	50.6	70.000	1.167	0.194	0.355
51656	88.4	94.000	1.567	0.339	0.477
51701	141.6	140.000	2.333	0.543	0.711
51704	232.9	67.241	1.121	0.893	0.341
51705	179.2	93.000	1.550	0.687	0.472
51709	174.7	84.000	1.400	0.670	0.426
51711	184.8	124.000	2.067	0.709	0.629
51716	79.2	84.000	1.400	0.304	0.426
51720	122.8	71.000	1.183	0.471	0.360
51730	61.3	75.000	1.250	0.235	0.381
51765	53.3	94.000	1.567	0.204	0.477
51777	53.1	115.000	1.917	0.204	0.584
51804	88.8	63.000	1.050	0.340	0.320
51810	102.3	64.000	1.067	0.392	0.325
51811	123.2	87.000	1.450	0.472	0.442
51818	91.8	104.000	1.733	0.352	0.528
51828	97.9	100.000	1.667	0.375	0.508
51839	48.5	74.000	1.233	0.186	0.376
51855	33.4	78.000	1.300	0.128	0.396
51931	70.9	72.000	1.200	0.272	0.365
52101	85.5	112.000	1.867	0.328	0.569
52112	83.9	134.000	2.233	0.322	0.680
52118	107.2	138.000	2.300	0.411	0.701
52203	79.7	78.000	1.300	0.306	0.396
52313	59	155.000	2.583	0.226	0.787

Figure 7. Joint distribution map of the total snowfall for the top ten maximum snowfall days and the accumulative wind scale for six days (including the day of snowfall) after the top ten maximum snowfall day at all stations in the Xinjiang Uygur Autonomous Region.

From Figure 7, we can draw four conclusions. (1) After heavy snowfalls, the wind is very strong in the northwestern Altay, the northwestern Tacheng Prefecture and Urumqi city. So, the snowdrift disasters that occur in these areas are very serious. (2) After a heavy snowfall, the wind is not very strong in the Kizilsu Kirghiz Autonomous Prefecture, the Bortala Mongol Autonomous Prefecture and northeastern Altay. So, the snowdrift disasters in these areas are not serious. (3) After a snowfall, the wind is very strong, but it does not snow very much in Hami city. So, it has potential to become an area with serious snowdrift disasters. (4) After snowfalls, the wind is very weak, and the snowfall is less than it is in other areas. So, the probability of the occurrence of snowdrift disasters is much lower than in the other areas.

7. Cumulative snowfall and wind scale in relation to the top ten maximum snowfall events in winter

In section 5, the authors analyzed the cumulative wind scale for the top ten maximum snowfall day at all stations in the Xinjiang Uygur Autonomous Region, without considering the total snowfall of the top ten maximum snowfall events in winter. Therefore, the authors extracted the wind scales of the days on which the top ten maximum snowfall events happened, and the wind scales of the five days after these snowfall events in winter in the Xinjiang Uygur Autonomous Region by using the above data. In this data on the different days at all stations, one day's data at stations 51431, 51470, 51431, 51346 and 51704, and two days' data at stations 51433 is missing. For the missing

data at different stations, we used the average value of the last days. To compare the joint distribution of the total snowfall and the accumulated wind scale, these data are also normalized according to the Formulas 1 and 2. After these processes, the average wind scale, maximum wind scale, normalized total snowfall, and normalized accumulated wind scale are shown in Table 9.

Table 9. Accumulative snowfall and accumulative wind scale taken from the top ten maximum snowfall events at each station in the Xinjiang Uygur Autonomous Region.

Station ID	Total snowing days (d)	Accumulative amount of snowfall (mm)	Accumulative wind scale	Average daily wind scale	Maximum wind scale	Normalized accumulative snowfall	Normalized accumulative wind scale
51053	46	256.1	283	2.948	6	0.519	0.990
51059	48	242.3	212	2.163	6	0.491	0.741
51060	45	212.8	256	2.695	5	0.431	0.895
51068	39	114.1	162	1.820	4	0.231	0.566
51076	59	301.8	151	1.385	3	0.612	0.528
51087	56	365.8	118	1.113	3	0.741	0.413
51133	45	426.8	137	1.442	3	0.865	0.479
51156	31	114.2	130	1.605	4	0.231	0.455
51186	46	304.9	84	0.875	2	0.618	0.294
51232	51	116.9	286	2.832	8	0.237	1.000
51238	28	204.2	98	1.256	2	0.414	0.343
51241	51	171	183	1.812	3	0.347	0.640
51243	33	111.4	131	1.578	5	0.226	0.458
51288	38	146.4	188	2.136	5	0.297	0.657
51330	43	166.1	123	1.323	5	0.337	0.430
51334	48	118.2	99	1.010	3	0.240	0.346
51346	29	222	116.08	1.469	3	0.450	0.406
51358	49	257.8	112	1.131	2	0.522	0.392
51365	24	125.8	120	1.622	3	0.255	0.420
51367	38	173.7	155	1.761	3	0.352	0.542
51379	29	138.2	157	1.987	3	0.280	0.549
51431	53	493.5	143.22	1.390	3	1.000	0.501
51433	53	426.6	134.82	1.309	5	0.864	0.471
51437	46	181.3	142	1.479	3	0.367	0.497
51463	33	263.5	133	1.602	3	0.534	0.465
51467	17	51.2	102	1.522	3	0.104	0.357
51470	37	245.4	197	2.264	4	0.497	0.689
51477	24	53.7	239	3.230	6	0.109	0.836
51495	17	21.9	160	2.388	8	0.044	0.559
51526	16	63.4	100	1.515	4	0.128	0.350
51542	44	85.9	172	1.830	5	0.174	0.601
51567	23	91.8	96	1.315	3	0.186	0.336
51573	13	72.1	66	1.048	2	0.146	0.231
51581	17	81.4	79	1.179	2	0.165	0.276

51628	30	177.2	114	1.425	3	0.359	0.399
51633	41	170.2	86	0.945	3	0.345	0.301
51639	29	100.9	93	1.177	2	0.204	0.325
51642	20	127.1	85	1.214	2	0.258	0.297
51644	31	117.1	124	1.531	2	0.237	0.434
51655	15	58.9	75	1.154	3	0.119	0.262
51656	26	76.1	110	1.447	4	0.154	0.385
51701	61	245.5	255	2.297	5	0.497	0.892
51704	28	249.1	97.13	1.245	3	0.505	0.340
51705	32	193	123	1.500	5	0.391	0.430
51709	30	201.9	108	1.350	2	0.409	0.378
51711	29	182	147	1.861	3	0.369	0.514
51716	18	72.8	87	1.279	3	0.148	0.304
51720	32	130.6	102	1.244	3	0.265	0.357
51730	24	60	82	1.108	2	0.122	0.287
51765	12	42.9	96	1.548	3	0.087	0.336
51777	21	64.7	128	1.803	4	0.131	0.448
51804	28	103.8	73	0.936	2	0.210	0.255
51810	29	100.6	89	1.127	2	0.204	0.311
51811	29	144.7	98	1.241	3	0.293	0.343
51818	32	130.9	115	1.402	3	0.265	0.402
51828	29	114.3	131	1.658	4	0.232	0.458
51839	31	50.3	93	1.148	3	0.102	0.325
51855	20	40.1	87	1.243	3	0.081	0.304
51931	29	72.3	89	1.127	2	0.147	0.311
52101	25	83.9	141	1.880	5	0.170	0.493
52112	25	93.4	152	2.027	4	0.189	0.531
52118	23	73.3	169	2.315	5	0.149	0.591
52203	20	97.4	90	1.286	3	0.197	0.315
52313	20	65.9	190	2.714	5	0.134	0.664

Based on the above normalized snowfall and wind scale data, a joint distribution map of the accumulative snowfall and wind scales at each station in the Xinjiang Uygur Autonomous Region was generated (Figure 8). It can be seen from this map that, after the top ten heavy snowfall events, the accumulative wind scale is high and the cumulative snowfall is large in the Altay Prefecture, the Bortala Mongol Autonomous Prefecture, the Tacheng Prefecture of the northern Xinjiang, and the Kizilsu Kirghiz Autonomous Prefecture of the western Xinjiang. However, in regard to the Altay Prefecture, the accumulative wind scale is higher in its northwestern quadrant than its southeastern counterpart. The cumulative wind scale is high, but the accumulative snowfall is small in the Hami Prefecture. In other areas, both are small.

Figure 8. Joint distribution map of the accumulative snowfall and accumulative wind scale from the top ten maximum snowfall events at each station in the Xinjiang Uygur Autonomous Region.

The maximum wind scale after a heavy snowfall is of great concern because it determines the intensity of the snow drifting disaster. The authors generated the joint distribution map of the accumulative snowfall and maximum wind scale from the top ten maximum snowfall events at each station in the Xinjiang Uygur Autonomous Region, which is shown in Figure 9. From this, we can know that the maximum wind scale after a heavy snowfall is high in the northwestern Altay Prefecture, the western Bortala Mongol Autonomous Prefecture, the western Kizilsu Kirghiz Autonomous Prefecture, the eastern Ili Kazakh Autonomous Prefecture, the eastern Hami city, and the eastern Urumqi. Among the above regions, the snowfall in the Altay region, Ili Kazakh Autonomous Prefecture and Tacheng Prefecture is very heavy. So, these three regions are the hardest hit by the snow drifting disasters. The snowfall is not very heavy in the western Bortala Mongol Autonomous Prefecture, the western Kizilsu Kirghiz Autonomous Prefecture, the eastern section of Hami city, and the eastern neighborhoods in Urumqi. However, if heavy snowfall occurs, which provides plenty of snow particles, it's very easy to pose a snow drifting disaster. These are the key regions in terms of snow drifting disasters and deserve our focus. Other areas are not as peculiarly prone to this type of disaster.

Figure 9. Joint distribution map of the accumulative snowfall and max wind scale from the top ten maximum snowfall events at each station in the Xinjiang Uygur Autonomous Region.

8. Conclusions

This study assessed the spatial distribution characteristics of wind and snowfall in the Xinjiang Uygur Autonomous Region of China based on the “Daily Data Sets of Climate Data for China International Surface Exchange Station (V3.0)”. The following conclusions can be drawn from this analysis.

(1) The weighted accumulated wind scale is large in the northwest of Altay, the east and west of Hami and at the junction of Urumqi and Turpan. However, it’s relatively small in other regions.

(2) The distribution map of the basic wind speed in winter at the 50-year reoccurrence period in the Xinjiang Uygur Autonomous Region was generated, which can be used in building structure design.

(3) The snowfall areas of the Xinjiang Uygur Autonomous Region are mainly in the north, and the heaviest snowfalls are in the Yili Kazak Autonomous and the northwestern Tacheng Prefectures.

(4) After heavy snowfall, the wind is very strong in northwestern Altay, the northwestern section of the Tacheng Prefecture and Urumqi city. So, the snowdrift disasters are very serious if they occur in these areas.

(5) After the top ten heavy snowfall events, the accumulative wind scale is high and the accumulative snowfall is large in the Altay Prefecture, the Bortala Mongol Autonomous Prefecture, and the Tacheng Prefecture of the northern Xinjiang, and the Kizilsu Kirghiz Autonomous Prefecture of the western Xinjiang.

Acknowledgement

The research reported herein was sponsored by the National Natural Science Foundation of China

(Grant No. 51778381) and the Hebei Province Natural Science Fund (Grant No. D2016210008 and Grant No. E2018210044). We would like to thank TopEdit (www.topeditsci.com) for English language editing of this manuscript.

Competing interests

The authors declare no competing interests.

Reference

- National Meteorological Centre (2012). GB/T 28591-2012 Wind Scale. Beijing, Standards Press of China. **GB/T 28591-2012**.
- National Meteorological Centre (2017). GB/T 21984-2017 Short-Range Weather Forecast. Beijing, Standards Press of China. **GB/T 21984-2017**.
- Chen, H. and X. Fan (2009). "Some Extreme Events of Weather, Climate and Related Phenomena in 2008." Climatic and Environmental Research **14**(3): 329-340. (in Chinese with English Abstract)
- Li, X., J. Zhu, Y. Jia, J. Liu, N. Li and F. Li (2007). "Formation process of extraordinarily serious snowstorm and its induced damage in 2007 in Liaoning Province." Chinese Journal of Ecology(8): 1250-1258. (in Chinese with English Abstract)
- Liu, J., X. Yao, Z. Han, P. Xie, L. Liu and P. Liu (2008). "Analysis on the cause and type of wind blowing snow on Highways in Xinjiang." Highway(1): 25-27. (in Chinese)
- Liu, W., X. Liu and H. Chen (2010). "On the Successful Shaping of Local Government Image in Blizzard Disaster: an Example of Xinjiang Blizzard Disaster in 2010." Journal of Institute of Disaster Prevention **12**(2): 119-121+136. (in Chinese with English Abstract)
- China Academy of Building Research (2012). Load code for the design of building structures, China Architecture & Building Press. **GB 50009-2012**.
- Su, G. (2017). Research on Numerical simulation of Wind blowing snow at railway cut and disaster prevention and control engineering Master, Lanzhou Jiaotong University. (in Chinese with English Abstract)
- Wang, T. (2009). Study on numerical simulation and prevention measures of the drifting snow disaster along railway Master, Lanzhou University. (in Chinese with English Abstract)
- Ying, C. (2007). Research on the Countermeasures of the Blowing Snow Hazards on Highways Doctor, Jilin University. (in Chinese with English Abstract)
- Zha, M., Z. Yu and H. Li (2012). "Study on the prevention against snow hazard for the highway between Karamay and Tacheng." Journal of China & Foreign Highway **32**(4): 67-71. (in Chinese)

Figures

Figure 1

Meteorological station distribution map in the Xinjiang Uygur Autonomous Region. Note: The designations employed and the presentation of the material on this map do not imply the expression of any opinion whatsoever on the part of Research Square concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. This map has been provided by the authors.

Distribution map of the weighted accumulative wind scale in the Xinjiang Uygur Autonomous Region

Figure 2

Distribution map of the weighted accumulative wind scale in the Xinjiang Uygur Autonomous Region

Note: The designations employed and the presentation of the material on this map do not imply the expression of any opinion whatsoever on the part of Research Square concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. This map has been provided by the authors.

Figure 3

Maximum wind direction rose diagrams for some stations in the Xinjiang Uygur Autonomous Region

Distribution map of the basic wind speed in winter at the 50-year reoccurrence period in the Xinjiang Uygur Autonomous Region

Figure 4

Distribution map of the basic wind speed in winter at the 50-year reoccurrence period in the Xinjiang Uygur Autonomous Region Note: The designations employed and the presentation of the material on this map do not imply the expression of any opinion whatsoever on the part of Research Square concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. This map has been provided by the authors.

Distribution map of weighted accumulative snowfall grade in the Xinjiang Uygur Autonomous Region

Figure 5

Distribution map of weighted accumulative snowfall grade in the Xinjiang Uygur Autonomous region

Note: The designations employed and the presentation of the material on this map do not imply the expression of any opinion whatsoever on the part of Research Square concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. This map has been provided by the authors.

Distribution map of the accumulative wind scale from the top ten maximum snowfall days at all stations in the Xinjiang Uygur Autonomous Region

Figure 6

Distribution map of the accumulative wind scale from the top ten maximum snowfall days at all stations in the Xinjiang Uygur Autonomous Region Note: The designations employed and the presentation of the material on this map do not imply the expression of any opinion whatsoever on the part of Research Square concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. This map has been provided by the authors.

Figure 7

Joint distribution map of the total snowfall for the top ten maximum snowfall days and the accumulative wind scale for six days (including the day of snowfall) after the top ten maximum snowfall day at all stations in the Xinjiang Uygur Autonomous Region Note: The designations employed and the presentation of the material on this map do not imply the expression of any opinion whatsoever on the part of Research Square concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. This map has been provided by the authors.

Figure 8

Joint distribution map of the accumulative snowfall and accumulative wind scale from the top ten maximum snowfall events at each station in the Xinjiang Uygur Autonomous Region. Note: The designations employed and the presentation of the material on this map do not imply the expression of any opinion whatsoever on the part of Research Square concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. This map has been provided by the authors.

Figure 9

Joint distribution map of the accumulative snowfall and max wind scale from the top ten maximum snowfall events at each station in the Xinjiang Uygur Autonomous Region. Note: The designations employed and the presentation of the material on this map do not imply the expression of any opinion whatsoever on the part of Research Square concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. This map has been provided by the authors.