

# The impact of atrial fibrillation on one-year mortality in patients with severe lower extremity arterial disease

Min-I Su

Taitung Mackay Memorial Hospital

Cheng-Wei Liu (✉ [issac700319@gmail.com](mailto:issac700319@gmail.com))

Tri-Service General Hospital Songshan Branch

---

## Research Article

**Keywords:** Afib, LEAD, MALEs, PAD, CHADS2

**Posted Date:** February 25th, 2021

**DOI:** <https://doi.org/10.21203/rs.3.rs-243651/v1>

**License:**  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

**Version of Record:** A version of this preprint was published at Journal of Clinical Medicine on March 31st, 2022. See the published version at <https://doi.org/10.3390/jcm11071936>.

# Abstract

## Background

and Aims: Atrial fibrillation (Afib) is associated with the incidence of lower extremity arterial disease (LEAD), but its effect on severe LEAD prognosis remains unclear. We investigated the association between Afib and clinical outcomes.

## Methods and Results

We retrospectively enrolled consecutive severe LEAD patients receiving percutaneous transluminal angioplasty between 2013/1/1 and 2018/12/31. Patients were divided by a history of any type of Afib and followed for at least one year. The primary outcome was all-cause mortality. Secondary outcomes were cardiac-related mortality, major adverse cardiovascular events (MACEs), and major adverse limb events (MALEs). The study included 222 patients aged  $74 \pm 11$  years (54% male), and 12.6% had acute limb ischemia. The Afib group had significantly higher rates of all-cause mortality (42.9% vs. 20.1%,  $P = 0.014$ ) and MACEs (32.1% vs. 14.4%,  $P = 0.028$ ) than the non-Afib group. After adjustment for confounders, Afib was independently associated with all-cause mortality (adjusted HR: 2.153, 95% CI: 1.084–4.276,  $P = 0.029$ ) and MACEs (adjusted HR: 2.338, 95% CI: 1.054–2.188,  $P = 0.037$ ).

## Conclusions

Afib was significantly associated with increased risks of one-year all-cause mortality and MACEs in severe LEAD patients. Future studies should investigate whether oral anticoagulants benefit these patients.

## Introduction

Atrial fibrillation and peripheral artery disease (PAD) are common atherosclerotic consequences that interact with each other. Patients with atrial fibrillation have a higher incidence of PAD than those without atrial fibrillation<sup>1</sup>. Conversely, PAD is also associated with an increased risk of developing atrial fibrillation<sup>2</sup>. A nationwide cohort study showed that PAD was independently associated with new-onset atrial fibrillation<sup>3</sup>, and another study showed that patients with atrial fibrillation had a higher incidence of PAD<sup>1</sup>. One type of severe complications of atrial fibrillation is thromboembolic events, which are frequently diagnosed in PAD patients. These episodes are not limited to cerebrovascular events and include peripheral artery thromboembolic complications, such as splenic, renal, and mesenteric artery infarction, and acute limb ischemia<sup>4</sup>. Overall mortality and cardiac-related mortality were also found to be elevated in patients with atrial fibrillation and symptomatic PAD independent of age and comorbidities<sup>5</sup>. Lower extremity arterial disease (LEAD), defined by an ankle-brachial index  $< 0.9$ , was found to be associated with two-fold higher risks of myocardial infarction and vascular death in patients with atrial fibrillation<sup>6</sup>. Baseline atrial fibrillation was reported to be associated with an increased risk of incident

acute limb ischemia in a large-scale case-control study<sup>7</sup>. Patients with LEAD without a medical history of atrial fibrillation at baseline had an increased incidence of atrial fibrillation in a population-based cohort study<sup>8</sup>. Regarding atrial fibrillation and LEAD, it remains unclear which comes first<sup>3,9</sup>, but it is known that patients with concomitant atrial fibrillation and PAD have a synergistically elevated risk of cardiac mortality<sup>10</sup>. To date, the effect of atrial fibrillation on prognosis has rarely been reported in patients with severe LEAD. Therefore, we conducted the present study to comprehensively investigate mortality and cardiovascular outcomes in patients with severe LEAD and atrial fibrillation.

## Results

We initially enrolled 225 patients who presented with severe LEAD and excluded three patients who refused surgical amputation. The study cohort consisted of 222 patients aged 74±11 years (54% male), and 12.6% had acute limb ischemia. The patients with atrial fibrillation were significantly older than those without atrial fibrillation (80.6±9.1 vs. 72.6±11.5 years,  $P = 0.001$ ), were more likely to have a history of ischemic stroke (28.6% vs. 13.9%,  $P = 0.056$ ), were more likely to have acute limb ischemia (25.0% vs. 10.8%,  $P = 0.035$ ), had higher CHADS2 scores (2.9±1.3 vs. 2.2±1.2,  $P = 0.004$ ) and higher serum uric acid levels (6.8±2.9 vs. 5.7±2.2 mg/dl,  $P = 0.022$ ). The patients with atrial fibrillation had lower prevalences of chronic kidney disease and end-stage renal disease and lower serum creatinine levels than those without atrial fibrillation (1.8±1.1 vs. 3.7±3.5 mg/dl,  $P < 0.001$ ). Table 1 describes the details of the baseline characteristics of the study population.

During the study period, the patients with atrial fibrillation had higher incidences of all-cause mortality (42.9% vs. 20.1%,  $P = 0.014$ ), cardiac-related mortality (21.4% vs. 10.3%,  $P = 0.111$ ), MACEs (32.1% vs. 14.4%,  $P = 0.028$ ), MALEs (17.9% vs. 14.9%,  $P = 0.778$ ) and in-hospital mortality (14.3% vs. 6.7%,  $P = 0.242$ ) than those without atrial fibrillation. In univariate logistic regression analyses, atrial fibrillation was significantly associated with increased risks of all-cause mortality (crude hazard ratio (cHR): 2.435, 95% CI: 1.274-4.654,  $P = 0.007$ ) and MACEs (cHR: 2.479, 95% CI: 1.169-5.257,  $P = 0.018$ ). There was a tendency towards a significant association between atrial fibrillation and cardiac-related mortality (cHR: 2.436, 95% CI: 0.977-6.072,  $P = 0.056$ ). No significant associations of atrial fibrillation with MALEs (cHR: 1.280, 95% CI: 0.495-3.307,  $P = 0.610$ ) and in-hospital mortality (cHR: 2.120, 95% CI: 0.691-6.503,  $P = 0.189$ ) were found. Acute limb ischemia and Rutherford stage were both associated with all-cause mortality (cHR: 4.133, 95% CI: 2.257-7.567,  $P = 0.003$  and cHR: 2.073, 95% CI: 1.330-3.233,  $P = 0.001$ ); acute limb ischemia was also associated with cardiac-related mortality (cHR: 3.769, 95% CI: 1.579-8.998,  $P = 0.003$ ) and MACEs (cHR: 2.592, 95% CI: 1.222-5.497,  $P = 0.013$ ), and Rutherford stage was associated with MALEs (cHR: 4.227, 95% CI: 2.339-7.636,  $P < 0.001$ ). We show the other confounders associated with the study outcomes in Table 2.

In multivariate logistic regression analyses, atrial fibrillation remained significantly associated with all-cause mortality (adjusted HR (aHR): 2.193, 95% CI: 1.109-4.336,  $P = 0.024$ ) and MACEs (aHR: 2.322, 95% CI: 1.045-5.158,  $P = 0.039$ ), independent of acute limb ischemia and Rutherford stage. We did not find a significant association between atrial fibrillation and cardiac-related mortality (aHR: 2.146, 95% CI: 0.819-

5.626,  $P = 0.120$ ) or between atrial fibrillation and MALEs (aHR: 1.757, 95% CI: 0.664-4.789,  $P = 0.271$ ) after adjusting for the confounders. Figure 1 shows a Kaplan-Meier curve. Acute limb ischemia (aHR: 2.872, 95% CI: 1.491-5.533,  $P = 0.002$ ) and Rutherford stage (aHR: 1.918, 95% CI: 1.186-3.103,  $P = 0.008$ ) were significantly associated with all-cause mortality but not cardiac-related mortality or MACEs. Rutherford stage was significantly associated with MALEs (aHR: 5.577, 95% CI: 2.780-11.19,  $P < 0.001$ ). Heart rate at presentation was significantly associated with all-cause and cardiac-related mortality and MACEs (Table 2).

In the sensitivity analyses, we adjusted for kidney function and body mass index, although these factors were not significantly associated with the study outcomes in univariate logistic regression analyses. The significant association between atrial fibrillation and all-cause mortality remained unchanged after adjustment for kidney function, represented by the creatinine level, creatinine clearance, and the estimated glomerular filtration rate. After we adjusted for body mass index and kidney function in various statistical models, atrial fibrillation was still associated with one-year mortality (aHR: 2.158, 95% CI: 1.012-4.605,  $P = 0.047$  for creatinine clearance and aHR: 2.209, 95% CI: 1.036-4.710,  $P = 0.040$  for the estimated glomerular filtration rate), but we found only a tendency toward a significant association when the creatinine level was used (aHR: 2.080, 95% CI: 0.950-4.555,  $P = 0.067$ ). Table 3 shows the results of the sensitivity analyses.

## Discussion

In the present study, we showed that in addition to acute limb ischemia and Rutherford stage, atrial fibrillation was an independent predictor of all-cause mortality. The patients with atrial fibrillation had lower creatinine levels but a higher incidence of all-cause mortality than the patients without atrial fibrillation. In the various statistical models, atrial fibrillation doubled the risk of all-cause mortality, and the significant associations were fairly consistent, although only a tendency toward significance was found after we adjusted for body mass index in addition to the other cofounders. In our study, we did not find a significant difference in body mass index between the patients with and without atrial fibrillation, and we therefore did not initially adjust for it as a confounder in our reduced statistical models. The effect of body mass index on mortality remains controversial in patients with LEAD<sup>11,12</sup>. Body mass index was found to have a complex inverse association with the three-year survival rate in patients with critical limb ischemia undergoing endovascular treatment for infrapopliteal lesions in a previous study: overweight patients had the highest one-year survival rate (86.1%), followed by normal-weight patients (78.6%), and underweight patients had the lowest mortality rate (68.4%)<sup>12</sup>. In contrast, the Atherosclerosis Risk in Communities (ARIC) study showed that body mass index was positively associated with the incidence of peripheral artery disease requiring hospitalization, and the risk was relatively higher in patients with critical limb ischemia<sup>11</sup>. We believe that a low body mass index value ( $< 20 \text{ kg/m}^2$ ) indirectly reflects a poor nutritional status and compromised wound healing in patients with severe LEAD, whereas a high body mass index value ( $> 25 \text{ kg/m}^2$ ) in this population is generally associated with a

greater prevalence of cardiovascular comorbidities, such as hypertension, diabetes, and dyslipidemia. Therefore, we suggest that it is beneficial for patients with severe LEAD to maintain a normal weight<sup>11</sup>.

In this study, we enrolled patients with severe LEAD, and all of our patients with atrial fibrillation had CHA<sub>2</sub>DS<sub>2</sub>Vsc scores  $\geq 3$ . According to the current guidelines, oral anticoagulants are strongly recommended for patients with nonvalvular atrial fibrillation and a CHA<sub>2</sub>DS<sub>2</sub>Vsc score  $\geq 2$ <sup>13,14</sup>. To date, no randomized controlled trial has been published to determine which antithrombotic regimen is best in patients with simultaneous atrial fibrillation and severe LEAD. The ROCKET-AF trial showed that patients with atrial fibrillation and LEAD did not have an elevated risk of stroke or systemic embolization, and a higher risk of major bleeding or nonmajor clinically relevant bleeding was found in patients treated with rivaroxaban than in those treated with warfarin<sup>15</sup>. The ARISTOTLE trial showed that patients with or without atrial fibrillation did not have a significantly different risk of stroke or systemic embolization, and the benefits of apixaban and warfarin were similar<sup>16</sup>. However, the two studies enrolled patients with atrial fibrillation and performed a subgroup analysis of patients with LEAD<sup>15,16</sup>, which is quite different from a study that enrolled LEAD patients with or without atrial fibrillation, based on the different natures of atrial fibrillation and LEAD. Although these two diseases have concomitant risk factors and share the characteristics of an increased inflammatory burden and a prothrombotic state, endothelial dysfunction and atherosclerosis play major roles in LEAD but not atrial fibrillation<sup>17</sup>. The VOYAGER trial showed that very-low-dose rivaroxaban in addition to aspirin compared with aspirin alone significantly reduced the incidence of acute limb ischemia in patients with peripheral artery disease after revascularization, but all-cause mortality did not differ significantly between the two groups. Notably, the VOYAGER trial excluded patients with atrial fibrillation and patients with major tissue loss<sup>18</sup>, but we did not exclude patients with chronic ulcers or major tissue loss of the lower extremities; instead, we enrolled all-comers. Based on the similar pathophysiology of coronary artery disease and LEAD, patients with atrial fibrillation who received percutaneous coronary stenting could be included in our study, although the percentage of patients with LEAD was not reported in previous studies<sup>19–22</sup>. The current evidence from four major randomized controlled trials showed that direct oral anticoagulants (DOACs) plus P2Y<sub>12</sub> inhibitors can prevent bleeding complications in patients with atrial fibrillation who receive coronary stent(s) without increasing the risk of stent thromboses. Although the selection of the proper antithrombotic regimen for patients with atrial fibrillation and severe LEAD is far beyond the scope of this study, our results showed that atrial fibrillation increased the risk of mortality, and it is worth investigating the effect of antithrombotic treatment in this population in the future.

This study had several limitations. First, we enrolled patients with severe LEAD, so the number of patients in our study population was relatively small. Therefore, we lacked adequate power to identify a significant association between atrial fibrillation and the other study outcomes after we adjusted for confounders. However, in the ROCKET-AF trial and the ARISTOTLE trial had 884 and 839 atrial fibrillation patients with a medical history of LEAD, respectively<sup>15,16</sup>. We did not find a significant association between atrial fibrillation and MACEs, including cardiac-related mortality, nonfatal myocardial infarction, and nonfatal stroke. However, a systematic review and meta-analysis showed that atrial fibrillation was associated

with an increased risk of MACE<sup>23</sup>. Second, our study enrolled all-comers. We noted that a relatively small proportion of the patients had received secondary prevention for dyslipidemia before they presented at our hospital. In addition, the percentage of the patients who had been prescribed oral anticoagulants seemed low in the present study. In our study, 53% of the patients with atrial fibrillation had a creatinine clearance value less than 30 ml/min, and the net clinical benefits should be balanced between antithrombotic treatment and bleeding complications in these patients. We will try to enroll additional patients with atrial fibrillation and prospectively conduct a study to investigate the effect of antithrombotic treatment on the study outcomes.

In conclusion, our retrospective cohort study showed that patients with severe LEAD and atrial fibrillation had higher incidences of all-cause and cardiac-related mortality and MACEs than those without atrial fibrillation. We showed the adverse effect of atrial fibrillation on all-cause mortality; acute limb ischemia and Rutherford stage were also associated with all-cause mortality. Future studies should investigate the effect of antithrombotic treatment on mortality in this study population.

## Methods

We retrospectively enrolled consecutive patients with severe LEAD who received percutaneous transluminal angioplasty at our hospital between 2013/1/1 and 2018/12/31. The present study enrolled all-comers; we excluded three patients with critical limb ischemia and unsalvageable limbs who refused surgical amputation. The patients' baseline characteristics, laboratory data, procedural details, and outcomes were collected from the medical records. We followed the patients until 2019/12/31. The study was approved by the Mackay Memorial Hospital under Institutional Review Board number 20MMHIS034e. The informed consent was waived by the Mackay memorial hospital institutional review board because the present study is a low-risk retrospective cohort study. We conducted the present study in accordance with the Declaration of Helsinki.

We performed emergency and urgent percutaneous transluminal angioplasty on patients with acute and chronic limb ischemia, respectively; elective percutaneous transluminal angioplasty was reserved for those patients with Rutherford stage IV disease. Physicians could decide whether to use antegrade or retrograde vascular access based on the type of lesion. During the procedure, we routinely administered heparin to maintain an active clotting time between 250 and 300 seconds and dual antiplatelet agents, namely, aspirin and clopidogrel, according to the current guidelines. Generally, we treated iliac and femoropopliteal lesions with balloon angioplasty, a drug-coated balloon or a stent, and we treated below-knee lesions with only balloon angioplasty. The decision to use a drug-coated balloon or stent was mainly influenced by the patient's economic status and left to the physicians' discretion. After the procedure was performed, we continued to administer dual antiplatelet agents for at least one month except in cases of clinically relevant bleeding complications. Because cilostazol has minimal effects on limb salvage and mortality, we did not routinely prescribe it to severe LEAD patients.

We defined severe LEAD based on the Rutherford classification system as follows: resting pain (stage IV), tissue loss (stage V), and gangrene (stage VI)<sup>24</sup>. The primary study outcome was all-cause mortality at the one-year follow-up. The secondary outcomes were cardiac-related mortality, major adverse cardiac events (MACEs) and major adverse limb events (MALEs) at the one-year follow-up and in-hospital mortality. We defined MACEs as the composite of cardiac-related mortality, nonfatal myocardial infarction, and nonfatal stroke and MALEs as amputation due to a vascular event above the forefoot, acute limb ischemia and clinically driven target vessel revascularization.

We present continuous variables as numbers and standard deviations and binary variables as numbers and percentages. We used independent t-tests and chi-squared tests to evaluate differences in continuous variables and binary variables, respectively. We first used univariate logistic regression analyses to identify potential confounders. Then, we adjusted for these confounders in multivariate logistic regression analyses of the relationships between the presence of atrial fibrillation and the study outcomes. In the statistical models, we adjusted for the presence of acute limb ischemia and Rutherford classification because these factors obviously contributed to the study outcomes. In the sensitivity analyses, we additionally adjusted for kidney function and body mass index if they were not associated with the study outcomes in multivariate logistic regression. Kidney function was represented by the serum creatinine level, creatinine clearance as determined with the Cockcroft-Gault formula, or the estimated glomerular filtration rate as determined by the Modification of Diet in Renal Disease study equation, as in our previous study<sup>25</sup>. Body mass index was divided by the thresholds of 20 and 25 into groups as follows: <20,  $\geq 20$  and <25, and  $\geq 25$ <sup>11,26</sup>. We calculated two-tailed P values, and we considered values of 0.05 or lower to be significant. We performed all statistical analyses with Statistical Package for the Social Sciences software (SPSS software, version 20.0).

### **Data Availability**

We can provide the anonymous study data if the corresponding author judges a request as reasonable.

## **Declarations**

### **Acknowledgments**

None.

### **Author Contributions**

Min-I Su designed the study and collected the data. Cheng-Wei Liu performed the statistical analyses, wrote the manuscript, and is responsible for correspondence.

### ***Competing Interests***

The authors declare no competing interests.

## References

1. Chang, C. J. *et al.* Atrial fibrillation increases the risk of peripheral arterial disease with relative complications and mortality: a population-based cohort study. *Medicine (Baltimore)***95**, e3002 (2016).
2. Bekwelem, W. *et al.* Association of peripheral artery disease with incident atrial fibrillation: the ARIC (Atherosclerosis Risk in Communities) study. *J. Am. Heart Assoc.***7**, e007452 (2018).
3. Lin, Y. S. *et al.* Peripheral arterial disease and atrial fibrillation and risk of stroke, heart failure hospitalization and cardiovascular death: a nationwide cohort study. *Int. J. Cardiol.***203**, 204-211 (2016).
4. Wasilewska, M. & Gosk-Bierska, I. Thromboembolism associated with atrial fibrillation as a cause of limb and organ ischemia. *Adv. Clin. Exp. Med.***22**, 865-873 (2013).
5. Winkel, T. A. *et al.* Prognosis of atrial fibrillation in patients with symptomatic peripheral arterial disease: data from the REduction of Atherothrombosis for Continued Health (REACH) Registry. *Eur. J. Vasc. Endovasc. Surg.***40**, 9-16 (2010).
6. Violi, F. *et al.* Ankle-Brachial index and cardiovascular events in atrial fibrillation. The ARAPACIS study. *Thromb. Haemost.***115**, 856-863 (2016).
7. Hess, C. N. *et al.* Acute limb ischemia in peripheral artery disease. *Circulation***140**, 556-565 (2019).
8. Griffin, W. F., Salahuddin, T., O'Neal, W. T. & Soliman, E. Z. Peripheral arterial disease is associated with an increased risk of atrial fibrillation in the elderly. *Europace***18**, 794-798 (2016).
9. O'Neal, W. T. *et al.* Peripheral arterial disease and risk of atrial fibrillation and stroke: the multi-ethnic study of atherosclerosis. *J. Am. Heart Assoc.***3**, e001270 (2014).
10. Goto, S. *et al.* Prevalence, clinical profile, and cardiovascular outcomes of atrial fibrillation patients with atherothrombosis. *Am. Heart J.***156**, 855-863, 863.e2 (2008).
11. Hicks, C. W. *et al.* Associations of obesity with incident hospitalization related to peripheral artery disease and critical limb ischemia in the ARIC study. *J. Am. Heart Assoc.***7**, e008644 (2018).
12. Murata, N. *et al.* Complex relationship of body mass index with mortality in patients with critical limb ischemia undergoing endovascular treatment. *Eur. J. Vasc. Endovasc. Surg.***49**, 297-305 (2015).
13. Imamura, T. & Kinugawa, K. Novel rate control strategy with landiolol in patients with cardiac dysfunction and atrial fibrillation. *ESC Heart Fail.***7**, 2208-2213 (2020).
14. January, C. T. *et al.* 2019 AHA/ACC/HRS focused update of the 2014 AHA/ACC/HRS guideline for the management of patients with atrial fibrillation: a report of the American college of cardiology/American heart association task force on clinical practice guidelines and the heart rhythm society in collaboration with the society of thoracic surgeons. *Circulation***140**, e125-e151 (2019).
15. Jones, W. S. *et al.* Efficacy and safety of rivaroxaban compared with warfarin in patients with peripheral artery disease and non-valvular atrial fibrillation: insights from ROCKET AF. *Eur. Heart J.***35**, 242-249 (2014).

16. Hu, P. T. *et al.* Efficacy and safety of apixaban compared with warfarin in patients with atrial fibrillation and peripheral artery disease: insights from the ARISTOTLE trial. *J. Am. Heart Assoc.***6**, e004699 (2017).
17. Proietti, M. Is there a relationship between atrial fibrillation and peripheral arterial disease? <https://www.escardio.org/Journals/E-Journal-of-Cardiology-Practice/Volume-16/Is-there-a-relationship-between-atrial-fibrillation-and-peripheral-arterial-disease> (2018).
18. Bonaca, M. P. *et al.* Rivaroxaban in peripheral artery disease after revascularization. *N. Engl. J. Med.***382**, 1994-2004 (2020).
19. Cannon, C. P. *et al.* Dual antithrombotic therapy with dabigatran after PCI in atrial fibrillation. *N. Engl. J. Med.***377**, 1513-1524 (2017).
20. Gibson, C. M. *et al.* Prevention of bleeding in patients with atrial fibrillation undergoing PCI. *N. Engl. J. Med.***375**, 2423-2434 (2016).
21. Lopes, R. D. *et al.* Antithrombotic therapy after acute coronary syndrome or PCI in atrial fibrillation. *N. Engl. J. Med.***380**, 1509-1524 (2019).
22. Vranckx, P. *et al.* Edoxaban-based versus vitamin K antagonist-based antithrombotic regimen after successful coronary stenting in patients with atrial fibrillation (ENTRUST-AF PCI): a randomised, open-label, phase 3b trial. *Lancet***394**, 1335-1343 (2019).
23. Odutayo, A. *et al.* Atrial fibrillation and risks of cardiovascular disease, renal disease, and death: systematic review and meta-analysis. *BMJ***354**, i4482 (2016).
24. Shishehbor, M. H. *et al.* Critical limb ischemia: an expert statement. *J. Am. Coll. Cardiol.***68**, 2002-2015 (2016).
25. Liu, C. W. *et al.* The dose-response effects of uric acid on the prevalence of metabolic syndrome and electrocardiographic left ventricular hypertrophy in healthy individuals. *Nutr. Metab. Cardiovasc. Dis.***29**, 30-38 (2019).
26. Global BMI Mortality Collaboration *et al.* Body-mass index and all-cause mortality: individual-participant-data meta-analysis of 239 prospective studies in four continents. *Lancet***388**, 776-786 (2016).

## Tables

**Table 1. Baseline and procedural characteristics and laboratory data in patients with severe lower extremity arterial diseases**

	No Afib		With Afib		P
	N = 194		N = 28		
Age (years)	72.6	(11.5)	80.6	(9.1)	0.001
Male gender	107	55.2%	12	42.9%	0.232
Body mass index (kg/m <sup>2</sup> )	23.8	(4.2)	22.8	(3.9)	0.252
Heart rate at baseline (beats per minute)	87.4	(17.2)	90.7	(19.6)	0.349
Systolic BP at baseline (mm Hg)	148.9	(31.1)	141.8	(27.8)	0.253
Diastolic BP at baseline (mm Hg)	75.6	(14.3)	73.6	(13.1)	0.472
Current/past smoker	49	25.3%	4	14.2%	0.275
Alcohol intake	62	31.9%	7	25.0%	0.678
Family history of premature CAD	2	1%	0	0%	1.000
History of hypertension	124	63.9%	21	75.0%	0.293
History of diabetes mellitus	133	68.6%	17	60.7%	0.398
History of insulin use	27	13.9%	1	3.6%	0.218
History of dyslipidemia	42	21.6%	4	14.3%	0.461
History of kidney disease					0.011
Normal kidney function	117	60.3%	24	85.7%	
Chronic kidney disease	32	16.5%	4	14.3%	
End-stage renal disease	45	23.2%	0	0	
History of CAD	81	41.8%	9	32.1%	0.412
History of myocardial infarction	12	6.2%	2	7.1%	0.692
History of chronic heart failure					0.684
NYHA class I	8	4.1%	2	7.1%	
NYHA class II	10	5.2%	2	7.1%	
NYHA class III	12	6.2%	3	10.7%	
NYHA class IV	5	2.6%	0	0%	
History of carotid artery stenosis	3	1.5%	0	0%	1.000

History of ischemic stroke	27	13.9%	8	28.6%	0.056
History of any cancer	10	5.2%	3	10.7%	0.216
History of amputation	13	6.7%	0	0%	0.949
Above-knee amputation	6	3.1%	0	0%	
Below-knee amputation	3	1.5%	0	0%	
Forefoot amputation	4	2.1%	0	0%	
Presented with acute ischemic limb	21	10.8%	7	25.0%	0.035
Rutherford classification					0.663
Class III	23	11.9%	2	7.1%	
Class IV	45	23.2%	9	32.1%	
Class V	114	58.8%	16	57.1%	
Class VI	12	6.2%	1	3.6%	
CHADS2 score	2.2	(1.2)	2.9	(1.3)	0.004
Laboratory data					
Total cholesterol (mg/dl)	160	(47.0)	148.0	(37.0)	0.204
High-density lipoprotein cholesterol (mg/dl)	40.0	(16.0)	42.0	(27.0)	0.815
Low-density lipoprotein cholesterol (mg/dl)	95.0	(37.0)	93.0	(30.0)	0.902
Triglyceride (mg/dl)	158.0	(125.0)	105.0	(67.0)	0.033
Fasting glucose (mg/dl)	176.0	(99.0)	177.0	(116.0)	0.965
Glycosylated hemoglobin (%)	7.3	(1.9)	7.9	(2.1)	0.257
Creatinine (mg/dl)	3.7	(3.5)	1.8	(1.1)	<0.001
Creatinine clearance (ml/min)	34.1	(31.0)	33.3	(19.4)	0.852
Estimated glomerular filtration rate (ml/min/1.73 <sup>2</sup> )	41.6	(37.0)	46.2	(28.7)	0.448
Alanine transaminase (IU/L)	22.0	(19.0)	30.0	(31.0)	0.150
Uric acid (mg/dl)	5.7	(2.2)	6.8	(2.9)	0.022
White blood cell count 10 <sup>3</sup> /μl	9645	(4970)	8593	(4623)	0.029
Neutrophil ratio (%)	70.5	(13.3)	67.7	(17.4)	0.330

Lymphocyte ratio (%)	17.1	(10.1)	17.0	(8.4)	0.969
NLR	7.9	(10.0)	6.3	(6.4)	0.421
<b>Medication use at baseline</b>					
Aspirin	70	36.1%	12	42.9%	0.533
Cilostazol	93	47.9%	15	53.6%	0.687
Clopidogrel	57	29.4%	4	14.3%	0.115
Oral anti-coagulants	0	0%	7	25.0%	<0.001
Pentoxifylline	8	4.1%	3	10.7%	0.148
ACEI or ARB	9	4.6%	2	7.1%	0.634
Beta-blocker	33	17.0%	6	21.4%	0.596
Calcium channel blocker	39	20.1%	9	32.1%	0.149
Statin	42	21.6%	2	7.1%	0.080
Urate lowering therapy	3	1.5%	2	7.1%	0.121

Values are expressed as numbers (standard deviation) or numbers and percentages.

Afib = atrial fibrillation; CAD = coronary artery disease; NYHA = New York Heart Association; BP = blood pressure; NLR = neutrophil-to-lymphocyte ratio; ACEI = angiotensin-converting enzyme inhibitor; ARB = angiotensin receptor blocker

**Table 2. Variables associated with study outcomes in patients with critical limb ischemia in logistic regression analyses**

	Crude hazard ratio	95% confidence interval		P	Adjusted hazard ratio	95% confidence interval		P
All-cause mortality at one year								
Atrial fibrillation	2.435	1.274	4.654	0.007	2.193	1.109	4.336	0.024
Age (years)	1.033	1.006	1.060	0.016	1.018	0.991	1.046	0.200
Male gender	0.678	0.390	1.177	0.167	0.693	0.389	1.235	0.214
Body mass index (kg/m <sup>2</sup> )	0.918	0.850	0.990	0.027				
Smoking	0.925	0.558	1.532	0.762				
Heart rate (beats per minute)	1.026	1.011	1.042	0.001	1.019	1.003	1.034	0.017
Systolic blood pressure (mm Hg)	0.995	0.986	1.005	0.325				
Diastolic blood pressure (mm Hg)	1.013	0.992	1.033	0.226				
Hypertension (yes or no)	0.675	0.388	1.175	0.164				
Dyslipidemia (yes or no)	0.777	0.440	1.370	0.383				
Chronic heart failure (yes or no)	1.021	0.512	2.039	0.952				
Ischemic stroke (yes or no)	1.361	0.681	2.717	0.383				
Acute ischemic limb (yes or no)	4.133	2.257	7.567	<0.001	2.872	1.491	5.533	0.002
Rutherford classification (IV~VI)	2.073	1.330	3.233	0.001	1.918	1.186	3.103	0.008
Creatinine (mg/dl)	1.018	0.943	1.099	0.640				
Triglyceride (mg/dl)	0.997	0.993	1.001	0.095				
Uric acid (mg/dl)	1.017	0.888	1.164	0.810				
NLR	1.030	1.016	1.044	<0.001	1.032	1.015	1.050	<0.001
CHADS2 score	1.016	0.804	1.285	0.891				

Cardiac-related mortality at one year								
Atrial fibrillation	2.436	0.977	6.072	0.056	2.146	0.819	5.626	0.120
Age (years)	1.031	0.994	1.069	0.100	1.017	0.980	1.056	0.373
Male gender	0.703	0.325	1.520	0.371	0.712	0.319	1.588	0.407
Body mass index (kg/m <sup>2</sup> )	0.917	0.826	1.016	0.098				
Smoking	1.027	0.525	2.009	0.938				
Heart rate (beats per minute)	1.039	1.018	1.061	<0.001	1.032	1.011	1.053	0.003
Systolic blood pressure (mm Hg)	0.999	0.986	1.013	0.928				
Diastolic blood pressure (mm Hg)	1.021	0.993	1.049	0.151				
Hypertension (yes or no)	0.694	0.319	1.511	0.358				
Dyslipidemia	1.246	0.524	2.963	0.619				
Chronic heart failure (yes or no)	1.551	0.652	3.690	0.321				
Ischemic stroke (yes or no)	0.737	0.221	2.455	0.619				
Acute ischemic limb (yes or no)	3.769	1.579	8.998	0.003	2.247	0.862	5.856	0.098
Rutherford classification (IV~VI)	1.346	0.778	2.327	0.288	1.189	0.668	2.114	0.557
Creatinine (mg/dl)	1.007	0.902	1.125	0.897				
Triglyceride (mg/dl)	0.997	0.993	1.002	0.239				
Uric acid (mg/dl)	1.040	0.860	1.257	0.688				
NLR	1.027	1.006	1.049	0.011	1.031	1.004	1.059	0.023
CHADS2 score	1.029	0.741	1.428	0.867				
Major adverse cardiac events at one year								
Atrial fibrillation	2.479	1.169	5.257	0.018	2.322	1.045	5.158	0.039
Age (years)	1.016	0.986	1.046	0.300	1.003	0.972	1.034	0.866

Male gender	0.805	0.422	1.534	0.510	0.858	0.446	1.649	0.645
Body mass index (kg/m <sup>2</sup> )	0.923	0.849	1.004	0.062				
Smoking	1.523	0.945	2.454	0.084				
Heart rate (beats per minute)	1.025	1.008	1.042	0.004	1.020	1.002	1.037	0.025
Systolic blood pressure (mm Hg)	0.998	0.988	1.009	0.715				
Diastolic blood pressure (mm Hg)	1.008	0.985	1.031	0.506				
Hypertension (yes or no)	0.753	0.391	1.452	0.398				
Dyslipidemia	1.318	0.638	2.723	0.456				
Chronic heart failure (yes or no)	1.654	0.800	3.416	0.174				
Ischemic stroke (yes or no)	1.044	0.436	2.503	0.923				
Acute ischemic limb (yes or no)	2.592	1.222	5.497	0.013	1.609	0.697	3.719	0.265
Rutherford classification (IV~VI)	1.257	0.806	1.960	0.314	1.133	0.711	1.805	0.600
Creatinine (mg/dl)	0.973	0.877	1.080	0.610				
Triglyceride (mg/dl)	0.999	0.995	1.002	0.434				
Uric acid (mg/dl)	1.047	0.905	1.213	0.536				
NLR	1.016	0.993	1.039	0.176	1.016	0.988	1.044	0.277
CHADS2 score	1.095	0.834	1.438	0.513				
<b>Major adverse limb events at one year</b>								
Atrial fibrillation	1.280	0.495	3.307	0.610	1.757	0.664	4.789	0.271
Age (years)	0.979	0.952	1.007	0.146	0.972	0.943	1.002	0.067
Male gender	1.145	0.582	2.253	0.696	1.709	0.851	3.433	0.132
Body mass index (kg/m <sup>2</sup> )	0.941	0.864	1.025	0.160				
Smoking	1.009	0.553	1.841	0.977				

Heart rate (beats per minute)	0.999	0.980	1.018	0.926				
Systolic blood pressure (mm Hg)	0.991	0.980	1.002	0.105				
Diastolic blood pressure (mm Hg)	0.984	0.960	1.008	0.196				
Hypertension (yes or no)	0.842	0.422	1.682	0.627				
Dyslipidemia	1.332	0.622	2.854	0.461				
Chronic heart failure (yes or no)	2.160	1.053	4.433	0.036	2.158	1.025	4.544	0.043
Ischemic stroke (yes or no)	1.151	0.476	2.779	0.755				
Acute ischemic limb (yes or no)	0.976	0.344	2.770	0.963	0.785	0.258	2.385	0.669
Rutherford classification (IV~VI)	4.227	2.339	7.636	<0.001	5.577	2.780	11.19	<0.001
Creatinine (mg/dl)	1.044	0.957	1.138	0.337				
Triglyceride (mg/dl)	1.000	0.998	1.003	0.844				
Uric acid (mg/dl)	0.906	0.765	1.073	0.251				
NLR	1.016	0.992	1.040	0.190	1.002	0.973	1.033	0.876
CHADS2 score	1.091	0.826	1.442	0.539				

a NLR as a continuous variable and NLR as a binary variable were not simultaneously adjusted in the multivariate logistic regression analyses.

<b>Table 3 The association between atrial fibrillation and the study outcomes in the sensitivity analyses</b>				
<b>The study outcomes</b>	<b>adjusted hazard ratio</b>	<b>95% confidence interval</b>		<b>P</b>
<b>All-cause mortality at one year</b>				
Model 1	2.416	1.204	-4.845	0.013
Model 2	2.141	1.047	-4.377	0.037
Model 3	2.533	1.304	-4.917	0.006
Model 4	2.080	0.950	-4.555	0.067
Model 5	2.158	1.012	-4.605	0.047
Model 6	2.209	1.036	-4.710	0.040
<b>Cardiac-related mortality</b>				
Model 1	2.276	0.853	-6.075	0.101
Model 2	2.176	0.801	-5.913	0.127
Model 3	2.483	0.979	-6.297	0.055
Model 4	2.022	0.708	-5.778	0.189
Model 5	2.169	0.789	-5.963	0.133
Model 6	2.169	0.792	-5.938	0.132
<b>Major adverse cardiac events at one year</b>				
Model 1	2.303	1.032	-5.140	0.042
Model 2	1.869	0.795	-4.393	0.151
Model 3	2.408	1.117	-5.191	0.025
Model 4	1.682	0.699	-4.048	0.246
Model 5	1.780	0.760	-4.172	0.184
Model 6	1.757	0.753	-4.099	0.192
Model 1 adjusted for creatinine in addition to age, gender, heart rate, the presence of acute limb ischemia, Rutherford classification stages, neutrophil-to-lymphocyte ratio.				
Model 2 adjusted for creatinine clearance in addition to heart rate, the presence of acute limb ischemia, Rutherford classification stages, neutrophil-to-lymphocyte ratio.				
Model 3 adjusted for estimated glomerular filtration rate in addition to heart rate, the presence of acute limb ischemia, Rutherford classification stages, neutrophil-to-lymphocyte ratio.				
Model 4 additionally adjusted for body mass index on the top of the model 1.				

Model 5 additionally adjusted for body mass index on the top of model 2.

Model 6 additionally adjusted for body mass index on the top of model 3.

## Figures




Figure 1

Kaplan-Meier curves showed that atrial fibrillation was significantly associated with an increased risk of one-year all-cause (1a) but not cardiac-related mortality (1b) in patients with severe lower extremity arterial disease after we adjusted for confounders. A significant association was found between atrial fibrillation and major adverse cardiac events (1c) other than major adverse limb events (1d).

## Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Table120210215.docx](#)
- [Table220200215.docx](#)
- [Table320210215.docx](#)