Interplay of Transcription Factor 7-Like-2 Polymorphisms with Type 2 Diabetes Mellitus in a Ghanaian Population

Christian Obirikorang

Kwame Nkrumah University of Science and Technology

Evans Asamoah Adu (✉ evansasamoahadu@gmail.com)

Kwame Nkrumah University of Science and Technology

Emmanuel Acheampong

Edith Cowan University

Enoch Odame Anto

Kwame Nkrumah University of Science and Technology

Lawrence Quaye

University for Development Studies

Brodrick Yeboah Amoah

University of Ghana

Max Efui Annani-Akollor

Kwame Nkrumah University of Science and Technology

Aaron Siaw Kwakye

Eastern Regional Hospital, Koforidua,

Foster Fokuoh

Eastern Regional Hospital, Koforidua,

Michael Appiah

Kwame Nkrumah University of Science and Technology

Eric Nana Yaw Nyarko

Kwame Nkrumah University of Science and Technology

Freeman Aidoo

Kwame Nkrumah University of Science and Technology

Eric Adua

Edith Cowan University

Ebenezer Afrifa-Yamoah

Edith Cowan University

Lois Balmer

Edith Cowan University

Wei Wang

Research Article

Keywords: TCF7L2, type 2 diabetes mellitus, cardiometabolic risk factors, single nucleotide polymorphisms

Posted Date: February 26th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-244726/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.
[Read Full License](#)

Abstract

Background

Type-2 diabetes mellitus (T2DM) have been strongly associated with single nucleotide polymorphisms (SNPs) in the TCF7L2 gene. This study investigated the interplay between rs12255372 and rs7903146 with T2DM in a Ghanaian population.

Methods

A case-control design was used for this study. A total of 106 T2DM patients and 110 apparently healthy control participants were selected. Basic data collected included body mass index, blood pressure and socio-demographics. Fasting blood samples were collected and used for serum lipid analysis, HbA1c, glucose estimation and DNA extraction. Common and allele-specific primers were designed for genotyping using the Modified Tetra-Primer Amplification assay. Associations were evaluated using logistic regression models.

Results

The rs7903146 risk variant was significantly associated with 2.16 vs 4.06 increased odds for T2DM in patients < 60 years vs ≥ 60 years. Both rs7903146 and rs12255372 were significantly associated with increased odds of T2DM in women, overweight/obese; negative T2DM family history and low-HDL-C. In a multivariate model, rs7903146 but not rs12255372 was significantly associated with 2.18, 5.01 and 2.25 increased odds of T2DM, under the codominant, recessive and additive model, respectively ($p < 0.05$). Haplotype analysis showed that rs7903146 and rs12255372 SNPs are weakly associated ($D' = 0.021$; $r^2 = 0.0001$).

Conclusion

The association between rs7903146 and rs12255372 with T2DM looks to be more highly associated in a subgroup- women and those without T2DM family history, yet have cardiometabolic risk.

Background

Diabetes continues to rise in prevalence across the globe. Alarmingly, 4 out of 5 people with diabetes live in low-middle-income countries (LMIC), where adequate resources for diabetes management are scarce [1]. According to the International Diabetes Federation (IDF) 2017, it is estimated that nearly half a billion of the world's population is living with diabetes [1]. Khan *et al.*, [2] reported that 462 million individuals are affected by diabetes, corresponding to 6.28% of the world's population. This figure is projected to reach 7079 individuals per 100,000 by 2030, reflecting a continued rise across all regions of the world [2]. Due

to the continues increase rate of diabetes worldwide, the 2030 projected rate is likely to be exceeded [2, 3]. This leads to high mortality from diabetic complications including cardiovascular disease [4]. The alarming situation is that 174.8 million (45.8%) of all diabetic cases, in adults, are undiagnosed with the most affected individuals from LMIC [5]. Thus, T2DM remains a threat to the world's population, even though there is increasing life-expectancy, which requires both public health and clinical care to prevent what is already a worldwide health crisis.

In Ghana, T2DM is prevalent among 6.46% [6] of the population, which is significantly greater than the 2017 International Diabetes Federation (IDF) estimation [1]. A number of factors including physical inactivity, poor diet, rapid urbanisation and ageing are common among people living with diabetes [7]. Besides these factors, T2DM has a strong genetic predisposition, taking the combination of inequalities across different populations, familial aggregation and concordance among monozygotic twins [8, 9]. Specifically, in T2DM, these factors are closely related to the metabolic abnormalities that underline the conditions of impaired insulin action, obesity, increased endogenous glucose output, and insulin secretory dysfunction [10]. It is well-known that genes control and regulate glucose and metabolic homeostasis, which influence the digestion, absorption and metabolic processes of lipids and carbohydrates can initiate the major mechanisms in T2DM [11].

Genome-wide association studies (GWAS) have identified more than 100 genetic variants in metabolic pathways which would provide new ways to understand the mechanism of T2DM [12, 13]. Among these risk variants, the transcription factor 7-like 2 (TCF7L2) represent the most identified and promising T2DM candidate gene [14].

It is well recognised as the gene that contains variants with the strongest association with T2DM, identified to date [14–19]. A common microsatellite in the TCF7L2 gene region DG10S478 (containing SNPs rs7903146 and rs12255372) was found to be associated with T2DM [20], the finding has been convincingly replicated in different population groups. Using non-coding SNPs in strong linkage disequilibrium with DG10S478 [16, 21, 22], an estimated population attributable risk of 17–28% has been established.

TCF7L2 encodes a high mobility group of box-containing transcription factors, which is involved in the Wnt signalling pathway [17]. The Wnt signalling pathway represents one of the pancreatic beta cell's important developmental and regulatory mechanism, which is critical in determining its fate, proliferation, polarity, and apoptosis during embryonic development and adult tissue homeostasis [20]. Risk alleles of TCF7L2 have been strongly associated with impaired insulin secretion/beta-cell function and T2DM [23–25]. Likewise, TCF7L2 has been shown to mediate regulation of glucagon-like peptide-1 expression, which represents a mechanism by which alteration of this gene influences susceptibility to T2DM [24]. There has been only a limited number of studies within the African Context, which makes it difficult to pool the effects and homogeneity of variants associated with T2DM [26]. Thus, replication studies are needed to understand the effect of these gene variants on T2DM. This study investigated the interplay between TCF7L2 gene polymorphisms and T2DM in a Ghanaian population.

Results

Table 1 shows the socio-demographic characteristics of the study participants. The proportion of participants aged 60 and above was significantly higher in the cases compared to the control group (31.1% vs 17.3%, $p = 0.017$). Family history of T2DM was more observed in the case group than the control group (42.5% vs 11.8%, $p < 0.001$). Aside from LDL-C ($p = 0.288$), the mean TC, TG, HDL-C and LDL-C significantly differed between the cases and control participants ($p < 0.05$). The proportion of cases vs controls with high TC (12.3% vs 5.6%, $p = 0.077$), overweight (40.6% vs 35.5%, $p = 0.441$) and obese (27.4% vs 19.1%, $p = 0.149$) did not differ significantly. High BP, atherogenic dyslipidaemia and Metabolic Syndrome (MetS) were higher in the cases than the controls (p -value < 0.001).

Table 1
Definition of metabolic syndrome

Component	Standard criteria [45]	Criteria used in the present study
Elevated waist circumference	Population- and country-specific definitions	BMI ≥ 30 kg/m ²
Elevated TG	TG ≥ 150 mg/dL (1.7 mmol/L), or drug treatment	Fasting TG ≥ 150 mg/dl (1.7 mmol/L)
Reduced HDL-C	HDL-C < 40 mg/dL (1.0 mmol/L) in males; HDL-C < 50 mg/dl (1.3 mmol/L) in females; or drug treatment	HDL-C < 40 mg/dl (1.0 mmol/L) in males HDL-C < 50 mg/dl (1.3 mmol/L) in females
Elevated blood pressure	Systolic ≥ 130 and/or diastolic ≥ 85 mmHg, or drug treatment	Systolic ≥ 130 and/or diastolic ≥ 85 mmHg
Elevated glucose	≥ 100 mg/dL in plasma, or drug treatment	Fasting blood glucose ≥ 100 mg/dL (5.5 mmol/L) and/or T2DM based on hospital records

Genotype and allele frequencies of TCF7L2 gene polymorphism

The genotype and minor allele frequencies (MAF) of rs7903146 and rs12255372 of the TCF7L2 gene are shown in Table 2. The MAF of rs7903146 was 50.5% and 33.3% among cases and control participants, with statistically significant differences ($p = 0.011$). The homozygote recessive allele (TT) of rs7903146, was more prevalent among cases than control participants (13.2% and 4.5%, $p = 0.024$). The MAF of rs12255372 was 24.1% among cases, which is significantly higher ($p = 0.013$) than that of controls (14.2%). The GG genotype ($p = 0.013$) and GT genotype ($p = 0.023$) but not the TT genotype ($p = 0.501$) of rs12255372 was disproportionately distributed between the cases vs control participants.

Table 2
Primers and Concentration of reagents

SNP	Primer name	Primer Sequence (5'→3')	Melting Temp	Product (BP)
rs7903146	FIP (T allele)	TAGAGAGCTAAGCACTTTTATAGAGAT	57.5	395
rs7903146	RIP (C allele)	GCCTCATACGGCAATTAAATTATAGAG	58.3	285
rs7903146	FOP	AAAACCTTTCCAATTTTTCACCTG	57.5	628
rs7903146	ROP	ATCCATCCACAAGCTAACCCT	59.1	628
rs12255372	FIP (T allele)	CCAGGAATATCCAGGCAAGATTG	55.8	171
rs12255372	RIP (C allele)	GGCCTGAGTAATTATCAGAATATGCTG	58.3	106
rs12255372	FOP	GTCCCTTGAGGTGTACTGCAA	56.7	228
rs12255372	ROP	TATTTGGCATTCAAATGGACGC	55.0	228
FIP- forward inner primer; RIP- reverse inner primer; FOP; forward outer primer; ROP: reverse outer primer; Tm- melting temperature; Ta- annealing temperature				

The linkage disequilibrium (LD) and pairwise LD coefficients r^2 and D' between rs12255372 and rs7903146 demonstrated that the alleles for the rs7903146 (C > T) and rs12255372 (G > T) SNPs are weakly associated ($D' = 0.021$; $r^2 = 0.0001$).

Association between TCF7L2 polymorphisms and T2DM

There is an association between rs12255372 and rs7903146 with T2DM (Table 3). Here rs7903146 was significantly associated with increased risk of T2DM in the codominant model (OR = 2.22, $p = 0.008$), recessive model (OR = 5.28, $p = 0.004$) and additive model (OR = 2.47, $p = 0.002$). After adjusting the model for age, sex, family history, high cholesterol and overweight/obesity, rs7903146 was still associated with T2DM; additive (OR = 2.25, $p = 0.003$), recessive (OR = 4.92, $p = 0.011$) and codominant (OR = 2.29, $p = 0.019$). rs12255372 was significantly associated with T2DM under the codominant (OR = 2.10, $p = 0.017$) and additive model (OR = 1.70, $p = 0.026$) but not the recessive model (OR = 1.86, $p = 0.309$). After adjustment for confounders, no association was found between rs12255372 and T2DM ($p > 0.05$). In a multivariate model, rs7903146 but not rs12255372 was significantly associated with 2.18, 5.01 and 2.25 increased odds of T2DM under the codominant, recessive and additive model respectively ($p < 0.05$).

Table 3

Comparison of demographic variables, anthropometric measures and biochemical profiles between the cases and controls

Variables	Cases (N = 106)	Controls (N = 110)	P-value
Sex			
Male	46 (43.4)	37 (33.6)	0.140
Female	60 (56.6)	73 (66.4)	
Family history of T2DM			
No	61 (57.7)	97 (88.2)	
Yes	45 (42.5)	13 (11.8)	< 0.001
Age (years)	53.90 ± 12.31	45.80 ± 12.14	< 0.001
< 60 years	73 (68.9)	91 (82.7)	
≥ 60 years	33 (31.1)	19 (17.3)	0.017
Anthropometric variables#			
BMI (Kg/m ²)	28.18 ± 5.53	26.10 ± 4.41	0.002
Systolic BP (mmHg)	133.35 ± 19.76	121.25 ± 18.41	< 0.001
Diastolic BP (mmHg)	83.68 ± 11.74	76.22 ± 11.17	< 0.001
Biochemical parameters#			
Triglycerides (mmol/l)	1.69 ± 0.64	1.29 ± 0.50	< 0.001
T. Chol (mmol/l)	5.25 ± 1.03	4.65 ± 1.06	< 0.001
HDL-C(mmol/l)	0.99 ± 0.24	1.23 ± 0.27	< 0.001
LDL-C (mmol/l)	3.44 ± 0.95	3.32 ± 0.72	0.288
FPG (mmol/l)	8.72 ± 4.66	4.85 ± 0.91	< 0.001
HbA1c (%)	6.99 ± 1.87	4.83 ± 0.60	< 0.001
Cardiometabolic factors			
High BP	36 (34.0)	10 (9.1)	< 0.001
High TC	13 (12.3)	6 (5.5)	0.077
High TG	61 (57.5)	16 (14.5%)	< 0.001
All values are presented as frequency and percentages unless stated. # Values are presented as mean and standard deviations. P-values highlighted are statistically significant (< 0.05)			

Variables	Cases (N = 106)	Controls (N = 110)	P-value
Low HDL-C	81 (76.4)	52 (47.3)	< 0.001
High LDL-C	64 (60.4)	36 (32.7)	< 0.001
Atherogenic dyslipidaemia	44 (41.6)	5 (4.5)	< 0.001
Overweight	43 (40.6)	39 (35.5)	0.441
Obesity	29 (27.4)	21 (19.1)	0.149
MetS	70 (66.0)	14 (12.7)	< 0.001
All values are presented as frequency and percentages unless stated. # Values are presented as mean and standard deviations. P-values highlighted are statistically significant (< 0.05)			

Table 4 demonstrates an association between rs12255372 and rs7903146 polymorphisms and T2DM within subcategories. The rs7903146 polymorphisms (C > T) was significantly associated with 2.16 vs 4.06 increased odds ratio of T2DM in persons aged below 60 years vs persons aged above 60-years. The rs7903146 (C > T) was significantly associated with increased odds ratio of T2DM in both normal weight (OR = 2.52, 95%CI: 1.02–6.26) and overweight/obese (OR = 2.15, 95%CI: 1.01–4.56) individuals; normal BP (OR = 2.93, 95%CI: 1.51–5.66); negative diabetes family history (OR = 2.71, 95%CI: 1.35–5.45); females (2.71, 95%CI: 1.30–5.65); elevated TG (OR = 3.87, 95%CI: 1.21–12.41) and low HDL-C (OR = 2.95, 95%CI: 1.40–6.24). On the other hand, rs12255372 (G > T) was significantly associated with excess risk of T2DM among patients aged 60 and above (OR = 52.45, 95%CI: 2.92–942.04); increased odds of T2DM among patients with low HDL-C (OR = 2.35, 95%CI: 1.10–5.00), overweight/obese (OR = 2.43, 95%CI: 1.19–4.98), negative family history (OR = 2.07, 95%CI: 1.04–4.12) and females (OR = 2.34, 95%CI: 1.11–4.95).

Table 4: Distribution of *TCF7L2* gene polymorphism in cases and control participants

SNP	Controls (n=110)		Cases (n=106)		P-value
	%	Allele frequency	N (%)	Allele frequency	
		C T		C T	
rs7903146		0.667 0.333		0.495 0.505	0.011
CC	49 (44.5)		26 (24.5)		0.002
CT	56 (50.9)		66 (62.3)		0.092
TT	5 (4.5)		14 (13.2)		0.024
		G T		G T	
rs12255372		0.857 0.143		0.759 0.241	0.068
GG	81 (73.6)		61 (57.5)		0.013
GT	24 (21.8)		38 (35.8)		0.023
TT	5 (4.5)		7 (6.6)		0.501

Highlighted P-values are statistically significant.

Individuals with the rs7903146 polymorphism comparatively had higher mean TG ($p = 0.048$) and high mean LDL-C in both the controls ($p = 0.001$) and cases ($p = 0.046$). Among the cases, rs7903146 SNP was associated with significantly higher mean systolic BP ($p = 0.015$) (Table S2). In table S2, those with rs12255372 SNPs, among the control group, had higher mean TG ($p = 0.003$), TC ($p = 0.048$), LDL-C ($p = 0.001$), FPG ($p = 0.005$) and HbA1c ($p < 0.001$). Among the cases, rs12255372 polymorphism was associated with higher BMI ($p = 0.044$) and systolic BP ($p = 0.003$).

In Table S3, rs7903146 was significantly associated with increased odds of high LDL among the controls (OR = 2.98, $p = 0.017$). Among the case group, individuals with the minor allele of rs7903146 were significantly associated with an increased odds ratio of high TG (OR = 2.95, $p = 0.023$) (Table S3). rs12255372 SNP was significantly associated with an increased odds ratio of high cholesterol (OR = 14.0, $p = 0.023$) and elevated LDC-C (OR = 3.12, $p = 0.027$) among the control group (Table S3). Among the case group, rs12255372 was significantly associated with increased odds of high BP (OR = 3.01, $p = 0.013$) and overweight/obesity (OR = 3.33, $p = 0.012$).

Discussion

Variants of TCFL2 has been consistently associated with T2DM in several studies among multiple ethnic groups [16, 21, 22]. This study demonstrated the transferability of the TCF7L2 gene polymorphisms (rs12255372 and rs7903146) and their association with T2DM among a Ghanaian population. However, the associations were observed to be relevant only in a subgroup- women, patients that do not have a family history of diabetes and those with dyslipidaemia with or without elevated blood pressure and increased BMI.

This study observed that rs7903146 and rs12255372 polymorphisms have a significant effect on T2DM, however, the effect was mediated by other diabetes risk factors. In a systematic review and Meta-analysis among a Sub-Saharan Africa (SSA) population [26], the effect of rs7903146 was 1.75 in a recessive model and 2.27 in an additive model, which is similar to our present findings. These effects are also consistent among the Caucasian, East Asian, South Asian and other ethnicities [21]. To the best of our knowledge, this study is currently among the few studies in SSA, that have demonstrated the transferability of rs12255372 [27, 28]. In contrast studies by Yako *et al.*, [29] and Mandour *et al.*, [30], found the SNP rs12255372 to not associated with T2DM, in a South African mixed-ancestry and Egyptian population, respectively.

From a Meta-analysis by Wang *et al.*, [22] among different ethnic groups, there are significant differences in the LD structure between rs12255372 and rs7903146. From the findings of this study, weak interaction between rs12255372 and rs7903146 for T2DM was reported (in both male and female participants). This result is similar to that observed among the Yoruba population ($D' = 0.075$, $r^2 = 0.001$) but unlike that of the Europeans ($D' = 0.948$, $r^2 = 0.746$) and Chinese ($D' = 0.479$, $r^2 = 0.114$) [22].

The MAF of rs7903146 observed in this study is similar to the findings of Danquah *et al.*, [15] among a Ghanaian population, where the MAF of rs7903146 was 30% in the control population and 36% in T2DM population. Among other SSA populations, the MAF of rs7903146 among T2DM vs controls has been reported at 9.5% vs 0% in Cameroonian population [31] and 32.2% vs 24.3% in South African mixed-ancestry [29]. Furthermore, from the Allele Frequency Aggregator (ALFA) HapMap and 1000 Genomes project [32] the MAF of rs7903146 were 29.2%, 24.1% and 22.8%, respectively, which is consistent with our findings. The race-specific risk allele frequency of rs7903146 is lower among the Asian population (range: 3.1%-4.7%) and higher among Europeans (29.2%), Africans (29.5%), African American (29.7%), Latin American (34.2%) and South Asians (31.8%) [32]. A wide population difference in the minor allele frequency of rs12255372 (Asians = 0.000 to 0.029, Europeans = 0.217-0.50 and SSA = 0.267–0.274) has been reported [17, 30, 33]. Moreover, from the results of the 1000 genomes project, the MAF of rs12255372 is 21.4% globally, 30.2% in Africa, 29.2% among Europeans and the Americans 21.9%, similar to our study.

From the findings of this study and related literature, the association between rs12255372 and rs7903146 with T2DM are similar among West African populations, Asians, Caucasians [34, 35] and other population groups [16]. This suggests that the association of rs7903146 and rs12255372 with T2DM is moderately strong in multiple ethnicities which are likely to project a vital contribution of differences in environmental factors that confer susceptibility to T2DM.

It appears that rs12255372 and rs7903146 are not associated with T2DM among West African populations (Ghanaians and Yoruba), unlike other population groups. This finding is essential in understanding the different types and aetiology of T2DM present in SSA [36], and the wider population differences in T2DM prevalence. For example, as indicated in this study the association between rs12255372 with T2DM was relevant among women, obese/overweight individuals, individuals with negative family history, low HDL-C and individuals aged above 60 years. Similarly, rs7903146 was relevant among females, individuals with negative family history, elevated TG and low HDL-C but normal blood pressure. This indicated that TCF7L2 risk variants may be relevant to a subgroup of individuals with single or multiple diabetes risk factors, other than those with a positive family history [37].

This study observed that the risk allele rs7903146 was associated with elevated FPG and HbA1c in both the diabetic and controls. In addition to FPG and HbA1c, the risk alleles of rs12255372 were associated with elevated BMI, blood pressure, TG and reduced LDL in T2DM and controls. This finding reflects the pleiotropic role TCF7L2 polymorphisms play in T2DM (adipogenesis, incretin and insulin expression as well as beta-cell growth and development). The rs7903146 risk alleles are associated with reduced proportional control of β -cell function among newly diagnosed T2DM patients [44]. Loos *et al.*, [38] reported that the T-allele of rs7903146 was associated with higher HbA1c concentrations and reduced β -cell function, assessed by the homeostasis model assessment of β -cell function. TCF7L2 protein was also found to be increased during adipogenesis in 3T3-L1 cells and primary adipocyte stem cells [46] and TCF7L2 expression was required for the regulation of Wnt [39]. High expression of TCF7L2 mRNA was identified with reduced TCF7L2 protein levels in individuals with the T-allele of rs7903146, which is

associated with a downregulation of GIP- and GLP-1 receptors and impaired beta-cell function Shu *et al.*, [40].

These findings imply that TCF7L2 polymorphisms rs7903146 and rs12255372 may confer T2DM susceptibility through the expression of adipocyte-insulin resistance, delayed insulin secretion (resulting from beta-cell dysfunction) and reduced intracellular energy homeostasis. Therefore, individuals with the risk alleles of TCF7L2 (rs7903146 and rs12255372) may present with obesity, elevated TG and reduced HDL-C. Also, the findings suggest that among individuals with multiple diabetes risk factors, those who do develop T2DM may have a genetic predisposition to beta-cell dysfunction which includes the SNPs rs12255372 and rs7903146 [18].

Some limitations of the study are worth mentioning when interpreting the findings of this study. The differences in population substructures may result in a different interpretation of risk alleles in candidate gene association studies. Also, the number of genetic roots underlying T2DM is still being investigated and could be numerous given its multifaceted nature. Therefore, the effect of the variants among the small population considered in this study may be unpredictably low, when we consider it from a larger population. Besides this limitation, the findings of this study are consistent with the recommended approach for conducting a candidate gene association study and present consistent findings with similar studies among other population groups.

Methods

This case-control study was conducted at the Begoro District Hospital (BDH) and two communities (Bosuso and Begoro) in the Eastern Region of Ghana. The study included participants aged 30–79 years. Cases were defined as clinically diagnosed T2DM patients attending the diabetic clinic at the BDH. Controls were defined as non-diabetic patients with normal fasting plasma glucose (FPG) and/or normal HbA1C that voluntarily participated in a health screening program conducted as part of the study in Bosuso or Begoro. Participants willing to have their body mass index (BMI), FPG, HbA1c and lipid profile checked and provided written informed consent during the health screening were recruited as controls. Control participants with a self-reported, previous or current, history of renal, liver, thyroid disorders or other endocrine diseases, chronic diseases, on hormone replacement therapy or corticosteroid therapy were excluded from the study. T2DM patients who self-managed or being managed for micro-vascular or macrovascular complications were excluded from the study.

Sample Size Calculation

Taking the minor allele frequency (MAF) of the lead SNP rs7903146 from a previous study [15], at a patient-to-control ratio of 1:1, at 95% CI, and 6% proportional differences ($P^* = (0.36 + 0.30)/2 = 0.33$; $(P_1 - P_2) = (0.36 - 0.30) = 0.06$, the minimum recommended sample size at 80% power replication under Kelsey method [41] were 100 cases and 100 controls. Overall, 106 T2DM and 110 non-diabetic controls met the inclusion criteria.

Recruitment Process

The control participants were recruited during a health screening program conducted in the two communities. Additional controls were sorted through community members that were informed via the various religious organizations. Initially, the intent, preparation and benefits of the health screening program was announced in all the religious-based organisations. The health screening process took place every morning (Monday, Wednesday and Friday) from 8:00 am to 11:30 am from May 2019 to November 2019. T2DM patients were recruited using a total sampling method. All patients that visited the BDH hospital during the study period and met the inclusion criteria were considered as a sample.

Blood pressure measurement was taken using an automated sphygmomanometer (Omron BP7200, Omron Healthcare, Inc., USA). Three different measurements were taken at 5-minute intervals and the average was used as the final reading. All blood pressure recordings were done in mmHg.

The Omron B511 (Omron Healthcare, Inc., USA) was used for weight measurement, following the manufacturer's instruction. Briefly, participants were asked to stand in upright on the main unit of the scale and without shoes. Portable Height Rod Stadiometers (QuickMedical, Warwick) were used for body height; the subject stood upright, with feet together and flat on the ground. BMI was calculated as the ratio of weight to height-squared (Kg/m^2).

Six millilitres (ml) of venous blood were taken in a fasting state and two millilitres each were placed into sterile ethylene diamine tetraacetic acid (EDTA) (BD Vacutainer®, USA), gel separator and Sodium Fluoride/Calcium oxalate tubes (BD Vacutainer®, USA). Fasting was defined as no caloric intake for at least 8 h, overnight. The EDTA samples were used for HbA1c analysis using standard A1cCare System (SD BIOSENSOR, republic of Korea). The remaining EDTA sample was used for DNA extraction and consequent SNPs genotyping. The gel separator tubes samples were processed by centrifugation at 3,000 rpm for 10 minutes and the serum was collected into 1.5 ml eppendorf tubes. The obtained sera were used for lipid profiling. The plasma obtained from the Sodium Fluoride/Calcium oxalate tube samples was used for the determination of Fasting Plasma Glucose (FPG).

Biochemical Measurements and Definition of Cardiometabolic risk factors

The Lipid and FPG analysis were conducted at the Eastern Regional Hospital's Clinical Chemistry laboratory, using the SELECTRAPRO M chemistry analyser (EliTechGroup B.V, Netherlands). All analysis was recorded in mmol/l. The lipid profile included HDL-C, low-density lipoprotein cholesterol (LDL-C) and TG and total cholesterol (TC)

Dyslipidaemia was defined as follows: high TC ($> 5.0\text{mmol/l}$), high LDL-C ($> 3.0\text{mmol/l}$), high TG ($> 1.7\text{mmol/l}$), low HDL-C ($< 1.0\text{mmol/l}$ for men and $< 1.2\text{mmol/l}$ for women) [42]. Atherogenic dyslipidaemia was defined as high TG levels, low HDL cholesterol levels and an increase in small dense LDL [43]. BMI was categorized into four groups according to the conventional WHO classification [44]: underweight ($<$

18.5 kg/m²), normal weight (18.5–24.9 kg/m²), overweight (25–29.9 kg/m²), and obese (≥ 30 kg/m²). Metabolic syndrome (MetS) was defined according to the National Cholesterol Education Program (NCEP) Adult Treatment Panel III (ATP III), without waist circumference [45]; BP over 130/85 mmHg, TG > 1.7 mmol/l, HDL-C levels less than 1.03 mmol/l (men) or 1.29 mmol/l (women) and fasting blood sugar over 5.5 mmol/l (Table 5).

Table 5
Association between rs12255372 and rs7903146 and T2DM

SNPs	Crude model		Adjusted model		Multivariate model	
	OR (95% CI)	P-value	OR (95% CI)	P-value	OR (95% CI)	P-value
rs7903146						
Codominant	2.22 (1.23–4.02)	0.008	2.29 (1.14–4.58)	0.019	2.18 (1.08–4.39)	0.030
Recessive	5.28 (1.71–16.27)	0.004	4.92 (1.45–16.72)	0.011	5.01 (1.44–17.44)	0.011
Additive	2.47 (1.38–4.42)	0.002	2.25 (1.32–3.83)	0.003	2.25 (1.31–3.84)	0.003
rs12255372						
Codominant	2.10 (1.14–3.87)	0.017	1.80 (0.92–3.52)	0.085	1.72 (0.86–3.43)	0.124
Recessive	1.86 (0.56–6.14)	0.309	1.13 (0.29–4.43)	0.859	1.26 (0.32–5.044)	0.742
Additive	1.70 (1.06–2.71)	0.026	1.40 (0.83–2.36)	0.206	1.39 (0.82–2.38)	0.222
OR- Odds ratio; CI- confidence interval. highlighted P-values are statistically significant.						

Genomic DNA Extraction

Genomic DNA was extracted using a modified, simplified non-enzymatic salting-out method (details are shown in supplementary material). Briefly, 300 μ l of whole blood was used from each of the 24 samples resulting in good quality of DNA with the concentration ranging between 60 to > 100 ng/ μ l being extracted. The results of 1% (w/v) Agarose gel and Thermo Scientific™ NanoDrop 2000 full spectrum of the product are presented in **Figure S1 and S2**.

Primer design and primer information

Based on the GenBank sequence of human TCF7L2 the sequence surrounding the SNPs rs7903146 (NG_012631.1:g.53341) and rs12255372 (NG_012631.1:g.10389) was obtained. Two sets of primers for modified tetra-primer amplification assay (MTPA) PCR were designed using web-based software (http://sci.ui.ac.ir/*rahgozar). Several pairs of primers were identified and further refined using Primer-BLAST (<http://www.ncbi.nlm.nih.gov/blast>) and Oligo-Analyzer 3.1 IDT (<https://www.idtdna.com/calc/analyzer>). The primers used are provided in Table 6.

Table 6
Association between rs12255372 and rs7903146 polymorphisms and T2DM while keeping other metabolic risk factors as fixed

Risk Factors	rs7903146	rs12255372
	OR (95% CI)	OR (95%CI)
Age		
< 60 years	2.16 (1.11–4.22)*	1.18 (0.62–2.26)
60 and above	4.06 (1.25–13.19)**	52.45 (2.92-942.04)**
Family history		
NFH	2.71 (1.35–5.45)**	2.07 (1.04–4.12)*
PFH	1.00 (0.25–3.98)	1.01 (0.31–3.36)
Sex		
Male	1.90 (0.75–4.84)	1.53 (0.64–3.68)
Female	2.71 (1.30–5.65) **	2.34 (1.11–4.95)**
BMI Status		
Normal weight	2.52 (1.02–6.26)*	1.28 (0.47–3.45)
Overweight/Obese	2.15 (1.01–4.56)*	2.43 (1.19–4.98)*
BP status		
Normal	2.93 (1.51–5.66)**	1.55 (0.81–2.95)
Hypertensive	1.35 (0.32–5.66)	2.81 (0.67–11.80)
Dyslipidaemia		
Normal TG	1.45 (0.70–2.99)	1.64 (0.76–3.55)
Elevated TG	3.87 (1.21–12.41)**	1.23 (0.42–3.61)
Normal HDL-C	1.57 (0.60–4.13)	1.48 (0.56–3.95)
Low HDL-C	2.95 (1.40–6.24)**	2.35 (1.10-5.00)*
OR- odds ratios, P-values are statistically significant.		
*- P-value < 0.05; **-P-values < 0.01		

Genotyping of TCF7L2 rs7903146 and rs12255372

Each 25 µL reaction mixture contained 1X of OneTaq® 2X Master Mix (New England Biolabs), 0.2 µM each of forward and reverse primers and > 50 ng of DNA. Incubation of the 25 µL reaction mixture was

completed at 94°C for 5 minutes, followed by 30 cycles of 30-sec denaturation (94°C), 1 min annealing (57.7°C for rs7903146 or 55.0°C for rs12255372), 30-sec extension (72°C) and additional 10 minutes extension at 72°C at the end of the 30 cycles. The final product was visualised with 1.5% (w/v) of agarose gel electrophoresis alongside a 100-base pair (bp) marker with Ethidium bromide stain (**Figure S3 and S4**). For quality control and troubleshooting see supplementary appendix (**Table S1**).

Data Analysis

Analysis of the data was conducted using SPSS version 25. Normality of continues data was checked using the Kolmogorov Simonov test. Descriptive statistics were reported using frequencies for categorical data and mean for continues data. Chi-square test was used to evaluate the independence of the categorical data and t-test for the continues data between cases and controls. To test for associations between T2DM phenotype and all SNPs, logistic regression models were fitted, in which each SNP was presented as a predictor variable whose values were equal to the number of copies of the minor allele (0, 1, 2) in an additive model, or presence of at least one copy of the minor allele (0, 1) in a codominant model or presence of two copies of the minor allele (0, 1) in a recessive model. Sex, age and family history were included as covariates in the fitted model. The structure of the model was represented as:

$$\text{Logit}[pr(D=1)] = \alpha + \beta_1 G + \beta_2 \text{sex}_{(male)} + \beta_3 \text{age}_{(\geq 60 \text{ years})} + \beta_{\text{positive family history}} + \beta_{\text{high cholesterol}} + \beta_{\text{overweight/obesity}}$$

To examine the associations between T2DM phenotype and various allelic configurations of rs7903146 and rs12255372, haplotype association analysis was performed. The statistical analyses for linkage disequilibrium (LD), haplotype frequencies, and multiple test corrections per 1000 random permutations were performed using the Haploview 4.2 software [46, 47]. The haplotype blocks were defined by the four-gamete test [48, 49] as implemented in Haploview. A weak interaction between rs7903146 and rs12255372 was observed, data not shown. Univariate models adjusted for age and gender, these were used to evaluate the association between the SNPs and cardiometabolic markers. P-values < 0.05 were considered statistically significant.

Conclusion

There is a relatively high prevalence of the risk alleles of TCF7L2 (rs7903146 and rs12255372) among the Ghana population with a modest association with T2DM. The association between rs7903146 and rs12255372 with T2DM may be relevant in a subgroup of individuals (women), particularly those with no family history of diabetes and those with cardiometabolic risk with or without elevated blood pressure. Therefore, genotyping of these SNPs is essential for future multi-ethnic association studies and clinically significant for classifying high-risk patients.

Declarations

Acknowledgements

The authors acknowledge the management of the Begoro District Hospital and the Eastern Regional Hospital, especially Mary Koryoe Tetteh (Head of Laboratory Department, BDH) and Aaron Siaw Kwakye, for their support and vital contribution to the work.

Authors Contribution

Conceptualization, CO, WW, LB; Methodology, CO, EAA, EA, MEAA, EO, LQ, BYA; Validation, LQ, WW, and LB; Formal Analysis, EAA, EAY; Investigation, CO, EAA, EA, ASK, FF.; Resources, MA, ENYN, FA, CO, EA, MEAA; Data Curation, EAA, ASK, FF; Writing – Original Draft Preparation, EAA.; Writing – Review & Editing, CO, EAA, EO, MEAA, EA, LQ, BYA, WW, LB; Supervision, CO. All authors read and approved the manuscript before submission.

Competing interests

The authors declare that they have no conflict of interest

Data Availability

All data supporting the conclusions of this study are summarized in Tables and supplementary materials. The raw data is available from the corresponding author upon reasonable request.

Ethics declarations

The study was conducted according to the guidelines of the Declaration of Helsinki and approved by the Committee on Human Research, Publications and Ethics (CHRPE), School of Medicine and Dentistry (SMD), Kwame Nkrumah University of Science and Technology (CHRPE/AP/034/20). Written informed consent was obtained from all subjects involved in the study.

References

1. IDF IDF. IDF Diabetes Atlas 8th Edition (2017). 2017.
2. Khan, M. A. B. *et al.* Epidemiology of Type 2 Diabetes - Global Burden of Disease and Forecasted Trends. *J Epidemiol Glob Health*.**10** (1), 107–111 Epub 2020/03/17 <https://doi.org/10.2991/jegh.k.191028.001> (2020).
3. da Rocha Fernandes, J. *et al.* IDF Diabetes Atlas estimates of 2014 global health expenditures on diabetes. *Diabetes Res Clin Pract*.**117**, 48–54 (2016).
4. Hudspeth, B. The burden of cardiovascular disease in patients with diabetes. *Am J Manag Care*.**24** (13 Suppl), S268–S272 (2018).
5. Beagley, J., Guariguata, L., Weil, C. & Motala, A. A. Global estimates of undiagnosed diabetes in adults. *Diabetes Res Clin Pract*.**103** (2), 150–160 (2014).

6. Asamoah-Boaheng, M., Sarfo-Kantanka, O., Tuffour, A. B., Eghan, B. & Mbanya, J. C. Prevalence and risk factors for diabetes mellitus among adults in Ghana: a systematic review and meta-analysis. *International Health*.**11** (2), 83–92 <https://doi.org/10.1093/inthealth/ihy067> (2018).
7. Issaka, A., Paradies, Y. & Stevenson, C. Modifiable and emerging risk factors for type 2 diabetes in Africa: a systematic review and meta-analysis protocol. *Systematic Reviews*.**7** (1), 139 <https://doi.org/10.1186/s13643-018-0801-y> (2018).
8. Meigs, J. B., Cupples, L. A. & Wilson, P. Parental transmission of type 2 diabetes: the Framingham Offspring Study. *Diabetes*.**49** (12), 2201–2207 (2000).
9. Poulsen, P., Kyvik, K. O., Vaag, A. & Beck-Nielsen, H. Heritability of type II (non-insulin-dependent) diabetes mellitus and abnormal glucose tolerance—a population-based twin study. *Diabetologia*.**42** (2), 139–145 (1999).
10. Zaman, G. S. *Pathogenesis of Insulin Resistance* (IntechOpen, Cellular Metabolism and Related Disorders, 2020).
11. Herman, M. A. & Kahn, B. B. Glucose transport and sensing in the maintenance of glucose homeostasis and metabolic harmony. *The Journal of clinical investigation*.**116** (7), 1767–1775 (2006).
12. Qi, L. *et al.* Genetic variants at 2q24 are associated with susceptibility to type 2 diabetes. *Hum Mol Genet*.**19** (13), 2706–2715 (2010).
13. Morris, A. P. *et al.* Large-scale association analysis provides insights into the genetic architecture and pathophysiology of type 2 diabetes. *Nat Genet*.**44** (9), 981 (2012).
14. Cauchi, S. *et al.* TCF7L2 is reproducibly associated with type 2 diabetes in various ethnic groups: a global meta-analysis. *J Mol Med*.**85** (7), 777–782 (2007).
15. Danquah, I. *et al.* The TCF7L2 rs7903146 (T) allele is associated with type 2 diabetes in urban Ghana: a hospital-based case–control study. *BMC Med Genet*.**14** (1), 96 (2013).
16. Dieudonne, N. *et al.* Association between the TCF7L2 rs12255372 (G/T) gene polymorphism and type 2 diabetes mellitus in a Cameroonian population: a pilot study. *Clinical and Translational Medicine*.**4**, <https://doi.org/10.1186/s40169-015-0058-1> (2015).
17. Huang, Z., Liao, Y., Huang, R. & Chen, J. Sun H-I. Possible role of TCF7L2 in the pathogenesis of type 2 diabetes mellitus. *Biotechnology & Biotechnological Equipment*.**32** (4), 830–834 (2018).
18. Katsoulis, K. *et al.* TCF7L2 gene variants predispose to the development of type 2 diabetes mellitus among individuals with metabolic syndrome. *Hormones*.**17** (3), 359–365 (2018).
19. Shu, L. *et al.* Transcription factor 7-like 2 regulates β -cell survival and function in human pancreatic islets. *Diabetes*.**57** (3), 645–653 (2008).
20. Grant, S. F. *et al.* Variant of transcription factor 7-like 2 (TCF7L2) gene confers risk of type 2 diabetes. *Nat Genet*.**38** (3), 320 (2006).
21. Ding, W. *et al.* Meta-analysis of association between TCF7L2 polymorphism rs7903146 and type 2 diabetes mellitus. *BMC Med Genet*.**19** (1), 38–38 <https://doi.org/10.1186/s12881-018-0553-5>

(2018).

22. Wang, J. *et al.* Association of rs12255372 in the TCF7L2 gene with type 2 diabetes mellitus: a meta-analysis. *Brazilian journal of medical and biological research = Revista brasileira de pesquisas medicas e biologicas*.**46** (4), 382–393 Epub 2013/04/12 <https://doi.org/10.1590/1414-431x20132677> (2013).
23. Lyssenko, V. *et al.* Mechanisms by which common variants in the TCF7L2 gene increase risk of type 2 diabetes. *The Journal of clinical investigation*.**117** (8), 2155–2163 (2007).
24. Ip, W., Chiang, Y-T-A. & Jin, T. The involvement of the wnt signaling pathway and TCF7L2 in diabetes mellitus: The current understanding, dispute, and perspective. *Cell Biosci*.**2** (1), 28–28 <https://doi.org/10.1186/2045-3701-2-28> (2012).
25. Wu, H-H. *et al.* TCF7L2 regulates pancreatic β -cell function through PI3K/AKT signal pathway. *Diabetol Metab Syndr*.**11** (1), 55 (2019).
26. Asamoah, E. A. *et al.* Heritability and Genetics of Type 2 Diabetes Mellitus in Sub-Saharan Africa: A Systematic Review and Meta-Analysis. *Journal of diabetes research*. 2020;2020.
27. Helgason, A. *et al.* Refining the impact of TCF7L2 gene variants on type 2 diabetes and adaptive evolution. *Nat Genet*.**39** (2), 218 (2007).
28. Nanfa, D. *et al.* Association between the TCF7L2 rs12255372 (G/T) gene polymorphism and type 2 diabetes mellitus in a Cameroonian population: a pilot study. *Clinical and translational medicine*.**4** (1), 17 (2015).
29. Yako, Y. Y. *et al.* Contribution of ENPP1, TCF7L2, and FTO polymorphisms to type 2 diabetes in mixed ancestry ethnic population of South Africa. *Afr Health Sci*.**15** (4), 1149–1160 (2015).
30. Mandour, I., Darwish, R., Fayez, R., Naguib, M. & El-Sayegh, S. TCF7L2 Gene Polymorphisms and Susceptibility to Type 2 Diabetes Mellitus, a Pilot Study. *Biomedical and Pharmacology Journal*.**11** (2), 1043–1049 (2018).
31. Guewo-Fokeng, M. *et al.* Contribution of the TCF7L2 rs7903146 (C/T) gene polymorphism to the susceptibility to type 2 diabetes mellitus in Cameroon. *Journal of Diabetes & Metabolic Disorders*.**14** (1), 26 (2015).
32. Clark, K. *et al.* *Nucleic Acids Res*.**44** (D1), D67–D72 <https://doi.org/10.1093/nar/gkv1276> (2016). Epub 2015/11/20
33. Lehman, D. M. *et al.* Haplotypes of transcription factor 7-like 2 (TCF7L2) gene and its upstream region are associated with type 2 diabetes and age of onset in Mexican Americans. *Diabetes*.**56** (2), 389–393 Epub 2007/01/30 <https://doi.org/10.2337/db06-0860> (2007).
34. Li, Y., Yang, X., Geng, H. & Gong, G. Type 2 diabetes mellitus and TCF7L2 gene rs12255372 G/T polymorphism: a meta-analysis involving 7990 subjects. *Int J Diabetes Dev Ctries*.**38** (1), 55–61 <https://doi.org/10.1007/s13410-017-0560-7> (2018).
35. Xi, X. & Ma, J. A meta-analysis on genetic associations between Transcription Factor 7 Like 2 polymorphisms and type 2 diabetes mellitus. *Genomics*.**112** (2), 1192–1196 (2020).

36. Kibirige, D. *et al.* Understanding the manifestation of diabetes in sub Saharan Africa to inform therapeutic approaches and preventive strategies: a narrative review. *Clinical Diabetes and Endocrinology*.**5** (1), 2 <https://doi.org/10.1186/s40842-019-0077-8> (2019).
37. Wunsch, C. *et al.* Lack of association between TCF7L2 gene variants and type 2 diabetes mellitus in a Brazilian sample of patients with the risk for cardiovascular disease. *Endocr Regul*.**53** (1), 1–7 (2019).
38. Loos, R. J. *et al.* TCF7L2 polymorphisms modulate proinsulin levels and β -cell function in a British Europid population. *Diabetes*.**56** (7), 1943–1947 (2007).
39. Chen, X. *et al.* The Diabetes Gene and Wnt Pathway Effector TCF7L2 Regulates Adipocyte Development and Function. *Diabetes*.**67** (4), 554–568 Epub 2018/01/11 <https://doi.org/10.2337/db17-0318> (2018).
40. Shu, L. *et al.* Decreased TCF7L2 protein levels in type 2 diabetes mellitus correlate with downregulation of GIP-and GLP-1 receptors and impaired beta-cell function. *Hum Mol Genet*.**18** (13), 2388–2399 (2009).
41. Kelsey, J. L., Whittemore, A. S., Evans, A. S. & Thompson, W. D. Methods in observational epidemiology: Monographs in Epidemiology and Biostatistics; 1996.
42. Graham, I. *et al.* European guidelines on cardiovascular disease prevention in clinical practice: executive summary: Fourth Joint Task Force of the European Society of Cardiology and Other Societies on Cardiovascular Disease Prevention in Clinical Practice (Constituted by representatives of nine societies and by invited experts). *Eur Heart J*.**28** (19), 2375–2414 (2007).
43. Semenkovich, C. F. Insulin resistance and atherosclerosis. *J Clin Invest*.**116** (7), 1813–1822 Epub 2006/07/11 <https://doi.org/10.1172/jci29024> (2006).
44. WHO EP. Executive summary of the clinical guidelines on the identification, evaluation, and treatment of overweight and obesity in adults. *Arch Intern Med*.**158**, 1855–1867 (1998).
45. Grundy, S. M. *et al.* Diagnosis and management of the metabolic syndrome: an American Heart Association/National Heart, Lung, and Blood Institute Scientific Statement. *Circulation*.**112** (17), 2735–2752 Epub 2005/09/15 <https://doi.org/10.1161/circulationaha.105.169404> (2005).
46. Barrett, J. C., Fry, B., Maller, J. & Daly, M. J. Haploview: analysis and visualization of LD and haplotype maps. *Bioinformatics*.**21** (2), 263–265 (2004).
47. Devlin, B. & Risch, N. A comparison of linkage disequilibrium measures for fine-scale mapping. *Genomics*.**29** (2), 311–322 (1995).
48. Hudson, R. R. & Kaplan, N. L. Statistical properties of the number of recombination events in the history of a sample of DNA sequences. *Genetics*.**111** (1), 147–164 (1985).
49. Wang, N., Akey, J. M., Zhang, K., Chakraborty, R. & Jin, L. J. T. A. J. H. G. Distribution of recombination crossovers and the origin of haplotype blocks: the interplay of population history, recombination, and mutation. *The American Journal of Human Genetics*.**71** (5), 1227–1234 (2002).

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Supplementarydata.docx](#)