

# Factors associated with malnutrition in children <5 years in western Kenya: A hospital-based unmatched case control study

**Edwin Gudu**

Moi Teaching and Referral Hospital

**Mark Obonyo**

Ministry of Health

**Victor Omballa**

Kenya Medical Research Institute

**Elvis Oyugi**

Ministry of Health

**Cecilia Kiilu**

Moi University

**Jane Githuku**

Ministry of Health

**Zeinab Gura**

Ministry of Health

**James Anthony Ransom** (✉ [james.ransom@piretpartners.com](mailto:james.ransom@piretpartners.com))

Piret Partners Consulting <https://orcid.org/0000-0001-6528-3994>

---

## Research article

### Keywords:

**Posted Date:** December 31st, 2019

**DOI:** <https://doi.org/10.21203/rs.2.11617/v2>

**License:**   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

**Version of Record:** A version of this preprint was published on July 29th, 2020. See the published version at <https://doi.org/10.1186/s40795-020-00357-4>.

# Abstract

**Background** Globally, under-nutrition accounts for >3 million deaths annually among children <5 years, with Kenya having ~35,000 deaths. We sought to identify factors associated with malnutrition in children aged <5 years in Western Kenya. **Methods** We conducted a hospital-based unmatched case-control study carried out between December 2016 and March 2017. Cases were defined as children aged 6-59 months with either z-score for weight-for-height,  $\leq -2SD$  or  $\geq +2SD$ ; weight-for-age,  $\leq -2SD$  or  $\geq +2SD$ ; or height-for-age of  $\leq -2SD$ . Controls were children aged 6-59 months with age-appropriate anthropometric measurements. Cases were consecutively recruited while systematic random sampling was used to select three controls. Unconditional logistic regression using backward elimination was used to identify independent associative factors using Epi-Info.

**Results** A total 94 cases and 281 controls were recruited. Of the cases, 86% (81/94) were under-nourished. Mother not having attended ante-natal clinic (OR=7.9; 95% CI: 1.5-41.2), deworming (OR=0.8; 95% CI: 0.4-1.2) and pre-lacteal feeding (OR=1.8; 95% CI: 1.1-3.0) were associated with under-nutrition. Delayed developmental milestones (AOR=13.9; 95% CI: 2.8-68.6); low birth weight (AOR=3.3; 95% CI: 1.4-7.6) and paternal lack of formal education (AOR=4.9; 95% CI: 1.3-18.9) were independently associated with under-nutrition. Urban residence (OR=5.4; 95% CI: 0.5-68.8) and male gender (OR=2.7; 95% CI: 0.7-11.8) were associated with over-nutrition with age <12 months (AOR=6.8; 95% CI: 1.6-29.7) and family income >5,000 Kenya shillings (AOR=4.6; 95% CI: 1.2-17.8) being independently associated.

**Conclusion** Proper pre-natal care, child feeding practices and deworming programs should be enhanced to reduce pediatric malnutrition.

## Introduction

Malnutrition refers to a state of either under-nutrition or over-nutrition. Under-nutrition occurs when the diet a person consumes does not meet their body's requirement for growth and development whereas over-nutrition occurs when a person consumes too many calories [1]. Good nutrition and feeding practices are critical to a child's growth and development especially during the first 2 years of life [2]. Under-nutrition has been known to impair a child's immunity leading to recurrent infections. It also impairs physical growth and cognitive development [3]. Long term effects of stunting on females also include obstructive labor once they reach the reproductive age group [3].

Under-nutrition is a great cause of morbidity and mortality especially in developing countries. Globally, malnutrition contributes to more than 3 million deaths among children <5 years annually [4]. UNICEF estimates that in Kenya, 239,446 children suffer from moderate acute malnutrition (MAM) and 2,600 children suffer from severe acute malnutrition (SAM) and are on clinical follow up. Under-nutrition also contributes to about 35,000 deaths among children <5 years each year in Kenya [4]. Stunting has also been linked to development of non-communicable diseases and lower adult productivity later in life. Children <5 years who are prone to recurrent infectious diseases such as diarrheal illnesses, respiratory

tract infections, tuberculosis and malaria usually have under-nutrition as a co-morbidity which if not treated promptly lead to high mortality rates [5].

According to Kenya Demographic Health Survey 2014; 26% of children <5 years are stunted, 4% are wasted and 11% are underweight. Malnutrition still remains to be a matter of public health concern in western Kenya. According to the survey 25.2% of children <5 years are stunted while 8.2% are severely stunted [6]. This means that 1 in 4 children suffer from chronic under-nutrition. Therefore, understanding of the factors associated with malnutrition (especially under-nutrition) is vital in preventing the development of long-term effects.

This study aimed to determine the clinical, demographic, and socio-economic factors associated with malnutrition in children <5 years by comparing those with malnutrition to those without. This study will contribute to the knowledge on prevention of malnutrition among children <5 years in Western Kenya by outlining the predisposing factors. This will by extension decrease morbidity and mortality related with the condition.

## **Materials And Methods**

### ***Study area***

The study was carried out at Alupe Sub-County Hospital. The hospital is a level 4 hospital located in Angorom ward, Teso South Sub-County in Busia County serving a catchment population of 34,321 persons [Figure 1] [7].

### ***Study Design***

We conducted a facility-based unmatched case control study carried out between May 2017 and June 2017. The study population consisted of all children below five years attending the child welfare clinic and the outpatient clinic within the hospital during the study period.

### ***Study hypothesis***

We hypothesized that improper infant and young child feeding practices as well as inconsistent child welfare clinic follow-up do not contribute a significant proportion of malnutrition in children <5 years attending Alupe Hospital in western Kenya, using the UNICEF conceptual framework for malnutrition [Figure 2].

### ***Case Definitions***

Under-nourished child was defined as a child aged 6-59 months attending the hospital as an inpatient or outpatient whose anthropometric measurements were not appropriate for their age with z-scores (Weight-for-Height WHZ, Weight-for-Age WAZ, Height-for-Age HAZ) of less than or equal to -2 SD. Weight for age Z-Scores from the WHO Charts were used to define presence of under-nutrition [8].

Over-nourished child was defined as a child aged 6-59 months attending the hospital as an inpatient or outpatient whose weight for height index (WHZ) and weight for age (WAZ) Z score is more than or equal to +2 SD.

A participant was classified as stunted if height-for-age z-score was less than -2SD and severely stunted if height-for-age z-score was less than -3SD. Wasting was defined as weight-for-height z-score of less than -2SD while severe wasting was weight-for-height z-score of less than -3SD. Any participant with weight-for-age z-score less than -2SD was said to be underweight while -3SD was said to be severely underweight. On the contrary, those with weight-for-age or weight for height z-scores more than +2SD were classified as overweight, while those with more than +3SD were obese.

### ***MUAC calculations***

For the MUAC cut-points to determine whether a child was under- or over-nourished, we used the cut-points of any child with MUAC <126 mm was classified as under-nourished and those with MUAC>179 mm were classified as over-nourished [8].

### ***Definition of controls***

Any child aged 6-59 months attending the hospital as an inpatient or outpatient whose anthropometric measurements are appropriate for their age with z-scores between -2SD and +2SD [9].

### ***Sample size determination***

The sample size was calculated using statistical software Epi Info® version 7.2.0. The study assumed a 95% confidence interval, 80% power, 10% wasting among controls [10], an Odds ratio of 2.5 and the ratio of cases to controls of 1:3. Using these assumptions, the minimum sample size was 375 (94 cases and 282 controls).

### ***Selection of cases and controls***

The cases were sampled consecutively given the low number seen each day for eligible children enrolled for nutritional support in welfare clinic. The controls were selected through systematic random sampling. The average number of children below five years visiting the outpatient section of the child welfare clinic on a daily basis was used as a sampling frame. This was determined by obtaining the number of children visiting the out-patient clinic between April and June of three preceding years before the study. The study was conducted during weekdays within the duration of the study period hence the number of controls to be enrolled in the study on any single day was pre-determined. Using the average number of patients seen each day at the clinic and number of controls to be enrolled in the study each day, a sampling interval was determined and the first control was picked randomly between one and the sampling interval. The sampling interval was then added to enroll the remaining controls. Any eligible participant whose legal parent/guardian did not give consent was replaced by the next available participant whose legal parent/guardian consented to the study.

## ***Data Collection***

Triage was carried out by the hospital staffs as is the norm and all critically and severely ill patients were urgently attended to by the hospital clinicians as per procedures and guidelines of the hospital. The weight was measured using electronic digital weighing scale (Seca®). For height/length, children <2 years were measured lying down (recumbent length) while those who were >2 years were measured standing up. For mid-upper arm circumference (MUAC) and head circumference, a non-stretch tape was used.

A pre-tested interviewer-administered questionnaire was used for each study participant to obtain demographic, clinical, nutritional, social and economic information. Each patient was de-identified by a unique code to ensure their privacy and maintenance of confidentiality.

## ***Data management***

Data entry, cleaning, validation and analysis was done using Microsoft Excel (Microsoft, Seattle, WA, USA), and Epi info version 7 (CDC, Atlanta, GA, USA). Anthropometric data was analyzed using WHO Anthro® software version 3.2.2 (WHO Anthro®) to assess nutritional indicators like weight-for-length/weight-for-height (wasting), weight-for-age (underweight or overweight), length-for-age/height-for-age (stunting), MUAC-for-age and HC-for-age. The software then provided the z-scores based on gender, age and the anthropometric measurements. We calculated measures of central tendency and dispersion for the continuous variables and proportions for categorical variables. For univariable analysis, we calculated odds ratios (OR), 95% confidence intervals (CI), chi-square statistics and p-values. Variables with p-value  $\leq 0.05$  were considered to be statistically significant. We carried out unconditional logistic regression with variables that had p-values of <0.2 at univariable analysis. A backward elimination stepwise method was used to identify independent factors associated with malnutrition. During model building, any variable that caused an insignificant increase in deviance on removal from the model were left out of the model while the variable that caused a significant increase in deviance on removal were retained in the model. All variables removed from the model when a backward stepwise method was performed and those known to be potential cofounders or factors associated with malnutrition from previous studies were tested for confounding, any of the mentioned variables that had a more than ten percentage change (>10%) between the crude and adjusted odds ratio was considered as a confounder. The final model after testing for all biologically and statistically plausible interactions had only variables with p-value of  $\leq 0.05$ .

# **Results**

## ***Description of the study participants***

There were 375 participants (94 cases and 281 controls). The median age of all the study participants was 16 months (IQR 10, 22). Males comprised 51% (191/375) with females accounting for 49%

(184/375). Of the cases, males accounted for 57% (54/94) while the females were 43% (40/94). Of the controls, females accounted for 51% (144/281) while the males accounted for 49% (137/281).

### ***Nutritional status of cases***

Of the cases, 84% (79/94) were under-nourished and 16% (15/94) over-nourished. Among those that were under-nourished, by assessing the WHZ score, 20% (16/79) were wasted while 9% (7/79) were severely wasted. Among the same group, using WAZ score, 39% (31/79) were underweight while 29% (23/79) were severely underweight. Using the HAZ score among the under-nourished, 46% (36/79) were stunted while 38% (30/79) were severely stunted. Among the over-nourished, 7% (1/15) were stunted while 20% (3/15) were severely stunted (**Table 1**).

### ***Univariable and multivariable analysis of factors associated with under-nutrition***

On univariable analysis, socio-demographic factors like high birth order of five or more (OR= 2.3; 95% CI: 0.9 – 6.0) , living in urban areas (OR=1.9; 95% CI: 0.8 - 4.3), children whose mothers had no formal education (OR=2.0; 95% CI 0.9 - 4.4), those whose fathers had no formal education (OR=4.6; 95% CI: 1.4 – 15.0) and those who came from large family sizes of more than 6 occupants (OR=1.8; 95% CI: 1.1 – 3.0) had higher odds of developing under-nutrition.

Pre-natal maternal factors were also shown to increase odds of developing under-nutrition. These included: participants whose mothers' did not attend antenatal clinic (ANC) at least once (OR=7.9; 95% CI: 1.5 - 41.6), participants whose mothers who did not attend 4 ANC visits as recommended by WHO (OR=1.6; 95% CI: 0.9 - 2.7) and those whose mothers had illness during pregnancy (OR=1.7; 95% CI: 1.0 - 2.8). The participants who were born preterm (OR=2.0; 95% CI: 0.6 - 7.4) and those with low birth weight (OR=2.8; 95% CI: 1.2 – 6.2) had higher odds of under-nutrition compared to term babies and those with normal birth weights.

Post-natal factors such as failure to complete or not being up-to-date on immunizations as per the national immunization schedule (OR=2.2; 95% CI: 0.7-7.2) and human immunodeficiency virus (HIV) sero-exposure (OR=1.4; 95% CI: 0.6 - 3.5) and delayed developmental milestones (OR=18.9; 95% CI: 4.1- 87.5) also increased the odds of developing under-nutrition. The participants who were eligible for deworming and had been dewormed at least once were protected from under-nutrition (OR=0.8; 95% CI: 0.4 - 1.2). Infant and young child feeding practices also affected nutritional status of the participants. During the study period, 24% (89/375) of the participants were receiving pre-lacteal feeds increased their odds of under-nutrition (OR 1.8; 95% CI: 1.0 - 3.1). Exclusive breastfeeding for 6 months as recommended by the WHO was also widely practiced with 72% (271/375) of the participant's parents adhering to this guideline. Forty percent (149/375) of the participant's parents still used bottle with nipple for feeding, while 31% (116/375) ceased breastfeeding before the recommended 2 years of age. However, during the study period, there was no statistically significant association between duration of exclusive breastfeeding for the first six months of life (OR=1.1; 95% CI 0.6 – 2.0), cessation of breastfeeding at less

than 2 years (OR= 0.6; 95% CI 0.2 – 1.7) or bottle with nipple feeding (OR=0.9; 95% CI 0.5 – 1.5) and developing under-nutrition.

Economic factors of the families also affected the nutritional status of the participants. Those whose mothers were unemployed had higher chances of under-nutrition (OR=1.8; 95% CI 1.0 -3.1) whereas those families with an average monthly income of above 5,000 Kenya shillings (KES) were protected of under-nutrition (OR 0.7; 95% CI 0.4 - 1.2) (**Table 2**).

On multivariable analysis, delayed developmental milestones (AOR=13.9; 95% CI: 2.8 - 68.6); low birth weight (AOR=3.3; 95% CI: 1.4 – 7.6) and paternal lack of formal education (AOR=4.9; 95% CI: 1.3 – 18.9) were found to be independently associated with under-nutrition.

### ***Univariable and multivariable analysis of factors associated with over-nutrition***

On univariable analysis, socio-demographic factors like age less than 12 months (OR= 6.0; 95% CI: 1.5 – 25.3), being male (OR=2.7; 95% CI: 0.7 – 11.8), living in urban areas (OR=5.4; 95% CI: 0.5 – 68.8), children whose mothers had no education (OR=3.3; 95% CI 0.4 - 25.8) had higher odds of developing over-nutrition. Participants with high birth weight of more than 2500 grams (OR=3.3; 95% CI 0.4 - 25.8) and those who came from families with monthly income more than 5,000 KES (OR=3.3; 95% CI 0.4 - 25.8) also had higher odds of over-nutrition (**Table 3**).

On multivariable analysis, age  $\leq$ 12 months (AOR=6.8; 95% CI: 1.6 – 29.7) and monthly family income  $\geq$  5,000 KES (AOR=4.6; 95% CI: 1.2 – 17.8) were found to be independently associated with over-nutrition.

## **Discussion**

The study identified various factors affecting nutritional status among children <5 years which need to be adequately addressed. This included both pre-natal and post-natal factors as well as infant and young child feeding practices. Therefore, consistent follow-up of pregnant mothers from the antenatal period and post-natal care of the children <5 years needs to be enhanced.

Under-nutrition was the predominant form of malnutrition among the study participants. However, the rising number of cases of over-nutrition point to the growing trend of double burden of malnutrition. Among the undernourished, we found that stunting was the most common form of malnutrition, followed by children who were underweight and wasting being the least common among the study population. Stunting was common among cases of under-nutrition and over-nutrition alike. Stunting is a chronic form of malnutrition that results from prolonged non-adherence to proper dietary requirements to meet the body's physiological needs. These findings were similar to those of a demographic and health survey carried out in the Western Kenya in 2014 [6]. Other studies carried out in Burundi and Uganda also had similar findings [11, 12].

Deworming of children >1 year of age was also found to be protective of under-nutrition. This finding was in line with another study done in India among pre-school children which showed substantial weight gain

among children who were dewormed [13]. This is because intestinal nematodes affect absorption of both micro and macronutrients which are vital for a child's growth. However, current systematic reviews show little benefit is derived from mass deworming. They show that children found to be worm infested are the ones that gain weight more significantly compared to non-worm infested children [14, 15].

Proper breastfeeding practices for children are advocated for by WHO [16]. Children that are breastfed up to 2 years of age show quicker linear growth than those breastfed for shorter durations [2]. Feeding practices such as bottle with nipple feeding, breastfeeding within thirty minutes of delivery, exclusive breast feeding for 6 months and cessation of breastfeeding at 2 years were also assessed during the study. However, they were not statistically significantly associated with under-nutrition. In contrast, the giving of pre-lacteal feeds adversely affected nutritional status and predisposed the children to under-nutrition. This has also been shown by other studies [17] [18] [19]. This could be due to the fact that pre-lacteal feeding affects the quality and quantity of breastfeeding which in turn affects the nutritional intake by the child. As such, proper education on feeding practices during post-natal period should be enhanced.

Children with under-nutrition were also shown to be more likely to have delayed developmental milestones. This finding was consistent with other studies [20] [21, 22]. This could be because they lack the macro and micronutrients necessary for normal growth and development. Children with prematurity and low birth weight also had higher odds of under-nutrition. These findings were similar from a review done in several countries [23]. This could be due to the fact that they require more nutrients for catch-up growth which if not provided in adequate quantities leave them vulnerable to develop under-nutrition. These children should therefore be followed up more closely.

We also found that lack of parental formal education was linked to development of under-nutrition with paternal illiteracy being shown to have a greater influence. This finding concurred with other studies [24] [19]. This could be due to the fact that the community being a patriarchal society, the fathers control the family's resources. As such, lack of formal education could mean no formal employment and by extension no regular source of income to provide for their families.

Our study also showed that the cases of over-nutrition were also high, compared to findings of other studies in Kenya, despite the hospital serving a population of predominantly low socio-economic status [25]. This clearly points to the double burden of malnutrition that is supported by other literature based on a critical review done in other lower middle income countries [26]. This is a new development over the last couple of decades that needs to be further explored to halt and decrease the burden of cases of over-nutrition.

During the study period, children aged less than 12 months were more likely to be over-nourished. This finding was similar to another study carried out in Kenya in 2009 [27]. This could be due to the fact that younger children are more likely to receive more attention and feeding effort from their parents as compared to older children. Male gender was also positively associated with over-nutrition. A study carried out in Kenya in 2016 had similar findings [28]. This could probably be due to the value and

cultural preferences placed on the male child. As such, they are likely to be better fed as compared to the girl child. This has also been shown in other Sub-Saharan African countries [29]

Children who came from households in urban areas and those who came from families with higher average monthly income had higher odds of over-nutrition. This finding was similar to other studies [27]. This could be due to the higher levels of income which increase their ability to provide more than enough nutrition for their growing children.

Maternal lack of formal education also increased the chances of developing over-nutrition. This finding was contrary with other studies carried out in Sub-Saharan Africa [29]. High birth weight was also linked to increased chances of over-nutrition. This has also been shown by other studies [29, 30]. However, the exact mechanism of this link has not yet been clearly described.

The study had some limitations. It was a hospital-based case control study and as such, its findings cannot be generalized to the entire population of under-five children in Western Kenya. The data collected on some of the variables could be susceptible to recall bias more so if the child was brought in by a guardian.

## Conclusions And Recommendations

Proper pre-natal care, child feeding practices and deworming programs should be enhanced. As such, we recommend that close monitoring especially of children more likely to be malnourished should be enhanced. This can be done by providing job aids to providers to help them talk to parents about adherence to key recommended practices such as appropriate feeding, continuous auditing of patient outcomes, and better use of data for improved decision-making should be implemented at these facilities.

Proper infant and young child feeding practices and deworming should be emphasized. Provider advocacy and better health education to parents should be intensified in the region for better outcomes. The hospital in conjunction with Busia County Government should organize for regular outreach to the community targeting pregnant and lactating mothers, strengthen deworming programs for children >1 year and all children with delayed developmental milestones. It should also organize for health advocacy camps targeting the parents with children < 5 years to educate them on the proper infant and young child feeding practices.

## Abbreviations

MAM: Moderate Acute Malnutrition; MUAC: Mid Upper Arm Circumference; WHZ: Weight for Height Z-score; WAZ: Weight for Age Z-score; HAZ: Height for Age Z-score; SD: standard deviation; OR: Odds ratio; AOR: Adjusted odds ratio; CI: Confidence interval

## Declarations

## Acknowledgement

We would like to appreciate the participants whose data was used in this study. We also acknowledge Alupe Sub-County Hospital and Busia County Government. We also thank the Kenya Field Epidemiology and Laboratory Training Programme for the financial and technical assistance in the carrying out of this study.

## Funding

This study was fully funded by Kenya Field Epidemiology and Laboratory Training Program. The funding body of this study did not participate in the design or conclusion of the study.

## Availability of data and materials

The datasets used and/or analyzed during the current study are available as Additional file 1.

## Author's contributions

EG and MO conceived the study, EG collected data, EG and MO analyzed the data and drafted the manuscript. All the authors helped with the interpretation of the results, read, critically reviewed and approved the final manuscript.

## Ethics approval and consent to participate

Ethical clearance was sought from Institutional Review Ethics Committee Moi University/Moi teaching and Referral Hospital under reference FAN: IREC 1870. Permission for the study was also sought from the County health department and the hospital administration. During the interviews informed consent was obtained from parents/legal guardians of all study participants after explaining the objectives of the study. Measures were taken to assure confidentiality of the information provided during these interviews and codes were used to de-classify personal identifying information of study participants. Measures were taken to assure collected data were properly stored and secured and only accessible to the investigators

## Consent for publication

Not applicable.

## Competing interests

The authors declare that they have no personal or financial competing interests that may bias publication of this manuscript.

## References

1. UNICEF, *Progress for children*, in *UNICEF*. 2016.

2. Onyango, A.W., S.A. Esrey, and M.S. Krammer, *Continued breastfeeding and child growth in the second year of life: a prospective cohort study in western Kenya*. The Lancet, 1999. 354(9195): p. 2041–2045.
3. Services, J.C.o.I.C.s., *Orphan Nutrition*. 2017.
4. UNICEF, *Kenya*. 2016.
5. Intitute, I.F.P.R., *Global Nutrition Report*. 2015, International Food Policy Research Intitute: Washington DC.
6. Statistics, K.N.B.o., *2014 Kenya Demographic and Health Survey*. 2015, Kenya National Beaureau of Statistics: Nairobi, Kenya.
7. System, K.H.I., *DHIS2*, in *DHIS2*. 2016.
8. World Health Organization and World Health Organization. *Nutrition for Health, 2009. WHO child growth standards: growth velocity based on weight, length and head circumference: methods and development*. World Health Organization.
9. Organization, W.H. *World Health Organization (WHO) - Global Database on Child Growth and Malnutrition*. 2017.
10. Shikur, B., *The relationship between Malaria and Malnutrition among under five children in Adami Tulu District, south-central Ethiopia: a case-control study*. 2014, Addis Ababa: Addis Ababa University College of health sciences.
11. Peter, Y., *Child Growth and Determinant Factors: A Case Study of Burundi*, in *Bioscience Engineering*. 2016, Gent.
12. Gilbert, H., *An investigation on factors associated with malnutrition among underfive children in Nakaseke and Nakasongola districts, Uganda*. BMC Pediatrics, 2015. 15(134).
13. Shally, A., R. Peto, and V. Pande, *Effects of deworming on malnourished preschool children in India*. Plos: Neglected Tropical Diseases, 2008. 2(4): p. 223.
14. Taylor, R., *Deworming school children in developing countries*. 2015, Cochrane: Cochrane Library.
15. Tim, A., *Deworming delusions? Mass drug administration in East African Schools*. Journal of Biosocial Science, 2016. 48: p. 116-147.
16. Organization, W.H., *Indicators for assessing breastfeeding practices*. 1991, World Health Organization: Geneva, Switzerland.
17. Nayak, R. and D. Mallapur, *Determinants of nutritional status in under five children- A cross sectional study*. Annals of community health, 2014. 2(2): p. 26-29.
18. Wagaye, F. and A. Tariku, *Undernutrition and associated factors among children aged 6-59 months in East Belesa District, northwest Ethiopia: a community based cross-sectional study*. BMC Public Health, 2016. 16(506): p. 1-10.
19. Asfaw, M. and D. Lamessa, *Prevalence of undernutrition and associated factors among children aged between six to fifty nine months in Bule Hora district, South Ethiopia*. BMC Public Health, 2015. 15(41).

20. A., A., *Children at risk for developmental delay can be recognized by stunting, being underweight, ill health, little maternal schooling or high gravidity*. NIH Public Access, 2010. 51(6): p. 652-659.
21. Hyun, P., *The impact of nutritional status and longitudinal recovery of motor and cognitive milestones in internationally adopted children*. International Journal of Environmental Research and Public Health, 2010. 8: p. 105 - 116.
22. Deepti, D., *A Community-Based Study on Growth and Development of Under-Five Children in an Urbanized Village of South Delhi*. Journal of Tropical Pediatrics, 2016. 62: p. 446 - 456.
23. Christian, P. and S. Lee, *Risk of childhood undernutrition related to small-for-gestational age and preterm birth in low- and middle-income countries*. International Journal of Epidemiology, 2013. 42: p. 1340-1355.
24. Zaida, H. and S. Luis *Cross-Sectional Study of Malnutrition and Associated Factors among School Aged Children in Rural and Urban Settings of Fogera and Libo Kemkem Districts, Ethiopia*. 2014.
25. Hoffman, D., Cacciola, T., Barrios, P. and Simon, J., 2017. Temporal changes and determinants of childhood nutritional status in Kenya and Zambia. *Journal of Health, Population and Nutrition*, 36(1), p.27.
26. Emma, T., *Childhood dual burden of under- and overnutrition in low- and middle-income countries: A critical review*. Food and Nutrition Bulletin, 2014. 35(2).
27. Constance, G., *Childhood overweight and obesity among Kenyan pre-school children: association with maternal and early child nutritional factors*. Public Health Nutrition, 2009. 13(4): p. 496-503.
28. Tankoi, E.O., *Determinants of Malnutrition among Children Aged 6-59 Months in Trans-Mara East Sub-County, Narok County, Kenya*. International Journal of Public Health and Safety, 2016. 1(3).
29. Keino, S., *Determinants of stunting and overweight among young children and adolescents in sub-Saharan Africa*. Food and Nutrition Bulletin, 2014. 35(2): p. 167 - 178.
30. Qiao, Y., *Birth weight and childhood obesity: a 12-country study*. International Journal of Obesity Supplements, 2015. 5: p. 74 - 79.

## Tables

Table 1: Nutritional status of the study participants, Alupe Sub-County Hospital, 2017

Category/Variable	n =375	Proportion (%)	95% CI
<b>Weight-for-Height Z score (WHZ)</b>			
Normal WHZ (> -2 and < +2)	342	91.2%	87.6 - 93.8
Overweight ( $\geq +2$ )	10	2.7%	1.3 - 5.0
Wasted ( $\leq -2$ and < -3)	16	4.3%	2.5 - 7.0
Severely Wasted ( $\leq -3$ )	7	1.9%	0.8 - 4.0
<b>Weight-for-Age Z score (WAZ)</b>			
Normal WAZ (> -2 and < +2)	313	83.5%	79.5 - 87.0
Overweight ( $\geq +2$ )	8	2.1%	1.0 - 4.3
Underweight ( $\leq -2$ and < -3)	31	8.3%	5.8 - 11.7
Severely Underweight ( $\leq -3$ )	23	6.1%	4.0 - 9.2
<b>Height-for-Age Z score (HAZ)</b>			
Normal HAZ (> -2 and < +2)	304	81.1%	76.7 - 84.8
Stunted ( $\leq -2$ and < -3)	38	10.1%	7.4 - 13.8
Severely Stunted ( $\leq -3$ )	33	8.8%	6.2 - 12.3

Table 2: Factors associated with under-nutrition participants, Alupe Sub-County Hospital, 2017\*

Variable	Category	Cases n=79 (n/%)	Controls n=236 (n/%)	OR	95% CI	P value
Residence	Urban	69 (87.3)	219 (92.7)	1.9	0.8 - 4.3	0.21
	Rural	10 (12.7)	17 (7.2)			
ANC attendance	Yes	5 (6.3)	2 (<1)	7.9	1.5 - 41.6	0.02
	No	74 (93.7)	234 (99.1)			
Illness during pregnancy <sup>1</sup>	Yes	43 (54.4)	102 (43.2)	1.7	1.0 - 2.8	0.07
	No	33 (41.7)	129 (54.6)			
Gestation	Pre-term	4 (5.1)	6 (2.5)	2.0	0.6 - 7.4	0.46
	Term	75 (94.9)	230 (97.5)			
Low birth weight <sup>2</sup>	Low	13 (16.5)	14 (5.9)	2.8	1.2 - 6.2	0.02
	Normal	57 (72.2)	169 (71.6)			
Delayed developmental milestones	Delayed	11 (13.9)	2 (<1)	18.9	4.1 - 87.5	<0.001
	Normal	68 (86.1)	234 (99.1)			
Deworming	Yes	49 (62.0)	163 (69.1)	0.7	0.4 - 1.2	0.31
	No	30 (38.0)	73 (30.9)			
Pre-lacteal feeds <sup>3</sup>	Yes	25 (31.6)	50 (21.2)	1.7	1.0 - 3.1	0.08
	No	53 (67.1)	185 (78.4)			
Mother's education level <sup>4</sup>	None	11 (13.9)	18 (7.6)	2.0	0.9 - 4.4	0.14
	Other	67 (84.8)	218 (92.4)			
Father's education level <sup>5</sup>	None	7 (8.9)	5 (2.1)	4.6	1.4 - 15.0	0.02
	Other	69 (87.3)	227 (96.2)			
Average monthly income KES.	≥5,000	25 (31.6)	96 (40.7)	0.7	0.4 - 1.2	0.20
	<5,000	54 (68.4)	140 (59.3)			
Birth order	≥5	8 (10.1)	11 (4.7)	2.3	0.9 - 6.0	0.14
	<5	71 (89.9)	225 (95.3)			
Number of house occupants	≥6	41 (51.9)	88 (37.3)	1.8	1.1 - 3.0	0.03
	<6	38 (48.1)	148 (62.7)			

\*2 cases and 5 controls were brought by their guardians who did not know whether the child's mother had any illness during pregnancy; 6 cases and 52 controls had high birth weight while 3 cases and 1 control were brought by guardians who did not know their birth weight; 1 case and 1 control had never been breastfed since birth (were on formula feeds);

1 case was brought by a guardian who did not know the mother's educational level; 3 cases and 4 controls were brought by guardians who did not know the father's education levels

Table 3: Factors associated with over-nutrition among participants, Alupe Sub-County Hospital, 2017

Variable	Category	Cases n=15 (n/%)	Controls n=45) (n/%)	OR	95% CI	P value
Age	≤12 months	12 (80.0)	18 (40.0)	6.0	1.5 - 25.3	0.02
	>12 months	3 (20.0)	27 (60.0)			
Gender	Male	10 (66.7)	19 (42.2)	2.7	0.7 - 11.8	0.14
	Female	5 (33.3)	26 (57.8)			
Residence	Urban	3 (20.0)	2 (4.4)	5.4	0.5 - 68.8	0.09
	Rural	12 (80.0)	43 (95.6)			
High birth weight* (>2500grams)	Yes	5 (33.3)	10 (22.2)	2.1	0.6 - 8.0	0.44
	No	8 (53.3)	34 (75.6)			
Dewormed	Yes	2 (13.3)	26 (57.8)	0.2	0.01 - 2.5	0.54
	No	13 (86.7)	19 (42.2)			
Mother's education level	None	2 (13.3)	2 (4.4)	3.3	0.4 - 25.8	0.52
	Primary education and above	13 (86.7)	43 (95.6)			
Average family income (KES.)	≥5,000	10 (66.7)	15 (33.3)	4.0	1.0 - 17.4	0.04
	<5,000	5 (33.3)	30 (66.7)			
Exclusive breastfeeding**	≥6months	10 (66.7)	38 (84.4)	0.5	0.1 - 1.8	0.48
	<6months	4 (26.7)	7 (15.6)			

\* 2 cases and 1 control had low birth weight; \*\* 1 of the cases was still being exclusively breastfed as at the time of the study.

# Figures




Figure 1

Geographical location of study site in Busia County, Kenya, indicating the location of the Alupe Sub-county hospital (red dot)




Figure 2

Shows conceptual framework of the study adapted from UNICEF conceptual framework for malnutrition, 1990