

Birt-Hogg-Dubé Syndrome in Chinese Patients: A Literature Review of 120 Families

Guofeng Zhang

The First Affiliated Hospital of USTC: Anhui Provincial Hospital

Xianmeng Chen

The First Affiliated Hospital of USTC: Anhui Provincial Hospital

Kaifeng Xu

Peking Union Medical College Hospital

Xiaowen Hu (✉ hu.xiaowen@hotmail.com)

The First Affiliated Hospital of USTC: Anhui Provincial Hospital <https://orcid.org/0000-0002-3727-6049>

Research Article

Keywords: Birt-Hogg-Dubé syndrome, FLCN gene, pneumothorax, skin lesion, renal tumor

Posted Date: February 26th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-247951/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Objective

To clarify the epidemiological and clinical features of Birt-Hogg-Dubé syndrome (BHDS) in Chinese patients.

Methods

We identified reports on Chinese patients with BHDS by searching the China Academic Journals Database, Wanfang Chinese Database, and PubMed databases, either in Chinese or English languages published from January 1, 2006 to December 31, 2020. Studies without sufficient clinical data were excluded and cases under 18 years old were excluded.

Results

Twenty papers were included and comprised 120 families with 221 cases. Most families with BHDS were reported from institutions in Beijing (66.7%) and Jiangsu Province (15.8%); 80.8% of cases were reported within the past five years. The average duration from clinical presentation to diagnosis was 9.6 years. The average age was 47.0 ± 13.9 years (range, 18-84 years) and the ratio of male to female was 1:1.6. The most common manifestations of BHDS were multiple pulmonary cysts (92.4%), spontaneous pneumothorax (71.0%), skin lesions (18.1%) and renal tumors (3.6%). Pulmonary cysts were predominantly distributed in the lower lobe on chest CT imaging. Family history of spontaneous pneumothorax was identified in 84.7% of the families and average number of pneumothoraxes was 1.8 (range, 1-6). The *FLCN* gene mutation c.1285dupC/delC in exon 11 was the most frequent mutation observed (17.4% of patients). The recurrence rate of pneumothorax after conservative treatment (including tube thoracostomy) was 29/41 (71%) while the pneumothorax recurred after surgical treatment (pulmonary bullectomy or pleurodesis) in only 4/37 (11%).

Conclusions

Although BHDS has been increasingly reported in the recent years, only minority of families were reported from institutions outside of Beijing and Jiangsu Province. The dominant clinical manifestations were pulmonary cysts associated with recurrent pneumothorax, while skin lesions and renal tumors were less commonly reported. Delayed diagnosis along with suboptimal management appear to represent critical challenges for Chinese patients with BHDS.

Background

Birt-Hogg-Dubé syndrome (BHDS, OMIM#135150) is a rare autosomal dominant inherited disorder, first described in 1977, featured by lung cysts, spontaneous pneumothorax, skin lesions (fibrofolliculomas and trichodiscomas) and renal tumors¹. It is caused by germline mutations in the folliculin (*FLCN*) gene, which is located on chromosome 17p11.2 and encodes a 579-amino acid protein with tumor suppressor

function, and interacts with the mammalian target of rapamycin (mTOR) and adenosine monophosphate-activated protein kinase (AMPK) signaling pathways²⁻⁵. Over 600 families with BHD have been reported worldwide in the past 40 years, mainly from Europe and USA. To date, over 200 *FLCN* gene variants have been identified to cause BHDS according to the Leiden Open Variation Database (<https://www.lovd.nl/>).

Multiple bilateral pulmonary cysts are the most common manifestation in both Asian and Caucasian individuals with BHDS. Fibrofolliculomas are highly specific manifestation occurring in more than 80% of Caucasian individuals with BHDS. Renal tumors develop in up to one-third of individuals with BHDS, with an average age of onset of 50 years reported in Caucasian individuals⁶. In East Asia, Japanese authors reported over 100 families with less frequent skin and renal manifestations⁷.

In 2008, the first detailed report of BHDS in Chinese patients focused on the gene mutations identified among patients manifesting spontaneous pneumothorax in Jiangsu province⁸. In 2017, Liu et al provided a systematic genetic and radiological description of a BHDS case series in the Chinese population⁹. Until recently, data on clinical features of Chinese patients with BHD were limited to case reports or small series, mainly with pulmonary features. Extrapulmonary abnormalities, including skin and kidney manifestations, are uncommonly reported in Chinese individuals of BHDS. With an increasing number of reports published in recent decades, it seemed warranted to summarize the prevalence and clinical characteristics associated with BHDS in the Chinese population in order to clarify differences between Chinese and Caucasian individuals with this disorder.

The purposes of this study are to fill this knowledge gap by exploring the epidemiology and disease characteristics of BHDS reported in China, focusing on clinical, radiological, genetic, diagnostic, and therapeutic aspects to improve recognition of this rare disease in the Chinese population.

Material And Methods

Study design and data collection

Firstly, the following keywords “Birt-Hogg-Dubé syndrome” or “BHD” or “BHDS” [All Fields] combined “China” or “Chinese” in the China Academic Journals Full-text Database (CAJ), Wanfang Chinese Database and PUBMED English Database were searched by Dr. Zhang in September 2020, and then updated in December 2020. This was supplemented with manual search to identify other relevant documents as conducted by two authors (Drs. Hu and Zhang). Articles were restricted to those published and available as full texts in either Chinese or English languages; conference abstracts were excluded. We included BHDS cases reported in China from January 1, 2008 to December 31, 2020; the diagnosis of BHDS was based on the criteria proposed by the European BHD consortium. BHDS could be diagnosed by fulfilling one major criterion or two minor criteria. The major criteria included: (1) at least five fibrofolliculomas or trichodiscomas (at least one histologically confirmed, of adult onset); (2) pathogenic *FLCN* germline mutation. The minor criteria included: (1) multiple lung cysts (bilateral basally located lung cysts with no other apparent cause, with or without spontaneous pneumothorax); (2) renal cancer

(early onset before age 50 or multifocal or bilateral, or mixed chromophobe and oncocytic histology); (3) a first-degree relative with BHDS¹⁰. Patients must have been living in China (including from Taiwan, Hong Kong, or Macau) and the papers must have provided enough information about the clinical features, including medical history, imaging data, gene results and therapeutic strategy. Exclusion criteria included duplicate reports, basic science research, and clinical studies without sufficient clinical data. Cases under 18 years old were excluded from the final analysis.

The study was waived by the institutional review board of the First Affiliated Hospital of University of Science and Technology of China because the data were obtained from published reported and were analyzed anonymously and retrospectively.

Clinical information including sex, age, family history, pulmonary cysts, detailed history of pneumothorax, skin lesion, renal tumor, gene testing, diagnosis, treatment, and outcome was collected.

Statistic methods

All data was expressed in the form of means and standard deviations ($\bar{x} \pm s$). SPSS 25.0 was used for data analysis and independent sample t-test was used to compare continuous variables. Chi-square test was used to compare categorical variables. Differences with a value of $P < 0.05$ was considered statistically significant.

Results

Demographic data

There were 49 BHDS-related papers in English or Chinese languages by Chinese authors between January 1, 2008 and December 31, 2020 (Fig. 1). Using the study criteria outlined above, 221 cases from 120 families with BHDS were included in the final analysis, which were reported in 20 papers (Table 1)^{8,9,11–28}. In 2008, the first 10 families with spontaneous pneumothorax and positive *FLCN* mutations in the Chinese population were described by a research team at Nanjing University⁸. As shown in Fig. 2, 97 (80.8%) of 221 cases were reported in the past five years. Furthermore, 60 (50%) families were reported in the past two years, and was equal to the number of cases reported in the preceding ten years. Although the reporting regions included 9 Chinese provinces (Table 2), most of the cases came from Beijing (80 families, 66.7%), followed by Jiangsu province (19 families, 15.8%). Among 221 patients, 38.9% were male and the ratio of male to female was 1:1.6. The average age at the time of diagnosis was 47.0 ± 13.9 years (range, 18–84 years).

Table 1
The features of enrolled papers of BHDS in the final study

Author	Province	Year of Publication	Number of Families	Number of Cases
Ren H-Z	Nanjing	2008	10	23
ShunYang So	Hong Kong	2009	1	1
GeeGwo Yang	Taiwan	2013	1	1
Zhichun Lin	Beijing	2014	1	3
Zhibo Liu	Beijing	2014	1	1
Yibing Ding	Nanjing	2015	9	40
Li Dong	Tianjin	2016	2	2
JF Zhu	Wuhan	2017	1	1
Lv Liu	Hunan	2017	2	8
Yaping Liu	Beijing	2017	27	27
Teng Li	Beijing	2017	2	2
Shengyu Hao	Shandong	2017	1	1
Xiaocan Hou	Hunan	2018	2	8
Fei Xie	Beijing	2019	1	1
Keqiang Liu	Beijing	2019	31	38
Qun Hu	Fujian	2019	1	1
Chunming Zheng	Beijing	2019	1	10
Yanguo Liu	Beijing	2020	16	37
Dandan Zong	Hunan	2020	1	7
Ting Guo	Hunan	2020	9	9
Total			120	221

Table 2
BHDS according reporting provinces in
China

Province	Family (%)	Case (%)
Beijing	80(66.7)	119(53.9)
Jiangsu	19(15.8)	63(28.6)
Hunan	14 (11.8)	32(14.6)
Tianjin	2 (1.7)	2 (0.9)
Shandong	1 (0.8)	1 (0.5)
Hubei	1 (0.8)	1 (0.5)
Fujian	1 (0.8)	1 (0.5)
Hong Kong	1 (0.8)	1 (0.5)
Taiwan	1 (0.8)	1 (0.5)
Total	120	221

Clinical manifestations

Spontaneous pneumothorax was observed in 157 patients; the characteristics of pneumothorax was described in 78 of these patients, including the age at initial pneumothorax, number of episodes, and side of pneumothorax. The average age at first pneumothorax episode was 39.2 ± 12.7 years (range, 18 to 71). The average number of pneumothorax per patient was 1.8 ± 1.1 (range, 1 to 6). Among these, 74 and 4 patients experienced unilateral and bilateral pneumothorax, respectively. Family history of pneumothorax was identified in 84.7% patients.

Among 221 patients, 40 (18.1%) patients presented with skin lesions, including 6 fibrofolliculomas, 1 with sarcoma, 11 with papules, and 22 cases without specific descriptions. Eight patients had renal cancers (3.6%) of whom two cases of renal cell carcinoma were not described in detail. One patient was diagnosed with bilateral renal carcinomas including chromophobe cell carcinoma on the left and clear cell carcinoma on the right. The remaining 5 patients had unilateral renal carcinoma. Thyroid carcinoma was found in two patients from two different families and uterine myoma was discovered in one case from Hunan Province.

Imaging findings

There were 195 patients (92.4%) with multiple pulmonary cystic lesions on chest CT (Table 3). The detailed locations of lung cysts were described in 75 cases; the cystic lesions were located bilaterally in 52 cases (69.3%), and most of them in the lower lung zones. Most of the cysts ranged 0.5-6cm in diameter; the largest cystic lesion measured 8.5cm.

Table 3
Comparison of characteristics among patients with BHDS between China, Japan and USA

Characteristic	China#	Japan ⁺	USA ^{\$}	P value
Pneumothorax (n, %)	157(71.0%)	230(73.7%)	79(76.0%)	0.616
Cysts on chest CT (n, %)	195(92.4%)	309(99.0%)*	88(85.0%)	≪0.01
Skin lesion (n, %)	40(18.1%)	76(48.7%)*	74(71.0%)*	≪0.01
Renal tumor (n, %)	8(3.6%)	40(34.8%)*	34(33.0%)*	≪0.01
*P≪0.05, #Current study; +data from Clin Genet 2016,90: 403–412; \$data from Ann Am Thorac Soc 2017,14 (5): 706–713				

FLCN mutations

All 221 patients were diagnosed by genetic testing and the results in detail were available for 144 of them. The mutation in exon was found in 134 patients, including almost all exons 4–14. The most frequent mutation (17.4%, 25/144) was the single deletion, duplication of cytosine in codon 1285 of exon 11 (c.1285del, c.1285dup), follow by the mutation of c.1579_1580ins in exon 14 (10.4%,15/144), as was reported in previous studies, and a three base pair deletion in exon 6 (c.924_926del) (9.0%,13/144) was the third most common mutation Other mutations occurred much less frequently. Mutation c.924_926del was associated with a 39% risk (5 of 13 patients) for pneumothorax, which increased to 60% for mutation c.1285dup (9 of 15 patients) and 73% for mutation c. 1579_1580ins (11 of 15 patients). However, the differences in the risk of pneumothorax associated with these three most common mutations was not statistically significant (patients must have experienced at least one pneumothorax episode) (χ^2 , 3.513; P value, 0.173). The mutant gene in 6 patients with fibrofolliculomas included exon 11, 9, 13; 8 renal carcinoma patients manifested mutations in exon 7, 9, 10, 11, 13, 14.

Diagnosis

All 221 patients were confirmed to have BHDS by detection of pathogenic variant in the *FLCN* gene. Detailed information was available for 32 families that were initially misdiagnosed. Almost all initial presentations involved pneumothorax, which was often misdiagnosed as primary spontaneous pneumothorax. One case reported from Fujian Province was misdiagnosed as lymphangioleiomyomatosis (LAM) and had been on rapamycin (sirolimus) treatment for several weeks. The average duration of misdiagnosis was 9.6 ± 10.8 years (range from 0 to 36 years) prior to the eventual diagnosis of BHDS.

Treatment

The detailed information regarding management of pneumothorax was available in 57 patients. Pneumothorax occurred in 37 patients with BHDS who received interventional treatment. Among them, 29 patients with pneumothorax were treated by bullectomy with pleurodesis; pneumothorax recurred in three cases. The remaining of pneumothorax were treated by pleurodesis alone and the other 5 by bullectomy.

Pneumothorax recurred in one patient after pleurodesis. The recurrence rate of pneumothorax after conservative treatment (including tube thoracostomy) was 29/41 (71%) while the pneumothorax recurred after surgical treatment (pulmonary bullectomy or pleurodesis) in only 4/37 (11%). Partial nephrectomies were performed in all renal cancer patients but the additional following-up was lacking in all cases.

Prognosis

The follow-up information for patients with BHDS in China were rarely described in the published reports. Among 43 patients with recurrent pneumothoraces, ipsilateral and contralateral recurrences were described in 67% and 33%, respectively.

Only one patient in Beijing was followed up after open partial nephrectomy and there was no new abnormality in the kidneys at three months.

Discussion

To the best of our knowledge, this study comprises the largest number of families of BHDS in the Chinese population. We found most BHDS families were reported from Beijing and Jiangsu Province, and over 80% of them were reported in only the past five years. In comparison to USA and Japanese BHDS cohorts, the dominant manifestations of Chinese patients were pulmonary cysts associated with recurrent pneumothorax, while skin lesions and renal tumors were less prevalent. Delayed diagnosis and suboptimal medical management presented common challenges for Chinese patients with BHDS.

BHDS is a rare inherited autosomal dominant disorder caused by germline mutations in the tumor suppressor gene *FLCN* that predisposes affected individuals to develop benign skin tumors (fibrofolliculomas), renal neoplasms, and pulmonary cysts with a risk of spontaneous pneumothorax. Since a research team from Nanjing University firstly explored the *FLCN* gene mutations on patients with spontaneous pneumothorax history in 2008, there has been more than 20 families with BHDS in China up to 2017 according to the BHD foundation statistics. Our study included 221 cases from 120 families that were reported between January 2008 and December 2020 in China. More than 80% of families were described in the past five years, half of them in the past two years. The increasing number of BHDS in China likely reflects the increasing awareness and recognition rather than increasing incidence of disorder. In recent decades, a fascinating progress has been achieved with respect to the diagnosis and management of rare diseases in China. A registry study on rare respiratory diseases including another diffuse cystic lung disease, LAM, has been ongoing nation-wide since 2016. Therefore, BHDS is also being increasingly diagnosed by clinicians in China. Nonetheless, over two-thirds of BHDS families were managed in Beijing. This huge imbalance of diagnosis and management on patients with rare diseases between Beijing and the rest of China suggests urgent need for improvements in the recognition of rare diseases nation-wide.

As in prior reports, respiratory system was the most frequently affected in Chinese patients with BHDS with 92.4% of our cohort manifesting lung cysts (Table 3). A study of 312 Japanese BHDS patients

reported that almost all patients had lung cysts⁷. Similarly, a study of Korean BHDS patients reported all cases had lung cysts²⁹. On the contrary, it was reported that the prevalence of lung cysts of BHDS in USA was 85%, which was lower compared to the Japanese cohort^{7,30}. This might indicate that Asian patients were more prone to pulmonary cysts than Caucasian patients. However, most of cases with BHDS in our study was diagnosed by respiratory physicians which may have introduced a bias. The widespread use of high-resolution computed tomography in recent decades has provided better detection of cystic lung diseases. As a result, cystic lung diseases such as BHDS have been better recognized and distinguished from emphysema.

Due to the high prevalence of lung cysts in adult patients with BHDS, spontaneous pneumothorax has been found to be a common presentation³. In our study, 71% of confirmed patients had at least one episode of pneumothorax and nearly 85% of them had a family history of pneumothorax. This was consistent with the features of patients with BHDS reported from other Asian countries²⁶. A recent survey of 104 American patients with BHDS reported a similar rate of spontaneous pneumothorax during their lifetime³⁰. BHDS-related pneumothorax seems to have a higher recurrence rate after conservative therapy (observation, chest drainage and chemical pleurodesis) compared to surgical management. Liu et al reported only 9% of patients had recurrence of pneumothorax after surgical treatment (pulmonary bullectomy and/or pleurodesis) while 52% of recurrence rate was observed after conservative treatment²⁴. Gupta et al reported pleurodesis (chemical or surgical) to reduce the ipsilateral recurrence rate by half (33% versus 63%)³⁰. Another diffuse cystic lung disease LAM is also associated with high recurrence rate of pneumothorax. In 2017, American Thoracic Society/Japanese Respiratory Society Guideline suggested that patients with LAM be offered ipsilateral pleurodesis after their initial pneumothorax rather than waiting for a recurrent pneumothorax before intervening with a pleural symphysis procedure³¹.

Renal cell cancer has been found to be the main manifestation associated with the poor prognosis of BHDS. Zbar et al found a 7-fold increase in the risk of renal tumors for BHD-affected family members when adjusted for age³². Subsequent study revealed that the prevalence of renal tumors in patients with BHD was 34% with a mean age at diagnosis of approximately 50 years³³. Chromophobe renal carcinoma was the most common histological type of renal cancer, followed by hybrid oncocytic/chromophobe tumor^{3,7}. Similar findings were reported in a recent largest study including 312 Japanese cases from 120 different families⁷. Renal carcinoma was detected in 8 (3.6%) patients in this study while that in Japan was 34.8% and 25% in Korea, respectively^{7,29}. The differences between these countries might attribute to less awareness and evaluation for renal manifestations of BHDS among Chinese physicians or ethnic differences; further studies are needed on this issue. In addition, lack of follow-up evaluations might have contributed to the low prevalence of renal cancer in the Chinese cohort. Thus, a long-term prospective observation study is needed to explore the true incidence of renal tumors among BHDS-affected families in the Chinese population.

Skin lesions were a common feature of BHDS in Caucasians, in whom it is found in about 80%~90%^{3,34}. Characteristic lesions fibrofolliculomas and trichodiscomas are important clues in recognizing BHDS, but the patients in Asian countries seem to have lower prevalence of skin lesions^{3,7,29}. Previous study demonstrated skin lesions developed in less than 30% in Chinese patients²⁶, similar to the current study. Only 6 patients were histologically diagnosed to have fibrofolliculomas. However, this probably represents an underestimation due to limited diagnostic capacity of this rare disease, especially among general pulmonologists. The incidence of skin lesions among patients with BHDS reported from Peking Union Medical College Hospital, a national referral center for rare respiratory diseases, significantly increased from 11% (3/27) in 2017 to 47.2% (17/36) in 2019^{9,21}. The awareness of characteristic skin manifestations was likely an important factor in the increased diagnosis rate over the interval. Similarly, dermatologist Chikako Iwabuchi and colleagues reported the increasing diagnosis rates of skin lesions in 31 Japanese individuals³⁵. Skin lesions were recognized in over 80% of BHDS patients, including almost 75% of them histologically confirmed by skin biopsies³⁵. Therefore, skin lesions appear to be more prevalent than previously reported in east Asian population, which would be warrant further assessment in clinical practice through cooperation with an experienced multidisciplinary team.

The *FLCN* mutations in exon 11 were found to be the most frequently detected among Caucasian population, mainly c.1285dup/del and c.1300G > C³⁶. Our study also showed that the most frequent mutation in Chinese BHD patients was the single deletion, duplication of cytosine in codon 1285 of exon 11. The correlation between genotype and phenotype has been explored in the past decades. There was rather high incidence of pneumothorax associate with *FLCN* hotspot mutation c.1285dup (60% risk) in this cohort. However, a cohort from Germany reported less than 50% of 197 patients with BHDS had pneumothorax. Significantly increased pneumothorax risks were observed for mutations c.1300G > C (59%) and c.250-2A > G (77%) compared with *FLCN* hotspot mutation c.1285dup (37% risk)³⁶. We are not able to explain the discrepant rates of pneumothorax associated with the mutation c.1285dup in Chinese patients compared to the Germany cohort. It would need further research on large-scale population to explore the correlation between these mutation and phenotype.

To date, there is no effective treatment for BHDS, but the benefits of an early diagnosis include appropriate screening for renal tumors among BHDS individuals, patient education and family counseling. An average delay from first symptom to diagnosis was up to 13 years in 104 patients reported from USA³⁰. The misdiagnosis was quite common. Previous results have demonstrated lower prevalence of skin lesions and renal tumors among of BHDS patients in China^{9,26}. Thus, the presenting manifestation of pneumothorax associated with characteristic lung cysts are important clues to achieve an early diagnosis. A study in Japan established a scoring system that has a high degree of sensitivity and specificity for distinguishing suspected BHDS from primary spontaneous pneumothorax³⁷. Several radiological studies also found appropriate evaluation of chest CT images to be helpful in differentiating BHDS from other diffuse cystic lung diseases³⁸⁻⁴². Therefore, the most common presentation pneumothorax in setting of characteristic cystic lung lesions or a family history of pneumothorax should suggest the need for *FLCN* gene testing to make timely diagnosis.

There were several limitations of this study. Firstly, this was a retrospective study, and majority of reported cases were from tertiary referral centers in Beijing. Secondly, some cases lacked data regarding the *FLCN* gene mutations as well as aspects of management and follow-up. However, this study collected all the consecutively cases reported in the Chinese population in recent decades and fully described the real-life diagnosis and managements of BHDS which should be valuable for better understanding this rare disease in the coming years.

Conclusion

Although BHDS has been increasingly reported in the recent years in China, only minority of the cases were managed in the regions beyond Beijing and Jiangsu Province. The dominant clinical manifestation was pulmonary cysts associated with recurrent pneumothorax, while skin lesions and renal tumors were less prevalent. Lack of disease awareness, delayed diagnosis along with suboptimal management present critical challenges for Chinese BHDS patients.

Declarations

Acknowledgments

We thank Prof. J.H. Ryu, Mayo Clinic Rochester, USA, for the unique contribution he gave revising this manuscript. We also want to thank Prof. Yaping Liu, Chinese Academy of Medical Sciences, for her contribution in genetic analysis of *FLCN* mutations.

Author contributions

Guofeng Zhang and Xiaowen Hu conceived the idea, designed, and supervised the study, drafted the manuscript, and had full access to all the data and took responsibility for the integrity of the data. Xianmeng Chen analyzed data and performed statistical analysis. All the authors reviewed and approved the final version of the manuscript.

Funding

The National Key Research and Development Program of China (2016YFC0901502).

Availability of data and materials

The datasets generated and analyzed for this study are not publicly available due to participant privacy but are available from the corresponding author upon reasonable request.

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing interest

The authors declare that they have no conflict of interest.

Author details

¹ Department of Pulmonary and Critical Care Medicine, the First Affiliated Hospital of USTC, Division of Life Sciences and Medicine, University of Science and Technology of China, Hefei, Anhui, China.² WanNan Medical College, Wuhu, Anhui, China.³ Department of Pulmonary and Critical Care Medicine, Peking Union Medical College Hospital, Beijing, China

References

1. Birt AR, Hogg GR, Dube WJ. Hereditary multiple fibrofolliculomas with trichodiscomas and acrochordons. *Arch Dermatol*. 1977;113(12):1674-7.
2. Nickerson ML, Warren MB, Toro JR, Matrosova V, Glenn G, Turner ML, et al. Mutations in a novel gene lead to kidney tumors, lung wall defects, and benign tumors of the hair follicle in patients with the Birt-Hogg-Dube syndrome. *Cancer Cell*. 2002;2(2):157-64.
3. Daccord C, Good JM, Morren MA, Bonny O, Hohl D, Lazor R. Birt-Hogg-Dube syndrome. *Eur Respir Rev*. 2020;29(157).
4. Kennedy JC, Khabibullin D, Boku Y, Shi W, Henske EP. New Developments in the Pathogenesis of Pulmonary Cysts in Birt-Hogg-Dube Syndrome. *Semin Respir Crit Care Med*. 2020;41(2):247-55.
5. Schmidt LS, Linehan WM. FLCN: The causative gene for Birt-Hogg-Dube syndrome. *Gene*. 2018;640:28-42.
6. Gupta N, Sunwoo BY, Kotloff RM. Birt-Hogg-Dube Syndrome. *Clin Chest Med*. 2016;37(3):475-86.
7. Furuya M, Yao M, Tanaka R, Nagashima Y, Kuroda N, Hasumi H, et al. Genetic, epidemiologic and clinicopathologic studies of Japanese Asian patients with Birt-Hogg-Dube syndrome. *Clin Genet*. 2016;90(5):403-12.
8. Ren HZ, Zhu CC, Yang C, Chen SL, Xie J, Hou YY, et al. Mutation analysis of the FLCN gene in Chinese patients with sporadic and familial isolated primary spontaneous pneumothorax. *Clin Genet*. 2008;74(2):178-83.
9. Liu Y, Xu Z, Feng R, Zhan Y, Wang J, Li G, et al. Clinical and genetic characteristics of chinese patients with Birt-Hogg-Dubé syndrome. *Orphanet Journal of Rare Diseases*. 2017;12(1):104.
10. Menko FH, van Steensel MA, Giraud S, Friis-Hansen L, Richard S, Ungari S, et al. Birt-Hogg-Dube syndrome: diagnosis and management. *Lancet Oncol*. 2009;10(12):1199-206.
11. Lin Z, Gong K, Pang B, Zeng C, Zhang D. Birt-Hogg-Dube syndrome with clear cell renal cell carcinoma in a Chinese family. *Intern Med*. 2014;53(24):2825-8.

12. Liu Z, Xu KF, Hu C, Chen RR, Duan J, Shi YZ, et al. Use of whole-exome sequencing for the diagnosis of atypical Birt-Hogg-Dube syndrome. *J Genet Genomics*. 2014;41(8):449-51.
13. Ding Y, Zhu C, Zou W, Ma DH, Min HY, Chen BF, et al. FLCN intragenic deletions in Chinese familial primary spontaneous pneumothorax. *Am J Med Genet A*. 2015;167A(5):1125-33.
14. Dong L, Gao M, Hao WJ, Zheng XQ, Li YG, Li XL, et al. Case Report of Birt-Hogg-Dube Syndrome: Germline Mutations of FLCN Detected in Patients With Renal Cancer and Thyroid Cancer. *Medicine*. 2016;95(22):e3695.
15. Zhu JF, Shen XQ, Zhu F, Tian L. Novel folliculin (FLCN) mutation and familial spontaneous pneumothorax. *QJM*. 2017;110(1):23-6.
16. Liu L, Yang K, Wang X, Shi Z, Yang Y, Yuan Y, et al. Detection of *Folliculin* Gene Mutations in Two Chinese Families with Birt-Hogg-Dube Syndrome. *Biomed Res Int*. 2017;2017:8751384.
17. Li T, Ning X, He Q, Gong K. Novel germline mutations in FLCN gene identified in two Chinese patients with Birt-Hogg-Dube syndrome. *Chin J Cancer*. 2017;36(1):4.
18. Hao S, Long F, Sun F, Liu T, Li D, Jiang S. Birt-Hogg-Dube syndrome: a literature review and case study of a Chinese woman presenting a novel FLCN mutation. *BMC Pulm Med*. 2017;17(1):43.
19. Hou X, Zhou Y, Peng Y, Qiu R, Xia K, Tang B, et al. Birt-Hogg-Dube syndrome in two Chinese families with mutations in the FLCN gene. *BMC Med Genet*. 2018;19(1):14.
20. Xie F, Mao QZ. A Case of Birt-Hogg-Dube syndrome with renal hybrid oncocytic/chromophobe tumors. *Medical Journal of Peking Union Medical College Hospital*. 2019;10(2):162-5.
21. Liu K, Xu W, Tian X, Xiao M, Zhao X, Zhang Q, et al. Genotypic characteristics of Chinese patients with BHD syndrome and functional analysis of FLCN variants. *Orphanet J Rare Dis*. 2019;14(1):223.
22. Hu Q, Lai GX, Xu LY, Xu C, Qu LJ, Zhang L. Recurrent pneumothorax with diffuse lung cysts. *Chin J Tuberc Respir Dis*. 2019;42(6):464-7.
23. Zheng CM, Hu XX, Gao YL, Miao JB, Li H. Recurrent primary spontaneous pneumothorax in a large Chinese family: a clinical and genetic investigation. *Chin Med J*. 2019;132(20):2402-7.
24. Liu Y, Xing H, Huang Y, Meng S, Wang J. Familial spontaneous pneumothorax: importance of screening for Birt-Hogg-Dube syndrome. *Eur J Cardiothorac Surg*. 2020;57(1):39-45.
25. Zong D, Li J, Liu X, Guo T, Ouyang R. Identification of a Novel Pathogenic Folliculin Variant in a Chinese Family With Birt-Hogg-Dube Syndrome (Hornstein-Knickenberg Syndrome). *Front Genet*. 2020;11:565566.
26. Guo T, Shen Q, Ouyang R, Song M, Zong D, Shi Z, et al. The clinical characteristics of East Asian patients with Birt-Hogg-Dube syndrome. *Ann Transl Med*. 2020;8(21):1436.
27. So SY. Spontaneous pneumothorax due to Birt-Hogg-Dube syndrome in a Chinese family. *Respirology*. 2009;14(5):775-6.
28. Yang GG, Chiang CC. Recurrent spontaneous pneumothoraces. *Am J Respir Crit Care Med*. 2013;188(1):e1-2.

29. Lee JH, Jeon MJ, Song JS, Chae EJ, Choi JH, Kin GH, et al. Birt-Hogg-Dube syndrome in Korean: clinicoradiologic features and long term follow-up. *Korean J Intern Med.* 2019;34(4):830-40.
30. Gupta N, Koprass EJ, Henske EP, James LE, El-Chemaly S, Veeraraghavan S, et al. Spontaneous Pneumothoraces in Patients with Birt-Hogg-Dube Syndrome. *Ann Am Thorac Soc.* 2017;14(5):706-13.
31. Gupta N, Finlay GA, Kotloff RM, Strange C, Wilson KC, Young LR, et al. Lymphangioliomyomatosis Diagnosis and Management: High-Resolution Chest Computed Tomography, Transbronchial Lung Biopsy, and Pleural Disease Management. An Official American Thoracic Society/Japanese Respiratory Society Clinical Practice Guideline. *Am J Respir Crit Care Med.* 2017;196(10):1337-48.
32. Zbar B, Alvord WG, Glenn G, Turner M, Pavlovich CP, Schmidt L, et al. Risk of renal and colonic neoplasms and spontaneous pneumothorax in the Birt-Hogg-Dube syndrome. *Cancer Epidemiol Biomarkers Prev.* 2002;11(4):393-400.
33. Toro JR, Wei MH, Glenn GM, Weinreich M, Toure O, Vocke C, et al. BHD mutations, clinical and molecular genetic investigations of Birt-Hogg-Dube syndrome: a new series of 50 families and a review of published reports. *J Med Genet.* 2008;45(6):321-31.
34. Steinlein OK, Ertl-Wagner B, Ruzicka T, Sattler EC. Birt-Hogg-Dube syndrome: an underdiagnosed genetic tumor syndrome. *J Dtsch Dermatol Ges.* 2018;16(3):278-83.
35. Iwabuchi C, Ebana H, Ishiko A, Negishi A, Mizobuchi T, Kumasaka T, et al. Skin lesions of Birt-Hogg-Dube syndrome: Clinical and histopathological findings in 31 Japanese patients who presented with pneumothorax and/or multiple lung cysts. *J Dermatol Sci.* 2018;89(1):77-84.
36. Sattler EC, Syunyaeva Z, Mansmann U, Steinlein OK. Genetic Risk Factors for Spontaneous Pneumothorax in Birt-Hogg-Dube Syndrome. *Chest.* 2020;157(5):1199-206.
37. Ebana H, Mizobuchi T, Kurihara M, Kobayashi E, Haga T, Okamoto S, et al. Novel clinical scoring system to identify patients with pneumothorax with suspicion for Birt-Hogg-Dube syndrome. *Respirology.* 2018;23(4):414-8.
38. Xu W, Xu Z, Liu Y, Zhan Y, Sui X, Feng R, al. Characterization of CT scans of patients with Birt-Hogg-Dube syndrome compared with those of Chinese patients with non-BHD diffuse cyst lung diseases. *Orphanet J Rare Dis.* 2020;15(1):176.
39. Park HJ, Park CH, Lee SE, Lee GD, Byun MK, Lee S, et al. Birt-Hogg-Dube syndrome prospectively detected by review of chest computed tomography scans. *Plos One.* 2017;12(2):e0170713.
40. Lee JE, Cha YK, Kim JS, Choi JH. Birt-Hogg-Dube syndrome: characteristic CT findings differentiating it from other diffuse cystic lung diseases. *Diagn Interv Radiol.* 2017;23(5):354-9.
41. Lee E, Sayyoub M, Haggerty JE, Kazerooni E, Agarwal PP. Role of Radiologists in the Diagnosis of Unsuspected Birt-Hogg-Dube Syndrome in a Tertiary Clinical Practice. *AJR Am J Roentgenol.* 2019;213(4):792-7.
42. Escalon JG, Richards JC, Koelsch T, Downey GP, Lynch DA. Isolated Cystic Lung Disease: An Algorithmic Approach to Distinguishing Birt-Hogg-Dube Syndrome, Lymphangioliomyomatosis, and Lymphocytic Interstitial Pneumonia. *AJR Am J Roentgenol.* 2019:1-5.

Figures




Figure 1

The flow diagram of this study Abbreviations: CAJ = China Academic Journals Full-text Database (CAJ), Wanfang = Wanfang Chinese Database.




Figure 2

Number of families and cases of BHDS in China distributed by year of report