

Oxalate nephropathy in an elderly patient with newly diagnosed celiac disease – a case report

Hendrik Wietse Zijlstra (✉ rik.zijlstra@gmail.com)

University Medical Center Groningen

Coen A. Stegeman

University Medical Center Groningen

Case Report

Keywords: Oxalate, nephropathy, celiac disease

Posted Date: February 6th, 2023

DOI: <https://doi.org/10.21203/rs.3.rs-2499613/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Additional Declarations: No competing interests reported.

Version of Record: A version of this preprint was published at BMC Nephrology on June 27th, 2023. See the published version at <https://doi.org/10.1186/s12882-023-03241-y>.

Abstract

Oxalate nephropathy has widely been described in gastrointestinal disease,. This is only the second publication of oxalate nephropathy in newly diagnosed celiac disease.

A 72-year-old Caucasian male presented to the hospital with abdominal discomfort and acute renal insufficiency with a creatinine of 290 $\mu\text{mol/L}$.

The clinical course, laboratory results and urinalysis was suspect for tubular injury. Renal biopsy showed calcium oxalate depositions. Elevated serum and urine oxalate levels established the diagnosis oxalate nephropathy.

The abdominal complaints, with steatorrhea and positive anti-tissue transglutaminase antibodies were diagnosed as celiac disease, which was confirmed after duodenal biopsies.

Treatment with prednisone, gluten-free, low oxalate and normal calcium diet, lowered the serum oxalate levels and improved his renal function.

Decreased absorption of free fatty acids can lead to increased free oxalate in the colon, due to the binding of free fatty acids to calcium, preventing the formation of the less absorbable calcium oxalate in the colon.

Oxalate dispositions in the kidney can lead to acute tubular injury and chronic renal insufficiency. Celiac disease is therefore one of the intestinal diseases that can lead to hyperoxaluria and oxalate nephropathy.

Introduction

Oxalate nephropathy is an uncommon cause of renal failure. The deposition of oxalate crystals in the tubules and interstitium can result in tubular injury and loss of kidney function. Hyperoxaluria can be primary, due to congenital defects in metabolism of glyoxylate and oxalate [1] or secondary caused by increased absorption, increased intake or decreased intestinal oxalate degradation [2]. Due to increased oxalate uptake in the digestive system, different intestinal diseases are known risk factors for development of nephrolithiasis and nephrocalcinosis.

We present a case of oxalate nephropathy caused by secondary hyperoxaluria in a 72 years old patient with newly diagnosed celiac disease, without nephrolithiasis.

Case Report

A 72-year-old Caucasian male, with a past medical history of eczema, for which he was recently treated with methotrexate, presented with a 3 month history of abdominal discomfort and diarrhea with 5 kg weight loss and fatigue. Colonoscopy in a different medical center did not show any pathology in colon

and distal ileum. He had discontinued the methotrexate he used for eczema three months ago because of these abdominal complaints, without clinical improvement. At presentation in our hospital he did not have a fever (99.3 °F (37,4 °C)), mild hypertension (167/83 mmHg) and a normal heart rate (62/min).

He did not use any medication at the time of presentation.

Laboratory tests showed acute renal failure with a creatinine of 290 µmol/L with a normal anion gap metabolic acidosis and no proteinuria. Five months prior to presentation his creatinine was 102 µmol/L. C-reactive peptide was 19 mg/L, with leukocytes of $7.2 \times 10^9/L$. The urine dipstick was negative for erythrocyturia or leucocyturia. Renal ultrasound did not show evidence of hydronephrosis.

He was admitted for further evaluation of renal impairment and abdominal complaints.

Fluid resuscitation did not improve the renal function, which made dehydration due to diarrhea as a cause of renal impairment unlikely. Microscopic urinalysis showed leucocyturia of approximately 10 leukocytes per high-power field, with possible leucocyte casts, and no evidence of glomerular erythrocyturia, consistent with tubular injury. Oxalate crystals were not seen. Remarkably, the initial screening with urine dipstick seems to be falsely negative for leucocyturia.

A renal biopsy was done because the renal function did not improve after rehydration and normalization of defecation pattern.

3 of 13 glomeruli were globally sclerosed and did not show any abnormalities otherwise. The tubulointerstitium contained multiple zones with atrophy, and fibrosis was present in approximately 20% of the cortical area. Various tubules contained calcium oxalate crystals and showed patchy tubulointerstitial nephritis with subtle lymphocytic infiltrates (Fig. 1).

Because tubulointerstitial nephritis (TIN), without known causing agent, likely contributed to the renal failure prednisone 30 mg once daily was initiated after the renal biopsy with results pending.

Meanwhile, the abdominal pain was found to be primarily after ingestion of gluten-rich diets, the anti-tissue transglutaminase antibodies were positive (42 U/ml) and there was documented steatorrhea (13.6 grams of total fecal fat per day). It was therefore thought to be caused by celiac disease. He was treated with a gluten-free diet, with improvement of the abdominal pain and normalization of fecal pattern.

In the outpatient setting the diagnosis was confirmed after duodenal biopsies showed partial villous atrophy and increased intraepithelial lymphocytes, consistent with celiac disease (Marsh 3a).

An increased serum oxalate (2.54 mg/L [N 0.0-0.45 mg/L]) and urinary oxalate excretion (1,23 mmol/24 hr [N 0,07–0,72 mmol/24 hr] and 117,23 mmol/mol creatinine [N 8,30–49,00 mmol/mol creatinine], with low glycolic acid and glyceric acid) were evidence for secondary hyperoxaluria (Table 1). The combination of calcium oxalate crystals, secondary hyperoxaluria and newly diagnosed celiac disease made oxalate nephropathy in celiac disease the most likely diagnosis. It was treated with a gluten-free,

low oxalate and normal calcium diet and high-volume intake. Because of the initial improvement the prednisone was continued.

Table 1
clinical renal function and oxalate levels at presentation and during outpatient follow-up

	At presentation	8 weeks after presentation	4 months after presentation	6 months after presentation
Serum creatinine (µmol/L)	290	152	178	179
Creatinine clearance (ml/min)	35	50	52	48
Serum oxalate (mg/L)	2.54	0.79	0.62	0.82
24h-urine oxalate (mmol/mol creatinine)	117.23	67.10	41.97	51.99
24h-urine oxalate (mmol/24hr)	1.23	0.83	0.56	0.63

There was a remarkable improvement of renal function, as well as oxalate levels in serum and urinary oxalate excretion. The prednisone was discontinued. The patient's renal function did improve, yet stabilized at a creatinine of 150 µmol/L (Table 1).

Discussion

First presentation of celiac disease in elderly people is more common than usually expected. Patel et al. showed that fifteen percent of the patients with newly diagnosed celiac were older than 65 years old[3]. The prevalence in a Finnish study showed that approximately 2% of patients between 52 and 74 have biopsy proven celiac disease, which is higher than in the general population[4].

There are few case reports describing methotrexate induced sprue-like syndrome with villous atrophy after initiation of methotrexate.[5] Because in this patient the anti-tissue transglutaminase antibodies were positive and because there was no mucosal healing four months after discontinuation of methotrexate, this was unlikely.

Even though intestinal diseases, like inflammatory bowel disease [6, 7] and bariatric surgery [8] are known risk factors for urinary stone disease and nephrocalcinosis, and even though celiac disease has been associated with an increased risk of nephrolithiasis, due to hyperoxaluria, this is only the second case report of acute renal failure caused by oxalate nephropathy in new onset celiac disease in a patient with previously normal renal function and without nephrolithiasis.

Kohler et al. described a case of new onset celiac disease resulting in oxalate nephropathy[9] and Capolongo et al. described oxalate nephropathy in a patient with subclinical celiac disease, developing after a kidney transplantation[10].

Intestinal diseases with fat malabsorption (e.g. pancreatic insufficiency [11], small bowel resection [8] or inflammatory bowel disease [6, 7]) can lead to increased serum oxalate levels[12]. Fat malabsorption leads to an increase of free fatty acids in the intestinal lumen. The free fatty acids bind to calcium, reducing the availability of calcium to bind to dietary oxalate. In normal functionality of the intestines, the intestinal absorption of calcium oxalate is less efficient than absorption of free oxalate. The increased concentration of free oxalate in the colon therefore leads to higher serum concentration. Additionally, free fatty acids and bile salts also increase the permeability of the colon for oxalate, contributing to increased oxalate absorption[13].

Increased serum oxalate concentration can lead to nephrolithiasis with calcium oxalate stones or the oxalate crystals can precipitate in the renal interstitium and tubules leading to tubular injury and interstitial inflammation, causing oxalate nephropathy.

Lumlertgul et al.[2] described in their review the different biopsy findings that can be found. They found that 71% (95% CI: 44–89) had acute tubular injury. Tubular damage and atrophy were found in 69% (95% CI: 43–87). 72% (95% CI 45.0–89.0) showed interstitial mononuclear cell infiltration, and glomerular changes were described in 59% (95% CI: 40–76) of which mostly mesangial cellular proliferation.

There are no randomized controlled trials on therapy for oxalate nephropathy. High volume intake, normal calcium and low oxalate diet, and if possible, treatment for the underlying cause of increased oxalate absorption are important strategies.

Oxalate nephropathy in general has a poor prognosis with studies showing 58% of patients requiring dialysis and only partial, if any recovery of renal function[2].

In this case the fast improvement of renal function can not be attributed to dissolving of oxalate crystals in the kidney. Most likely the oxalate crystals caused interstitial inflammation, which contributed to the decline in kidney function. This could explain the positive effect of prednisone and the initial increase in renal function. Incomplete recovery may be due to the remaining oxalate crystal deposits in the kidney. Possibly it is comparable to intratubular casts in multiple myeloma, which also results in interstitial inflammation with diffuse fibrosis and tubular atrophy, in which the casts dissolve in weeks.[14]

Conclusion

Oxalate nephropathy is a uncommon cause of acute renal failure. In various intestinal diseases, fat malabsorption can result in hyperoxaluria, leading to calcium oxalate depositions in the kidneys, causing renal insufficiency. Even though there are numerous reports on oxalate nephropathy in intestinal disease, this is only the second report on new onset celiac disease as a cause of oxalate nephropathy.

Because celiac disease is more common later in life than what is usually assumed, in patients in any age group with a combination of known celiac disease or clinical suspicion, and acute renal failure, physicians should be aware of oxalate nephropathy secondary to celiac disease.

Abbreviations

TIN
tubulointerstitial nephritis

Declarations

Ethics approval and consent to participate

Not applicable

Consent for publication

Informed consent obtained from the patient.

Availability of data and materials

Not applicable

Competing interests

There we no competing interest

Funding

Not applicable

Authors' contributions

HWZ writing of the essay; CAS supervision and mentorship. Each author contributed important intellectual content during manuscript drafting or revision, accepts personal accountability for the author's own contributions, and agrees to ensure that questions pertaining to the accuracy or integrity of any portion of the work are appropriately investigated and resolved.

Acknowledgements

Not applicable

Authors' information (optional)

Not applicable

References

1. Harambat J, Fargue S, Bacchetta J, Acquaviva C, Cochat P. Primary Hyperoxaluria. *International Journal of Nephrology*. 2011; doi:10.4061/2011/864580
2. Lumlertgul N, Siribamrungwong M, Jaber BL, Susantitaphong P. Secondary Oxalate Nephropathy: A Systematic Review. *Kidney Int Rep*. 2018 Nov; 3(6): 1363–1372.
3. Patel D, Kalkat P, Baisch D, Zipser R. Celiac Disease in the Elderly. *Gerontology*. 2005;51:213–214
4. Vilppula A, Collin, P, Mäki, M, et al. Undetected coeliac disease in the elderly. A biopsy-proven population-based study. *Digestive and Liver Disease*. 2008;40:809–813
5. Boscá M, Añón R, Mayordomo E, et al. Methotrexate induced sprue-like syndrome. *World J of Gastroenterology*. 2008;14(45):7009–7011.
6. Mandell I, Krauss E, Millan JC. Oxalate-induced acute renal failure in Crohn's disease. *Am J Med*. 1980;69:628–632.
7. Gelzayd EA, Breier RI, Kirsner JB. Nephrolithiasis in inflammatory bowel disease. *Am J Dig Dis*. 1968;13:1027–1034.
8. Asplin JR, Coe FL. Hyperoxaluria in kidney stone formers treated with modern bariatric surgery. *J Urol*. 2007;177(2):565.
9. Kohler GD, Gaut JP, Morrison AR. Diarrhea, weight loss, electrolyte abnormalities, and renal failure. *Kidney International*. 2015; 88:421–422.
10. Capolongo G, Abul-Ezz S, Moe OW, Sakhaee K. Subclinical Celiac Disease and Crystal-Induced Kidney Disease Following Kidney Transplant. *Am J Kidney Dis*. 2012; 60(4): 662–667.
11. Demoulin N, Issa Z, Crott R, et al. Enteric hyperoxaluria in chronic pancreatitis. *Medicine (Baltimore)*. 2017;96:e6758.
12. Nazzal L, Puri S, Goldfarb DS. Enteric hyperoxaluria: an important cause of end-stage kidney disease. *Nephrol Dial Transplant*. 2016;31(3):375–382.
13. Dobbins JW, Binder HJ. Effect of bile salts and fatty acids on the colonic absorption of oxalate. *Gastroenterology*. 1976;70:1096–1100.
14. Basnayake K, Hutchison C, Kamel D *et al*. Resolution of cast nephropathy following free light chain removal by haemodialysis in a patient with multiple myeloma: a case report. *J Med Case Reports*. 2008;doi:10.1186/1752-1947-2-380

Figures




Figure 1

Renal biopsy findings containing calcium oxalate crystals in the tubules, with crystals being birefringent under polarized light.