

Dysnatremia in Covid -19 Disease Characteristics of Patients, Clinical Manifestation and Outcome in Pakistan.

Faiza Saeed (✉ faiza_saeed_1@hotmail.com)

The Indus Hospital

Ashar Alam

The Indus Hospital

Shoukat Memon

The Indus Hospital

Javeria Chughtai

The Indus Hospital

Shahzad Ahmed

The Indus Hospital

Sobia Tariq

The Indus Hospital

Beena Salman

Kidney Centre

Salman Imtiaz

The Indus Hospital

Research Article

Keywords: Dysnatremias, Hyponatremia, Hypernatremia, covid 19, Pakistan. Electrolytes imbalance

Posted Date: March 2nd, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-251233/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background.

Imbalance of serum sodium in the form of hypo and hypernatremia has bad prognostic impact, and linked with increase morbidity and mortality. Covid 19 is a deadly viral infection can be complicated with either of the two state. This increases the complications associated with the covid 19 infection.

Material and method.

This is the observational cohort study, conducted at The Indus Hospital Karachi, from March 2020 to September 2020. All patients were divided in to three groups on the basis of their serum sodium level, and distribution of variables was observed in all these three groups. To observe the effect of dysnatremias on outcome, binary logistic regression was run and odds ratios with 95% confidence interval were obtained.

Results.

We included 655 patients, in which 70.7% were male, while 29.3% were female. Mean age was 54 ± 15.5 with minimum of 1 year and maximum of 95 year. Hyponatremia was more common 154(23.5%) than hypernatremia 79(12.1%) patients. Dysnatremia were common in patients of age group of 51 – 65 years (Hyponatremia 72(46.8%) hypernatremia 36(45.6%). Hyponatremia didn't show any significant effect on the clinical manifestation, critical illness and outcome of the patients except drowsiness ($p=0.022$). On the other hand hypernatremia had highly significant effect on clinical manifestation, critical illness and outcome of the patients. The patients who had hypernatremia died 16.8 times more than patients with normal sodium ($p= 0.001$).

Conclusion.

Dysnatremias has profound impact on the outcome of covid patients. We found development of hypernatremia has devastating effect on patient survival.

Background:

Water imbalance leading to Dysnatremia is the most common electrolyte disturbances observed in clinical medicine. Two different pathways control this tightly regulated mechanism, these include osmoregulation via ADH and volume regulation via RAAS and natriuretic hormones. Water and sodium balance are regulated independently by these pathways which prevents large changes in plasma osmolality and the effective circulating volume. Severe sodium disorders are related with increased morbidity and mortality (1), even mild hyponatremia, especially when it complicates conditions, such as heart failure and sepsis (2).

Hyponatremia is much common (50%) amongst hospitalized COVID-19 patients than expected (3),and a recent study further suggested that serum sodium concentration was inversely correlated with IL-6, and

association of hyponatremia with a more severe outcome of COVID-19 disease (4). Similarly, Hypernatremia in patients treated in medical intensive care units (ICUs) varied between 6% and 26% (5). During the treatment of patients with severe coronavirus disease 2019 (COVID-19), there was a frequent development of treatment-resistant hypernatremia possibly due to diuretics and steroids use as a part of covid – 19 infection treatment(6).

The impact of covid – 19 is different all over the world, Pakistan has relatively low prevalence of disease as up till now 530,818 cases have been detected, while total number of closed cases, 495,755 had an outcome out, of which 484,508(98%) recovered and 11,247(2%) died. In contrast to France with 3,011,257 cases detected, with total number of 288,469 closed case which had an outcome, 215,822 (75%) Recovered 72,647 (25%) patients died (7). Covid-19 is highly infective organism now is equipped with mutational characteristics. It can lead to a potentially lethal pneumonia which can cause mortality in hospitalized patients as high as 20.3–27.9% (8, 9). The association of hyponatremia and hypernatremia with viral infection and sepsis needs to be explicate and its relation with poor prognosis in patients admitted with Covid – 19 need to be unravel.

The primary objective of this study was to find out, whether dysnatremias are associated with age, comorbid conditions, mechanical ventilation, vasopressor support and death.

Till date very sparse data is available regarding sodium disorders associated with Covid-19. We conducted this study to determine the varying degree of sodium imbalance among patients with Covid-19 and identify the outcome associated with it in our region.

Material And Method:

This is an observational cohort study of all patients admitted in Indus Hospital Covid ICU with a confirmed diagnosis of Covid pneumonia by RT-PCR from 1st March 2020 to 30th Sep 2020. Patients who do not have complete data were excluded. Permission from the institutional ethical review committee was taken prior to conduction of the study. History, clinical examination, laboratory investigations were acquired from the Health Management Information System (HMIS) record of the patients admitted in Covid ICU of The Indus Hospital Korangi, Karachi. Data was collected on a structured proforma which included variables like age, gender, comorbid conditions, clinical presentation, hemodynamic status and lab parameters. Outcome of the patients like discharge or death was noted.

Statistical analysis:

The data was entered and analyzed on IBM, SPSS version 21. Cleaning and ordering of data was done prior to analysis. Mean \pm STD were computed for continuous data while frequency with percentage were obtained for categorical data. All patients were divided in three groups according to their sodium status (Eunatremia, hyponatremia and hypernatremia) and distribution of variables was observed in these three groups. Age of the patients was also categorized in three groups, and age group of 17–50 years was taken as reference category for other age groups in logistic regression. To observe the effect of hypo and

hypernatremia on outcome, critical illness of the patients, drowsiness, and confusion, binary logistic regression was run and odds ratio with 95% confidence interval were obtained. Similarly, logistic regression was also run to calculate the amount of effect of different variable like age, acute kidney injury (AKI), co morbid conditions and volume status on sodium. P value of ≤ 0.05 was considered significant.

Results:

There were total 655 patients in our study in which 463(70.7%) were male, while 192(29.3%) were female. Mean age was 54 ± 15.5 with minimum of 1 year and maximum of 95 year. In terms of clinical presentation, most of our patients with dysnatremias were euvolumic and asymptomatic Fig. 1A & 1B.

The patients who had hypernatremia were 79(12.1%), while hyponatremia was present in 154(23.5%) patients, on the other hand 422(64.4%) patients had normal range of sodium. Dysnatremia were common in patients of age group of 51–65 years (Hyponatremia 72(46.8%) hypernatremia 36(45.6%), as compared to other age groups. Among co morbid conditions hypertension (HTN) was the most prevalent comorbid 380(58%) and hypertensive patients had more hyponatremia and hypernatremia, as compared to normotensive patients. Similarly the diabetic patients also had more hypo and hypernatremia than non-diabetics. Multi organ dysfunction is a manifestation of severe covid and in our cohort AKI was the most occurring complication, encountered by 209(31.9%) patients and hypernatremia was widespread in this group of the patients 53(67.1%) as compared to non AKI patients 26(32.9%) Table 1a.

Critically ill patients who needed ventilatory and vasopressor support were 182(27.8%) and 194(29.6%) respectively, and hypernatremia was more common in these patients as compared to non-critical patients. (Ventilatory support 53[(67.1 %) vs 26(32.9%) and vasopressor support 57 (72.2%) vs 22(27.8%)]. Likewise hypernatremia was more frequent in patients who died 69(87.3%) than patients who recovered 10(12.7%).Table 1b

Effect of demographic and comorbid variables on Hyponatremia

Age, Diabetes mellitus (DM) and ischemic heart disease (IHD) were risk factors for developing hyponatremia in covid – 19 patients. The patients in age group of 51–65 years suffered hyponatremia 1.6 time more than patients of age group of 17–50 years ($p < 0.039$). Similarly, patients of age > 65 years had 1.9 times more at risk for hyponatremia than 17–50 years of age group ($p = 0.015$). The risk of hyponatremia in diabetics was 1.6 times than non-diabetics ($p = 0.009$). Similarly the patients with IHD developed hyponatremia 1.9 times more than patients without IHD ($p = 0.005$). Table 2

Effect of demographic and comorbid variables on Hypernatremia

In parallel with hyponatremia, hypernatremia was not only affected by the older age and DM, but with HTN, AKI and Thromboembolic event (TEE) also. The age group of 51–65 years had greater risk of hypernatremia about 2.2 times than the age group of 17–50 years ($p = 0.017$). Same as patients of > 65 years of age had 3.7 times more hypernatremia than patients of age of 17–50 years ($p < 0.001$). Hypertensive patients suffered with hypernatremia 1.9 times higher than normotensive ($p = 0.015$). Similarly, DM also 1.8 times more effect on hypernatremia than non-diabetic ($p = 0.014$). Presence of AKI in covid patients had 5.5 times more at risk for development of hypernatremia than non AKI patients ($p < 0.001$). Likewise patient with TEE had 3.8 times more hypernatremia than patients without TEE ($p = 0.001$). Patients with hypovolemic status had 3.9 times more hypernatremia than patients with euvolemia ($p < 0.001$) Table 2.

Effect of Dysnatremia on clinical status and outcome of the patients.

Hyponatremia didn't show any significant effect on the clinical manifestation, critical illness and outcome of the patients except drowsiness, as patients who had hyponatremia developed drowsiness 2 times more than the eunatremic patients ($p = 0.022$). (Eunatremia was the reference category)

On the other hand hypernatremia had highly significant effect on clinical manifestation, critical illness and outcome of the patients. The patients who had hypernatremia died 16.8 times more than patients with normal sodium ($p = 0.001$). Similarly, patients with hypernatremia needed vasopressors support 7.7 times more than the eunatremic patients ($p < 0.001$). Patients with hypernatremia required 6.4 times more ventilatory support as compared to the patients with normal level of sodium (< 0.001). Likewise, hypernatremic patients were also 6.6 times more drowsy than eunatremic patients ($p < 0.001$). Confusion was also 7.6 times higher in hypernatremia than eunatremic patients ($p < 0.001$) Table 3.

Discussion:

This is the first study from this country, reported the sodium disorder in Covid – 19 patients admitted in Covid-19 facility at Indus Hospital Karachi.

This study showed that the higher age, HTN, DM, IHD, AKI, TEE and Hypovolemia had devastating impact on sodium balance. We found that hyponatremia was more prevalent than hypernatremia 23% vs 12.1%, but hypernatremia has catastrophic effect on patient's survival as compared with hyponatremia. There are reason for it, as we know that higher age impairs water metabolism, there is relative as well as absolute reduction in total body water, which makes elderly population distinctly inclined to developed dysnatremias (10). Ageing, not only impair the thirst mechanism (11), but it also reduce the concentrating capability of kidneys which increases the risk of dehydration and hypernatremia (12). On the other hand their ability to excrete water load is also hamper, this predisposition contributes to the commonly

observed episode of hyponatremia in elderly hospitalized patients who received hypotonic fluid without proper monitoring (13).

We found significantly higher number of DM patients with both conditions in our population like other study in which, DM is a well-known cause of dysnatremias (14). In another study of 113 patients Liamis et al found poorly controlled sugar level was the major factor in the development of hypernatremia in one third (34%) of their patients. The possible explanation of this observation is the glycosuria induced osmotic diuresis, which tends to raise the serum sodium levels. Steroids are the mainstay of covid 19 therapy, which has dual effect on the generation of hypernatremia, steroids tends to raise the blood sugar level and make it difficult to control, causes osmotic diuresis, concomitantly, they have mineralocorticoid activity which retain sodium and causes hypernatremia. Similarly, hyponatremia can also develop in DM patients by various mechanisms. Firstly, hyperglycemia causes water shift from intracellular to extracellular space causes translocational hyponatremia. Secondly, excessive thirst causes increase water intake and in the background of covid pneumonia, an increase surge of antidiuretic hormone can cause hyponatremia. (15).

In our study, hypertension was significantly associated with hypernatremia, although hyponatremia was also common but didn't reach to a significant level. Ruiz and Cuesta in HOPE-COVID-19 registry analysis also found significant association of hypernatremia in hypertensive patients, the mechanism of this finding is not evident and it will need further evaluation (16). The hyponatremia on the other hand is also reportedly associated with covid hypertensive patients. The underlying mechanism is supposed to be an increased proinflammatory cytokine release like interleukin (IL)-6 induces antidiuretic hormone (ADH) production by direct non osmotic stimulation of ADH. Indirect stimulation of IL-6 is mediated by an injury to the alveolar basement membrane, resulting in activation of hypoxic pulmonary vasoconstriction pathway that lead ADH release⁶¹ (17). On the contrary, IHD was significantly associated with hyponatremia than hypernatremia. This association is also reported in HOPE –COVID-19 registry (16)

In our population, sodium metabolism was affected by the complications of covid-19 infection, like AKI and TEE. We found that, the patients who had AKI developed hypernatremia 5.5 times more as compared to those who did not have AKI. Sjöström and Rysz examined the value of sodium as a prognostic marker for length of hospitalization, AKI and mortality in a cohort of 223 critically ill patients with covid 19. They found patients who developed AKI developed severe hypernatremia along with high mortality (18). Similarly, TEE are well known complication of Covid 19 infection and we found this complication in those who have hypernatremia. The underlying pathophysiology of this macro and micro thrombi in small vessels is not clear and it needs further study (19). Multiple small studies in which critically ill patients with COVID-19 were routinely screened for thrombotic disease demonstrated high rates of thrombotic complications in these patients ranging from 69–85% despite thrombo prophylaxis (20, 21, 22). Similar to this we also observed hypernatremia led to critical illness like need of vasopressor and ventilatory support and high mortality. In a case series of covid patients with hypernatremia Zimmer et al found similar observation of increased ventilatory support along with increased ICU stay (5). On the other hand we didn't find any significant effect of hyponatremia on severity of disease. However in a registry analysis

Sanchez et al found hyponatremia as an independent risk factor for an increase in mortality as well as for an increase in the development of sepsis and requirement for intensive therapy. The reason of their finding might be due to relatively larger sample size of their study (16).

In conclusion, we found dysnatremias are frequent feature of covid 19 infection. The risk factors for this are older age, comorbid conditions and development of complications. Among the dysnatremias, hypernatremia had distressing impact on severity of disease and outcome.

Declarations:

- Ethics approval and consent to participate was taken from IRD-IRB (Reference #- IRD_IRB_2020_04_017).
- All methods were carried out in accordance with relevant guidelines and regulations.
- All experimental protocols were approved by Institutional Review Board (IRB) and Interactive Research and Development (IRD).
- Complete waiver of informed consent was taken because the study was a chart review and there was no patient interaction.
- Consent for publication – not applicable
- Availability of data and materials - The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.
- Competing interests- The authors declare that they have no competing interests
- Funding – no funding
- Authors' contributions:
 1. FS: conception and design of the study along with acquisition of data.
 2. AA : Reviewed the article.
 3. SM: Reviewed the article.
 4. JC: Acquisition of data and reviewed the article
 5. SA: Acquisition of data and reviewed the article
 6. ST: Acquisition of data and reviewed the article
 7. BS: Analysis and interpretation of Data
 8. SI: Drafting, conception, final review of the article along with interpretation and analysis of data.
- Acknowledgement- We acknowledged Dr Jahanzaib Khan (Senior Medical Officer- Dialysis) for helping us in collecting and arranging the Data and manuscript preparation.

References:

1. Reynolds RM, Padfield PL, Seckl JR (2006) Disorders of sodium balance. *BMJ (Clin Res Ed)* 332(7543):702–705. <https://doi.org/10.1136/bmj.332.7543.702>
2. Adrogué HJ, Madias NE (2000) Hyponatremia. *N Engl J Med* 342(21):1581–1589. <https://doi.org/10.1056/nejm200005253422107>
3. Aggarwal S, Garcia-Telles N, Aggarwal G, Lavie C, Lippi G, Henry BM. Clinical features, laboratory characteristics, and outcomes of patients hospitalized with coronavirus disease 2019 (COVID-19): Early report from the United States. *Diagnosis (Berl)*. 2020;7(2):91-6
4. Berni A, Malandrino D, Parenti G, Maggi M, Poggesi L, Peri A (2020) Hyponatremia, IL-6, and SARS-CoV-2 (COVID-19) infection: may all fit together? *J Endocrinol Invest* 43(8):1137–1139. <https://doi.org/10.1007/s40618-020-01301-w>
5. Zimmer MA, Zink AK, Weißer CW, Vogt U, Michelsen A, Priebe HJ, Mols G. Hyponatremia—a manifestation of COVID-19: a case series. *A&A Practice*. 2020 Jul;14(9):e01295.
6. Lindner G, Funk GC. Hyponatremia in critically ill patients. *J Crit Care*. 2013;28:216.e211–216.e220.
7. <https://www.worldometers.info/coronavirus/country/pakistan/> visited on 23-1 2021
8. Rosenberg ES, Dufort EM, Udo T, Wilberschied LA, Kumar J, Tesoriero J, et al. Association of Treatment With Hydroxychloroquine or Azithromycin With In-Hospital Mortality in Patients With COVID-19 in New York State. *JAMA* (2020) 323:2493. doi: 10.1001/jama.2020.8630.
9. Núñez-Gil IJ, Estrada V, Fernández-Pérez C, Fernández-Rozas I, Martín-Sánchez FJ, Macaya C. The COVID-19 curve, health system overload, and mortality. *Emergencias Rev la Soc Esp Med Emergencias* (2020) 32:293–5.
10. Ayus JC, Arieff AI. Abnormalities of water metabolism in the elderly. In *Seminars in nephrology* 1996 Jul 1 (Vol. 16, No. 4, pp. 277-288).
11. Phillips PA, Rolls BJ, Ledingham JG, Forsling ML, Morton JJ, Crowe MJ, et al. Reduced thirst after water deprivation in healthy elderly men. *N Engl J Med*. 1984;311:753–9.
12. Rowe JW, Shock NW, DeFronzo RA. The influence of age on the renal response to water deprivation in man. *Nephron*. 1976;17:270–8.
13. Beck LH. Changes in renal function with aging. *Clin Geriatr Med*. 1998;14:199–209.
14. Liamis G, Rodenburg EM, Hofman A, Zietse R, Stricker BH, Hoorn EJ. Electrolyte disorders in community subjects: prevalence and risk factors. *The American journal of medicine*. 2013 Mar 1;126(3):256-63.
15. Liamis G, Tsimihodimos V, Doumas M, Spyrou A, Bairaktari E, Elisaf M. Clinical and laboratory characteristics of hypernatraemia in an internal medicine clinic. *Nephrology Dialysis Transplantation*. 2008 Jan 1;23(1):136-43.
16. Ruiz-Sánchez JG, Núñez-Gil IJ, Cuesta M, Rubio MA, Maroun-Eid C, Arroyo-Espliguero R, Romero R, Becerra-Muñoz VM, Uribarri A, Feltes G, Trabattoni D. Prognostic impact of hyponatremia and hypernatremia in COVID-19 pneumonia. A HOPE-COVID-19 (Health Outcome Predictive Evaluation for COVID-19) registry analysis. *Frontiers in endocrinology*. 2020;11.

17. Lim JH, Jung HY, Choi JY, Park SH, Kim CD, Kim YL, Cho JH. Hypertension and Electrolyte Disorders in Patients with COVID-19. *Electrolytes & Blood Pressure: E & BP*. 2020 Dec;18(2):23.
18. Sjöström A, Rysz S, Sjöström H, Höybye C. Hyponatremia is common in patients with severe COVID-19 and indicates a poor prognosis. DOI:[10.21203/rs.3.rs-81965/v1](https://doi.org/10.21203/rs.3.rs-81965/v1).
19. Spyropoulos, A.C. et al. Scientific and standardization committee communication: clinical guidance on the diagnosis, prevention and treatment of venous thromboembolism in hospitalized patients with COVID-19. *J. Tromb. Haemost.* <https://doi.org/10.1111/jth.14929> (2020).
20. Llitjos, J.F. et al. High incidence of venous thromboembolic events in anticoagulated severe COVID-19 patients. *J. Tromb. Haemost.* <https://doi.org/10.1111/jth.14869> (2020).
21. Nahum, J. et al. Venous thrombosis among critically ill patients with coronavirus disease 2019 (COVID-19). *JAMA Netw. Open* 3, e2010478 (2020).
22. Ren, B. et al. Extremely high incidence of lower extremity deep venous thrombosis in 48 patients with severe COVID-19 in Wuhan. *Circulation* <https://doi.org/10.1161/CIRCULATIONAHA.120.047407> (2020).

Tables:

Table 1(a)

Distribution of demographic and clinical parameters of patient according to sodium levels n(%)=655

Variables		Eunatremic 422(64.4)	Hyponatremic 154(23.5)	Hypernatremic 79(12.1)	Total 655(%)
Age(years)	≤ 50	171(76)	40(26)	14(17.7)	225(34.4)
	51-65	175(41.5)	72(46.8)	36(45.6)	283(43.2)
	>65	76(18)	42(27.3)	29(36.7)	147(22.4)
Gender	Male	299(70.9)	102(66.2)	62(78.5)	463(70.7)
	Female	123(29.1)	52(33.8)	17(21.5)	192(29.3)
Comorbid of patients					
HTN	Yes	225(53.5)	99(64.3)	56(70.9)	380(58)
	No	197(46.7)	55(35.7)	23(29.1)	275(42)
DM	Yes	167(39.6)	84(54.5)	46(58.2)	297(45.3)
	No	225(60.4)	70(45.5)	33(41.8)	358(54.7)
IHD	Yes	58(13.7)	39(25.3)	19(24.1)	116(17.7)
	No	364(86.3)	115(74.7)	60(75.9)	539(82.3)
Complications of Covid 19 infection					
AKI	Yes	103(24.4)	53(34.4)	53(67.1)	209(31.9)
	No	319(75.6)	101(65.6)	26(32.9)	446(68.1)
CCF	Yes	17(4)	8(5.2)	5(6.3)	30(4.6)
	No	405(96)	146(94.8)	74(93.7)	625(95.4)
TEE	Yes	13(3.1)	8(5.2)	10(12.7)	31(4.7)
	No	409(96.9)	146(94.8)	69(87.3)	624(95.3)
HTN Hypertension, DM Diabetes mellitus, IHD Ischemic heart disease, AKI Acute kidney injury, CCF Congestive cardiac failure, TEE Thromboembolic event					

Table 1(b)

Distribution of clinical parameters of patients according to sodium levels n(%)=655

Variables		Eunatremic 422(64.4)	Hyponatremic 154(23.5)	Hypernatremic 79(12.1)	Total 655
Volume status	Euolumic	382(90.5)	128(83.1)	61(77.2)	571(87.2)
	Hypovolemic	13(3.1)	13(8.4)	12(15.2)	38(5.8)
	Hypervolumic	27(6.4)	13(8.4)	6(7.6)	46(7)
Drowsiness	Yes	31(7.3)	21(13.6)	27(34.2)	79(12.1)
	No	391(92.7)	133(86.4)	52(65.8)	576(87.9)
Confusion	Yes	33(7.8)	19(12.3)	31(39.2)	83(12.7)
	No	389(92.2)	135(87.7)	48(60.8)	572(87.3)
Vomiting	Yes	31(7.3)	18(11.7)	6(7.6)	55(8.4)
	No	391(92.7)	136(88.3)	73(92.4)	600(91.6)
Ventilatory support	Yes	102(24.2)	27(17.5)	53(67.1)	182(27.8)
	No	320(75.8)	127(82.5)	26(32.9)	473(72.2)
Vasopressor support	Yes	106(25.1)	31(20.1)	57(72.2)	194(29.6)
	No	316(74.9)	123(79.9)	22(27.8)	461(70.4)
Outcome	Died	123(29.1)	35(22.7)	69(87.3)	227(34.7)
	Recovered	299(70.7)	119(77.3)	10(12.7)	428(65.3)

Table 2

Effect of age and clinical variables on hypo and hypernatremia in covid 19 infection

Variables		HYPONATREMIA			HYPERNATREMIA		
		Odds Ratio	95% CI Lower-Upper	p value	Odds Ratio	95% CI Lower-Upper	p value
Age(years)	≤ 50	1	1		1	1	
	51-65	1.6	1.0 - 2.4	0.039	2.2	1.2 - 4.2	0.017
	>65	1.9	1.1 - 3.1	0.015	3.7	1.9 - 7.3	<0.001
HTN	Yes	1.4	0.97- 2.1	0.072	1.9	1.1- 3.2	0.015
DM	Yes	1.6	1.13 - 2.3	0.009	1.8	1.1 - 2.9	0.014
AKI	yes	1.2	0.79 - 1.7	0.446	5.5	3.3 - 9.1	<0.001
IHD	Yes	1.9	1.2 - 2.9	0.005	1.6	0.9 - 2.7	0.118
CCF	Yes	1.2	0.5 -2.7	0.677	1.5	0.55 - 4	0.431
TEE	Yes	1.1	0.49 - 2.6	0.78	3.8	1.7 - 8.5	0.001
Volume status	Euvolemia	1	1		1	1	
	Hypovolemia	1.8	0.89 - 3.6	0.099	3.9	1.9 - 8.0	<0.001
	Hypervolumeia	1.4	0.69 - 2.7	0.365	1.3	0.511 - 3.1	0.62
HTN Hypertension, DM Diabetes mellitus, AKI Acute kidney injury, IHD Ischemic heart disease, CCF Congestive cardiac failure, TEE Thromboembolic event							

Table 3

Effect of Dysnatremia on disease severity in covid 19 infection

Variables		Odds Ratio	95% CI Lower-Upper	p vale
Hyponatremia	Death	0.72	0.46-1	0.127
	Vasopressor need	0.75	0.48-1.2	0.214
	Ventilatory support	0.67	0.42-1.1	0.092
	Drowsiness	2	1.1- 3.6	0.022
	Confusion	1.7	0.9 - 3.0	0.097
Hypernatremia	Death	16.77	8.4-33.6	0.001
	Vasopressor need	7.7	4.5-13.2	<0.001
	Ventilatory support	6.4	3.8-10.8	<0.001
	Drowsiness	6.6	3.6-11.8	<0.001
	Confusion	7.6	4.3 - 13.5	<0.001

Figures

Figure 1A




Figure 1B




Figure 1

In terms of clinical presentation, most of our patients with dysnatremias were euvolumic and asymptomatic