Sensitization to cat and dog components and prediction of symptoms in cat-sensitized children.

Katarzyna Smejda

Uniwersytet Medyczny w Łodzi

Joanna Jerzyńska (✉ joanna.jerzynska@umed.lodz.pl)

Uniwersytet Medyczny w Łodzi <https://orcid.org/0000-0001-9504-2673>

Daniela Podlecka

Uniwersytet Medyczny w Łodzi

Agnieszka Brzozowska

Uniwersytet Medyczny w Łodzi

Research

Keywords: children, components, cat, dog, allergy, symptoms, sensitization

Posted Date: July 14th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-25143/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background: Sensitization to cat and/or dog allergens during childhood represents a risk factor for the development of allergic diseases later in life.

Material and Methods: The study investigated the association of patterns of sensitization to cat and dog allergen components with clinical symptoms of allergy to these furry animals among cat-sensitized children. It included 50 children aged 5-17 years who showed sensitization to cat in the skin prick test. They were considered to have cat allergy if they suffered from one or more allergy symptoms when being exposed to contact with a cat. The children were evaluated for the presence of bronchial asthma, atopic dermatitis and allergic rhinitis. Their mothers completed a questionnaire on pet exposure at home. Levels of serum IgE cat epitopes Fel d (1, 2, 4), as well as dog components Can f (1, 2, 3, 5) were measured in all the studied children.

Results: Respiratory symptoms following exposure to the cat allergen were most common in children with Fel d 2 epitope ($p = 0.041$). After contact with a dog, respiratory symptoms were most common in children with Can f 1 epitope ($p = 0.042$), eczema in children with sensitization to both Can f 1 ($p = 0.009$) and Can f 2 ($p = 0.002$), whereas eye symptoms occurred mostly in children with Can f 3 ($p = 0.039$).

Conclusions: Molecular diagnosis in patients with pet allergy may help clinicians to predict clinical symptoms and their severity.

Introduction

Cats and dogs are the most common household pets. According to the latest research, over half (52%) of households in Poland own a pet. Forty-two percent of Polish citizens declare that they have a dog, 26% report to keep a cat, and 5% possess other animals [1]. In the ECAP study of the Polish population, an average frequency of cat allergy was 13.5%, and in the case of dog allergy it was 9.7% [2]. The dog and cat allergens are a common cause of allergic sensitization and

a factor triggering respiratory symptoms worldwide.

Molecule-based diagnosis of allergy is an important advancement in improving sensitivity and specificity [3]. The major cat allergen, in up to 96% cases, is Fel d 1, a uteroglobin produced in the skin and salivary glands [4, 5, 6]. Other important cat allergens are albumin of Fel d 2 serum [7] and lipocalin Fel d 4 [8] which cross-reacts with the corresponding proteins from other animals [9]. Commonly described dog allergens represent lipocalins (Can f 1, Can f 2, Can f 4 and Can f 6) or albumins (Can f 3) [10], prostatic kallikrein (Can f 5) which is an androgen-regulated protein expressed in the prostate and detectable only in male dogs [11]. Due to the low degree of sequence identity among them, lipocalins were considered as species-specific allergy markers.

Sensitization to Fel d 1 and Can f 1 in childhood and polysensitization to either cat or dog allergen molecules predict allergy significantly better than cat or dog IgE extracts [12]. Fel d 7 is a common allergen in the cat-sensitized population that cross-reacts with Can f 1 and may contribute to symptoms in both cat-allergic and dog-allergic patients [13].

The study investigated the association of patterns of sensitization to cat and dog allergen components with clinical symptoms of allergy to these furry animals among cat-sensitized children, considering their prenatal/postnatal exposure to the pets.

Material And Methods

The study included 50 children of both sexes, aged 5–18 17 years, patients of our allergy clinic, seen between January and September of 2018, who showed sensitization to cat allergen based on the skin prick test. They were considered to have allergy to cat if they suffered from one or more allergy symptoms when exposed to contact with a cat, which was confirmed by positive medical history). The allergy was manifested among others by ocular symptoms (tearing, redness, pruritus), nasal symptoms (sneezing, nasal itch, runny nose), eczema, symptoms of the lower respiratory tract (cough, shortness of breath, wheezing). The children were also evaluated for the presence of bronchial asthma, atopic dermatitis, and allergic rhinitis. Their mothers completed a questionnaire on pet exposure at home: a) during pregnancy, b) in the first two years of the child's life, and c) after the second year of the child's life. Since a majority of the mothers reported having a dog rather than a cat, additional analyses of sensitization to dog allergens were performed. Children were considered to have allergy to dog if they suffered from one or more allergy symptoms that occurred when having contact with a dog, and at the same time presented Can f level (1, 2, 3 or 5) of least grade 1.

Blood samples were collected from all the studied children to measure the IgE serum levels for the cat epitopes, Fel d 1, Fel d 2, Fel d 4, as well as for the dog components, Can f 1, Can f 2, Can f 3, Can f 5. IgE serum levels > 0.35 IU/ml were regarded as a positive result. The serum samples were analyzed for IgE antibodies using the ImmunoCAP® system according to the manufacturer's guidelines (Phadia AB, Uppsala, Sweden). Serum samples of venous blood were used. The samples were stored at -20 °C. The ImmunoCAP Specific IgE detects IgE antibodies in the range from 0 to 100 kUA /l where "A" stands for allergen-specific antibodies. The result is read quantitatively. In clinical practice, a level of 0.35 kUA /l has commonly been accepted as a cut-off. The clinical performance is expressed as sensitivity, ranging from 84% to 95%, and specificity, ranging from 85–94%.

The study was approved by the Bioethics Committee of the Medical University of Lodz, Poland, RNN/39/14/KE; a written informed consent was obtained from all the subjects before the assays.

Statistical analysis

The investigated traits were described based on the measures of location – arithmetic or geometric mean, along with measures of dispersion – standard deviation, 95% confidence interval as well as

minimum and maximum values. The categorical variables were depicted by using absolute numbers and percentages.

Multivariate logistic regression models (for binary dependent variables such as allergy symptoms or having an animal at home) were performed in order to test statistical relationships. When dealing with non-normally distributed variables, robust standard errors (*i.e.* sandwich estimators) were used within a specific regression model. Some variables, which revealed the perfect prediction (*i.e.*, when each study participant or none of them displayed a specific trait), were excluded from a regression equation. All the regression models were controlled for the studied patients' age and gender. A level of $P < 0.05$ was considered statistically significant. All the statistical computations were carried out by means of Stata/Special Edition, release 14.2 (StataCorp LP, College Station, Texas, USA).

Results

Baseline characteristics of the studied population are presented in Table I.

Table I. Baseline characteristics of the studied patients (n = 50)
(presented as mean, standard deviation, 95% confidence interval, and min.-to-max. values for numerical traits; absolute numbers and percentages for discrete variables)

Investigated trait	Statistical parameter			
	<i>M</i>	<i>SD</i>	<i>95% CI</i>	<i>Min. - max.</i>
Age (years)	11.7	3.32	10.8-12.7	6-18
	Yes (n)	%	No (n)	%
Gender (female)	18	36.0	32	64.0
Symptoms after contact with the cat	38	76.0	12	24.0
Symptoms after contact with the dog	12	24.0	38	76.0
Keeping a cat in any period of life	16	32.0	34	68.0
Keeping a dog in any period of life	26	52.0	24	48.0
Presence of other pets	7	14.0	43	86.0
Asthma	45	90.0	5	10.0
Allergic rhinitis	50	100.0	0	0.0
Atopic dermatitis	6	12.0	44	88.0

Cat allergen components and allergy symptoms

Thirty-eight children were allergic to cats as determined by positive anamnesis, whereas twelve patients exhibited clinically non-relevant sensitization as indicated by negative anamnesis (Table II). Forty-seven of all the children (94.00%) had at least one of the *Fel d* epitopes tested, 34 children had only one epitope (72.34%), eight children had two epitopes (17.02%), four children had three epitopes (8.51%). There was no correlation between the number of epitopes and symptoms of allergy to cats. The most common epitope was *Fel d* 1 (90% of the whole study group); relevant results are displayed in Fig. 1. Respiratory symptoms following contact with a cat allergen were most frequent in the children with *Fel d* 2 epitope ($p = 0.041$) (Table III). There was no relationship between other (*i.e.*, ocular, nasal, cutaneous) symptoms of allergy to cats and the type of epitope (Table III). A correlation was found between the occurrence of *Fel d* 2 epitope and any symptoms appearing after contact with a cat ($p = 0.048$) (Table III).

Table II. Detailed characteristics of the studied patients concerning symptoms due to exposure to animal allergens (n = 50)
(presented as absolute numbers and percentages)

Investigated trait	Statistical parameter			
	Yes (n)	%	No (n)	%
Symptoms after contact with a cat	38	76.0	12	24.0
Ocular	31	62.0	19	38.0
Nasal	32	64.0	18	36.0
In the lower airways	10	20.0	40	80.0
Eczema	12	24.0	38	76.0
Symptoms after contact with a dog	12	76.0	38	24.0
Ocular	4	8.0	46	92.0
Nasal	7	14.0	43	86.0
In the lower airways	5	10.0	45	90.0
Eczema	7	14.0	43	86.0

Table III. Allergic symptoms after contact with a cat and/or a dog and the prevalence of selected *Fel d* and/or *Can f* epitopes

in the studied patients (n = 50)
(statistical significance of relationships)

Allergic symptoms after contact with the cat	<i>Fel d</i> 1	<i>Fel d</i> 2	<i>Fel d</i> 4	<i>Can f</i> 1	<i>Can f</i> 2	<i>Can f</i> 3	<i>Can f</i> 5
Any symptoms	NS*	p = 0.048	NS	NS	NS	NS	NS
Ocular	NS	NS	NS	p = 0.042	NS	p = 0.039	NS
Nasal	NS	NS	NS	NS	NS	NS	NS
In the lower airways	NS	p = 0.041	NS	NS	NS	NS	NS
Eczema	NS	NS	NS	p = 0.009	p = 0.002	NS	NS

NS – [statistically] not significant

Dog allergen components and allergy symptoms

Twelve children were allergic to dogs as confirmed by positive anamnesis and thirty-eight exhibited clinically non-relevant sensitization as indicated by negative anamnesis (Table II). Twenty nine of the fifty study participants (58.00%) had at least one of the Can f epitopes tested, sixteen children had only one epitope (55.17%), ten children had two epitopes (34.48%), one patient had three epitopes (3.45%), and one child had four epitopes (3.45%). There was no relationship between the number of epitopes and symptoms of allergy to dogs. The most common epitope was Can f 1 (32% of the study sample); the results are shown in Fig. 1. Eczema developing after contact with a dog showed a strong correlation with sensitization to both Can f 1 (p = 0.009) and Can f 2 (p = 0.002) allergen components (Table III). The presence of allergic symptoms in the eyes showed a significant relationship with sensitization to Can f 1 and Can f 3 (p = 0.042 and p = 0.039, respectively) (Table III). Additionally, the presence of Can f 1 and Can f 2 significantly conditioned the occurrence of eczema after contact with a dog (p = 0.009 and p = 0.002, respectively) (Table III).

Pet components and allergic disorders

There was no relationship between diagnosed asthma and sensitization to any pet components. Monosensitization to Can f 1 significantly underpinned the prevalence of atopic dermatitis in the studied children (p = 0.036).

Cross-reactivity

Table IV depicts the co-occurrence of selected Fel d and/or Can f epitopes in the studied children. Fifteen children (30%) showed sensitivity to Fel d 1 and Can f 1, whereas thirteen patients (26%) to Fel d 1 and Can f 5.

Table IV. Overlapping of selected *Fel d* and/or *Can f* epitopes in the studied patients (n = 50)

(presented as absolute numbers)

Epitope	<i>Fel d 1</i>	<i>Fel d 2</i>	<i>Fel d 4</i>	<i>Can f 1</i>	<i>Can f 2</i>	<i>Can f 3</i>	<i>Can f 5</i>
<i>Fel d 1</i>		4	11	15	6	5	13
<i>Fel d 2</i>	4		5	3	1	5	1
<i>Fel d 4</i>	11	5		6	4	5	2
<i>Can f 1</i>	15	3	6		6	1	6
<i>Can f 2</i>	6	1	4	6		1	2
<i>Can f 3</i>	5	5	5	3	1		1
<i>Can f 5</i>	13	1	3	6	2	1	

Prenatal/postnatal exposure to pets

Keeping a cat during pregnancy, in early childhood, or at any time during a child's life showed no relationship to sensitization to any feline epitopes tested. Having a dog in any period of pre-natal and post-natal life was associated with sensitization to Can f 1 epitope ($p = 0.033$) (Table V). Keeping a dog and a cat at the same time – regardless of the life stage, prenatal or postnatal, did not reveal any statistically significant relationship with sensitization to the epitopes tested.

Table V. Keeping a dog in the domicile and prevalence of selected *Can f* epitopes in the studied patients (n = 50)
(statistical significance of relationships)

Keeping a dog	<i>Can f 1</i>	<i>Can f 2</i>	<i>Can f 3</i>	<i>Can f 5</i>
Overall	$p = 0.033$	NS	NS	NS
During pregnancy	$p = 0.042$	NS	NS	NS
Once in the past	$p = 0.033$	NS	NS	NS
At the age of 0-2	$p = 0.048$	NS	NS	NS
At the age of over 2	$p = 0.021$	NS	NS	NS
Currently	NS	NS	NS	NS

NS - [statistically] not significant

Discussion

In the last few years, component resolved diagnostics (CRD) has been developing rapidly, and both singleplex and multiplex assays are used in relation to over one hundred recombinant or purified native allergen components [14]. Molecular allergy is based on identification, characterization and subsequent

use of single allergens, being components of complex allergen sources like pollen, mites, furred animals, foods or insect venoms [15, 16, 17]. Thus, carefully defined allergen molecules serve as a useful complement to the reagents available to date, and optimize IgE determination and detection of specific sensitization in the context of allergy diagnosis [16, 17]. Our study showed that respiratory symptoms after contact with a cat and a dog have a strong relationship with presence of Fel d 2 and Can f 1 components, respectively. The occurrence of eczema after contact with a dog reveals a strong correlation with the presence of Can f 1 and Can f 2 epitopes. The presence of ocular symptoms showed a significant relationship with Can f 3 epitope.

Gent et al. found that the risk for wheezing in children sensitized and exposed to pet allergen significantly increased (by 39% and 53% for Fel d 1 > 0.12 µg/g, Can f 1 > 1.2 µg/g, respectively) [18]. The most frequent symptoms of cat owners suffering from allergy when exposed to contact with their pets were rhinitis (sneezing, rhinorrhea, nasal obstruction and itching, postnasal drip) (80.0%), followed by conjunctivitis (73.3%), cutaneous symptoms (skin itching, skin rash, angioedema, urticaria) (33.3%), and lower respiratory symptoms (sputum, dyspnea, chest discomfort and pain, wheezing) (13.3%). Dog owners with allergy to their pets suffered from cutaneous symptoms more frequently than cat owners with cat allergy (53.3% vs. 33.3%, $P < 0.01$), especially in terms of urticaria (35.9% vs. 15.6%, $P < 0.01$) as well as from lower respiratory symptoms (33.3% vs 13.3%, $P < 0.05$), especially cough (23.3% vs 6.7%, $P < 0.05$) [19].

In our study there was no statistically significant relationship between diagnosed asthma and sensitization to any pet components. Monosensitization to Can f 1 was related to the diagnosis of atopic dermatitis. Uriarte et al. showed that Can f1 was associated with persistent rhinitis, Can f 2 with asthma diagnosis, Can f3 with severity of rhinitis and asthma, and Can f 5 both to persistence and severity of rhinitis. Sensitization to several allergens in patients (1, 2, 3 or 4) was associated with persistent moderate asthma or rhinitis. Direct contact with dogs was connected both with persistence and severity of rhinitis [20, 21]. Fel d2 was associated with severity of rhinitis and asthma. Fel d4 was associated with the presence of asthma symptoms. Direct contact with cats was associated both with persistence and severity of rhinitis [22]. Bjerg et al. reported a higher prevalence of asthma in children co-sensitized to Fel d1 and Fel d 4 than in children sensitized to Fel d 1 alone [23]. They also found a higher prevalence of asthma (55%) in patients with IgE reactivity, not only towards Fel d 1, but also towards Fel d 2 and/or rFel d 4. Wisniewski et al. described an association between high levels of IgE antibodies to nFel d 2 and rFel d 4 and atopic dermatitis in cat-allergic children [24].

The exposure to animal allergens constitutes a relevant risk factor for the development of allergic sensitization. Some authors suggest that early contact with dogs, before the first year of age, may have a protective effect in terms of allergic sensitizations [25, 26]. Similarly, conflicting studies have been reported for early cat exposures [27]. So far, no consensus has been reached on the potential association between animal exposure and prevention of onset of asthma or other allergic diseases in later life [28]. Conclusions presented in various studies also differ on the impact of canine or feline allergens as being those most protective ones [29]. In our study, keeping a dog at any time, in any period of prenatal and

postnatal life, was associated with the occurrence of IgE to Can f 1 epitope and allergy symptoms. While our study has not shown any relationship between keeping a cat during pregnancy, in early childhood, or at any time during a child's life, keeping a cat during pregnancy had no relation to the occurrence of any of the cat epitopes tested either. Many authors have found no relationship between possession of animals and exposure to asthma and other allergic diseases [30, 31]. In a meta-analysis of eleven European birth cohorts including > 22 thousand children, there was no relationship between keeping any furry animal (mainly a cat and dog) at home in the first two years of a child's life and occurrence of asthma at 6–10 years of age [32]. However, there are very reliable studies that suggest a protective effect of having a pet. Takkouche et al. showed that exposure to dogs slightly increased the risk of asthma, but exposure to cats played a protective role [33]. Researchers from Sweden found that keeping a dog in the first year of a child's life is associated with a significantly lower risk of asthma in children aged 6 years [34]. Another study, involving 696 Swedish children, reports that current asthma and asthma symptoms following contact with cats were associated with cosensitization to Fel d 1 and Fel d 4. Asthma was related to higher levels of component sensitization (Fel d 1 $\geq$ 15 ISU) while sensitization to more than one component from the same animal conferred the greatest risk [35]. Based on the evidence from the past decade it may be assumed that early exposure to dogs before the first year of life may have a protective effect in preventing allergic sensitizations [36, 37].

The limitation of our study is the small sample size and a restricted number of assayed components. All the studied children were diagnosed with allergic rhinitis, which is why the disease was not included in multifactor statistical analyses.

Conclusions

To sum up, our article described the pattern of sensitization to pet IgE components and its association with clinical symptoms. Our study showed that respiratory symptoms in children allergic to cats or dogs demonstrate a strong relationship with the presence of Fel d 2 and Can f 1 components, respectively. Eczema after contact with a dog showed a strong correlation with the presence of Can f 1 and Can f 2 epitopes. The presence of allergic eye symptoms showed a statistically significant relationship with the presence of the Can f 3 epitope. However, Fel d 1 is the most important cat allergen component, yet the functional role of this molecule remains unknown. Molecular diagnosis in patients with allergy to pets may help clinicians to predict clinical symptoms and their severity.

Declarations

Corresponding author:

Jerzynska J MD, PhD, Prof, Department of Pediatrics and Allergy, Medical University of Lodz, Poland, Copernicus Memorial Hospital, Korczak Pediatric Center, Piłsudskiego 71 Str., 90-329 Lodz, Poland, Tel ++48422074726, Fax ++48426776358, E-mail: alergol@kopernik.lodz.pl

Conflict of interest:

The authors declare no conflict of interest regarding publication of this article.

Financial information:

This study was funded by grant 503/2-056-01/503-21-001 from the Medical University of Lodz, Poland.

References

1. Kantar Public, may 2017, K.021/17.
2. Sybilski AJ, Raciborski F, Lipiec A, et al. Epidemiology of atopic dermatitis in Poland according to the Epidemiology of Allergic Disorders in Poland (ECAP) study. *J Dermatol*. 2015;42(2):140-147.
3. Canonica GW, Ansotegui IJ, Pawankar R, et al. A WAO - ARIA - GA²LEN consensus document on molecular-based allergy diagnostics. *World Allergy Organ J*. 2013;6:17.
4. Kaiser L, Velickovic TC, Badia-Martinez D, et al. Structural characterization of the tetrameric form of the major cat allergen Fel d 1. *J Mol Biol*. 2007;370:714–727.
5. Bonnet B, Messaoudi K, Jacomet F, et al. An update on molecular cat allergens: Fel d 1 and what else? Chapter 1: Fel d 1, the major cat allergen. *Allergy Asthma Clin Immunol*. 2018; 14: 14.
6. van Ree R, van Leeuwen WA, Bulder I, Bond J, Aalberse RC. Purified natural and recombinant Fel d 1 and cat albumin in in vitro diagnostics for cat allergy. *J Allergy Clin Immunol*. 1999;104:1223–1230.
7. Virtanen T, Kinnunen T. Mammalian allergens. *Clin Allergy Immunol*(2008) 21:201–1810.
8. Smith W, et al. Fel d 4, a cat lipocalin allergen. *Clin Exp Allergy*. 2004;34(11):1732–1738
9. Hilger C, Swiontek K, Arumugam K, Lehnert C, Hentges F. Identification of a new major dog allergen highly cross-reactive with Fel d 4 in a population of cat- and dog-sensitized patients. *J Allergy Clin Immunol*. 2012;129(4):1149–1151.e1142.
10. Hentges F, Leonard C, Arumugan K, Hilger C. Immune response to mammalian allergens. *Front Immunol*. 2014 May 21;5:234.
11. Mattsson L, Lundgren T, Everberg H, Larsson H, Lidholm J. Prostatic kallikrein: A new major dog allergen. *J Allergy Clin Immunol*. 2009;123:362–8.
12. Patelis A, Dosanjh A, Gunnbjörnsdóttir M, et al. New data analysis in a population study raises the hypothesis that particle size contributes to the pro-asthmatic potential of small pet animal allergens. *Ups J Med Sci*. 2016;121:25–32
13. Apostolovic D, Sánchez-Vidaurre S, Waden K, et al. The cat lipocalin Fel d 7 and its cross-reactivity with the dog lipocalin Can f 1. *Allergy*. 2016 Oct;71(10):1490-5.
14. Kleine-Tebbe J, Jakob T. Molecular allergy diagnostics using IgE singleplex determinations: methodological and practical considerations for use in clinical routine: Part 18 of the Series Molecular Allergology. *Allergo J Int*. 2015;24:185-197.

15. Kleine-Tebbe J, Jappe U. Molecular allergy diagnostic tests: development and relevance in clinical practice. *Allergol Select*. 2017 Aug 4;1(2):169-189.
16. Cui Y, Yu L, Teng F et al. Expression of recombinant allergen, Der f 1, Der f 2 and Der f 4 using baculovirus-insect cell systems. *Arch Med Sci*. 2018 Oct;14(6):1348-1354.
17. Petrosino MI, Scaparrotta A, Marcovecchio ML, et al. Usefulness of molecular diagnosis in egg allergic children. *Arch Med Sci*. 2018 Jan;14(1):132-137.
18. Gent JF, Kezik JM, Hill ME, Tsai E, Li DW, Leaderer BP. Household mold and dust allergens: exposure, sensitization and childhood asthma morbidity. *Environ Res*. 2012 Oct;118:86-93.
19. Yang MS, Lee SP, Kwon YJ, Lee SM. Dog and Cat Allergies and Allergen Avoidance Measures in Korean Adult Pet Owners Who Participated in a Pet Exhibition. *Allergy Asthma Immunol Res*. 2018 Mar; 10(2): 155–164.
20. Uriarte SA, Sastre J. Clinical relevance of molecular diagnosis in pet allergy. *Allergy*. 2016 Jul;71(7):1066-8.
21. Uriarte SO, Sastre J. Clinical impact of molecular diagnosis in dog allergy *Clin Transl Allergy*. 2014; 4(Suppl 2): P52.
22. Uriarte SO, Sastre J. Clinical impact of molecular diagnosis in cat allergy. *Clin Transl Allergy*. 2014; 4(Suppl 2): P51.
23. Bjerg A, Winberg A, Berthold M, Mattsson L, Borres MP, Rönmark E. A population-based study of animal component sensitization, asthma and rhinitis in schoolchildren. *Pediatr Allergy Immunol*. 2015;26(6):557–563.
24. Wisniewski JA, Agrawal R, Minnicozzi S, et al. Sensitization to food and inhalant allergens in relation to age and wheeze among children with atopic dermatitis. *Clin Exp Allergy*. 2013;43(10):1160–1170.
25. Perzanowski MS, Rönmark E, Platts-Mills TA, Lundbäck B. Effect of cat and dog ownership on sensitization and development of asthma among preteenage children. *Am J Respir Crit Care Med*. 2002;166:696–702.
26. Takkouche B, González-Barcala FJ, Etminan M, Fitzgerald M. Exposure to furry pets and the risk of asthma and allergic rhinitis: a meta-analysis. *Allergy*. 2008;63:857–864.
27. Celedón JC, Litonjua AA, Ryan L, Platts-Mills T, Weiss ST, Gold DR. Exposure to cat allergen, maternal history of asthma, and wheezing in first 5 years of life. *Lancet*. 2002;360:781–782.
28. Flohr C, Yeo L. Atopic dermatitis and the hygiene hypothesis revisited. In: Shiohara T, editor. *Pathogenesis and management of atopic dermatitis*. Basel: Karger; 2011. pp. 1–34.
29. Mandhane PJ, Sears MR, Poulton R, et al. Cats and dogs and the risk of atopy in childhood and adulthood. *J Allergy Clin Immunol*. 2009;124:745–750.
30. Chen CM, Tischer C, Schnappinger M, et al. The role of cats and dogs in asthma and allergy – a systematic review. *Int J Hyg Environ Health* 2010;213:1-31.
31. Medjo B, Atanaskovic-Markovic M, Nikolic D, et al. Association between pet-keeping and asthma in school children. *Pediatr Int* 2013;55:133-7.

32. Lødrup Carlsen KC, Roll S, Carlsen KH, et al. Does pet ownership in infancy lead to asthma or allergy at school age? Pooled analysis of individual participant data from 11 European birth cohorts. *PLoS One*. 2012;7:e43214.
33. Takkouche B, González-Barcala FJ, Etminan M, et al. Exposure to furry pets and the risk of asthma and allergic rhinitis: a meta-analysis. *Allergy* 2008;63:857-64.
34. Fall T, Lundholm C, Örtqvist AK, et al. Early Exposure to Dogs and Farm Animals and the Risk of Childhood Asthma. *JAMA Pediatr* 2015;169:e153219.
35. Bjerg A, Winberg A, Berthold M, Mattsson L, Borres MP, Rönmark E. A population-based study of animal component sensitization, asthma, and rhinitis in schoolchildren. *Pediatr Allergy Immunol*. 2015;26:557–563.
36. Perzanowski MS, Rönmark E, Platts-Mills TA, Lundbäck B. Effect of cat and dog ownership on sensitization and development of asthma among preteenage children. *Am J Respir Crit Care Med*. 2002;166:696–702.
37. Takkouche B, González-Barcala FJ, Etminan M, Fitzgerald M. Exposure to furry pets and the risk of asthma and allergic rhinitis: a meta-analysis. *Allergy*. 2008;63:857–864.

Figures

Figure 1

.Presence of selected epitopes - referring to allergy to cat dander - in the studied patients (n = 50)
(presented as percentages)

Figure 2

Presence of selected epitopes - referring to allergy to dog dander - in the studied patients (n = 50)
(presented as percentages)