

# Clinical Characteristics of Magnetic Foreign Body Mis-ingestion in Children—A Big Issue for Small Children

**Yu Kun Huang**

Children's Hospital of Fudan University Xiamen Branch

**Shao Xian Hong**

Children's Hospital of Fudan University Xiamen Branch

**I Hsin Tai**

China Medical University Hospital

**Kai Sheng Hsieh** (✉ [kshsieh@hotmail.com](mailto:kshsieh@hotmail.com))

Taipei Medical University-Shuang Ho Hospital

---

## Research Article

**Keywords:** magnetic foreign bodies, mis-ingestion in children, necrosis of intestinal wall, intestinal perforation

**Posted Date:** March 3rd, 2021

**DOI:** <https://doi.org/10.21203/rs.3.rs-251786/v1>

**License:**  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

# Abstract

**Objective:** Magnetic foreign body mis-ingestion (MFBM) is now occurring more frequently. it may cause remarkable mortality and morbidity in children.

**Methods:** A retrospective analysis of the clinical data of children admitted to Xiamen Children's Hospital between March 2017 and July 2020 due to accidental MFBM.

**Results:** A total of 14 children who had MFBM were collected, the proportion between urban and rural areas was 8:6, and the ratio of male to female was 6:1. The age ranged from 1.2 to 8.9 years (median 4.6 years). The number of magnetic foreign bodies ingested by mistake is 1 to 17 (average 6.5) Magnetic foreign objects are divided into magnet (3 cases) + magnetic beads (11 cases). About 40% (5/14) of this patient series showed no available mis-ingestion history. Management includes: 4 cases of open surgery (including 1 case of laparoscopic transfer to operation), 3 cases of laparoscopic surgery, 2 cases of gastroscopy, 5 cases of conservative treatment of foreign bodies discharged through the anus. Of the 7 surgical cases, 6 cases presented with intestinal obstruction and intestinal perforation (at least 1 intestinal perforation and at most 5). Abdominal sonography has limitations in the detection of magnetic foreign bodies in the digestive tract. The proportion of laparoscopic surgery in the 7 surgical cases is nearly half. All surgical cases recovered smoothly after treatment.

**Conclusions:** Our experience shows that MFBM is a big issue for the small children! The early symptoms of MFBM are often atypical especially among young children and MFBM may lead to severe adverse events. We proposed a management strategy for MFBM in children. We advise pediatricians/emergency physicians, parents/children's guardians and society should raise the collaborated alertness of MFBM. Global awareness of risk prevention of magnetic material accidental ingestion cannot be overemphasized.

## Introduction

Foreign bodies in the digestive tract of children have become one of important accidental injuries in children. The first case of foreign bodies in the digestive tract of children was reported in 1692<sup>1</sup>. It was the 4-year-old Prince Brandenburg who swallowed a leather shoe fastener. Since then, similar situations are not uncommon, and the consequences and complications caused by this are quite different. It mainly depends on the shape, nature, and quantity of the foreign body. Its outcomes ranged from the spontaneous discharge, intestinal perforation, obstruction, bleeding, and even sepsis, septic shock, etc<sup>2,3</sup>. Some of the complications are life-threatening. To clarify clinical status of MFBM in children, we therefore conducted this study.

## Materials And Methods

We respectively retrieved and reviewed the clinical data of all children admitted to Xiamen children's Hospital (XMCH), including age, gender, clinical manifestations, medical history and signs, imaging

examinations, treatment, and outcome, etc.

The diagnostic criteria for magnetic foreign bodies were based on the history of accidental swallowing of magnetic foreign bodies, clinical manifestations, abdominal signs, and abdominal imaging examination (abdominal X-ray and CT examination). The relevant data were analyzed and presented. The study was approved by the institutional review board of Xiamen children's Hospital. The authors have identified the institutional and/or licensing committee approving the experiments, including any relevant details; confirming that all experiments were performed in accordance with relevant guidelines and regulations. Informed consent was waived by institutional board of Xiamen children's hospital as retrospective study is performed.

## Results

### 1. **General information**

Between March 2017 and July 2020, 14 cases of children with MFBM were admitted to Xiamen Children's Hospital, including 12 males and 2 females. The ratio of urban to rural areas is 8:6, and the ratio of males to females is 6:1. The age span is between 1.2 years old and 8.9 years old, and the median age is 4.6 years old. The number of ingested magnetic foreign bodies ranged from 1 to 17, with an average of 6.5. The properties of magnetic foreign bodies are divided into 3 cases of magnets and 11 cases of magnetic beads.

### 2. **Clinical symptoms**

The main manifestations of 14 children were abdominal pain, bloating, vomiting, fever, etc. Among them, 6 cases were complicated by intestinal obstruction and intestinal perforation (1-5 locations). Some of them had no clinical symptoms due to the short time of ingesting foreign bodies. Of the 14 cases, 9 cases were able to provide a history of ingesting foreign bodies at the time of medical visit, and the other 5 children and their families could not provide a history of ingesting foreign bodies. The clinical characteristics of these children without history of MFBM are shown in Table 1 and Table 2 showed clinical characteristics of these children with traceable history of MFBM. The cases in group 1 had a complicated clinical course and longer hospital stay (median 13 days, range 10-22 days) than those in group 2 (median 3 days, range 2-12 days) as shown in Table 3.

### 3. **Image analysis:**

(1). The abdominal X-rays of all 14 children (Figure 1) clearly showed foreign bodies in the digestive tract. In cases No. 5 and 9 it appeared that the location of the magnetic foreign body was in the stomach. However, subsequent gastroscopy showed there was no foreign body in the stomach. The mis-ingested magnetic foreign bodies showed various shapes of stacks of magnets.

(2). The abdominal sonography before surgery did not indicate the presence of foreign bodies in the digestive tract in cases No. 1, No. 2, 5, 7, and 8. Neither had it showed magnetic foreign bodies in case 1 who had sought medical help in another hospital prior to arrival at XMCH.

#### 4. **Management and outcome**

Gastroscopy was performed on 2 cases (case 5 and case 9) but failed to show the magnets in the stomach. In case 5, laparotomy was performed to retrieve the magnets. While in case 9, conservative care was made and spontaneous discharge occurred 3 days later. Seven of the 14 cases underwent surgical treatment to remove foreign bodies (including 4 cases of open surgery and 3 cases of laparoscopic surgery); of the 7 cases underwent operation, 6 cases had complicated intestinal obstruction and intestinal perforation (at least 1 intestinal perforation, at most 5 Place). Two of the 14 cases showed that the location of the foreign body was in the stomach by the abdominal X-ray, and the gastroscopy did not show the foreign body in the stomach. The proportion of laparoscopy in the 7 surgical cases is nearly half, including case 2 who initially had laparoscopy but subsequently switched to open laparotomy suggesting that laparoscopic exploration can be used as the preferred treatment for intestinal perforation and/or intestinal obstruction caused by magnetic foreign bodies in the digestive tract. All surgical cases recovered smoothly after treatment, and no complications occurred during the longest follow-up period of six months.

## **Discussion**

Foreign bodies in the digestive tract are not uncommon in children. In recent years, related reports have gradually increased, especially the reports of magnetic foreign bodies. This is related to the wide application of magnetic materials in children's toys. Among the 14 cases in this group, 12 were boys and 2 were girls. The ratio of urban to rural areas was 8:6. The age span was between 1.2 and 8.9 with a median of 4.6. The proportion of boys is much higher than that of girls, accounting for 85.7%, which is related to boys' active and naughty nature; in terms of age proportions, there are 3 infants, 9 preschoolers, 2 schooler children, indicating 64% preschoolers. This is probably related to the strong curiosity of preschool children, the lack of relevant safety knowledge, and the weak sense of alertness of parents/guardians.

Compared with other foreign bodies, mis-ingestion of magnetic foreign bodies in the digestive tract is unique and need to be paid attention to. In contrast to other kind of foreign bodies, the magnetic foreign bodies (such as magnets, magnetic beads) are easily attracted to each other (if the mis-ingested number is 2 or more) in the digestive tract, as reflected in the abdominal X-rays of these patients. The aggregation of magnets may thus cause the compression of the luminal wall of the gastrointestinal tract resulting in ischemic necrosis, which can cause single or multiple intestinal perforation, intestinal obstruction, peritonitis, intussusception, and hemorrhage<sup>4-6</sup>. In severe cases, it can lead to sepsis, septic shock and life-threatening.<sup>3</sup>; Of the 14 cases in this series, 2 cases ingested single magnetic foreign body while the remaining 12 cases mis-ingested multiple magnetic foreign bodies, which accounted for as high as

85.7% among the 14 children with MFBM. This is a higher proportion compared to the 67% of children mis-ingested multiple magnets or additional metal foreign bodies reported by Sola et al.<sup>1</sup> Therefore, in view of the occurrence rate of multiple magnetic foreign bodies and its possible serious complication, the proportion of surgical operations is higher than other kind foreign bodies. Compared with the damage caused by multiple magnetic foreign bodies, a single magnetic foreign body can often be expelled along with feces by itself, and the outcome is relatively better.

In this group of 14 children, 9 cases clearly provided medical history at the time of medical visit, 5 (35.7%) cases failed to provide adequate medical history, and all these 5 cases had intestinal obstruction and intestinal perforation (1–5 locations). One case even complicated with sepsis. Thus, the clinical medical history often is unclear for foreign body mis-ingestion in children. In view of the possible serious complications for children with MFBM, it is therefore particularly important that for children with unexplained abdominal symptoms, foreign body mis-ingestion, particularly magnetic foreign body mis-ingestion should be considered because the children who mis-ingested foreign body often could not express the fact and abdominal plain X-ray should be taken accordingly.

Abdominal sonography, although with many advantages such as non-radiation, high portability, and non-invasiveness etc., may not be helpful in children with MFBM. In our series, 4 of them underwent abdominal sonography in our hospital and another 1 had abdominal sonography in other hospital prior to admission to our hospital. In all 5 cases the abdominal sonography failed to obtain positive information. Therefore, until more experience is accumulated, abdominal sonography is not considered for diagnosis of MFBM in children.

There were 2 children underwent abdominal X-ray examination which indicated that the foreign body was in the stomach, but the emergency gastroscopy results showed that there was no foreign body in the stomach. This might be due to time lag between the abdominal X-ray and gastroscopy or due to an orthogonal plane of abdominal X-ray may be needed to localize the exact location of the magnetic foreign body. In another case, the abdominal X-ray (anterior position) showed the illusion of "1 magnet" at the time of presentation, but the lateral radiograph showed "2 magnets", again suggesting that at least orthogonal X-ray examination should be performed for cases with mis-ingestion of magnetic or metal foreign bodies in the digestive tract. Kircher et al.<sup>10</sup> have reported related similar situations.

There is currently no consensus on how to deal with a single mis-ingested magnetic foreign body. At present, Clinicians generally agree that conservative observation and dynamic follow-up of abdominal X-rays. Most single magnetic foreign bodies can be expelled with feces by themselves. Two of the 14 cases in this series mis-ingested single magnetic foreign bodies, which were all conservatively managed. The magnetic foreign bodies were eventually discharged through the anus on their own without complications such as intestinal perforation or obstruction. Scholar Sun Jun et al.<sup>11,12</sup> suggested that children who accidentally ingested multiple magnets should be actively treated with surgery, regardless of their clinical manifestations. Among the 14 cases in this series, multiple magnetic foreign bodies accounted for 12 cases, but 7 cases were removed by surgery and 2 cases were removed by gastroscopy. In this group of

cases, 6 out of 7 undergoing surgery showed complicated intestinal perforation (the perforation sites were up to 5), but in all cases of intestinal perforation, the abdominal X-ray did not show obvious free gas under the diaphragm. However, it was confirmed that the gastrointestinal perforation was present, all with multiple site of perforation, during the operation. This is probably due to the surrounding effect by adhesive tissue. Therefore, it should be noted that the absence of free gas does not rule out absence of intestinal perforation. Three cases were managed conservatively.

Intestinal perforation caused by multiple magnetic foreign bodies is often located in multiple locations, scattered and hidden. Therefore, regardless of the choice of laparotomy or laparoscopic exploration, the entire digestive tract (from stomach to rectum) should be explored to ensure that no digestion is missed. Tract perforation and no missing magnetic foreign body, if necessary, X-ray should be rechecked during the operation to avoid undetected magnet(s); in this group of cases, there are 3 cases of multiple foreign bodies (the number of foreign bodies is 4, 8, 11) foreign bodies were expelled spontaneously after conservative management. Among them, the child who mis-ingested 11 magnetic foreign bodies expelled them in 13 days.

In addition, a total of 4 patients had clinical followed up, the follow-up time was 1 month, 1 month, 5 months, and 6 months. The remaining cases were not followed up due to reasons such as residence. All these cases had all paroxysmal abdominal pain, the degree and frequency of abdominal pain varied; there was no accompanied vomiting, diarrhea, and intestinal obstruction. The abdominal pain relieved spontaneously. The follow-up radiologic imaging all showed no obvious abnormal lesions.

Based on this series, we thus suggest a protocol in handling mis-ingested multiple magnetic foreign bodies is as follows: 1. Children with multiple MFBM within 4 hours and confirmed their presence in the stomach region on the orthogonal abdominal X-ray, an emergent gastroscopy should be done immediately /as soon as possible to decrease the time interval for gastroscopy. 2. For children who mis-ingest multiple magnetic foreign bodies or for a long time (more than 4 hours) or accompanied by abdominal pain, vomiting, or bloating, fever, etc., we recommend emergency surgery. The indications for surgery cannot be judged solely on the results of abdominal X-ray examination and other signs of intestinal perforation, such as free air under the diaphragm. Regarding the choice of surgical methods, laparotomy or laparoscopic exploration can be selected according to the situation of the child and surgeon's preference. Laparoscopy may be attempted first. 3. For children who mis-ingested multiple magnetic foreign bodies, but the foreign body has passed through the stomach and duodenum and cannot be removed by the gastroscope, present no abdominal pain or vomiting, no abdominal distension, no fever and no other symptoms, and show normal stools, and abdominal X-rays did not indicate intestinal perforation, intestinal obstruction, intestinal motility changes, can continue to observe and follow-up.

In summary, pediatricians/emergency physicians should be vigilant about the hazards caused by mis-ingestion of magnetic foreign bodies in children. They should make great effort to make differential diagnosis among children with gastrointestinal symptoms and to recognize magnetic foreign bodies as

early as possible, actively manage them, reduce complications, and strengthen publicity. For magnetic foreign bodies in the digestive tract, abdominal X-ray examination has a higher diagnostic value than ultrasound, but additional orthogonal plane may be attempted to better localize the location of the magnetic foreign bodies in the abdomen. Also, the management planning should be judged according to the number and size of the foreign body, the number of mis-ingestion magnets and the condition of the children. For children who mis-ingested multiple magnetic foreign bodies, or mis-ingested it multiple times, and who presented with intestinal obstruction or pneumoperitoneum, active surgical operations are required. For management, laparoscopy may be the first choice.

Our experience shows we encountered 14 cases of MFBM in about 2.5 years. To the best of our knowledge, this is the largest series of MFBM in a single center in such a short time span. Thus MFBM is really a big issue for the small children! In view of the harm caused by magnetic foreign bodies in the digestive tract, we hereby call for:(1) Children's parents/ guardians and the whole society should be repeatedly educated through multiple channels to let them alert the seriousness of the accidental MFBI, strengthen guardianship, take responsibility on site, and reduce the incidents of MFBM; (2) Currently, Many countries probably have various existing laws/regulations on toy production and quality, but in fact there are deficiencies to earnestly inform toy risks and related conditions and to implement more effective measures to prevent children, especially small children from being exposed to the risk of MFBM, It is a serious big global issue on the further focus of relevant laws and regulations to strictly supervise small children's magnetic toys and eliminate them from the source; the family and society's awareness of the risk of accidental ingestion of magnetic materials should always be emphasized.

## **Declarations**

### **Author contributions:**

YKH has made substantial contributions to the conception of the work and drafted the work; approved the submitted version the acquisition; agreed both to be personally accountable for the author's own contributions and to ensure that questions related to the accuracy or integrity of any part of the work

SXH has made substantial contributions to the conception, acquisition, analysis, and interpretation of data of the work. approved the submitted version the acquisition; agreed both to be personally accountable for the author's own contributions and to ensure that questions related to the accuracy or integrity of any part of the work

IHT has substantively revised the manuscript; approved the submitted version the acquisition; agreed both to be personally accountable for the author's own contributions and to ensure that questions related to the accuracy or integrity of any part of the work

KSH has made substantial contributions to the conception; the acquisition, analysis, and interpretation of data; approved the submitted version the acquisition; agreed both to be personally accountable for the

author's own contributions and to ensure that questions related to the accuracy or integrity of any part of the work

## Additional Information (including a Competing Interests Statement)

Competing Interests Statement: The author(s) declare no competing interests.

## References

1. Ying Fang. Endoscopic treatment of foreign bodies in the digestive tract of children. *Chinese Journal of Digestive Endoscopy*. **34** (2), 80–82 (2017).
2. Tang, Y. & Xu, J. Yuanjun Hu, Hefeng Wang. Perforation of gastrointestinal tract caused by multiple magnetic foreign bodies. *Chinese Journal of Digestive Endoscopy*. **36** (1), 54–56 (2019).
3. Cox, S. *et al*. The risks of gastrointestinal injury due to ingested magnetic beads[J]. *S Afr Med J*, 2014, 104(4): 277–278.
4. Chen, C., Wang, A. & Chen, X. *et al*. Multiple intestinal perforation caused by a child eating a magnet by mistake. A case report. *Chinese Journal of Pediatric Surgery*. **31** (1), 80 (2010).
5. Cox, S. *et al*. The risks of gastrointestinal injury due to ingested magnetic beads. *S Afr Med J*. **104** (4), 277–278 (2014).
6. Arslan, S., Basuguy, E. & *et al* Zeytun. H., Jejunoleal perforation and volvulus caused by multiple magnet ingestion. *Acta Clin Croat*, 2015, 54(1): 96–98.
7. Corduk, N. & Odabas, S. E. A Sarioglu-Buke. intestinal perforation caused by multiple magnet ingestion. *Afr J Paediatr Surg*. **11** (1), 84–86 (2014).
8. Fuzhong Liu, S. & Wang, Y. Chen. Intussusception caused by multiple magnets A case report. *Chinese Journal of Women and Children's Clinical Medicine (Electronic Edition)* 2012, 8(3): 395.
9. Corduk, N., Odabas, S. E. & Sarioglu-Buke, A. intestinal perforation caused by multiple magnet ingestion. *Afr J Paediatr Surg*. **11** (1), 84–86 (2014).
10. Sola, R. *et al*. Magnet Foreign Body Ingestion: Rare Occurrence but Big Consequences. *J Pediatr Surg*, 2018, 53(9): 1815–1819.
11. Kircher, M. F., Milla, S. & Callahan, M. J. Ingestion of magnetic foreign bodies causing multiple bowel perforations. *Pediatr Radiol*, 2007, 37(9): 933–936.
12. Sun, J. *et al*. of multiple magnetic foreign bodies in the gastrointestinal tract. Reports of 3 cases and literature review. *Chinese Journal of Applied Pediatrics and Clinical Medicine*. **31** (17), 1339–1342 (2016).
13. Yan, X. P. *et al*. Operation or non-operation: strategy of managing multi-magnets in digestive tract. *Int J Colorectal Dis*, 2016, 31(4): 945–946.

# Tables

Due to technical limitations, tables are only available as a download in the Supplemental Files section.

# Figures




Figure 1

Abdominal X-rays of the 14 children with different number and different locations of the mis-ingested magnets/magnetic beads (please refer to table 1 and table 2 in detail)

## Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Table1.png](#)
- [Table2.png](#)
- [Table3.png](#)