

# Dietary Habits and its Association with Metabolic Syndrome in a large sample of Iranian adults: a population-based study

**Elnaz Lorzadeh**

Shahid Sadoughi University of Medical Sciences and Health Services

**Zohreh Sadat Sangsefidi**

Shahid Sadoughi University of Medical Sciences and Health Services

**Masoud Mirzaei**

Shahid Sadoughi University of Medical Sciences and Health Services

**Mahdieh Hosseinzadeh** (✉ [hoseinzade.mahdie@gmail.com](mailto:hoseinzade.mahdie@gmail.com))

<https://orcid.org/0000-0001-7482-2494>

---

## Research article

**Keywords:** Metabolic Syndrome, Dietary habits, Diet

**Posted Date:** May 3rd, 2020

**DOI:** <https://doi.org/10.21203/rs.3.rs-25601/v1>

**License:**   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

# Abstract

**Background** Central obesity, insulin resistance, dyslipidemia and hypertension are the core components of metabolic syndrome (MetS) which is coincident with unhealthy dietary habits in the Middle-Eastern countries. The aim of this study was to explore the association between dietary habits and MetS of adult population living in Yazd Greater Area, Iran.

**Methods** This is a cross-sectional study which uses the data of a population-based cohort study on Iranian adults, known as Yazd Health Study (YaHS). The relationship between dietary habits and metabolic syndrome among adults ( $n = 2896$ ) were analyzed using multiple logistic regression method.

**Results** The prevalence of the MetS among the participants was 32.2%. Outcomes from logistic regression examining show that breakfast consumption has a significant inverse effect on the occurrence of MetS after adjustment for age, education level, physical activity status, history of chronic diseases and smoking (odds ratio (OR) = 0.38, 95% confidence interval (CI) = 0.14, 0.97). This effect remains significant even after adjustment for body mass index (BMI) and reveals that odds of having MetS is 69% lower in breakfast consumers in contrast to non-consumers (OR = 0.31, 95% CI = 0.11, 0.87). However, no significant relationship was observed between other dietary habits including consumption of sweetened drinks, sugar cubes, and fast foods and MetS after adjustment for all potential confounders

**Conclusions** This study revealed that eating breakfast has an inverse relationship with metabolic syndrome. Finding out stronger evidence in relation between dietary habits and MetS, more researches especially population-based cohort studies are needed to be conducted.

## Background

With increased risk of diabetes mellitus (DM) and cardiovascular disease (CVD), which are mostly derived from metabolic syndrome (MetS) [1], urgent strategies to prevent its global epidemic are demanded [2]. Central obesity, insulin resistance, dyslipidemia and hypertension are the core components of metabolic syndrome [3]. According to the international diabetes federation (IDF), the rapid increase of MetS has been estimated to be 20–25% globally [4] which is parallel to the growing rate of DM, hypertension, cardiovascular disease and obesity [2]. The prevalence of MetS in different studies varies, depending majorly on the used criteria of different definitions and their compositions (such as sex, age, race and ethnicity) [5]. For example results of recent systematic review and meta-analysis of studies revealed the overall estimation of MetS prevalence in Iran to be 36.9% based on the Adult Treatment Panel guidelines III (ATP III) criteria and 34.6% according to IDF [6, 7]. However, regardless of the criteria used, it has been reported that the prevalence of metabolic syndrome around the world is rising drastically [8, 9].

Unhealthy diet and lifestyle are some of the risk factors contributed to MetS [10, 11]. Several studies have been conducted to explore the relationship between dietary habits and MetS [12–19]. It has been reported that dietary habits that are more westernized can increase the chances of MetS [19], whereas diets rich in

Loading [MathJax]/jax/output/CommonHTML/jax.js ction effect.

Considering what has been mentioned above, emerging evidence reveals that higher consumption of sugar sweetened beverages consist of energy-containing sweeteners such as sucrose, high fructose corn syrup or fruit-juice concentrates [20] is associated with increasing risks of MetS [21] not only in the US diet [22] but also, in developing countries such as India and China [23]. These findings also have been confirmed by a meta-analyses of cohort studies [24].

Moreover, studies suggests that consumption of out-of-home meals and energy dense fast foods [25] and snacks which contain highly processed meat, total fat (saturated and trans fatty acids) and refined carbohydrates [26] as well as sodium, can cause postprandial metabolic disorders such as dyslipidemia, subclinical inflammatory and oxidative stress and thereby increase occurrence of MetS [27] [28, 29]. All of which lead to elevated fasting insulin levels and metabolic syndrome in adults [27]. In regard to breakfast habits and MetS, there are some controversial results. For example, those who habitually do not consume breakfast are at higher risk for skipping other meals, snacking, having a sedentary lifestyle and being obese or overweight [30, 31]. Eating breakfast is associated with higher overall diet quality [32] and since it is satiating, it can reduce the total energy intake of the day and cause beneficial impact on improving metabolic parameters [33, 34]. On the contrary, a study suggested that skipping breakfast alone did not have any significant association with MetS prevalence [35].

In despite previous studies that have investigated metabolic syndrome's association with dietary habits, there are only a limited number of them conducted in the Middle-eastern countries where nutritional transition to westernized eating habits has caused an increase in the prevalence of overweight and obesity. This study aims to explore the association between dietary habits and MetS of a large population of adults living in Yazd Greater Area in Iran.

## Materials And Methods

### Study population and data collection

Yazd Health Study (YaHS) recruitment phase data was used for this cross-sectional study. YaHS was initiated among the adult population living in Yazd Greater Area, with 10000 participants between the ages of 20-70 years old). Established in November 2014 and containing two phases, YaHS at first, randomly selected 200 clusters according to the city post codes. Then, the interviewers set up a meeting time with the residences at each of their assigned addresses. At last, as the interviewers visited the neighborhood of the first addressee based on the study protocol to fill in the questionnaire forms [36]. A validated questionnaire consists of 300 questions including the following sections were filled by trained interviewers; a) demographics, b) physical activity, c) sleep quality and quantity, d) mental health, e) past medical history of chronic disease and surgical operations, f) dental health, g) history of accidents, h) dietary habits, i) occupation and social life, j) traditional medicine, k) smoking habit and drug addiction, l) women's health and m) quality of life [36]. In the present research, out of 3748 available cases with data on dietary intakes and main variables associated with metabolic syndrome, 2896 participants were

considered according to exclusion criteria (including having a history of cardiovascular disease, diabetes and cancer).

Demographics data, anthropometrics (weight, height and waist circumference), socioeconomic status, education, physical activity, tobacco smoking and dietary habits in addition to biochemical assessments including; fasting blood glucose, triglyceride and HDL cholesterol levels were also carried out.

The present study was approved by the Ethics Committee of Shahid Sadoughi University of Medical Science (IR.SSU.MEDICINE.REC.1395.287). Written informed consents were obtained from all the participants.

In general, all the required information on participants' dietary habits, anthropometric indices measurements, blood test results, blood pressure measurement and confounding factors consisting of socioeconomic status, history of chronic diseases and physical activity status, were extracted and merged from YaHS database [37].

## **Dietary Assessment**

We aimed at obtaining accurate information on some dietary habits noted by participants via a questionnaire [37]. These items were as follows; i) sweetened drinks (fruit juices, artificially or homemade sweetened beverages) with frequency consumption of not at all, less than once per week, once or more per week, ii) fast foods consumption with frequency of not at all or few times per year, 1-3 times per months, once or more per week, iii) breakfast consumption as not at all in frequency, once per week or more than once per week, iv) sugar cubes with the serving intake of not at all, 1-2 cubes per day and more than 2 cubes per day.

## **Diagnosis of metabolic syndrome**

National Cholesterol Education Program and Adult Treatment Panel III criteria (NCEP) present the definition of MetS in the current study [38-40]. To be diagnosed with at least 3 risk factors out of the five following cases, means a participant has metabolic syndrome if: i) Triglyceride (TG) above 150 mg/dl or consuming triglyceride lowering agents (hyperglyceride); ii) high density lipoprotein-cholesterol (HDL-C) level of less than 40 mg/dl in men and less than 50 mg/dl in women or any kind of pharmacological treatment; iii) Systolic blood pressure above 130 mmHg and diastolic blood pressure above 85 mmHg (hypertension); iv) Fasting blood glucose above 100 mg/dl or usage of pharmacological treatment as a control from blood sugar; and v) and waist circumference (WC) above 91.5 cm in men and 85.5 cm in women (adopted for only Iranian population) [41].

Loading [MathJax]/jax/output/CommonHTML/jax.js

## Evaluation of anthropometric indices

Weight was recorded with participants wearing lightweight clothing and no shoes using a digital scale (Model BF511, Omron Co. Karada Body Scan, Osaka, Japan) with the precision of 0.1 kg. A non-stretchable tape meter with the precision of 0.5 cm was used to measure height with lightweight clothing and no shoes and with their heels, hip, shoulder and head touching the wall. Body mass index (BMI; weight (kg) / height (m<sup>2</sup>)) was calculated from measuring height and weight. Waist circumference was also measured by a tape in the horizontal plane midway between the iliac crest and the rib cage with the precision of 0.1 cm.

## Blood pressure measurement

Systolic and diastolic blood pressures were measured in the sitting position by an automatic digital blood pressure monitor (Reister Germany). Each measurement was repeated for two times after every 5-minute interval.

## Laboratory data

Data on serum level of fasting blood glucose (FBG) (mg / dl), total cholesterol, triglycerides, low density lipoprotein-cholesterol (LDL-C) and high density lipoprotein-cholesterol (HDL-C) were measured and collected. The equipment used in this study included a calibrated Ciba Corning auto-analyzer device (Ciba Corp., Basle, Switzerland (and Pars Azmoon company kits to assess fasting blood glucose and triglyceride after centrifuging and bionic kits as an analyzer for HDL cholesterol.

## Physical activity assessment

For ascertainment physical activities the Iranian Short Version of International Physical Activity Questionnaire (IPAQ) was given to the participants [42, 43]. Physical activity levels of each participant were categorized as active and inactive based on the guideline of short form of IPAQ [44].

## Statistical analysis

All statistical analyses were performed using SPSS version 24 (SPSS Inc., Chicago, IL, USA) and to show the qualitative variables, they went under a frequency and percentage discretionary. Multiple-logistic regression was used to examine the association between dietary habits and MetS. The lowest frequency or serving indicates a reference for all the models and adjustments of the confounding factors which are as follows: Age (20-29, 30-39, 40-49, 50-59, 60-69 years), educational level (secondary school and lower, bachelors, masters and PhD), history of chronic diseases

(yes/no, including : hypertension, diabetes, cardiovascular disease, cancer, depression, dyslipidemia), smoking history (yes/no), physical activity level (inactive and active), and BMI (continuous).

## Results

Table 1 shows the general characteristics of study population. A majority of participants (53.2%) as illustrated were females. Furthermore, a high percent of the participants (24.7%) were between ages of 40–49 years old and had secondary school or lower education (53.5%). The smoking status also demonstrates that a high percent of the volunteers did not smoke (90.0%) and were married (86.9%). Moreover, 51.8% of the participants were active on their physical activity statue.

Table 1  
Distribution of all participants according to general characteristics.

<b>Variables</b>	<b>Total (n = 2896) N (%)</b>
<b>Sex</b>	
Male	1352 (46.8%)
Female	1534 (53.2%)
<b>Age (years)</b>	
20–29	541 (18.8%)
30–39	639 (22.2%)
40–49	713 (24.7%)
50–59	560 (19.4%)
60–69	429 (14.9%)
<b>Education level</b>	
Secondary school and lower	1534 (53.5%)
Diploma and Graduate diploma	877 (30.6%)
Bachelors	389 (13.6%)
Masters and PhD	66 (2.3%)
<b>Smoking</b>	
Yes	185 (6.6%)
Sometimes	55 (2.0%)
Quit	43 (1.5%)
No	2533 (90.0%)
<b>Marital status</b>	
Married	2505 (86.9%)
Single	274 (9.5%)
Widowed	89 (3.1%)

<b>Variables</b>	<b>Total (n = 2896)</b>
	<b>N (%)</b>
Divorced	15 (0.5%)
<b>Physical activity</b>	1396 (48.2%)
Inactive	1500 (51.8%)
Active	

The prevalence of the metabolic syndrome among the participants was 32.2% as demonstrated in Table 2. Additionally, the plurality of the participants (38.6%) were overweight (BMI = 25-29.9). Moreover, 38.8% met the criterion for hypertriglyceridemia (plasma TG higher than 150 mg/dl) and 29.3% had FBG higher than 100 mg/dl, while 65.8% had low levels of HDL-C ( $\leq$ 40 mg/dl in men and  $\leq$ 50 mg/dl in women). Furthermore, abdominal obesity which is considered as waist circumference  $>$ 91.5 cm for men and 85.5 cm for women afflicted 65.1% of the study population. It was also found that 45.5% of the subjects suffered from high blood pressure ( $\geq$  130/85).

Table 2

Distribution of participants according to metabolic syndrome (MetS) and its components.

<b>Variables</b>	<b>Total (n = 2896) N (%)</b>
<b>BMI (kg/m<sup>2</sup>)</b>	
Low weight (< 18.5 )	80 (2.9%)
Normal (18.5–24.9)	863 (31.3%)
Overweight (25-29.9)	1062 (38.6%)
Obesity ( 30-39.9)	671 (24.4%)
Extreme Obesity ( $\geq 40$ )	77 (2.8%)
<b>Triglyceride status (mg/dl)</b>	
Lower than 150	1768 (61.2%)
Equal to or higher than 150	1119 (38.8%)
<b>Fasting Plasma glucose status (mg/dl)</b>	
Lower than 100	2044 (70.7%)
Equal to or higher than 100	849 (29.3%)
<b>Waist Circumference status (cm)</b>	
91.5 > for men and 85.5 > for women	1001 (34.9%)
91.5 $\leq$ for men and 85.5 $\leq$ for women	1870 (65.1%)
<b>HDL status (mg/dl)</b>	
40 $\ll$ / $>$ <i>cript</i> $>$ for men and 50 $\ll$ / $>$ <i>cript</i> $>$ for women	1900 (65.8%)
40 $>$ for men and 50 $>$ for women	986 (34.2%)
<b>Hypertension status (mmHg)</b>	
No	1571 (54.5%)
Yes	1312 (45.5%)
<b>MetS</b>	
No	1940 (67.8%)
Yes	920 (32.2%)

Furthermore, in Table 3 after adjusting the confounders, results of the logistic regression examining the association among those who consume breakfast for once per week compared to those who completely skip breakfast shows a significant relationship with odds of metabolic syndrome, which means that these subjects have 62% lesser chance of MetS (odds ratio (OR) = 0.38, 95% confidence interval (CI) = 0.14 to 0.97). This effect remains significant even after adjustment for BMI and reveals that odds of MetS is 69% lower in breakfast consumption as once per week in contrast to non-consumption (OR = 0.31, 95% CI = 0.11 to 0.87).

Table 3

Multivariable-adjusted odds ratios (95% CI) for metabolic syndrome across different frequencies or servings for various dietary habits in a sample of Iranian adults.

<sup>1</sup> Dietary habits	Metabolic Syndrom			
	<sup>2</sup> Multivariable adjusted		<sup>3</sup> Multivariable + BMI	
	OR	95% CI	OR	95% CI
<b>Sweetened drinks</b>				
Not at all	Reference		Reference	
Lower than once per week	0.93	0.71–1.22	0.97	0.74–1.28
Once or more per week	0.98	0.76–1.27	1.07	0.82–1.40
<b>Fast foods</b>				
Not at all or few times per year	Reference		Reference	
1–3 times per month	1.02	0.81–1.28	0.93	0.73–1.18
Once or more per week	0.92	0.62–1.36	0.88	0.58–1.31
<b>Breakfast</b>				
Not at all	Reference		Reference	
Once per week	0.38*	0.14–0.97*	0.31*	0.11–0.87*
More than once per week	0.74	0.40–1.37	0.36	0.36–1.32
<b>Sugar cubes</b>				
Not at all	Reference		Reference	
1–2 cubes per day	1.01	0.71–1.45	1.03	0.71–1.49
More than 2 cubes per day	1.05	0.76–1.45	1.09	0.78–1.52

<sup>1</sup> Dietary habits for all items were presented as frequency of consumption except for sugar cubes which presented as serving of intake.

<sup>2</sup> Adjusted for age (20–29, 30–39, 40–49, 50–59, 60–69 years), education level (Secondary school and lower, High school, Diploma and Graduate diploma, Bachelors, Masters and PhD), physical activity level (active and inactive), history of chronic diseases (hypertension, diabetes, cardiovascular disease, cancer, depression and dyslipidemia), smoking (yes/no).

<sup>3</sup> Adjusted for age (20–29, 30–39, 40–49, 50–59, 60–69 years), education level (Secondary school and lower, High school, Diploma and Graduate diploma, Bachelors, Masters and PhD), physical activity level (active and inactive), history of chronic diseases (hypertension, diabetes, cardiovascular disease, cancer, depression, and dyslipidemia), smoking (yes/no) and BMI.

\* .05.

Nevertheless, no significant association has been witnessed between other dietary habits including consumption of sweetened drinks, sugar cubes, and fast foods with MetS after adjusting for potential confounders.

## Discussion

In the current study, consumption of breakfast can significantly contribute to low chance of MetS. Conversely, consumptions of sweetened beverages, fast foods and sugar cubes have been shown to have non-significant impact on the occurrence of MetS. These relations between dietary habits and MetS were maintained after adjusting for potential cofounders such as age, education level; and physical activity level; history of chronic diseases; smoking and BMI.

Many studies suggest that eating breakfast can associate with an overall better diet quality and healthy lifestyle [45–47]. It has been also reported that individuals who eat breakfast frequently have a lower risk of an array of metabolic outcomes comparing to their peers who infrequently or never eat breakfast [48]. Many studies suggest that breakfast intake is beneficial to metabolic health which is in the same line as our findings [49, 50]. This, as provided by a spectrum of research evidence could be for the reason that eating breakfast plays an important role in factors related to appetite and glucose, insulin and lipid metabolism [50–52]. It has been discussed in some studies that poor breakfast habits can be a part of unhealthy lifestyle which predicts metabolic syndrome [53]. Several evidences suggest that breakfast intake beneficial effects depend on the type of breakfast [54–56]. It has also been demonstrated that healthy breakfast (such as fruit smoothie or milk breakfast) may influence subsequent dietary intake [57]. Iranian breakfast usually consists of bread, cheese or butter or egg and drinks such as tea and milk [58] which can have a positive effect on metabolic factors. As it has been reported in recent studies whole-grains [59] and low fat dairy [60] consumption as common components of healthy breakfast can be beneficial.

On the contrary, controversial outcomes report that a recommendation to eat or skip breakfast may have no effect on body weight [61]. It has been reported that breakfast intake can potentially improve energy balance along with insulin and glucose level thus leading to satiety and lesser weight [62]. Breakfast consumers are also more likely to have better overall dietary quality along with following dietary recommendations more than non-consumers [63]. Obesity and high fasting glucose, the components of the metabolic syndrome and higher levels of total cholesterol (TC) and LDL-C are also evident after skipping breakfast completely compared to regular breakfast eaters [64]. According to a population-based cohort study, MetS prevalence grows with the continuation of omitting breakfast or poor breakfast habits [53].

Furthermore, we found no association between consumptions of sweetened beverages and sugar cubes and MetS in our population. In consistent to our outcomes, a few studies suggest no significant association between sweetened beverages consumption and higher risk of metabolic syndrome and its components [65–67]. A prospective study also, showed that sweetened drinks consumption was not

associated with incident of metabolic syndrome in a middle-aged population [68]. This, however, is in contrast with several other studies that report sugar-sweetened drinks intake has significant effect on metabolic syndrome [69–71]. Sweetened beverages that were included in this study consist of artificially or homemade beverages. Sweetened beverages in Iran are usually of plant origin which contain beneficial compounds such as polyphenols which are inversely associate with MetS and its components [72]. This issue as well as different in cofounders adjusted in analysis could be a source of contradiction in this study and others, because some studies did not adjust for BMI which is an important variable in relation between dietary habits and MetS.

Additionally we could not find any significant association between fast foods intake and metabolic syndrome. In contrast with our findings, some studies suggest that regular fast foods consumption has irreparable effect on general health and can increase the risk of obesity, insulin resistance and other metabolic abnormalities [73–75]. However, there is not a definitive agreement on ‘fast foods’ definition [76] and differences in ingredients of the fast foods in the different studies could have caused discrepancies in the outcomes. Moreover, fast foods consumption in our study was generally low (most of the study population (60.6%) did not consume fast foods at all or the intakes were only few times per year.

This study contains a few limitations. First of all, the design of this study is cross-sectional which does not provide a liable source of causality. Second, the data on dietary assessments were based on self-reported questionnaires. And third, the impact of confounders could not be fully controlled due to unknown or unmeasured factors. Strength points suggest that this study investigates the impact of dietary habits on metabolic syndrome among a relatively large sample in a Middle Eastern country. A large extent of confounders that might affect the metabolic syndrome or its components have been also considered and controlled.

## Conclusions

As a conclusion, this study provides evidence that there is in fact an inverse relationship between breakfast intake and metabolic syndrome. No significant association has been seen between other dietary habits and MetS. However, to find out stronger evidence in relation between dietary habits and metabolic syndrome, more studies especially population-based cohort researches need to be conducted.

## Abbreviations

MetS: metabolic syndrome; YaHS: Yazd Health Study; OR: odds ratio; CI: confidence interval; DM: diabetes mellitus; CVD: cardiovascular disease; IDF: international diabetes federation; ATP III: Adult Treatment Panel guidelines III; NCEP: National Cholesterol Education Program and Adult Treatment Panel III; HDL-C: high density lipoprotein-cholesterol; TG: Triglyceride; WC: waist circumference; BMI: Body mass index; FBG: fasting blood glucose; LDL-C: low density lipoprotein-cholesterol; IPAQ: International Physical Activity Questionnaire; TC: total cholesterol

Loading [MathJax]/jax/output/CommonHTML/jax.js

# Declarations

## Ethics approval and consent to participate

The present research was approved by the Ethics Committee of Shahid Sadoughi University of Medical Science (IR.SSU.MEDICINE.REC.1395.287). Written informed consents were obtained from all the participants.

## Consent for publication

Not applicable

## Availability of data and material

None

## Competing interests

The authors declare that they have no conflict of interest.

## Funding

No funding was received regarding to the present study.

## Authors' contributions

1. M and M. H designed the study.
2. S. S and M. H conducted the statistical analysis.
3. S. S and E. L wrote the draft of manuscript.
4. M and M. H critically revised the manuscript and confirmed the final version of it to submit.

All authors read and approved the final version of manuscript.

## Acknowledgements

The authors appreciate Shahid Sadoughi University of Medical Sciences, Yazd, Iran to support this research.

## References

1. Grundy SM, Hansen B, Smith SC Jr, Cleeman JI, Kahn RA. Clinical management of metabolic syndrome: report of the American Heart Association/National Heart, Lung, and Blood Institute/American Diabetes Association conference on scientific issues related to management. *Arteriosclerosis, thrombosis, and vascular biology*. 2004;24:e19-24.
2. Zimmet P, Alberti KG, Shaw J. Global and societal implications of the diabetes epidemic. *Nature*. 2001;414:782–7.
3. Kirby MG, Wagg A, Cardozo L, Chapple C, Castro-Diaz D, De Ridder D, et al. Overactive bladder: is there a link to the metabolic syndrome in men? *Neurourol Urodyn*. 2010;29:1360–4.
4. O'Neill S, O'Driscoll LJO. Metabolic syndrome: a closer look at the growing epidemic and its associated pathologies. 2015;16:1–12.
5. Cornier M-A, Dabelea D, Hernandez TL, Lindstrom RC, Steig AJ, Stob NR, et al. The metabolic syndrome. 2008;29:777–822.
6. Dalvand S, Bakhshi E, Zarei M, Asl MT, Ghanei RJMSNJ. Prevalence of Metabolic Syndrome in Iran: A systematic review and meta-analysis. 2017;5:1–14.
7. Amirkalali B, Fakhrzadeh H, Sharifi F, Kelishadi R, Zamani F, Asayesh H, et al. Prevalence of metabolic syndrome and its components in the Iranian adult population: a systematic review and meta-analysis. 2015;17.
8. Aguilar M, Bhuket T, Torres S, Liu B, Wong RJJJ. Prevalence of the metabolic syndrome in the United States, 2003–2012. 2015;313:1973–4.
9. Prasad DS, Kabir Z, Dash AK, Das BC. Prevalence and risk factors for metabolic syndrome in Asian Indians: A community study from urban Eastern India. *Journal of cardiovascular disease research*. 2012;3:204–11.
10. Organization WH. Global action plan for the prevention and control of noncommunicable diseases 2013–2020. 2013.
11. Cox JL, Vallis TM, Pfammatter A, Szpilfogel C, Carr B, O'Neill BJJCJoC. A novel approach to cardiovascular health by optimizing risk management (ANCHOR): behavioural modification in primary care effectively reduces global risk. 2013;29:1400–7.
12. Azadbakht L, Mirmiran P, Esmailzadeh A, Azizi FJTAjocn. Dairy consumption is inversely associated with the prevalence of the metabolic syndrome in Tehranian adults–. 2005;82:523–30.

13. Elwood PC, Pickering JE, Fehily AMJJoE, Health C. Milk and dairy consumption, diabetes and the metabolic syndrome: the Caerphilly prospective study. 2007;61:695–8.
14. Esmailzadeh A, Kimiagar M, Mehrabi Y, Azadbakht L, Hu FB. Willett WCJTAjocn. Fruit and vegetable intakes, C-reactive protein, and the metabolic syndrome. 2006;84:1489–97.
15. Lutsey PL, Steffen LM, Stevens J. Dietary intake and development of the metabolic syndrome: The atherosclerosis risk in communities study. *Circulation: LIPPINCOTT WILLIAMS & WILKINS 530 WALNUT ST, PHILADELPHIA, PA 19106 – 3621 USA*; 2007. p. E220-E.
16. Panagiotakos DB, Pitsavos C, Skoumas Y, Stefanadis CJJotADA. The association between food patterns and the metabolic syndrome using principal components analysis. *The ATTICA Study*. 2007;107:979–87.
17. Ruidavets J-B, Bongard V, Dallongeville J, Arveiler D, Ducimetière P, Perret B, et al. High consumptions of grain, fish, dairy products and combinations of these are associated with a low prevalence of metabolic syndrome. 2007;61:810–7.
18. Snijder MB, van der Heijden AA, van Dam RM, Stehouwer CD, Hiddink GJ, Nijpels G, et al. Is higher dairy consumption associated with lower body weight and fewer metabolic disturbances? *The Hoorn Study*. 2007;85:989–95.
19. Wirfält E, Hedblad B, Gullberg B, Mattisson I, Andrén C, Rosander U, et al. Food patterns and components of the metabolic syndrome in men and women: a cross-sectional study within the Malmö Diet and Cancer cohort. 2001;154:1150–9.
20. Duffey KJ, Popkin BM. High-fructose corn syrup: is this what's for dinner? *The American journal of clinical nutrition*. 2008;88:1722 s-32 s.
21. Malik VS, Popkin BM, Bray GA, Despres JP, Hu FB. Sugar-sweetened beverages, obesity, type 2 diabetes mellitus, and cardiovascular disease risk. *Circulation*. 2010;121:1356–64.
22. Popkin BM. Patterns of beverage use across the lifecycle. *Physiol Behav*. 2010;100:4–9.
23. Bray GA. *The Battle of the Bulge: A History of Obesity*: Dorrance Publishing Company; 2007.
24. Malik VS, Popkin BM, Bray GA, Després J-P, Willett WC. Hu FBJDc. Sugar sweetened beverages and risk of metabolic syndrome and type 2 diabetes: a meta-analysis. 2010.
25. Bahadoran Z, Mirmiran P, Hosseini-Esfahani F, Azizi FJEjocn. Fast food consumption and the risk of metabolic syndrome after 3-years of follow-up: Tehran Lipid and Glucose Study. 2013;67:1303.
26. Esmailzadeh A, Azadbakht LJEjocn. Dietary energy density and the metabolic syndrome among Iranian women. 2011;65:598.
27. Mendoza JA, Drewnowski A, Christakis DAJDc. Dietary energy density is associated with obesity and the metabolic syndrome in US adults. 2007;30:974–9.
28. Devaraj S, Wang-Polagruto J, Polagruto J, Keen CL, Jialal IJM. High-fat, energy-dense, fast-food–style breakfast results in an increase in oxidative stress in metabolic syndrome. 2008;57:867–70.
29. Marangoni F, Martini D, Scaglioni S, Sculati M, Donini LM, Leonardi F, et al. Snacking in nutrition and health. *Int J Food Sci Nutr*. 2019;70:909–23.

30. Rampersaud GC, Pereira MA, Girard BL, Adams J, Metz J. Breakfast habits, nutritional status, body weight, and academic performance in children and adolescents. *Journal of the American Dietetic Association*. 2005;105:743 – 60; quiz 61 – 2..
31. Utter J, Scragg R, Mhurchu CN, Schaaf D. At-home breakfast consumption among New Zealand children: associations with body mass index and related nutrition behaviors. *J Am Diet Assoc*. 2007;107:570–6.
32. Carson TA, Siega-Riz A, Popkin B. importance of breakfast meal type to daily nutrient intake: differences by age and ethnicity. *Cereal Foods World*. 1999.
33. Ruxton C, Kirk TR. Breakfast: a review of associations with measures of dietary intake, physiology and biochemistry. *Br J Nutr*. 1997;78:199–213.
34. de Castro JM. The time of day of food intake influences overall intake in humans. *J Nutr*. 2004;134:104–11.
35. Kutsuma A, Nakajima K, Suwa K. Potential Association between Breakfast Skipping and Concomitant Late-Night-Dinner Eating with Metabolic Syndrome and Proteinuria in the Japanese Population. *Scientifica*. 2014;2014:253581-.
36. Mirzaei M, Salehi-Abargouei A, Mirzaei M, Mohsenpour MA. Cohort Profile: The Yazd Health Study (YaHS): a population-based study of adults aged 20–70 years (study design and baseline population data). *International journal of epidemiology*. 2018;47:697-8 h.
37. Mirzaei M, Salehi-Abargouei A, Mirzaei M, Mohsenpour M. Cohort Profile: The Yazd Health Study (YaHS): a population-based study of adults aged 20–70 years (study design and baseline population data). *International journal of epidemiology*. 2017.
38. Chackrewarthy S, Gunasekera D, Pathmeswaren A, Wijekoon C, Ranawaka U, Kato N, et al. A Comparison between Revised NCEP ATP III and IDF Definitions in Diagnosing Metabolic Syndrome in an Urban Sri Lankan Population: The Ragama Health Study.
39. Grundy SM, Cleeman JI, Daniels SR, Donato KA, Eckel RH, Franklin BA, et al. Diagnosis and management of the metabolic syndrome: an American Heart Association/National Heart, Lung, and Blood Institute scientific statement. *Circulation*. 2005;112:2735–52.
40. Huang PL. A comprehensive definition for metabolic syndrome. *Dis Models Mech*. 2009;2:231–7.
41. Esteghamati A, Ashraf H, Rashidi A, Meysamie A. Waist circumference cut-off points for the diagnosis of metabolic syndrome in Iranian adults. *Diabetes Res Clin Pract*. 2008;82:104–7.
42. Hu FB. Dietary pattern analysis: a new direction in nutritional epidemiology. *Curr Opin Lipidol*. 2002;13:3–9.
43. Moghaddam MB, Aghdam FB, Jafarabadi MA, Allahverdipour H, Nikookheslat SD, Safarpour S. The Iranian Version of International Physical Activity Questionnaire (IPAQ) in Iran: content and construct validity, factor structure, internal consistency and stability. *World applied sciences journal*. 2012;18:1073–80.
44. Committee IR. Guidelines for data processing and analysis of the International Physical Activity

45. Ruxton CH, Kirk TR. Breakfast: a review of associations with measures of dietary intake, physiology and biochemistry. *Br J Nutr.* 1997;78:199–213.
46. de Castro JM. The time of day of food intake influences overall intake in humans. *J Nutr.* 2004;134:104–11.
47. Song WO, Chun OK, Obayashi S, Cho S, Chung CE. Is consumption of breakfast associated with body mass index in US adults? *J Am Diet Assoc.* 2005;105:1373–82.
48. Odegaard AO, Jacobs DR, Steffen LM, Van Horn L, Ludwig DS, Pereira MA. Breakfast Frequency and Development of Metabolic Risk. *Diabetes Care.* 2013;36:3100–6.
49. Cayres SU, Junior IF, Barbosa MF, Christofaro DG, Fernandes RA. Breakfast frequency, adiposity, and cardiovascular risk factors as markers in adolescents. *Cardiol Young.* 2016;26:244–9.
50. Odegaard AO, Jacobs DR Jr, Steffen LM, Van Horn L, Ludwig DS, Pereira MA. Breakfast frequency and development of metabolic risk. *Diabetes Care.* 2013;36:3100–6.
51. Klok MD, Jakobsdottir S, Drent ML. The role of leptin and ghrelin in the regulation of food intake and body weight in humans: a review. *Obesity reviews: an official journal of the International Association for the Study of Obesity.* 2007;8:21–34.
52. Schlundt DG, Hill JO, Sbrocco T, Pope-Cordle J, Sharp T. The role of breakfast in the treatment of obesity: a randomized clinical trial. *Am J Clin Nutr.* 1992;55:645–51.
53. Wennberg M, Gustafsson PE, Wennberg P, Hammarström A. Poor breakfast habits in adolescence predict the metabolic syndrome in adulthood. *Public Health Nutr.* 2015;18:122–9.
54. Cho S, Dietrich M, Brown CJ, Clark CA, Block G. The effect of breakfast type on total daily energy intake and body mass index: results from the Third National Health and Nutrition Examination Survey (NHANES III). *J Am Coll Nutr.* 2003;22:296–302.
55. Deshmukh-Taskar PR, Radcliffe JD, Liu Y, Nicklas TA. Do breakfast skipping and breakfast type affect energy intake, nutrient intake, nutrient adequacy, and diet quality in young adults? NHANES 1999–2002. *J Am Coll Nutr.* 2010;29:407–18.
56. O'Neil CE, Nicklas TA, Fulgoni VL III. Nutrient intake, diet quality, and weight/adiposity parameters in breakfast patterns compared with no breakfast in adults: National Health and Nutrition Examination Survey 2001–2008. *Journal of the Academy of Nutrition Dietetics.* 2014;114:27–43.
57. McCartney D, Langston K, Desbrow B, Khalesi S, Irwin C. The influence of a fruit smoothie or cereal and milk breakfast on subsequent dietary intake: a pilot study. *Int J Food Sci Nutr.* 2019;70:612–22.
58. Ahadi Z, Kelishadi R, Qorbani M, Zahedi H, Aram M, Motlagh ME, et al. Association of Breakfast Intake with Psychiatric Distress and Violent Behaviors in Iranian Children and Adolescents: The CASPIAN- IV Study. *Indian J Pediatr.* 2016;83:922–9.
59. Tieri M, Ghelfi F, Vitale M, Vetrani C, Marventano S, Lafranconi A, et al. Whole grain consumption and human health: an umbrella review of observational studies. *International Journal of Food Sciences and Nutrition.* 2020:1–10.

60. Godos J, Tieri M, Ghelfi F, Titta L, Marventano S, Lafranconi A, et al. Dairy foods and health: an umbrella review of observational studies. *Int J Food Sci Nutr*. 2020;71:138–51.
61. Dhurandhar EJ, Dawson J, Alcorn A, Larsen LH, Thomas EA, Cardel M, et al. The effectiveness of breakfast recommendations on weight loss: a randomized controlled trial. *Am J Clin Nutr*. 2014;100:507–13.
62. Freitas Júnior IF, Christofaro DGD, Codogno JS, Monteiro PA, Silveira LS, Fernandes RA. The Association between Skipping Breakfast and Biochemical Variables in Sedentary Obese Children and Adolescents. *The Journal of Pediatrics*. 2012;161:871–4.
63. Rampersaud GC, Pereira MA, Girard BL, Adams J, Metz J. Breakfast habits, nutritional status, body weight, and academic performance in children and adolescents. *J Am Diet Assoc*. 2005;105:743–60.
64. Farshchi HR, Taylor MA, Macdonald IA. Beneficial metabolic effects of regular meal frequency on dietary thermogenesis, insulin sensitivity, and fasting lipid profiles in healthy obese women. *Am J Clin Nutr*. 2005;81:16–24.
65. Johnson L, Mander AP, Jones LR, Emmett PM, Jebb SA. Is sugar-sweetened beverage consumption associated with increased fatness in children? *Nutrition*. 2007;23:557–63.
66. Valente H, Teixeira V, Padrao P, Bessa M, Cordeiro T, Moreira A, et al. Sugar-sweetened beverage intake and overweight in children from a Mediterranean country. *Public Health Nutr*. 2011;14:127–32.
67. Khosravi-Boroujeni H, Sarrafzadegan N, Mohammadifard N, Alikhasi H, Sajjadi F, Asgari S, et al. Consumption of sugar-sweetened beverages in relation to the metabolic syndrome among Iranian adults. *Obes Facts*. 2012;5:527–37.
68. Lutsey PL, Steffen LM, Stevens J. Dietary intake and the development of the metabolic syndrome: the Atherosclerosis Risk in Communities study. *Circulation*. 2008;117:754–61.
69. Rivera JA, Muñoz-Hernández O, Rosas-Peralta M, Aguilar-Salinas CA, Popkin BM, Willett WC. Consumo de bebidas para una vida saludable: recomendaciones para la población mexicana. *Boletín médico del Hospital infantil de México*. 2008;65:208–37.
70. Woodward-Lopez G, Kao J, Ritchie L. To what extent have sweetened beverages contributed to the obesity epidemic? *Public Health Nutr*. 2011;14:499–509.
71. Malik VS, Schulze MB, Hu FB. Intake of sugar-sweetened beverages and weight gain: a systematic review–. *Am J Clin Nutr*. 2006;84:274–88.
72. Sohrab G, Hosseinpour-Niazi S, Hejazi J, Yuzbashian E, Mirmiran P, Azizi F. Dietary polyphenols and metabolic syndrome among Iranian adults. *Int J Food Sci Nutr*. 2013;64:661–7.
73. Rudolph TK, Ruempler K, Schwedhelm E, Tan-Andresen J, Riederer U, Böger RH, et al. Acute effects of various fast-food meals on vascular function and cardiovascular disease risk markers: the Hamburg Burger Trial. *Am J Clin Nutr*. 2007;86:334–40.
74. Pereira MA, Kartashov AI, Ebbeling CB, Van Horn L, Slattery ML, Jacobs DR Jr, et al. Fast-food habits, weight gain, and insulin resistance (the CARDIA study): 15-year prospective analysis. *The lancet*. 2005;365:36–42.

75. Duffey KJ, Gordon-Larsen P, Steffen LM, Jacobs DR Jr, Popkin BM. Regular consumption from fast food establishments relative to other restaurants is differentially associated with metabolic outcomes in young adults. *J Nutr.* 2009;139:2113–8.
76. Glanz K, Basil M, Maibach E, Goldberg J, Snyder D. Why Americans eat what they do: taste, nutrition, cost, convenience, and weight control concerns as influences on food consumption. *J Am Diet Assoc.* 1998;98:1118–26.