

Optimal Parity Cut-off Values for Predicting Postpartum Hemorrhage in Vaginal Deliveries and Cesarean Sections

Yasuhiro Miyoshi (✉ yasuhiro.miyoshi@gmail.com)

Research article

Keywords: Zambia, postpartum hemorrhage, risk factor, high parity, cut-off value

Posted Date: May 8th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-25706/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Pan African Medical Journal on January 1st, 2020. See the published version at <https://doi.org/10.11604/pamj.2020.37.336.24065>.

Abstract

Background High parity is a major public health concern in developing countries and it is a risk factor for postpartum hemorrhage (PPH). The aim of this study was to analyze the optimal parity cut-off values for predicting PPH in vaginal deliveries and cesarean sections in a rural Zambian setting.

Methods All women who delivered at Zimba Mission Hospital in 2017 were reviewed in this retrospective survey. Those whose records were missing data on parity and those with risk factors for developing PPH (e.g. birth weight $\geq 4,000$ g, multiple pregnancy, assisted vaginal delivery, and placenta previa) were excluded. We analyzed the association between parity and PPH using multiple logistic regression and ROC curve analyses.

Results Among the 1,555 women included in the study, 72 (4.6%) women experienced PPH. The optimal cut-off values for parity in vaginal deliveries and cesarean sections were para 7 and 3, respectively. Using these cut-off values, the adjusted odds ratios (95% confidence intervals) were 3.26 (1.15, 9.21) and 8.28 (2.25, 30.5), respectively.

Conclusions Proper preparation is required for vaginal deliveries in women with a history of ≥ 7 births and cesarean sections in women with a history of ≥ 3 births.

Background

Postpartum hemorrhage (PPH) is the leading cause of maternal mortality, accounting for 27.1% of all maternal deaths [1]. Although maternal mortality rates have declined greatly in developed countries, PPH remains a serious problem in developing countries, where more than 99% of maternal mortality due to PPH occurs [1]. Thirty-four percent of maternal mortality in Zambia is attributable to PPH [2].

An understanding of risk factors is important to properly prepare for deliveries. High parity is a risk factor for PPH [3–5]. The prevalence of grand multiparity (para 5 and above) [6] is as low as 3–4% in developed countries, while it is 19.3% in developing countries [7–9]. Associated factors include a high rate of unmet contraceptive needs, compromised or inadequate antenatal and delivery care, and low socio-economic status [10]. The national referral guideline in Zambia states that grand multiparous women should deliver in hospitals rather than health centers [11].

However, what qualifies as high parity, and whether para 5 is the optimal cut-off value remain to be elucidated. Additionally, no studies have investigated whether the cut-off value changes according to the mode of delivery.

The aim of this study was to analyze the optimal parity cut-off values for predicting PPH in vaginal deliveries and cesarean sections in a rural Zambian setting.

Methods

Study Design, Population, And Setting

This is a secondary analysis on data from a previously published study on PPH [12]. This retrospective cohort study was conducted at Zimba Mission Hospital in Southern Province, Zambia. This district hospital is located 400 km south of Lusaka, the capital of Zambia. It accepts patients referred from as many as 10 health centers in the catchment area, which has a population of 98,000. The study population included all women who delivered at the hospital between January 1 and December 31, 2017. Those whose records were missing data on parity and those with previously reported risk factors for developing PPH [13] (e.g. birth weight \geq 4,000 g, multiple pregnancy, assisted vaginal delivery [by vacuum or forceps]), and placenta previa were excluded.

Data Collection And Definitions

Demographic, clinical, and outcome data were extracted from the admission, delivery, and operation registers for all patients during the study period. Blood loss was estimated after delivery by the attending clinician (a midwife or doctor). PPH was defined as \geq 500 ml blood loss in the 24 h after vaginal delivery or \geq 1,000 ml blood loss in the 24 h after cesarean section [13].

Active management of the third stage of labor (AMTSL), consisting of the intramuscular injection of oxytocin (10 IU), controlled cord traction, and uterine massage, was performed for all patients who delivered vaginally. Additional oxytocin was given and bimanual uterine compression was performed in cases of PPH. Patients undergoing cesarean section also received oxytocin (10 IU). The management of the third stage of labor and PPH were based on the guideline proposed by University Teaching Hospital, Lusaka, Zambia [14].

Statistical analysis

Data were entered in Microsoft Excel (version 14.1.0; Microsoft®, Redmond, WA) and exported to EZR (version 3.1.2; Saitama Medical Center, Jichi Medical University, Saitama, Japan), which was used to perform the statistical analysis. The Mann Whitney U test was used to analyze continuous variables, and Pearson's chi-squared test was used to analyze qualitative variables. A multiple logistic regression analysis was performed to identify independent variables. Factors with p values $<$ 0.05 according to the Mann Whitney U test or Pearson's chi-squared test were entered into the multivariate analysis. P values of $<$ 0.05 were considered to indicate statistical significance. The Optimal parity cut-off values for predicting PPH in vaginal deliveries and cesarean sections were analyzed by multiple logistic regression and ROC curve analyses.

Results

A total of 1,704 women were reviewed and 1,555 women were included in the current study after excluding 149 cases for the following reasons: missing data on parity (n = 10), birth weight \geq 4,000 g (n =

47), multiple pregnancy (n = 47), assisted vaginal delivery (n = 40), and placenta previa (n = 9) (Fig. 1). The maternal and neonatal characteristics are listed in Table 1. Women with PPH tended to be older (p = 0.002), and have higher parity (p < 0.001). The rate of previous cesarean section (p = 0.017) and the cesarean section at the current pregnancy (p < 0.001) was higher in the PPH group. Birth weights tended to be greater in the PPH group (p = 0.010) (Table 1). The multiple logistic regression analysis demonstrated that cesarean section at the current pregnancy was the only factor significantly associated with PPH (p < 0.001) and the association between parity as a continuous variable and PPH lost significance after adjusting for confounders (Table 2).

Table 1
Maternal and neonatal characteristics

Characteristic	No PPH (n = 1,483)	PPH (n = 72)	p
Maternal age (years)*	24 (19–33)	30 (22–36)	0.002
Parity*	1 (0–4)	3 (1–6)	< 0.001
Previous cesarean section*	64 (4.3)	8 (11.1)	0.017
HIV seropositivity	73 (4.9)	4 (5.6)	> 0.999
HDP	32 (2.2)	3 (4.2)	0.474
Gestational age of delivery (weeks)	39 (37–40)	39 (38–40)	0.876
Cesarean section*	231 (15.6)	30 (41.7)	< 0.001
Birth weight (kg)*	3.05 ± 0.47	3.14 ± 0.58	0.010
Values are shown as the median (25th- 75th percentile) or number (percentage).			
PPH, postpartum hemorrhage; HIV, human immunodeficiency virus; HDP, hypertensive disorders of pregnancy.			
*p < 0.05			

Table 2
Multiple logistic regression analysis for PPH

Characteristic	Adjusted OR (95% CI)	<i>p</i>
Maternal age (years)	1.01 (0.96, 1.07)	0.697
Parity	1.13 (0.96, 1.34)	0.139
Previous cesarean section	1.26 (0.54, 2.97)	0.595
HIV seropositivity	1.02 (0.36, 2.95)	0.966
HDP	1.90 (0.55, 6.57)	0.311
Gestational age of delivery (weeks)	0.96 (0.86, 1.06)	0.378
Cesarean section*	3.83 (2.22, 6.61)	< 0.001
Birth weight (kg)	1.12 (0.64, 1.93)	0.697
PPH, postpartum hemorrhage; HIV, human immunodeficiency virus; HDP, hypertensive disorders of pregnancy; OR, odds ration; CI, confidence interval.		
* <i>p</i> < 0.05		

To investigate the optimal parity cut-off values for the prediction of subsequent PPH, we analyzed the association between parity and PPH separately in vaginal deliveries and cesarean sections. The incidence of PPH and the relative odds ratios (OR) for developing PPH were calculated (Tables 3 and 4). The statistical analysis revealed that, in vaginal deliveries, para ≥ 7 was associated with a significantly greater risk of PPH [Odds ratio 3.26; 95% confidence interval (95% CI) 1.15, 9.21; *p* = 0.025]. For para ≥ 7 , the positive predictive value (PPV), negative predictive value (NPV), sensitivity and specificity were 7.5%, 97.2%, 19.0% and 92.3%, respectively.

Table 3
 Incidences and odds ratios for PPH in vaginal deliveries in subgroups divided by different number of parity

Parity	Incidence of PPH (below vs. above)	Adjusted OR (95% CI)	<i>p</i>
1	3.5% vs. 3.1%	0.48 (0.19–1.23)	0.127
2	3.5% vs. 3.0%	0.38 (0.14–1.02)	0.055
3	2.9% vs. 3.8%	1.23 (0.45–3.36)	0.685
4	2.7% vs. 4.3%	2.05 (0.73–5.74)	0.172
5	2.9% vs. 4.4%	1.70 (0.62–4.65)	0.302
6	2.9% vs. 4.8%	1.69 (0.62–4.61)	0.305
7*	2.9% vs. 7.6%	3.26 (1.15–9.21)	0.025
8	3.2% vs. 4.0%	1.08 (0.23–5.04)	0.927
9	3.2% vs. 6.2%	1.77 (0.21–14.80)	0.598
10	3.5% vs. 0%	NA	NA
PPH, postpartum hemorrhage; OR, odds ration; CI, confidence interval; NA, not applicable.			
* <i>p</i> < 0.05			

Table 4
Incidences and odds ratios for PPH in cesarean sections in subgroups divided by different number of parity

Parity	Incidence of PPH (below vs. above)	Adjusted OR (95% CI)	<i>p</i>
1*	0.9% vs. 19.5%	28.8 (3.02–275)	0.004
2*	2.6% vs. 24.1%	10.0 (2.38–42.2)	0.002
3*	3.9% vs. 28.0%	8.28 (2.25–30.5)	0.001
4	7.2% vs. 23.9%	1.18 (0.36–3.90)	0.784
5	7.8% vs. 25.0%	1.06 (0.34–3.34)	0.921
6	9.4% vs. 23.7%	0.74 (0.24–2.24)	0.592
7	10.1% vs. 26.1%	0.76 (0.22–2.62)	0.660
8	10.9% vs. 23.1%	0.41 (0.07–2.33)	0.312
9	11.7% vs. 0%	NA	NA
10	11.6% vs. 0%	NA	NA
PPH, postpartum hemorrhage; OR, odds ratio; CI, confidence interval; NA, not applicable.			
* <i>p</i> < 0.05			

In cesarean sections, parity cut-off values of 1–3 were significantly associated with PPH. The ROC curve analysis showed that para 3 was the optimal cut-off value for predicting PPH with AUC of 0.787 (Fig. 2). Using para 3 as the cut-off value (OR 9.58; 95% CI 3.91, 23.5; *p* < 0.001), the PPV, NPV, sensitivity and specificity were 28.1%, 96.1%, 76.7% and 74.5%, respectively. All 4 cases of uterine rupture in patients without a history of cesarean section or uterine operation occurred in women with a history of ≥ 3 births (para 3, para 5, para 5 and para 6, respectively). All of these women were referred from health centers during labor and cesarean section was performed at the hospital.

Discussion

This study demonstrated that optimal parity cut-off values for predicting PPH in vaginal and cesarean section deliveries were para 7 and para 3, respectively.

To the best of our knowledge, this is the first study to demonstrate the optimal parity cut-off values for indicating the risk of PPH according to the mode of delivery. Previous studies [3–5] have shown an association between high parity and PPH. However these studies did not demonstrate the optimal cut-off

values. Some previous studies showed that grand multiparity was not associated with PPH [7, 10]. However this might be because para 5 is not the optimal cut-off.

As parity increase, a women's myometrial muscular strength may decrease due to a reduction of collagen fibers [5]. Therefore when parity increases, the probability of experiencing PPH increases. The discordancy of parity cut-off values between vaginal delivery and cesarean section indicates that cesarean section might affect the function of collagen fibers.

The association between parity as continuous variable and PPH lost significance after adjustment for confounders. However, parity as a categorical valuable, when using the specified cut-off value, was significantly associated with PPH. This suggests that the association between parity and PPH is not completely linear. Our result is consistent with a previous study in Australia [15].

The study findings suggest that healthcare workers should be well prepared to deal with the women with high parity. Several health professionals should attend vaginal delivery for women with a history of ≥ 7 births. Sufficient blood for transfusion should be prepared before performing cesarean section for the women with a history of ≥ 3 births because more blood loss is expected in comparison to vaginal delivery [12]. Due to the risk of uterine rupture, it is advisable for women with a history of ≥ 3 births to deliver at a hospital.

The present study was associated with some limitations. The diagnosis of PPH was based on estimated (rather than measured) blood loss. In addition, the gestational age data may not have been accurate because ultrasound is usually unavailable during early pregnancy and expected due dates are determined based on the last menstrual period, which is subject to memory bias. Parity may not have been completely accurate because Zambian medical practitioners tend to omit stillbirths and child deaths when calculating parity. Thus, parity may have been higher than documented in some cases. Furthermore, the registers did not contain data on the previous history of PPH or the body mass index, which prevented the evaluation of the effects of these known risk factors for PPH in the present study [16, 17]. Additionally, the socioeconomic factors of the women such as the place of residence, household income and number of prenatal checkups were not evaluated for the same reason. Finally, the data were collected at one district hospital. Thus, it might be difficult to generalize these data to the whole population of Zambia or other countries.

Conclusions

A history of ≥ 7 births in vaginal deliveries and ≥ 3 births in cesarean section deliveries was associated with PPH. During labor, providers caring for patients with these risk factors should be prepared to manage PPH.

Abbreviations

PPH, postpartum hemorrhage; HIV, human immunodeficiency virus; HDP, hypertensive disorders of pregnancy; OR, odds ratio; CI, confidence interval.

Declarations

Ethics approval and consent to participate

This study was approved by the research ethics committees of the University of Zambia Biomedical Research Ethics Committee (No. 001-03-19). The hospital administrator at Zimba Mission Hospital has granted permission to conduct the research after the ethical approval was obtained at an executive meeting. No ethical issue arose during this study, as it was retrospective and all data were anonymous.

Consent for publications

Not applicable

Availability of data and materials

The datasets during the current study are not publicly available to protect the participants' anonymity. But it can be freely available from the corresponding author on reasonable request.

Competing interests

The authors declare that they have no competing interests.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Authors' contributions

YM was involved in all processes of writing this paper.

Acknowledgements

Not applicable

References

1. Say L, Chou D, Gemmill A, Tunçalp Ö, Moller AB, Daniels J, et al. Global causes of maternal death: a WHO systematic analysis. *Lancet Glob Health*. 2014;2:e323-33.
2. Ministry of Community Development. Mother and Child Health, Ministry of Health. No Woman Should Die Giving Life Committing to Child Survival: A Promise Renewed. In: Roadmap for

- Accelerating Reduction of Maternal, Newborn and Child Mortality, 2013–2016. Ministry of Health; 2013.
3. Jaleel R, Khan A. Post-partum haemorrhage—a risk factor analysis. *Mymensingh Med J*. 2010;19(2):282–9.
 4. Habitamu D, Goshu YA, Zeleke LB. The magnitude and associated factors of postpartum hemorrhage among mothers who delivered at Debre Tabor general hospital 2018. *BMC Res Notes*. 2019;12(1):618.
 5. Gudeta TA, Kebede DS, Negeri GA, Dow MK, Hassen S. Magnitude of post-partum hemorrhage among women who received postpartum care at Bedele hospital south west, Ethiopia, 2018. *J Pregnancy Child Health*. 2018;5:396.
 6. Abu-Heija AT, Chalabi HE. Great grand multiparity: is it a risk? *J Obstet Gynaecol*. 1998;18(2):136–8.
 7. Muniro Z, Tarimo CS, Mahande MJ, Maro E, Mchome B. Grand multiparity as a predictor of adverse pregnancy outcome among women who delivered at a tertiary hospital in Northern Tanzania. *BMC Pregnancy Childbirth*. 2019;19(1):222.
 8. Births: Final Data for 2017
Martin JA, Hamilton BE, Osterman MJK, Driscoll AK, Drake P. Births: Final Data for 2017. *Natl Vital Stat Rep*. 2018; 67(8): 1–50.
 9. Al-Shaikh GK, Ibrahim GH, Fayed AA, Al-Mandeel H. Grand multiparity and the possible risk of adverse maternal and neonatal outcomes: a dilemma to be deciphered. *BMC Pregnancy Childbirth*. 2017;17(1):310.
 10. Ajong AB, Agbor VN, Simo LP, Noubiap JJ, Njim T. Grand multiparity in rural Cameroon: prevalence and adverse maternal and fetal delivery outcomes. *BMC Pregnancy Childbirth*. 2019;19:233.
 11. Republic of Zambia Ministry of Health. Zambia National Maternal and Neonatal Services Referral guidelines. Republic of Zambia Ministry of Health Web site. available at https://www.moh.gov.zm/?wpfb_dl=94. Accessed April 13, 2020.
 12. Miyoshi Y. Incidence, risk factors, treatment and outcomes of postpartum hemorrhage at a district hospital in Zambia. *Journal of International Health*. 2019;34(4):209–16.
 13. World Health Organization. WHO recommendations for the prevention and treatment of postpartum haemorrhage. WHO sexual and reproductive health. 2018.
 14. Department of Obstetrics & Gynaecology UT, Hospital, Lusaka, Zambia. Postpartum hemorrhage (PPH). In: *Obstetrics & Gynaecology Protocols and Guidelines*. Lusaka, Zambia. University of Zambia School of Medicine Medical Education Partnership Initiative; 2014.
 15. Bai J, Wong FW, Bauman A, Mohsin M. Parity and pregnancy outcomes. *Am J Obstet Gynecol*. 2002;186(2):274.
 16. Henry A, Birch MR, Sullivan EA, Katz S, Wang YA. Primary postpartum haemorrhage in an Australian tertiary hospital: a case-control study. *The Australian New Zealand Journal of Obstetrics Gynaecology*. 2005;45(3):233–6.

17. Sebire NJ, Jolly M, Harris JP, Wadsworth J, Joffe M, Beard RW, et al. Maternal obesity and pregnancy outcome: a study of 287,213 pregnancies in London. *International Journal of Obesity Related Metabolic Disorders*. 2001;25(8):1175–82.

Figures




Figure 1

Flow chart of the patients included in this study




Figure 2

Figure 2