

Primary and secondary aortopathy associated with adult congenital heart disease - retrospective study

Ingrid Schusterova

Vychodoslovensky Ustav Srdcovych a Cievnych Chorob AS

Alzbeta Banovcinova

Univerzita Pavla Jozefa Safarika v Kosiciach Lekarska fakulta

Marianna Vachalcova

Vychodoslovensky Ustav Srdcovych a Cievnych Chorob AS

Marta Jakubova

Vychodoslovensky Ustav Srdcovych a Cievnych Chorob AS

Panagiotis Artemiou (✉ panayiotisartemiou@yahoo.com)

Univerzita Komenskeho Lekarska fakulta

Research article

Keywords: primary/secondary aortopathy, aortic root, ascending aorta, congenital heart disease

Posted Date: May 11th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-25783/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published on September 10th, 2020. See the published version at <https://doi.org/10.1186/s13019-020-01283-z>.

Abstract

Background: Primary and secondary aortopathy are frequently encountered in patients with congenital heart disease. The aim of this study is to present our experience and the incidence of primary and secondary adult CHD-associated aortopathy.

Methods. The cohort is comprised of adult patients with congenital heart disease from the registry of the Eastern Slovakia Institute of Cardiovascular Diseases. Data from the last follow-up examinations are included in this study. In the primary and secondary aortopathy group were 35 and 12 patients respectively. As a control group were selected 64 patients with non aortopathy associated congenital heart disease (atrial and ventricular septal defect).

Results: Patients with primary and secondary aortopathy had larger ascending aorta/aortic root diameters than the control group (36.28 (26-49) mm vs 30.25 (21-41) mm $p=0.000113$, 33.82 (27-49) mm vs 29.03 (19-38)mm $p=0.000366$ and 42.1 (30-50) mm vs 30.25 (21-41) mm, $p=0.000106$, 35.67 (27-48) mm vs 29.03 (19-38) mm, $p=0.000119$ respectively). Moreover, patients with secondary aortopathy had statistically significant larger ascending aorta diameter compared to the patients with primary aortopathy (42.1 (30-50) mm vs 36.28 (26-49) mm $p=0.030$). During the follow-up period, were performed only in 2 patients (one from each group) operations on the aortic root and the ascending aorta due to aortic root or ascending aorta dilatation.

Conclusion: More patients with secondary aortopathy had dilated ascending aorta/ aortic root, as well as larger aortic diameters compare to the patients with primary aortopathy. Routine follow-up of these patients with attention to aortic diameter is necessary.

Background

Dilatation of the aortic root and the ascending aorta is frequently encountered in patients with congenital heart disease (CHD) at initial presentation and during follow-up.

Primary aortic dilatation is mainly associated with coarctation of the aorta (CoA), bicuspid aortic valve (BAV) and conotruncal abnormalities such as tetralogy of Fallot (TOF), pulmonary atresia with ventricular septal defect (PA/VSD) or truncus arteriosus (TAC). The evolution of the aortic size after birth will result from a combination of intrinsic pathology, hemodynamic factors, associated malformations, surgical or catheter interventions, and control of risk factors later in life (1).

Secondary dilatation of the aortic root and to a lesser extent of the ascending aorta, is seen after congenital cardiac surgery, when the original aortic root is replaced by a pulmonary autograft, as in Ross operation, or modified as in the arterial switch operation (ASO) or systemic outflow tract reconstruction in single ventricle (SV) patients. In these situations the neo-aortic root consists mainly of pulmonary arterial root tissue introduced in the high pressure left-sided system, often leading to dilatation in a time-dependent fashion (1).

The dilatation of the aorta or the neo-aortic root is not a stand-alone characteristic, but needs to be regarded as a part of the aorto-ventricular complex, compromising the systemic ventricle, the aortic valve, the aortic root, and the vascular wall. Each component of this complex may by itself influence the other components, thus introducing a dysfunction at multiple levels, often defined as aortopathy (2).

CHD-associated aortopathy shows histological and functional similarities with Marfan syndrome, degeneration of the aortic media (cystic medial necrosis) (3,4).

The aim of this study is to present our experience and the incidence of primary and secondary adult CHD-associated aortopathy.

Methods

Patients and methods

This is a retrospective study and the cohort is comprised of adult patients with congenital heart disease from the registry of the Eastern Slovakia Institute of Cardiovascular Diseases and all the relevant data were obtained from the medical records. Patients with CHD at the age of 18 years old are included in the adult CHD registry of the Eastern Slovakia Institute of Cardiovascular Diseases and they have annually follow-up examination with a transthoracic echocardiography (TTE) of the ascending aorta and the aortic root. Data from the last follow-up examinations which were done during the last year are included in this study. In the primary and secondary aortopathy group were included 35 patients with CoA (2 patients), BAV (26 patients), TOF (7 patients), and 12 patients that underwent Ross operation (4 patients), arterial switch operation due to transposition of great arteries (TGA) (6 patients) and Fontan procedure (2 patients) respectively. As a control group we selected 64 patients with non aortopathy associated CHD (atrial and ventricular septal defect). The diameter of the aortic root and ascending aorta were measured by TTE and all evaluations were done according the standard techniques recommended by the American Society of Echocardiography (5). The criteria for intervention on the aortic root and ascending aorta were according to the international guidelines and recommendations (3).

An informed consent from the patients and approval from the institutional review board were obtained in order to present this study.

Statistical analysis

All variables were expressed as median and the qualitative variables as numbers and percentages. A one-way ANOVA and the Tukey's range test paired were used to compare the variables. A p value of less than 0.05 was considered statistically significant. The statistical analyses were performed using the StatSoft, Inc (2007). STATISTICA (data analysis software system), version 8.0 www.statsoft.com

Results

The median age of the patients in the group of the primary and secondary aortopathy and the control group were 30.65 (18-60), 29.24 (18-44) and 36.35 (19-66) years old respectively (Table 1).

Patients with secondary aortopathy had statistically significant larger aortic root diameter compared to the control group (35.67 (27-48) mm vs 29.03 (19-38) mm, $p < 0.001$) (Graf 1, Table 1). Compared to the patients with primary aortopathy had larger aortic root diameter (35.67 (27-48) mm vs 33.82 (27-49) mm $p = 0.07$), but with no statistical significance (Graf 1, Table 1). Also patients with primary aortopathy had statistically significant larger aortic root diameter compared to the control group (33.82 (27-49) mm vs 29.03 (19-38) mm, $p < 0.001$) (Graf 1, Table 1).

Patients with secondary aortopathy had statistically significant larger ascending aorta diameter compared to the control group (42.10 (30-50) mm vs 30.25 (21-41) mm, $p < 0.001$) as well as to the patients with primary aortopathy (42.10 (30-50) mm vs 36.28 (26-49) mm, $p < 0.05$) (Graf 1, Table 1). Moreover, patients with primary aortopathy had also statistically significant larger ascending aorta diameter compared to the control group (36.28 (26-49) mm vs 30.25 (21-41) mm, $p < 0.001$) (Graf 1, Table 1).

Table 1 Characteristic of study groups

	Median age (years)	Median ascending aorta (mm)	Median aortic root (mm)
Primary aortopathy	30.65	36.28	33.82
Secondary aortopathy	29.24	42.10	35.65
Control group	36.35	30.25	29.03

In the primary aortopathy group, 8 patients (22.85%) had ascending aorta diameter between 35-40 mm, 9 patients (25.71%) had ascending aorta diameter ≥ 40 mm, 7 patients (20.00%) had aortic root diameter between 35-40 mm and 6 patients (17.14%) had aortic root aorta diameter ≥ 40 mm (Table 2).

In the secondary aortopathy group, 3 patients (25.00%) had ascending aorta diameter between 35-40 mm, and 6 patients (50.00%) had ascending aorta diameter ≥ 40 mm, 2 patients (16.67%) had aortic root diameter between 35-40 mm and 4 patients (33.33%) had aortic root aorta diameter ≥ 40 mm (Table 2).

Table 2 Characteristic of aortic root and ascending aorta diamerets in study groups- primary and secondary aortopathy

	Ascending aorta 35-40 mm	Ascending aorta ≥ 40 mm	Aortic root 35-40 mm	Aortic root ≥ 40 mm
Primary aortopathy	8 (22.85%)	9 (25.71%)	7 (20.00%)	6 (17.14%)
Secondary aortopathy	3 (25.00%)	6 (50.00%)	2 (16.67%)	4 (33.33%)

In the primary aortopathy group 9 patients (25.71%) had undergone at least one prior operation. One patient had repair of CoA and later reoperation due to subvalvular aortic stenosis, 1 patient with BAV had aortic valve replacement (AVR) and later redo-AVR, 1 patient with BAV had replacement of the ascending aorta due to aortic aneurysm, and 6 patients had repair of TOF, where 3 of them had multiple reoperations.

In the secondary aortopathy group, all the patients had undergone at least one prior operation. Four patients had Ross procedure, where 1 patient due to dilatation of the pulmonary homograft had a reoperation and replacement of the neo-aorta, 6 patients had switch operation due to TGA, and 2 patients had a Fontan procedure.

During the follow-up period, were performed only in 2 patients (one from each group) operations on the aortic root and the ascending aorta due to aortic root or ascending aorta dilatation. The rest of the patients are under follow-up screening and so far they do not meet indications for any procedure on the aortic root and the ascending aorta.

Discussion

Progressive proximal aortic (aortic root and ascending aorta) dilatation is frequently found in adults with unrepaired (primary aortopathy) or repaired (secondary aortopathy) CHD. In our study, in the primary aortopathy group 25.71% and 17.14% of the patients had dilated ascending aorta and aortic root respectively. Similar results are reported also by other authors. Steward et al (6) found dilated aortic root in 16%. There was a trend to a more abnormally widened aorta in patients that had surgery for repair of CoA later in life, and 5 of the patients in their series that died from aortic aneurysm rupture, had original surgery at a mean age of 19 years old, and were known of associated hypertension. It is unclear if early repair of CoA will always be able to prevent late aortic dilatation. Even neonatal intervention does not prevent the occurrence of late hypertension, which by itself may trigger aortic dilatation (7). Biopsies studies of the aortic wall found increased amount of collagen and a decreased smooth muscle content in the pre-stenotic region (8). In our study, no patient after CoA repair needed intervention on the aorta during the follow-up period due to dilatation of the ascending aorta or the aortic root.

Concomitant aortic dilatation is seen in 80% of patients with BAV (9). Studies showed that the dilatation results from a combination of intrinsic aortic wall modifications (genetic theory) and hemodynamic

changes induced by the bicuspid valve. The marked heterogeneity of BAV disease leads to different phenotypes, resulting in a large clinical variation of BAV patients (10,11). The strong association of BAV with CoA may indicate that BAV disease involves the ascending aorta and aortic arch extending to the ligamentum arteriosum. Dilatation of the ascending aorta occurs as a consequence of aortic medial degeneration (12). During the follow-up period only one patient with BAV from our study group underwent replacement of the ascending aorta due to aortic aneurysm.

Dilatation of the proximal aorta is a common feature in patients with unrepaired TOF. Corrective surgery has dramatically improved long-term prognosis, and nearly 90% of the patients surviving well into adulthood (13). However, persistent aortic root dilatation is increasingly reported in adult patients, years after the corrective surgery. In 1997 the first series of progressive aortic root dilatation was published, where a substantial cohort developed subsequent aortic valve incompetence, necessitating reoperation on the aortic root (14).

The underlying mechanism of the aortic dilatation in TOF are both hemodynamic and intrinsic wall abnormalities like cystic medial necrosis as in Marfan syndrome. Presence of right to left shunt, other congenital anomalies, complete repair at older age and even a genetic factor have been implicated in aortic dilatation in TOF (15-17). In a homogenous cohort of TOF repaired early in infancy, was found that the ascending aortic size decreases with growth of the patient during the first years after surgery, irrespective of the total histology score at surgery (18). These findings support the presumption that mitigation of the transaortic flow by early surgical repair of TOF triggers a remodeling process that may interrupt the progression of the limited histological alterations of the aortic root, thus preventing late aortic dilatation.

In a recent review by Mongeon et al (19), in adult patients 35 years after repair of TOF at a mean of 7 years of age, an aortic dimension of ≥ 40 mm was found in 29% and moderate to severe aortic regurgitation in 3.5% of the patients. Only 3 cases of aortic dissection later after TOF repair have been described, all in severely dilated aortas of ≥ 70 mm (1)

Moreover, in the secondary aortopathy group, 50% and 33.30% of the patients had dilated ascending aorta and aortic root respectively.

After Ross procedure, the neo-aortic root dilates mainly at the sinus portion and the sinotubular junction, and less at the neo-aortic annulus itself. Dilatation occurs rapidly within the first days after surgery, with a further increase during the first year of follow-up, without causing significant aortic regurgitation in the medium phase term (20). The postoperative distention of the pulmonary autograft leads to remodeling of the wall with intimal thickening, medial elastin fragmentation, hypertrophic smooth muscle cells and increased medial and adventitial fibrosis (21). Freedom from autograft reoperation has been reported between 74% and 93% at 10 years and between 65% to 82% at 15 years (1). In our study only one patient after the Ross procedure needed reoperation due to pulmonary autograft dilatation during the follow-up period.

In patients with TGA, after the arterial switch operation (ASO), the native pulmonary valve and root assume the role of systemic arterial valve and root. Progressive dilatation of the neo-aortic root exceeds somatic growth during a long follow-up period. A dilated neo-aortic root is seen in at least 50% of all patients after ASO (22). Risk factors that are associated with neo-aortic root dilatation after ASO include the presence of a ventricular septal defect, previous pulmonary artery banding, older age and surgical technical factors (1,22,23).

Also, adults patients with TGA after the atrial switch operation have a greater incidence of dilatation of both the pulmonary artery and aorta (24).

Freedom from aortic root reoperation was reported to be between 83% and 97% after ASO (1). Moreover, only a single case report of surgical repair of aneurysm of the ascending aorta after atrial switch operation has been described (25). In our study group, no patient after ASO or atrial switch operation needed reoperation on the ascending aorta.

Evidence of aortic dilatation has been reported in patients after the Fontan procedure. In a study with a median follow-up of 9 years neo-aortic root dilatation was observed in 98% of the patients (26). Histological analysis demonstrated findings seen in other forms of CHD- associated aortopathies, such as fragmentation of elastic fibers and deposition of myxoid material (27). Aortic dissection in patients after the Fontan procedure was reported in two patients with dilated aortic root (1). In our study group, no patient after the Fontan procedure needed reoperation on the ascending aorta or the aortic root.

One of the main findings of our study is that patients with secondary aortopathy had larger ascending aorta and aortic root dimensions than the patients with primary aortopathy. In the existing literature, so far there are no studies with direct comparison of aortic dimensions between the primary and secondary aortopathy. In our opinion, secondary aortopathy in contrast to the primary aortopathy, the neo-aorta consists mainly of pulmonary artery tissue introduced in the high pressure left-sided system, often leading to more severe dilatation in a time-dependent fashion than in the primary aortopathy.

Regarding aortopathy-associated CHD other than BAV, dissection risk is low (28). Since, there are no specific guidelines recommendations (3), and no reports suggesting high risk of aortic complications at neo-aortic root/ascending aortic diameters \geq 55 mm, there is a pragmatic attitude to these patients with a restrictive policy following general aortic disease guidelines (55 mm) (3, 29). On the other hand, because the time factor is the principal determinant of late neo-aorta dilatation and some cases of quickly progressive diameter increase or fatal complications have been described, a close observation is warranted (3, 30). In these patients, aortic surgery was performed at lower aortic diameters, particularly if surgery was indicated for aortic valve dysfunction (28).

Regarding BAV the life risk of aortic dissection has been reported to be 9 times higher than that of the general population (1). Moreover, Kuijpers JM et al (28), reported 10-year dissection incidence of 0.3%. This low aortic-dissection risk in BAV is also reported in population-based and post- AVR BAV cohorts (28). Current guideline recommended aortic diameters thresholds for prophylactic surgery in BAV is 55

mm or 50 mm with risk factors, and 45 mm at the time of AVR for dysfunctional BAV (28). Surgical treatment for ascending aortic dilatation in CoA may be considered when the diameter is ≥ 55 mm (≥ 27 mm/m²) or if rapid progression (31). Moreover, the close association between BAV and CoA imply strategies established for BAV may be appropriate (3).

In conclusion, progressive aortic root and/or ascending aorta dilatation is frequently found in adults with repaired or unrepaired CHD. Primary aortopathy is associated with BAV, CoA and conotruncal abnormalities, where secondary aortopathy is after congenital heart surgery, by which the original aortic root/ ascending aorta is replaced by a pulmonary autograft, as in Ross operation or modified as in ASO of Fontan procedure. It was observed that, more patients with secondary aortopathy had dilated ascending aorta and aortic root, as well as larger aortic diameters compare to the patients with primary aortopathy. Routine follow-up of these patients with attention to aortic diameter is necessary.

Conclusions

More patients with secondary aortopathy had dilated ascending aorta/ aortic root, as well as larger aortic diameters compare to the patients with primary aortopathy. Routine follow-up of these patients with attention to aortic diameter is necessary.

Abbreviations

CHD - congenital heart disease

CoA - coarctation of the aorta

BAV- bicuspid aortic valve

TOF - tetralogy of Fallot

PA -pulmonary atresia

VSD - ventricular septal defect

TAC - truncus arteriosus

SV - single ventricle

TTE - transthoracic echocardiography

TGA - transposition of great arteries

AVR - aortic valve replacement

Declarations

Ethics approval and consent to participate

An informed consent from the patients and approval from the institutional review board were obtained in order to present this study.

Consent for publication

An informed consent from the patients and approval from the institutional review board were obtained in order to present this study.

Availability of data and materials

The datasets used and/or analysed during the current study are available from the corresponding author on request.

Competing interests

“The authors declare that they have no competing interests”.

Funding

This is a retrospective study and the cohort is comprised of adult patients with congenital heart disease from the registry of the Eastern Slovakia Institute of Cardiovascular Diseases and all the relevant data were obtained from the medical records.

Authors' contributions

IS- analyzed and interpreted the patient data, contributor in writing the manuscript

PA- contributor in writing the manuscript

AB- contributor in writing the manuscript

MV- *literature searching*, contributor in writing the manuscript

MJ – *literature searching*, contributor in writing the manuscript

All authors read and approved the final manuscript.

Acknowledgements

Not applicable

References

1. Francois K. Aortopathy associated with congenital heart disease: A current literature review. *Ann Pediatr Cardiol* 2015;8:25-36
2. Niwa K, Perloff JK, Bhuta SM, ET AL. Structural abnormalities of great arterial walls in congenital heart disease: light and electron microscopic analyses. *Circulation* 2001;103:393-400
3. Kuijpers JM, Mulder BJM. Aortopathies in adult congenital heart disease and genetic aortopathy syndromes: management strategies and indication for surgery. *Heart* 2017;103:952-966
4. Niwa K. Aortic dilatation in complex congenital heart disease. *Cardiovasc Diagn Ther* 2018;8(6):725-738
5. Goldstein SA, Evangelista A, Abbara S, et al. Multimodality imaging of diseases of the thoracic aorta in adults. From the American Society of Echocardiography and the European Association of Cardiovascular Imaging. *J Am Soc Echocardiogr* 2015;28:119-82
6. Steward AB, Ahmed R, Travill CM, Newman CG. Coarctation of the aorta life and health 20-44 years after surgical repair. *Br Heart J* 1993;69:65-70
7. Brouwer AB, Erasmus ME, Ebels T, Eijgelaar A. Influence of age on survival, late hypertension and recoarctation in elective aortic coarctation repair. Including long-term results after elective aortic coarctation repair with a follow-up from 22 to 44 years. *J Thorac Cardiovasc Surg* 1994;108:525-531
8. Sehested J, Baandrup U, Mikkelsen E. Different reactivity and structure of the prestenotic and poststenotic aorta in human coarctation. Implications for baroreceptor function. *Circulation* 1982;65:1060-1065
9. Friedman T, Mani A, Elefteriades JA. Bicuspid aortic valve: clinical approach and scientific review of a common clinical entity. *Exprt Rev Cardiovasc Ther* 2008;6:235-248
10. Della Corte A, Bancone C, Quarto C, et al. Predictors of ascending aortic dilatation with bicuspid aortic valves: A wide spectrum of disease expression. *Eur J Cardiothorac Surg* 2007;31:397-405
11. Girdauskas E, Borger MA, Secknus MA, Girdauskas G, Kuntze T. Is aortopathy in bicuspid aortic valve disease a congenital defect or a result of abnormal hemodynamics? A critical reappraisal of a one-sided argument. *Eur J Cardiothorac Surg* 2011;39:809-814
12. Tadros TM, Klein MD, Shapira OM. Ascending aortic dilatation associated with bicuspid aortic valve: Pathophysiology, molecular biology, and clinical implications. *Circulation* 2009;119:880-890
13. Murphy JG, Gersh BJ, Mair DD, et al. Long-term outcome in patients undergoing surgical repair of tetralogy of Fallot. *N Engl J Med* 1993;329:593-599
14. Dodds GA, Warnes CA, Danielson GK. Aortic valve replacement after repair of pulmonary atresia and ventricular septal defect of tetralogy of Fallot. *J Thorac Cardiovasc Surg* 1997;113:736-741
15. Tan JL, Davlouros PA, McCarthy KP, Gatzoulis MA, Ho SY. Intrinsic histological abnormalities of aortic root and ascending aorta in tetralogy of Fallot: Evidence of causative mechanism for aortic dilatation and aortopathy. *Circulation* 2005;112:961-968
16. Goldmuntz E, Clark BJ, Mitchell LE, et al. Frequency of 22q11 deletions in patients with conotruncal defects. *J Am Coll Cardiol* 1998;32:492-498

17. Chowdhury UK, Mishra AK, Balakrishnan P, et al. Role of fibrillin-1 genetic mutations and polymorphism in aortic dilatation in patients undergoing intracardiac repair of tetralogy of Fallot. *J Thorac Cardiovasc Surg* 2008;136:757-766
18. Francois K, Creytens D, De Groove K, et al. Analysis of the aortic root in patients with tetralogy of Fallot undergoing early repair: Form follows function. *J Thorac Cardiovasc Surg* 2014;148:1555-1559
19. Mongeon FP, Gurvitz MZ, Broberg CS, et al. Alliance for Adult Research in Congenital Cardiology (AARCCA). Aortic root dilatation in adults with surgically repaired tetralogy of Fallot: A multicenter cross-sectional study. *Circulation* 2013;127:172-179
20. Hokken RB, Boders AJ, Schiks-Berghourt MB, et al. Does the pulmonary autograft in the aortic position in adults increase in diameter? An echocardiographic study. *J Thorac Cardiovasc Surg* 1997;113:667-674
21. Schoof PH, Takkenberg JJ, van Suylen RJ, et al. Degeneration of the pulmonary autograft: An explants study. *J Thorac Cardiovasc Surg* 2009;132:1426-1432
22. Schwartz ML, Gauvreau K, del Nido P, et al. Long-term predictors of aortic root dilatation and aortic regurgitation after arterial switch operation. *Circulation* 2004;10:II:28-32
23. McMahan CJ, Ravekes WJ, Smith EO, et al. Risk factors for neo-aortic root enlargement and aortic regurgitation following arterial switch operation. *Pediatr Cardiol* 2004;25:329-335
24. Yurasek GK, Gauvreau K, Poewll AJ, et al. Great vessel root and artery dimensions in transposition of the great arteries with atrial switch operation. *Pediatr Cardiol* 2014;35:457-462
25. Ono M, Goerler H, Boething D, et al. Valve-sparing operation for aortic root aneurysm late after Mustard procedure. *Ann Thorac Surg* 2007;83:2224-2226
26. Cohen MS, Marino BS, McElhinney DB, et al. Neo-aortic root dilatation and valve regurgitation up to 21 years after staged reconstruction for hypoplastic left heart syndrome. *J Am Coll Cardiol* 2003;42:533-540
27. Okamoto RJ, Xu H, Kouchoukos NT, Moon MR, Sundt TM. The influence of mechanical properties on wall stress and distensibility of the dilated ascending aorta. *J Thorac Cardiovasc Surg* 2003;126:842-850
28. Kuijpers JM, Koolbergen DR, Groenink M, et al. Aortic dissection and prophylactic surgery in congenital heart disease. *Int J Cardiol* 2019;274:113-115
29. Stulak JM, Dearani JA, Burkhart HM, Sund TM, Connolly HM, Schaff HV. Does the dilated ascending aorta in an adult with congenital heart disease require intervention.? *J Thorac Cardiovasc Surg* 2010;140:S52-57
30. Artemiou P, Schusterova I, Tohatyova A, Cocherova J, Krchno P, Sabol F. Follow-up after the Ross procedure, how significant it is, case reports of three patients. *J Cardiothrac Surg* 2015;10:144 doi:10.1186/s13019-015-0369-8
31. Baumgartner H, Bonhoeffer P, De Groot NM, et al. Task force on the management of grown-up congenital heart disease of the European Society of Cardiology (ESC):ESC Committee for Practice

Figures




Figure 1

Aortic root and ascending aorta in study groups- primary and secondary aortopathy and control group