

# Body Size Dependent Dispersal Influences Stability in Heterogeneous Metacommunities

Kurt Anderson (✉ [kurt.anderson@ucr.edu](mailto:kurt.anderson@ucr.edu))

University of California, Riverside

Ashkaan Fahimipour

National Oceanic and Atmospheric Administration

---

## Research Article

**Keywords:** Body size, dispersal influences stability, heterogeneous metacommunities, biological processes

**Posted Date:** March 2nd, 2021

**DOI:** <https://doi.org/10.21203/rs.3.rs-257849/v1>

**License:** © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

# Body size dependent dispersal influences stability in heterogeneous metacommunities

Kurt E. Anderson<sup>1,\*</sup> and Ashkaan K. Fahimipour<sup>2,3</sup>

<sup>1</sup>*University of California Riverside, Department of Evolution, Ecology, & Organismal Biology*

<sup>2</sup>*National Oceanic and Atmospheric Administration, Southwest Fisheries Science Center*

<sup>3</sup>*University of California Davis, Department of Computer Science*

## Abstract

Body size affects key biological processes across the tree of life, with particular importance for food web dynamics and stability. Traits influencing movement capabilities depend strongly on body size, yet the effects of allometrically-structured dispersal on food web stability are less well understood than other demographic processes. Here we study the stability properties of spatially-arranged model food webs in which larger bodied species occupy higher trophic positions, while species' body sizes also determine the rates at which they traverse spatial networks of heterogeneous habitat patches. Our analysis shows an apparent stabilizing effect of positive dispersal rate scaling with body size compared to negative scaling relationships or uniform dispersal. However, as the global coupling strength among patches increases, the benefits of positive body size-dispersal scaling disappear. A permutational analysis shows that breaking allometric dispersal hierarchies while preserving dispersal rate distributions rarely alters qualitative aspects of metacommunity stability. Taken together, these results suggest that the oft-predicted stabilizing effects of large mobile predators may, for some dimensions of ecological stability, be attributed to increased patch coupling *per se*, and not necessarily coupling by top trophic levels in particular.

## Introduction

What allows large, complex ecosystems to be stable? May's analysis of randomly arranged communities of self-limiting populations challenged previous ecological thinking on this issue, showing that greater species richness and interaction connectivity tended to destabilize random communities rather than stabilize them [1, 2]. This and subsequent theory set the stage for decades of work analyzing the complicated relationship between diversity and dynamics that continues today [3, 4, 5, 6, 7]. In contrast to randomly-assembled communities, an emerging focus in modern biodiversity theory is

---

\*To whom correspondence should be addressed: kurt.anderson@ucr.edu

on the non-random structural features of ecological systems that impart stability [4, 8, 9]. For communities organized around feeding relationships (*i.e.*, food webs), two types of *structure* receiving extensive attention are allometric hierarchies, where larger species mostly eat smaller ones and populations experience other demographic rates dependent on body-size [5, 10, 11]; and dispersal among spatially discrete habitats [12, 13, 14, 15].

Body size-based food web topologies and allometric scaling of population demographic rates have both shown to be stabilizing for models of trophic interactions [5, 6, 10, 16, 17]. Simple mass-based hierarchical feeding rules — where species high in the feeding hierarchy are interpreted as large-bodied predators — can successfully reproduce realistic food web topologies [4, 5, 11, 18, 19, 20, 21] that are more likely to be dynamically stable than random network configurations [16, 17]. Likewise, allometric scaling of species' demographic rates such as handling times, conversion efficiencies, and biomass turnover, are predicted to stabilize food webs [5, 10, 22, 23, 24, 25]. This is particularly the case when the ratio of body masses between resources and consumers is large and consistent with values observed in natural food webs [5, 10, 26].

At larger scales, dispersal generates structure by linking spatially distinct food webs through the movement of individuals. The role of dispersal in population dynamics and community composition is a central focus in ecology, with early work emphasizing the colonization of islands by mainland species [27, 28] and the “rescue” of small populations in sink habitats [29]. Colonization-extinction dynamics in spatially subdivided habitats, originally examined in a population context [30, 31], extended these results and have been shown to promote higher regional food web diversity than can be supported in isolated well-mixed systems [15]. Dispersal can also stabilize species interactions locally by mimicking density-dependence in *per capita* growth rates [32, 33, 34, 35]. More recent work has linked metacommunity dynamics to May's original examination of species richness and connectance, showing that dispersal among spatially distinct subpopulations can be strongly stabilizing [36, 37]. Complexity-stability relationships in these cases are relaxed or even reversed relative to results seen in linear stability analyses of random matrices.

Much like trophic interaction rates, many traits that influence animal locomotion, movement speed, and potentially dispersal also vary with body size [38, 39, 40]. Spatial patterns of resource use, home ranges, and geographic range size also exhibit strong allometric relationships [24, 41, 42, 43, 44]. These patterns in turn create the potential for spatial coupling of habitat patches [45] and hence local food webs [16, 46, 47] that depends on body size. While many studies emphasize faster movements of large consumers, suggesting positive body-size dispersal relationships, large-bodied species many face greater dispersal limitation in some habitats [48, 49, 50], suggesting a range of potential relationships between body size and dispersal across ecosystems [15].

Theoretical and empirical evidence suggests an important role for body size-dispersal scaling in ecosystem dynamics, yet the full effects of dispersal variation among species in food webs have so far been difficult to systematize [15]. Some mathematical models indicate that the coupling of distinct food webs by consumer movement can be stabilizing when those webs represent different energy




Figure 1: Model metacommunities are composed of local food webs connected to one another by dispersal. **A.** Each local web inhabits a habitat patch that is part of a spatial network, generated as a random geometric graph. Species in each web have a body size that is larger at higher trophic levels. Food webs have the same number of species and topology in all patches, but interaction rates and other ecological parameters vary among habitats mimicking spatial environmental heterogeneity. **B.** Dispersal varies as either an increasing or decreasing function of body size.

channels or environmental conditions [12, 16, 47]. In other models, greater mobility of consumers is a key requirement for instability [51, 52, 53, 54]. Yet other examples have identified dispersal-driven instabilities for communities in which primary producers traverse space more rapidly than other species [13]. Overall, the effects of dispersal are complicated in ecological networks and a general understanding of how dispersal rules influence food web stability is lacking [13, 15].

Here, we examine how body size scaling of species' dispersal rates influences stability in model trophic metacommunities (Fig. 1). In particular, we ask whether body size-dependent variation in dispersal rates influences trophic metacommunity stability relative to rates that are either uniform or randomly varying among species. We examine landscapes of discrete habitat patches that include variation in local abiotic conditions, generating spatial heterogeneity in rates of primary production and trophic interactions among species. Clearly, the stability of such heterogeneous metacommunities will depend on the proportion of patches in the landscape with locally favorable conditions for stability. We show how the body size scaling of species' dispersal rates alters this relationship. Because general rules describing the dependence of dispersal on body size are lacking and likely vary among ecosystems [15], we consider both positive and negative relationships between dispersal rates and body sizes. Our results show strong effects of dispersal-body size scaling on metacommunity stability, largely due to increased connectivity among local webs with different stability properties.

## Model formulation

We model trophic metacommunities as copies of food webs consisting of  $S$  species, embedded in a set of  $N$  patches. We chose to represent spatial networks as random geometric graphs, which provide a reasonable approximation for real networks [15] of habitats and the dispersal connections between them (Fig. 1).

Food web topologies were generated using the niche model [18], which recapitulates realistic yet variable feeding relationships with only two input parameters, species richness  $S$  and connectance  $C$

[18]. Briefly, species are assigned a position on a 1-dimensional niche axis and feed on species over a range determined by  $C$ . The feeding range is centered below the species' niche positions, creating a trophic hierarchy where each species  $i$  has trophic position  $T_i$  defined by the path length to any basal producer.

Following [5, 7, 10, 11], we assigned body sizes assuming that the normalized mass  $M_i$  of each species  $i$  scales with their trophic position  $T_i$ , as  $M_i = R^{T_i}$ . For reported results, normalized producer body sizes are uniformly set to one while  $R = 42$ . This value of  $R$  represents the average predator-prey body mass ratio reported by [26], although our qualitative results are consistent across a wide range of realistic ratios.

Dynamics on the links defined by each niche model topology were represented as the set of ordinary differential equations of the form

$$\frac{dB_i^k}{dt} = G_{B_i^k}(B_i^k) + F_{B_i^k}(\mathbf{B}^k) - X_{B_i^k}(B_i^k) - \sum_j E_{B_j^k B_i^k}(\mathbf{B}^k) + \sum_l [D_{B_i^l}(B_i^l) - D_{B_i^k}(B_i^k)] \quad (1)$$

where  $B_i^k$  is the biomass density of species  $i$  in patch  $k$ . The non-specified function  $G$  is the growth rate of primary producers,  $F$  is the rate of biomass accumulation due to feeding on other species,  $X$  is rate of biomass loss due to respiration and mortality,  $E$  is the rate of biomass loss due to consumption by species  $j$ , and  $D$  is the dispersal rate between patches. While the functional forms are not explicitly specified, the general form of equation (1) admits the calculation of a Jacobian matrix that quantifies how species in the metacommunity respond to perturbations from steady state and therefore metacommunity stability [55, 56]. Using the generalized modeling method [6, 55, 57], the derivatives of functions  $G$ ,  $F$ ,  $X$ , and  $E$  that constitute the non-dispersal elements of the Jacobian matrix can be recast in terms of scale, branching, and elasticity parameters (see Methods). The generalized model parameters have clear ecological interpretations: scale parameters set the time scale of biomass turnover, while branching and elasticity parameters set the relative contributions of different processes to biomass gains and losses and the form of non-linearities, respectively [6, 55, 56, 57]. The range of generalized model parameter values studied here map to most commonly encountered functional forms (e.g., Lotka-Volterra-like and Holling type II and III functional responses) that link interaction rates with species' densities in conventional models, and follow ecologically-based arguments from prior work [6, 55, 56] (Table 1).

Trophic metacommunities were constructed as follows. First, a spatial structure of  $N = 10$  habitat patches was randomly generated as a geometric graph on the interval  $[0, 1]$  with a neighborhood radius of 0.32 (see Methods). Each patch  $k$  in the spatial network contains a copy of the same  $S$  species niche model [18] web. Heterogeneity in factors that influence primary production and trophic interactions were modeled as random variation among patches in branching and elasticity parameters [6, 55, 56, 57]. Values of these spatially variable parameters were drawn independently from uniform distributions defined by ecologically meaningful ranges (Table 1).

For simplicity, we assume that there is no dependence of *per capita* dispersal rates on patch identity or on interspecific densities (i.e. no cross-diffusion). We further assume that dispersal is a linear

Parameter	Description	Range or Value
<b>Scale</b>		
$\alpha_i$	Normalized turnover rate of species $i$	$M_i^{-1/4}$
<b>Branching</b>		
$\rho_i^k$	Fraction of biomass gains from predation, for species $i$ in patch $k$	0 or 1
$\sigma_i^k$	Fraction of biomass loss from predation, for species $i$ in patch $k$	[0, 1]
$\beta_{ij}^k$	Fraction of species $i$ 's loss from consumption by $j$ in patch $k$	[0, 1]
$\chi_{ij}^k$	Fraction of species $j$ 's gains from consumption of $i$ in patch $k$	[0, 1]
<b>Elasticity</b>		
$\phi_i^k$	Nutrient availability for producer $i$ in patch $k$	[0, 1]
$\psi_i^k$	Sensitivity of $i$ 's predation rate to its own density in patch $k$	[0.5, 1]
$\lambda_{ji}^k$	Sensitivity of $j$ 's foraging preferences to prey densities in patch $k$	[1]
$\gamma_i^k$	Sensitivity of $i$ 's predation rate to total prey density in patch $k$	[0.5, 1.5]
$\mu_i^k$	Sensitivity of species $i$ 's mortality to its own density in patch $k$	[1, 1.5]
<b>Niche model</b>		
$S$	Species richness	[10, 30]
$C$	Food web connectance	[0.12, 0.24]

Table 1 Parameter definitions and ranges used in metacommunity simulations.

function of local intraspecific density and allow for dispersal rates to vary among species, 120

$$D_{B_i^k}(B_i^k) = \delta_i^k B_i^k, D_{B_i^l}(B_i^l) = \delta_i^l B_i^l; \delta_i^k = \delta_i^l = \delta_i. \quad (2)$$

Each species  $i$  in a particular metacommunity scenario is assigned a species-specific dispersal rate  $\delta_i$  121  
that is dependent on its body size. We chose a power law relationship between body mass and dis- 122  
persal rate, as this general form captures many allometric scaling relationships related to locomotive 123  
capabilities and spatial habitat use [24, 38, 40, 58]. Specifically, 124

$$\delta_i = dM_i^z \quad (3)$$

where  $d$  is the global link strength of the spatial network and  $z$  is the body size scaling exponent 125  
for dispersal. Producers have a body size equal to one such that for producers,  $\delta_i = d$ . When  $z$  is 126  
positive, species with higher trophic positions and thus larger body sizes traverse the spatial network 127  
at a higher rate. This scenario potentially reflects terrestrial and pelagic food webs where larger 128  
animals have greater mobility and hence dispersal potential [12]. In other systems,  $z$  may be negative 129  
[13, 15, 40], giving species in lower trophic levels the fastest dispersal rates with lower rates for larger 130  
bodied species. 131

We examined metacommunity stability using linear stability analysis. This procedure was applied 132  
in classic work on community stability by May and others [1, 2] and has since been expanded to 133  
incorporate spatial processes [14, 36]. Linear stability is assessed by examining the eigenvalues of 134

the Jacobian matrix of the trophic metacommunity,  $\mathbf{J}$ . Each local food web has a corresponding local  
Jacobian  $\mathbf{J}_k$  that is derived from equation (1); these local Jacobians are collected and numerically  
arranged as blocks on the diagonal of the  $SN \times SN$  matrix  $\mathbf{P}$ . The local food web information is then  
used to calculate the metacommunity Jacobian  $\mathbf{J}$  using the equation [13, 14, 59]

$$\mathbf{J} = \mathbf{P} - \mathbf{L} \otimes \mathbf{D}, \quad (4)$$

where  $\otimes$  is the Kronecker product; the  $N \times N$  matrix  $\mathbf{L}$  is the Laplacian of the corresponding spatial  
network; and  $\mathbf{D}$  is an  $S \times S$  Jacobian-like diagonal matrix containing species-specific dispersal rates  
calculated from equation (2). Eigenvalues of local food web Jacobians  $\mathbf{J}_k$  and the metacommunity  
Jacobian  $\mathbf{J}$  were computed numerically; stability occurs when the real parts of all eigenvalues are  
negative. Additional details regarding this method of analysis are found in refs. [13, 14, 59, 60] and in  
the Methods.

Body size scaling could increase stability in metacommunities owing to simple increases in overall  
dispersal, rather than because of particular relationships between body size, trophic position and  
dispersal rate. To test this possibility, we compared our suite of model metacommunities with different  
dispersal rules to those with random variation in dispersal rates. For each metacommunity and each  
dispersal scenario, we computed Jacobians with 100 random permutations of species' dispersal rates  
(*i.e.*, the diagonal entries of  $\mathbf{D}$ , see eq. 3) and compared stability properties in these randomized  
metacommunities to their corresponding intact systems.

## Results

Linear stability of both local food webs and metacommunities varied dramatically over the multitude  
of food webs examined, depending on species richness, food web connectance, branching, elasticity  
[6], and dispersal parameters. The baseline dispersal rate  $d$  and body size scaling exponent  $z$  in par-  
ticular influenced metacommunity stability. Figure 2 presents the aggregated results of all numerical  
analyses organized by local food web and metacommunity stability. The proportion of stable local  
food webs gives the proportion of webs in a metacommunity that would be stable in the absence of  
any dispersal. The proportion of stable metacommunities gives the probability that a corresponding  
trophic metacommunity will be stable when those patches are then linked by dispersal. The pro-  
portion of stable local food webs has the intuitive effect of positively increasing the chance that the  
metacommunity they constitute will also be stable; a metacommunity composed entirely of locally  
stable food webs is always stable. However, the strength of this positive relationship relies critically  
on the rules that determine dispersal rates across trophic levels.

Metacommunities are most likely to be stable when larger-bodied species disperse through spatial  
networks of habitat patches faster than smaller-bodied ones (that is, for  $z > 0$ , Fig. 2). Thus, having  
more patches with conditions that promote stability and experiencing higher overall spatial coupling  
both stabilize metacommunities. For a given proportion of stable local food webs and for all but the




Figure 2: The relationship between local and metacommunity stability as influenced by dispersal. The baseline dispersal rate  $d$  and the body size scaling coefficient  $z$  determine the species-specific dispersal rate  $\delta_i = dM_i^z$ , where  $M_i$  is the mass of species  $i$ . All species have the same dispersal rate  $d$  when  $z = 0$ . When  $z$  is positive, larger bodied species have faster dispersal rates, whereas when  $z$  is negative, it is smaller bodied species that have faster dispersal rates. Other parameters vary across all simulations as indicated by Table 1. Error bars denote  $\pm 2$  S.E.M., and are too small to see.

strongest global coupling (*i.e.* large  $d$ ), metacommunities with a positive relationship between body size and dispersal rate are most likely to stable, while those with negative scaling are least likely to be stable.

The stabilizing effects of positive body size-dispersal scaling are most pronounced when the global link strength among habitat patches  $d$  is relatively low. When  $d = 1$ , metacommunity stability with size-dependent dispersal is nearly indistinguishable from the case of uniform dispersal. As overall large levels of dispersal provide a substantial stabilizing effect, positive body size-dispersal scaling therefore appears to be beneficial for stability because large consumers increase spatial connectivity above the baseline rate set by primary producers (*i.e.* it is guaranteed that  $\delta_i \geq d$ ). With negative scaling, these same consumers have lower dispersal rates than producers (*i.e.*  $\delta_i \leq d$ ), lowering the overall spatial coupling between metacommunity patches and reducing stability.

Correlations between key model parameters (see Methods) and metacommunity stability are shown in Fig. 3, confirming the importance of dispersal rules. Of all model parameters examined, the baseline dispersal rate  $d$  and the body-size scaling parameter  $z$  have by far the strongest stabilizing effects. In contrast, species richness  $S$  and food web connectance  $C$  have the greatest de-stabilizing effects, largely through their well-documented effects on local food webs [2, 3, 6]. No other parameter showed a notable correlation with stability, either positive or negative.

The results in Fig. 2 show how body size scaling of dispersal can stabilize metacommunities by increasing coupling among patches. In Fig. 4, we compare stability of metacommunities with body size-dispersal scaling to those where the same set of dispersal rates are randomly re-assigned to species (permuted metacommunities). With these comparisons we ask whether, for a given global link strength  $d$ , it is coupling by large-bodied species specifically or simply greater coupling overall that drives stability. The 1:1 line in Fig. 4 shows where the leading eigenvalue  $\lambda_1$  of a metacommunity is the same as median leading eigenvalues of the corresponding permuted metacommunities. Most eigenvalue comparisons fall near this line across the entire range of dispersal scaling exponents  $z$




Figure 3: The effects of key parameters on metacommunity stability. Stability is as defined in Fig. 2 and parameters are defined in Table 1. Correlations given are coefficients from the best-fitting generalized linear model  $\pm 2$  SEM. Positive correlations indicate that larger values of a parameter correspond to a higher probability that a randomly-assembled metacommunity will be stable.

and global link strengths  $d$  (Supplementary Fig. 1). Figure 5 presents these results categorized by their qualitative effects on stability. The sign of the leading eigenvalues for the metacommunities with permuted dispersal are typically the same as the intact ones, indicating that the median effect of shuffling dispersal rates among species is rarely a qualitative change in stability. In fact, there is no combination of body size scaling  $z$  and dispersal coupling  $d$  where a qualitative change in stability occurs in more than 25% of metacommunities (Supplementary Fig. 2). Yet despite the frequent preservation of qualitative dynamics, re-arranging which species exhibit the highest dispersal rates on a patch network does show a high potential to alter the magnitude of  $\lambda_1$  (Fig. 4; Supplementary Figs. 1 & 2) and therefore the rates at which metacommunities return to, or depart from, steady states following perturbation. We conjecture that dispersal hierarchies may be important for the transient dynamics of ecosystems, and suggest that an understanding of how dispersal rules impact different dimensions of ecological stability will be an important goal for future work.

## Discussion

Our model reveals an apparent stabilizing effect of positive body size scaling of dispersal in heterogeneous metacommunities (Fig. 2), consistent with some conceptual and mathematical theories [14, 16, 61]. However, by comparing metacommunities with allometric dispersal to those whose dispersal rates were random permutations of the original values (Figs. 4 & 5), we found evidence that increased overall connectivity, and not predator movement *per se*, was responsible for metacommunity stability. Increased connectivity manifests either through larger values of global patch coupling  $d$  or stronger positive body size-dispersal scaling  $z$ , and generates conditions in which fewer stable




Figure 4: The effect of dispersal variation on metacommunity stability for spatial link strength  $d = 0.1$ . The metacommunity is stable when the real part of the leading eigenvalue of the metacommunity Jacobian  $\lambda_1 < 0$ . Allometric dispersal is defined eq. 2. Permuted dispersal refers to cases where allometric dispersal rates were randomly reassigned to new species. Each unique metacommunity with allometric dispersal was compared to 100 randomly permuted counterparts, and are shown with 1:1 lines. Grey regions mark portions of the plot representing qualitative changes in stability.

patches are required to “rescue” unstable patches in a metacommunity. 214

There are many potential explanations for why random dispersal maintains roughly the same 215  
degree of metacommunity stability as positive body size scaling. The stabilizing effects of dispersal 216  
in randomly assembled metacommunities have been shown to operate when dispersal is both ho- 217  
mogeneous and variable among species [36]. We conjecture that, for the non-random food webs we 218  
examine here, the existence of stabilizing structural motifs could be enhanced by dispersal. Con- 219  
sumers play key roles in the stabilizing effects of food webs compartments [62], including linking 220  
separate energy channels [16, 63] and forming long interaction loops [64]. Higher dispersal by these 221  
consumers, even if they are not top predators, could substantially enhance metacommunity stability 222  
by connecting stabilizing subcommunities across habitats. Additionally, patterns of dispersal where 223  
low level consumers have the greatest dispersal connectivity have been shown to confer stability in 224  
some tri-trophic metacommunities [54]. Finally, differential dispersal among species at similar trophic 225  
levels could generate competition-colonization and fecundity-dispersal trade-offs (reviewed in [65]), 226  
which may enhance stability in simple metacommunities. How the above mechanisms operate in 227  
more complex spatial food webs remains an open question. 228

Negative body size-dispersal scaling relationships lead to lower stability in our model by lowering 229  
dispersal coupling relative to situations where all species in the food web share the same dispersal rate. 230  
These results are most likely relevant for systems where connectivity results from passive transport of 231  
smaller-bodied organisms, for example wind-dispersed plants or freshwater zooplankton, flooding- 232  
dispersed pond invertebrates, and marine organisms with planktonic larvae. Passive dispersal can 233  
generate stochastic variation in connectivity [66], which may have complicated effects on stability. 234  
Yet it can also lead to increased connectivity by large numbers of individuals [49, 67, 68]. Thus, while 235  
negative scaling could potentially equate to low stability, the overall levels of dispersal may be very 236  
high in systems where lower trophic levels disperse at greater rates, reducing the scaling effect and 237  
leading to higher stability. 238




Figure 5: Eigenvalues  $\lambda_1$  from fig. 4 categorized by qualitative effects on stability. Categories *Stability is gained* and *Stability is lost* correspond to cases where the median effect of permuting dispersal rates is a change in the sign of  $\lambda_1$ . *Stability is unaffected* indicates no sign change.

Spatial variation in model parameters were central to how dispersal connectivity influenced stability in our model. The positive relationship between stability and the dispersal parameters  $d$  and  $z$  was consistent throughout the parameter region we explored; additional numerical results not shown confirmed this relationship more broadly. This pattern contrasts with some other metacommunity models that exhibited a unimodal relationship between dispersal and stability [69, 70, 71, 72]. Unimodal dispersal-stability relationships occur when very high dispersal leads the system to behave as a single, well-mixed habitat. Species persistence and the variation-reducing effects of dispersal therefore no longer operate. Similar outcomes do not seem to occur in our model, at least at the parameter values examined, because rates of primary production and trophic interactions are forced to vary in space. A key effect of dispersal in the presence of such spatial variation is to reduce variation among interaction rates of the average, well-mixed food web, which actually increases stability [36, 37].

Homogenous systems could exhibit very different dynamics. Dispersal-induced instabilities are frequent in homogeneous space trophic models (e.g. [73, 13]), leading to spatially patterned steady-states or asynchronous oscillations even when local dynamics are stable. The presence of high-dispersing top predators in more complex food webs appears to generally stabilize spatially homogeneous systems [14], although faster consumer movement is also a necessary condition for spatially patterned steady states to arise in two- and three-species food chains [74, 73]. Dispersal-induced oscillatory instabilities in contrast are more likely when primary producers disperse much faster than primary and secondary consumers in simple food chains [73, 54]. Therefore, we might expect a generally stabilizing effect of positive body-size scaling and a generally destabilizing effect from negative scaling with oscillatory instabilities. However, these predictions could be complicated in more complex ecological networks, where spatial dynamics can be additionally influenced by consumers sharing a common prey (e.g. [75]) or prey being consumed by a shared predator (e.g. [73]). Furthermore, pattern formation and asynchronous oscillations could impart “stability” in a different sense by lowering population variation at larger scales [54, 76].

Our definition of stability pertains to metacommunity behavior near equilibrium and is therefore limited in describing non-equilibrium dynamics. In this class of models, instabilities can in-

deed include trajectories that tend to zero (i.e., species extinctions), but other outcomes including non-equilibrium co-existence with synchronous or asynchronous oscillations are also possible [77, 78, 79, 80]. The relationship between species persistence and other non-equilibrium dynamics is not always clear, particularly when species interactions are nonlinear. In metacommunities, regional persistence can occur even in the presence of local extinctions [69, 70, 72]. Unstable oscillations or even local extinctions may in fact drive spatially asynchronous dynamics that enhances regional persistence [70, 54, 76, 72]. In these cases, positive body size scaling could counteract such effects; just as coupling by large, mobile consumers stabilizes heterogeneous metacommunities in our model, predator movement could dampen the amplitude of asynchronous oscillations. Wide-ranging predators may also synchronize prey dynamics [81, 82, 83]. The rich non-equilibrium behavior possible in highly speciose food webs with nonlinearities, like we examine here, will likely require extensive investigation and generalizations may be challenging [15].

Consumers, especially large-bodied top predators, are being disproportionately lost from the world's ecosystems [61, 84, 85]. These losses have far-ranging implications for ecosystem structure and stability [86, 87, 6, 61, 84, 14]. A potential consequence of consumer extinction is loss of spatial coupling, which previous research and our results here suggest could lead to further regional instability [63, 12, 14], including increased variability and subsequent species losses. While positive body-size scaling had the greatest stabilizing effect in our model, we also found that similarly strong dispersal coupling by species in other trophic positions yielded equivalent stability. Thus, our results suggest that the conservation value of connectivity may not be lost when top consumers are. Instead, identifying alternative agents of connectivity and promoting their dispersal following top consumer loss may serve as a productive strategy. In practice, finding substitutes for large bodied consumers in some ecosystems may be difficult given the out-sized role in ecosystem coupling these species play. Yet, continued threats to top consumer persistence and the potential conservation value of dispersal among other species suggest the utility of planning for robust connectivity at the community rather than population level [88, 89].

## Methods

Steady states and stability of equation (1) were studied using the generalized modeling method [6, 55, 57]. The method assumes all populations in the food web possess a steady state, allowing us to re-cast population densities and functions as normalized proportions of the steady state. Thus, for each population  $B_i^k$  of  $i = 1, \dots, S$  species across  $k = 1, \dots, N$  patches there exists a steady state  $B_i^{k*}$  that allows us to define the normalized densities  $b_i^k = \frac{B_i^k}{B_i^{k*}}$ . The normalized equations for the non-dispersal

components of the metacommunity are therefore

298

$$\begin{aligned} \frac{db_i^k}{dt} = & \alpha_i [(1 - \rho_i^k)g_{B_i^k}(b_i^k) + \rho_i^k f_{B_i^k}(c_{B_i^k}, b_i^k) - (1 - \sigma_i^k)x_{B_i^k}(b_i^k) \\ & - \sigma_i^k \sum_j \beta_{ij}^k e_{B_j^k B_i^k}(\mathbf{b}^k)]; \quad c_{B_j^k} = \sum_i \chi_{ji} \tilde{f}_{B_j^k}(B_i^k). \end{aligned} \quad (5)$$

The functions  $g$ ,  $f$ ,  $x$ , and  $e$  represent the functions  $G$ ,  $F$ ,  $X$ , and  $E$  from equation (1) normalized by their values evaluated at steady state; the newly introduced variables  $c_{B_j^k}$  and  $\tilde{f}_{B_j^k}(B_i^k)$  represent the total amount of food available to species  $j$  and the contribution of species  $i$  to the food available to species  $j$ , respectively [56]. Additional scale and branching parameters that arise from the normalization procedure set the rates and strength of interactions in the local food web. The normalized turnover rate  $\alpha_i$  scales the biomass flow rates for each species in the food web with mass,  $\alpha_i = M_i^{-1/4}$ , while the branching parameters quantify the structure of these flows. Specific interpretations of the scale and branching parameters are given in Table 1 and provided in detail in [55].

299  
300  
301  
302  
303  
304  
305  
306

Determining stability of equation (4) relies on establishing the Jacobian matrix for the local food web,  $\mathbf{J}_k$ . The Jacobian is constructed from elements that describe the change in the dynamic equation for each species that occurs given a change in each component state variable and the functions of each state variable near steady state. The local food web Jacobian therefore is a square  $S \times S$  matrix where diagonal entries  $J_{i,i}$  describe the effects of a change in species  $i$  on itself and the non-diagonal entries  $J_{j,i}$  describe the effects of species  $i$  on species  $j$ . Quantifying the changes of functions of state variables near steady state is accomplished in the generalized modeling framework by defining the following exponent parameters [6, 13, 55, 56, 57],

307  
308  
309  
310  
311  
312  
313  
314

$$\begin{aligned} \phi_i^k &= \left. \frac{\partial}{\partial b_i^k} g_{B_i^k}(b_i^k) \right|_{x=x^*}, \quad \psi_i^k = \left. \frac{\partial}{\partial b_i^k} f_{B_i^k}(c_{B_i^k}, b_i^k) \right|_{x=x^*}, \quad \lambda_{ji}^k = \left. \frac{\partial}{\partial b_i^k} \tilde{f}_{B_j^k}(B_i^k) \right|_{x=x^*}, \\ \gamma_i^k &= \left. \frac{\partial}{\partial c_{B_i^k}} f_{B_i^k}(c_{B_i^k}, b_i^k) \right|_{x=x^*}, \quad \mu_i^k = \left. \frac{\partial}{\partial b_i^k} x_{B_i^k}(b_i^k) \right|_{x=x^*}. \end{aligned} \quad (6)$$

These exponent parameters can be interpreted formally as elasticities [56, 57]. Furthermore, they recapitulate effects of relevant non-linearities on ecological dynamics commonly employed in standard ecological models such as the amount of saturation in the functional response ( $\gamma_i^k$ ), the shape of the producer growth function ( $\phi_i^k$ ), the presence of intraspecific consumer interference ( $\psi_i^k$ ), and the density-dependence of consumer mortality rates ( $\mu_i^k$ ). The exponent parameter  $\lambda_{j,i}^k$  can be interpreted as the adaptability of consumer preferences to different prey items; following previous work [6] we assume constant preferences for simplicity.

315  
316  
317  
318  
319  
320  
321

Niche model food web topologies that specify the structure of interactions in equation (5) were generated following [18]. Each species  $i$  is assumed to exist on a niche axis between  $[0,1]$  and is assigned a niche value using a uniform distribution. The species then is assumed to consume all species over a range  $r_i$  that is near or below the position of species  $i$  on the niche axis, generating a trophic hierarchy. The location of the range is assigned using a beta function with expected value  $2C$ .

322  
323  
324  
325  
326

We chose input values for species richness  $S$  as integers between 10 and 30 inclusive, [10..30], and for connectance  $C$  as (0.12, 0.14, ..., 0.24). The latter range was chosen to encompass empirically observed connectance values [90]. Only webs with a single connected component were retained for analysis.

Spatial networks were generated as random geometric graphs (RGG). Networks of  $N$  patches were generated by first randomly assigning coordinates in 2 dimensional space to each patch drawn from a standard uniform distribution. patches were then connected if the Euclidean distance between their coordinates fell below a threshold  $n$ . Networks with unconnected patches were discarded. For results included here, we used  $N = 10$  and  $n = 0.32$ , yet simulations with greater numbers of patches, different values of  $n$ , and even different network configurations exhibited qualitatively similar results.

The spatial structure of the RGG is encoded in the Laplacian matrix  $L$ . The Laplacian  $L$  is an  $N \times N$  matrix where the diagonal entries  $L_{k=l}$  represent the number of dispersal connections (i.e. degree) of each patch  $k$ . The off-diagonal elements of  $L$  reflect dispersal connections between individual patches, where  $L_{k \neq l} = -1$  when patches  $k$  and  $l$  are connected and 0 otherwise. Linking the Laplacian to the dispersal matrix  $D$  yields the spatial structure of the dispersal network. The dispersal matrix is an  $S \times S$  matrix with species-specific dispersal rates on the diagonal  $D_{i=j} = \delta_i$  and all other entries zero. Using  $L \otimes D$ , with  $\otimes$  being the Kronecker product, yields the  $SN \times SN$  block matrix that describes the pattern of connections between all patches by all species.

Local food webs were embedded in the spatial structure of the metacommunity using equation (3). Each local web Jacobian  $J_k$  was numerically placed on the diagonal of the metacommunity food web matrix  $P$ . Because we assume environmental heterogeneity, the entries of each  $J_k$  in a given  $P$  vary, although the topologies remain fixed. Environmental heterogeneity is implemented as variation in branching and elasticity parameters; the value of each parameter for each species in a local web  $J_k$  was determined by drawing from uniform distributions in the appropriate range defined in Table 1.

We generated one hundred unique food web topologies for each combination of species richness  $S$  and connectance  $C$ , yielding 7,700 unique metacommunity topologies. We explored metacommunity dynamics over five values of dispersal coupling  $d$  and seven values of dispersal allometry  $z$ , yielding 192,500 unique metacommunities. Dispersal structure of each unique metacommunity was then permuted 100 times; the stability of the metacommunity was compared with the median stability of the random dispersal metacommunities.

Associations between different parameters and metacommunity stability were quantified using generalized linear models (GLMs) with binomial errors and a logit link function, with an information criterion-based model selection scheme. Following White et al. [91] we use GLMs as a framework for partitioning variance and correlations between important model parameters and metacommunity stability, and not for assessing statistical significance. We first fitted a global model comprising linear combinations of fixed effects for local food web species richness  $S$ , web connectance  $C$ , the strength of spatial network links  $d$ , the body size scaling exponent for allometric dispersal  $z$ , and the spatial variances and means of parameters describing predator satiation, interaction strengths, and nutrient availability to primary producers (Table 1). We then computed Akaike's information criterion (AIC)

for all submodels comprising different combinations of fixed effects in the global model, and selected the model with the lowest AIC score as the best fitting. This model included four terms: species richness  $S$ , web connectance  $C$ , global spatial network link strength  $d$ , and the exponent of allometric dispersal  $z$  (Fig. 3).

## References

- [1] Gardner, M. R. & Ashby, W. R. Connectance of large dynamic (cybernetic) systems: critical values for stability. *Nature* **228**, 784–784 (1970).
- [2] May, R. M. Will a large complex system be stable? *Nature* **238**, 413–414 (1972).
- [3] McCann, K. S. The diversity–stability debate. *Nature* **405**, 228–233 (2000).
- [4] Dunne, J. A. The network structure of food webs. *Ecological networks: linking structure to dynamics in food webs* 27–86 (2006).
- [5] Williams, R. J., Brose, U. & Martinez, N. D. Homage to yodzis and innes 1992: scaling up feeding-based population dynamics to complex ecological networks. In *From energetics to ecosystems: the dynamics and structure of ecological systems*, 37–51 (Springer, 2007).
- [6] Gross, T., Rudolf, L., Levin, S. A. & Dieckmann, U. Generalized models reveal stabilizing factors in food webs. *Science* **325**, 747–750 (2009).
- [7] Fahimipour, A. K., Anderson, K. E. & Williams, R. J. Compensation masks trophic cascades in complex food webs. *Theoretical Ecology* **10**, 245–253 (2017).
- [8] Rooney, N. & McCann, K. S. Integrating food web diversity, structure and stability. *Trends in ecology & evolution* **27**, 40–46 (2012).
- [9] Jacquet, C. *et al.* No complexity–stability relationship in empirical ecosystems. *Nature communications* **7**, 12573 (2016).
- [10] Brose, U., Williams, R. J. & Martinez, N. D. Allometric scaling enhances stability in complex food webs. *Ecology letters* **9**, 1228–1236 (2006).
- [11] Martinez, N. D. Allometric trophic networks from individuals to socio-ecosystems: Consumer–resource theory of the ecological elephant in the room. *Frontiers in Ecology and Evolution* **8**, 92 (2020).
- [12] McCann, K. S., Rasmussen, J. & Umbanhowar, J. The dynamics of spatially coupled food webs. *Ecology letters* **8**, 513–523 (2005).
- [13] Brechtel, A., Gramlich, P., Ritterskamp, D., Drossel, B. & Gross, T. Master stability functions reveal diffusion-driven pattern formation in networks. *Physical Review E* **97**, 032307 (2018).
- [14] Brechtel, A., Gross, T. & Drossel, B. Far-ranging generalist top predators enhance the stability of meta-foodwebs. *Scientific reports* **9**, 1–15 (2019).
- [15] Gross, T. & *et. al.* Modern models of trophic meta-communities. *Phil. Trans. Roy. Soc. B* (in press).
- [16] Rooney, N., McCann, K., Gellner, G. & Moore, J. C. Structural asymmetry and the stability of diverse food webs. *Nature* **442**, 265–269 (2006).
- [17] Otto, S. B., Rall, B. C. & Brose, U. Allometric degree distributions facilitate food-web stability. *Nature* **450**, 1226–1229 (2007).
- [18] Williams, R. J. & Martinez, N. D. Simple rules yield complex food webs. *Nature* **404**, 180–183 (2000).
- [19] Cohen, J. E., Pimm, S. L., Yodzis, P. & Saldaña, J. Body sizes of animal predators and animal prey in food webs. *Journal of animal ecology* 67–78 (1993).

- [20] Petchey, O. L., Beckerman, A. P., Riede, J. O. & Warren, P. H. Size, foraging, and food web structure. *Proceedings of the National Academy of Sciences* **105**, 4191–4196 (2008). 404  
405
- [21] Cohen, J. E., Briand, F. & Newman, C. M. *Community food webs: data and theory*, vol. 20 (Springer Science & Business Media, 2012). 406  
407
- [22] Elton, C. S. *Animal ecology* (University of Chicago Press, 2001). 408
- [23] Lindeman, R. L. The trophic-dynamic aspect of ecology. *Ecology* **23**, 399–417 (1942). 409
- [24] Peters, R. H. & Peters, R. H. *The ecological implications of body size*, vol. 2 (Cambridge university press, 1986). 410
- [25] Riede, J. O. *et al.* Stepping in elton’s footprints: a general scaling model for body masses and trophic levels across ecosystems. *Ecology letters* **14**, 169–178 (2011). 411  
412
- [26] Brose, U. *et al.* Consumer–resource body-size relationships in natural food webs. *Ecology* **87**, 2411–2417 (2006). 413
- [27] MacArthur, R. H. & Wilson, E. O. *The theory of island biogeography*, vol. 1 (Princeton university press, 2001). 414
- [28] Simberloff, D. S. & Wilson, E. O. Experimental zoogeography of islands: the colonization of empty islands. *Ecology* **50**, 278–296 (1969). 415  
416
- [29] Brown, J. H. & Kodric-Brown, A. Turnover rates in insular biogeography: effect of immigration on extinction. *Ecology* **58**, 445–449 (1977). 417  
418
- [30] Levins, R. Some demographic and genetic consequences of environmental heterogeneity for biological control. *American Entomologist* **15**, 237–240 (1969). 419  
420
- [31] Gotelli, N. J. Metapopulation models: the rescue effect, the propagule rain, and the core-satellite hypothesis. *The American Naturalist* **138**, 768–776 (1991). 421  
422
- [32] Crowley, P. H. Dispersal and the stability of predator-prey interactions. *The American Naturalist* **118**, 673–701 (1981). 423
- [33] Reeve, J. D. Environmental variability, migration, and persistence in host-parasitoid systems. *The American Naturalist* **132**, 810–836 (1988). 424  
425
- [34] Murdoch, W. W. Population regulation in theory and practice. *Ecology* **75**, 271–287 (1994). 426
- [35] Briggs, C. J. & Hoopes, M. F. Stabilizing effects in spatial parasitoid–host and predator–prey models: a review. *Theoretical population biology* **65**, 299–315 (2004). 427  
428
- [36] Gravel, D., Massol, F. & Leibold, M. A. Stability and complexity in model meta-ecosystems. *Nature communications* **7**, 1–8 (2016). 429  
430
- [37] Mougi, A. & Kondoh, M. Food-web complexity, meta-community complexity and community stability. *Scientific reports* **6**, 24478 (2016). 431  
432
- [38] Domenici, P. The scaling of locomotor performance in predator–prey encounters: from fish to killer whales. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology* **131**, 169–182 (2001). 433  
434
- [39] Hirt, M. R., Laueremann, T., Brose, U., Noldus, L. P. & Dell, A. I. The little things that run: a general scaling of invertebrate exploratory speed with body mass. *Ecology* **98**, 2751–2757 (2017). 435  
436
- [40] Hirt, M. R., Jetz, W., Rall, B. C. & Brose, U. A general scaling law reveals why the largest animals are not the fastest. *Nature Ecology & Evolution* **1**, 1116–1122 (2017). 437  
438
- [41] Reiss, M. Scaling of home range size: body size, metabolic needs and ecology. *Trends in Ecology & Evolution* **3**, 85–86 (1988). 439  
440
- [42] Minns, C. K. Allometry of home range size in lake and river fishes. *Canadian Journal of Fisheries and Aquatic Sciences* **52**, 1499–1508 (1995). 441  
442
- [43] Jetz, W., Carbone, C., Fulford, J. & Brown, J. H. The scaling of animal space use. *Science* **306**, 266–268 (2004). 443

- [44] Hendriks, A. J., Willers, B. J., Lenders, H. R. & Leuven, R. S. Towards a coherent allometric framework for individual home ranges, key population patches and geographic ranges. *Ecography* **32**, 929–942 (2009). 444  
445
- [45] Hartfelder, J. *et al.* The allometry of movement predicts the connectivity of communities. *Proceedings of the National Academy of Sciences* **117**, 22274–22280 (2020). 446  
447
- [46] Vander Zanden, M. J. & Vadeboncoeur, Y. Fishes as integrators of benthic and pelagic food webs in lakes. *Ecology* **83**, 2152–2161 (2002). 448  
449
- [47] Wolkovich, E. M. *et al.* Linking the green and brown worlds: the prevalence and effect of multichannel feeding in food webs. *Ecology* **95**, 3376–3386 (2014). 450  
451
- [48] Lomolino, M. V. Immigrant selection, predation, and the distributions of *Microtus pennsylvanicus* and *Blarina brevicauda* on islands. *The American Naturalist* **123**, 468–483 (1984). 452  
453
- [49] Beisner, B. E., Peres-Neto, P. R., Lindström, E. S., Barnett, A. & Longhi, M. L. The role of environmental and spatial processes in structuring lake communities from bacteria to fish. *Ecology* **87**, 2985–2991 (2006). 454  
455
- [50] De Bie, T. *et al.* Body size and dispersal mode as key traits determining metacommunity structure of aquatic organisms. *Ecology Letters* **15**, 740–747 (2012). 456  
457
- [51] Kareiva, P. Population dynamics in spatially complex environments: theory and data. *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences* **330**, 175–190 (1990). 458  
459
- [52] Murray, J. *Mathematical biology II: spatial models and biomedical applications*, vol. 3 (Springer-Verlag, 2001). 460
- [53] Rietkerk, M. & Van de Koppel, J. Regular pattern formation in real ecosystems. *Trends in ecology & evolution* **23**, 169–175 (2008). 461  
462
- [54] Pedersen, E. J., Marleau, J. N., Granados, M., Moeller, H. V. & Guichard, F. Nonhierarchical dispersal promotes stability and resilience in a tritrophic metacommunity. *The American Naturalist* **187**, E116–E128 (2016). 463  
464
- [55] Gross, T. & Feudel, U. Generalized models as a universal approach to the analysis of nonlinear dynamical systems. *Physical Review E* **73**, 016205 (2006). 465  
466
- [56] Yeakel, J. D., Stiefs, D., Novak, M. & Gross, T. Generalized modeling of ecological population dynamics. *Theoretical Ecology* **4**, 179–194 (2011). 467  
468
- [57] Gross, T. & Feudel, U. Generalized models as a universal approach to the analysis of nonlinear dynamical systems. *Physical Review E* **73**, 016205 (2006). 469  
470
- [58] Hirt, M. R. *et al.* Bridging scales: allometric random walks link movement and biodiversity research. *Trends in ecology & evolution* **33**, 701–712 (2018). 471  
472
- [59] Segel, L. A. & Levin, S. A. Application of nonlinear stability theory to the study of the effects of diffusion on predator-prey interactions. In *AIP conference proceedings*, vol. 27, 123–152 (American Institute of Physics, 1976). 473  
474
- [60] Othmer, H. G. & Scriven, L. Non-linear aspects of dynamic pattern in cellular networks. *Journal of theoretical biology* **43**, 83–112 (1974). 475  
476
- [61] Estes, J. A. *et al.* Trophic downgrading of planet earth. *science* **333**, 301–306 (2011). 477
- [62] Krause, A. E., Frank, K. A., Mason, D. M., Ulanowicz, R. E. & Taylor, W. W. Compartments revealed in food-web structure. *Nature* **426**, 282–285 (2003). 478  
479
- [63] Post, D. M., Conners, M. E. & Goldberg, D. S. Prey preference by a top predator and the stability of linked food chains. *Ecology* **81**, 8–14 (2000). 480  
481
- [64] Neutel, A.-M., Heesterbeek, J. A. & de Ruiter, P. C. Stability in real food webs: weak links in long loops. *Science* **296**, 1120–1123 (2002). 482  
483
- [65] Leibold, M. A. *et al.* The metacommunity concept: a framework for multi-scale community ecology. *Ecology Letters* **7**, 601–613 (2004). 484  
485

- [66] Jenkins, D. G. *et al.* Does size matter for dispersal distance? *Global Ecology and Biogeography* **16**, 415–425 (2007). 486
- [67] Heino, J. *et al.* Metacommunity organisation, spatial extent and dispersal in aquatic systems: patterns, processes and prospects. *Freshwater Biology* **60**, 845–869 (2015). 487  
488
- [68] Siegel, D. *et al.* The stochastic nature of larval connectivity among nearshore marine populations. *Proceedings of the National Academy of Sciences* **105**, 8974–8979 (2008). 489  
490
- [69] Pillai, P., Loreau, M. & Gonzalez, A. A patch-dynamic framework for food web metacommunities. *Theoretical Ecology* **3**, 223–237 (2010). 491  
492
- [70] Pillai, P., Gonzalez, A. & Loreau, M. Metacommunity theory explains the emergence of food web complexity. *Proceedings of the National Academy of Sciences* **108**, 19293–19298 (2011). 493  
494
- [71] Plitzko, S. J. & Drossel, B. The effect of dispersal between patches on the stability of large trophic food webs. *Theoretical Ecology* **8**, 233–244 (2015). 495  
496
- [72] Guichard, F. Recent advances in metacommunities and meta-ecosystem theories. *F1000Research* **6** (2017). 497
- [73] Hata, S., Nakao, H. & Mikhailov, A. S. Dispersal-induced destabilization of metapopulations and oscillatory turing patterns in ecological networks. *Scientific reports* **4**, 3585 (2014). 498  
499
- [74] White, K. & Gilligan, C. Spatial heterogeneity in three species, plant–parasite–hyperparasite, systems. *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences* **353**, 543–557 (1998). 500  
501
- [75] Gibert, J. P. & Yeakel, J. D. Laplacian matrices and turing bifurcations: Revisiting levin 1974 and the consequences of spatial structure and movement for ecological dynamics. *Theoretical Ecology* **12**, 265–281 (2019). 502  
503
- [76] Fox, J. W., Vasseur, D., Cotroneo, M., Guan, L. & Simon, F. Population extinctions can increase metapopulation persistence. *Nature ecology & evolution* **1**, 1271–1278 (2017). 504  
505
- [77] Hastings, A. Food web theory and stability. *Ecology* **69**, 1665–1668 (1988). 506
- [78] Anderson, H., Hutson, V. & Law, R. On the conditions for permanence of species in ecological communities. *The American Naturalist* **139**, 663–668 (1992). 507  
508
- [79] Haydon, D. Pivotal assumptions determining the relationship between stability and complexity: an analytical synthesis of the stability-complexity debate. *The American Naturalist* **144**, 14–29 (1994). 509  
510
- [80] Chen, X. & Cohen, J. E. Global stability, local stability and permanence in model food webs. *Journal of Theoretical Biology* **212**, 223–235 (2001). 511  
512
- [81] Bjørnstad, O. N., Ims, R. A. & Lambin, X. Spatial population dynamics: analyzing patterns and processes of population synchrony. *Trends in Ecology & Evolution* **14**, 427–432 (1999). 513  
514
- [82] Ims, R. A. & Andreassen, H. P. Spatial synchronization of vole population dynamics by predatory birds. *Nature* **408**, 194–196 (2000). 515  
516
- [83] Sundell, J. *et al.* Large-scale spatial dynamics of vole populations in finland revealed by the breeding success of vole-eating avian predators. *Journal of Animal Ecology* **73**, 167–178 (2004). 517  
518
- [84] Ripple, W. J. *et al.* Status and ecological effects of the world’s largest carnivores. *Science* **343** (2014). 519
- [85] McCauley, D. J. *et al.* Marine defaunation: animal loss in the global ocean. *Science* **347** (2015). 520
- [86] Parsons, T. The removal of marine predators by fisheries and the impact of trophic structure. *Marine Pollution Bulletin* **25**, 51–53 (1992). 521  
522
- [87] Baum, J. K. & Worm, B. Cascading top-down effects of changing oceanic predator abundances. *Journal of Animal Ecology* **78**, 699–714 (2009). 523  
524
- [88] Albert, C. H., Rayfield, B., Dumitru, M. & Gonzalez, A. Applying network theory to prioritize multispecies habitat networks that are robust to climate and land-use change. *Conservation Biology* **31**, 1383–1396 (2017). 525  
526

- [89] Schiesari, L. *et al.* Towards an applied metaecology. *Perspectives in ecology and conservation* **17**, 172–181 (2019). 527
- [90] Vermaat, J. E., Dunne, J. A. & Gilbert, A. J. Major dimensions in food-web structure properties. *Ecology* **90**, 278–282 (2009). 528  
529
- [91] White, J. W., Rassweiler, A., Samhouri, J. F., Stier, A. C. & White, C. Ecologists should not use statistical significance tests to interpret simulation model results. *Oikos* **123**, 385–388 (2014). 530  
531

### **Acknowledgments**

532

K.E.A. was supported by a grant from the National Science Foundation #DEB-1553718. A.K.F. was supported by a fellowship from the National Research Council Research Associateship Program.

533

534

### **Authors contributions**

535

K.E.A. and A.K.F. conceptualized the study, wrote the manuscript, and contributed analyses.

536

### **Competing interests**

537

The authors declare no competing interests.

538

# Figures




Figure 1

Model metacommunities are composed of local food webs connected to one another by dispersal. A. Each local web inhabits a habitat patch that is part of a spatial network, generated as a random geometric graph. Species in each web have a body size that is larger at higher trophic levels. Food webs have the same number of species and topology in all patches, but interaction rates and other ecological parameters vary among habitats mimicking spatial environmental heterogeneity. B. Dispersal varies as either an increasing or decreasing function of body size.





Figure 2


Please see the Manuscript PDF file for the complete figure caption.

## Correlation with metacommunity stability




**Figure 3**

The effects of key parameters on metacommunity stability. Stability is as defined in Fig. 2 and parameters are defined in Table 1. Correlations given are coefficients from the best-fitting generalized linear model  $\pm 2$  SEM. Positive correlations indicate that larger values of a parameter correspond to a higher probability that a randomly-assembled metacommunity will be stable.



**Figure 4**

The effect of dispersal variation on metacommunity stability for spatial link strength  $d=0.1$ . The metacommunity is stable when the real part of the leading eigenvalue of the metacommunity Jacobian  $\lambda_1 < 0$ . Allometric dispersal is defined eq. 2. Permuted dispersal refers to cases where allometric dispersal rates were randomly reassigned to new species. Each unique metacommunity with allometric dispersal was compared to 100 randomly permuted counterparts, and are shown with 1:1 lines. Grey regions mark portions of the plot representing qualitative changes in stability.



**Figure 5**

Eigenvalues  $\lambda_1$  from fig. 4 categorized by qualitative effects on stability. Categories Stability is gained and Stability is lost correspond to cases where the median effect of permuting dispersal rates is a change in the sign of  $\lambda_1$ . Stability is unaffected indicates no sign change.

## Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Supplement.pdf](#)